



Publication Year	2017
Acceptance in OA @INAF	2020-12-01T15:38:05Z
Title	Central compact objects in supernova remnants
Authors	DE LUCA, Andrea
DOI	10.1088/1742-6596/932/1/012006
Handle	http://hdl.handle.net/20.500.12386/28604
Journal	JOURNAL OF PHYSICS. CONFERENCE SERIES
Number	932

PAPER • OPEN ACCESS

Central compact objects in supernova remnants

To cite this article: A De Luca 2017 *J. Phys.: Conf. Ser.* **932** 012006

View the [article online](#) for updates and enhancements.

Related content

- [Cosmic-ray astrophysicists close in on nearby supernova source](#)
- [Searching for gravitational waves from Cassiopeia A with LIGO](#)
- [What stars become supernovae?](#)

Recent citations

- [The Timing Behavior of the Central Compact Object Pulsar 1E 1207.45209](#)
E. V. Gotthelf and J. P. Halpern
- [The Proper Motion of the Central Compact Object RX J0822–4300 in the Supernova Remnant Puppis A, Revisited](#)
Martin Mayer *et al.*
- [NS 1987A in SN 1987A](#)
Dany Page *et al.*



IOP | ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Central compact objects in supernova remnants

A De Luca

INAF/IASF Milano, Via Bassini 15, I-20133 Milano, Italy

E-mail: deluca@iasf-milano.inaf.it

Abstract. Central Compact Objects (CCOs) are a handful of sources located close to the geometrical center of young supernova remnants. They only show thermal-like, soft X-ray emission and have no counterparts at any other wavelength. While the first observed CCO turned out to be a very peculiar magnetar, discovery that three members of the family are weakly magnetised Isolated Neutron Stars (INSs) set the basis for an interpretation of the class. However, the phenomenology of CCOs and their relationship with other classes of INSs, possibly ruled by supernova fall-back accretion, are still far from being well understood.

1. Introduction

A dozen of point-like, soft X-ray sources has been discovered over the last two decades, gently shining very close to the geometrical center of young (0.3–7 kyr) supernova remnants (SNR). Their spectra are thermal-like, usually well described by the sum of two blackbodies with high temperatures (0.2–0.5 keV) and very small emitting radii (ranging from 0.1 to a few km). Their emission is generally steady, with a luminosity of the order of 10^{33} erg s⁻¹, and pulsations are undetected in most cases. They have no counterparts at any wavelength. No associated structures of diffuse, non-thermal emission are seen. Such sources are clearly different from both standard rotation-powered pulsars and magnetars. They have been dubbed, as a class, “Central Compact Objects” (CCOs)¹ – indeed, a designation suggesting our rather poor understanding of their nature and physics. Discovery of fast pulsations and measurement of a tiny, positive period derivative in three sources set the basis of a framework for CCOs as young, Isolated Neutron Stars (INS) with a weak dipole field. In this short paper, I will focus on recent results and directions in CCO studies, including discovery of the magnetar nature of the first observed CCO – a possibly unique source. I will show that an explanation of the properties of these sources would be highly relevant for our overall understanding of NS production rate, physics of core-collapse, INS diversity and evolution. Previous reviews on CCOs were given by [2, 3, 4].

2. 1E 161348–5055: a unique, slowly-rotating magnetar.

1E 161348–5055 (1E 1613) in the 2 kyr-old SNR RCW103 has been the prototype for CCOs. It was the first candidate radio-quiet INS discovered in a SNR [5]. However, more recent observations clearly separated 1E 1613 from CCOs and settled the case for a truly unique phenomenology: 1E 1613 displays a dramatic long-term variability with large outbursts and a puzzling periodicity at 6.67 hours, together with a young age and lack of an optical/infrared counterpart [6, 7]. The nature of the source, possibly the first low-mass X-ray binary system

¹ The name has been used for the first time by G. Pavlov, referring to the central source in the Cas A SNR [1].



observed inside a SNR, or a very peculiar isolated, young magnetar with an extremely slow spin period, possibly braked by interaction with a surrounding disk [6], remained debated for a decade – even more exotic pictures were proposed [8, 9, 10, 11, 12, 13].

A partial answer to this puzzle came only very recently. Indeed, on 2016 June 22, the Burst Alert Telescope onboard *Swift* detected a magnetar-like, short X-ray burst from the direction of 1E 1613, with a spectrum well described by a blackbody model ($kT \sim 9$ keV) and a luminosity of $\sim 2 \times 10^{39}$ erg s $^{-1}$. A strong X-ray outburst was simultaneously observed from 1E 1613 with the *Swift* X-Ray Telescope (XRT), with the 0.5–10 keV flux a factor of 100 brighter than the quiescent level observed up to one month before [14, 15]. Follow-up observations performed with *Chandra* and *NuSTAR* unveiled (i) a dramatic change in the shape of the 6.67 hour modulation, two broad peaks per period replacing the sinusoidal shape that had been observed since 2005 [14]; (ii) a non-thermal spectral component extending up to ~ 30 keV (modulated up to ~ 20 keV), never detected before, well described by a hard power law with $\Gamma \sim 1.2$, superimposed to the thermal continuum dominating at lower energies [14]. Based on a series of *Swift*/XRT observations, the overall energy emitted in the outburst (impulsive plus persistent) was computed to be $\sim 2.6 \times 10^{42}$ ergs [14]. A likely counterpart in the near infrared was detected by [16] with the *Hubble Space Telescope* with AB magnitudes of 26.3 and 24.2 in the F110W and F160W filters, respectively (roughly corresponding to the J and H bands). The same source was detected with the ESO/VLT with $K_s=20.9 \pm 0.1$ (De Luca et al, in preparation). All results from such multiwavelength observations strongly point to a magnetar interpretation for 1E 1613 [14, 15, 16]. Indeed, 1E 1613 stands out as a unique source because of its rotation period, slower by three orders of magnitude with respect to any other known magnetar candidate.

Assuming that 1E 1613 was born with a spin period of a few hours is hardly conciliable with angular momentum conservation in the progenitor star’s collapsing core and also clashes with the commonly assumed dynamo mechanism for the generation of the gigantic magnetic field in magnetars, requiring a fast initial period (\sim millisecond) [17]. Disregarding such a possibility, one should explain how a fast-spinning NS could be slowed-down to a period of 6.67 hours in ~ 2 kyr. Standard magneto-dipole braking, coupled to magneto-thermal evolution, cannot account for such a slow rotation for any reasonable assumption on the newborn NS properties [14]. Braking of a strongly-magnetized NS by material/magnetic interaction with a low-mass companion star was discussed by [9]. However, near-infrared observations rule out any companion with a mass larger than an M8 dwarf [16] – it seems unlikely that a binary system with such an extreme mass ratio survive a supernova explosion². An alternative possibility is to invoke the role of supernova fallback material [18, 19]. The NS could have been braked by propeller interaction with a long-lived residual disk. Such a possibility had been considered by [6, 8] and, more recently, by [20, 21]. Although the formation and stability of fallback disks surrounding newborn NSs has been questioned by [22], it was shown that a very low-mass disk ($\sim 10^{-9} M_\odot$) could brake the NS to the observed spin rate in $\sim 1 - 3$ kyr, provided it is endowed by a dipole magnetic field of 5×10^{15} G [21], the largest in the magnetar family. As a further possibility, the NS could have been slowed-down in an earlier phase – propeller interaction starting at the onset of fallback accretion (with no need for formation and long-term survival of a disk). Such a picture was mentioned by [14, 23], but no detailed calculations have been presented so far.

3. CCO pulsars: neutron stars with a weak dipole magnetic field.

A key discovery has been the detection of fast pulsations from three objects of the class: 1E 1207.4–5209 in G296.5+10.0 ($P \sim 424$ ms, [24]), CXOU J185238.6+004020 in Kes 79 ($P \sim 105$ ms, [25]) and RX J0852.0–4622 in Puppis A ($P \sim 112$ ms, [26]). This proved with no doubt that these sources are NSs. Even more enlightening has been the measurement of their period

² The same argument, together with the difficulty in explaining the dramatic long-term variability in pulse shape and pulsed fraction, argues against an interpretation of the 6.67 hr cycle as the orbital period in a binary.

derivatives [4, 27], which turned out to be very small (indeed, very long observation campaigns were required). Implications are extremely interesting: (i) the dipole magnetic field inferred from standard magneto-dipole braking $B_{\text{dip}} = 3.2 \times 10^{19} (P\dot{P})^{1/2}$ is $10^{10} - 10^{11}$ G, remarkably smaller than the one of the bulk of the rotation-powered pulsar population; (ii) the characteristic age ($\tau_C = P/2\dot{P}$) is 4–5 orders of magnitude larger than the age of the host SNR (indeed, the CCO birth period is likely very close to the currently observed one); (iii) the spin-down energy loss is more than 10 times smaller than the X-ray luminosity. However, several properties of the CCO pulsars do not fit easily in the picture of weakly magnetised INs.

CXOU J185238.6+004020 displays an extremely high modulation (pulsed fraction $\sim 64\%$, with very little dependence on energy) of its thermal emission, consisting of the sum of two hot blackbodies with tiny emitting areas [27]. Reproducing the observed pulse shape and fraction with thermal emission models is challenging [28] and points to a peculiar thermal map, with a high temperature contrast between a small emitting region (a hot spot surrounded by a warmer region) and the remaining cooler and unobservable surface of the star. A highly elongated shape in the longitudinal direction (with respect to the spin axis) is required for the emitting region; a more conventional polar cap geometry requires the radiation to be highly beamed [29].

RX J0852.0–4622 too has a thermal continuum well described by two blackbodies. Its pulse shape is sinusoidal, with a very abrupt, 180° phase reversal at ~ 1.2 keV, where the hot and warm blackbodies switch dominance [30]. This is consistent with an anti-podal hot spot model, with two small emitting regions with different size and temperature located at opposite sides of the NS [30]. A spectral feature is also seen in the low-energy portion of the spectrum. It can be described either as an emission line at ~ 0.75 keV, or as a couple of absorption lines at ~ 0.46 keV and ~ 0.92 keV. Significant variability in the feature is seen between two observations performed in 2001 and 2009: assuming the emission line model, the central energy decreases from ~ 0.79 to ~ 0.71 keV [4, 31]. No further variability is seen after 2009 [4].

1E 1207.4–5209 displays multiple absorption features at harmonically spaced energies (0.7, 1.4, 2.1 and possibly 2.8 keV) superimposed to a thermal spectrum [32, 33, 34, 35]. After a long debate (see [3]), measurement of P and \dot{P} points to an interpretation of the features as due to electron cyclotron scattering close to the star’s surface, the feature at 0.7 keV being the fundamental. The magnetic field strength in the region where the lines are formed would be $\sim 8 \times 10^{10}$ G (accounting for a gravitational redshift $z = 0.3$), in broad agreement with the value inferred by magneto-dipole braking ($\sim 9.8 \times 10^{10}$ G). The relative strength of the harmonics was explained by taking into account resonances in the photospheric free-free opacity in the presence of the magnetic field [36, 37, 38]. The X-ray pulsation is dominated by the complex modulation of the spectral features as a function of the rotational phase [35]; the continuum, well described by the sum of two blackbody curves, being almost unpulsed. No simple constraints on the geometry of the thermally emitting regions could be set.

4. Other CCOs: homogeneous group or mixed bag?

The family of CCOs includes about ten more sources³. Thermal emission properties are pretty homogeneous, with no pulsations (rather deep upper limits have been set in a few cases), nor long-term variability (although multi-epoch coverage is limited in most cases). It seems reasonable to assume that such sources are INs similar to the three CCO pulsars, possibly with even smaller dipole fields.

The most famous CCO was discovered close to the center of the very young (~ 350 yr) Cas A SNR in the *Chandra* “first light” image [39]. An early claim of a magnetar nature for this CCO, based on detection of the possible infrared echo of a giant flare occurred around 1950 [40] was later retracted [41, 42]. Multi-epoch observations with *Chandra* point to long-term

³ see www.iasf-milano.inaf.it/~deluca/cco/main.htm for an updated list

evolution of the spectral shape and luminosity of this source, consistent with a $\sim 4\%$ temperature decrease in about 10 yr [43]. Such a direct observation of the NS cooling would have very important implications for our understanding of the properties of ultra-dense matter in the NS interior [44, 45, 46, 47, 48]. However, more recent analysis of *Chandra* data focused on possible systematics affecting the measurement (e.g., instrument calibration issues) and concluded that the reality and rate of the possible temperature decrease are uncertain [49, 50].

It was shown [51] that a carbon atmosphere NS model with low magnetic field provides a good description of the spectrum of this source and implies an emitting region consistent with the entire NS surface. This possibly solves the puzzling lack of pulsations coupled to the large temperature anisotropy (with uncomfortably small emitting areas) as derived from other spectral models (both blackbody curves and weakly-magnetised NS hydrogen atmosphere models), pointing to the picture of a weakly magnetized NS, with uniform emission from its surface and with active nuclear burning in its surface layers.

The same picture was proposed for other CCOs, as soon as high-quality X-ray spectra became available: the source in G353.6–0.7 [52, 53] (possibly the hottest known INS), as well as the source in G15.9+0.2 [54]. It was stressed that the carbon atmosphere models makes it possible to reconcile thermal luminosities with the current distance estimates and also to constrain the equation of state of the NS [53, 55]. As a matter of fact, however, such models cannot explain the phenomenology of all CCOs. For instance, the carbon atmosphere model would describe the emission of the CCO pulsar in Kes 79 as coming from the whole surface of the NS, but this is clearly at odds with the observed very high pulsed fraction [29]. Indeed, the current upper limits on the CCO pulsations are not stringent enough to rule out highly anisotropic surface temperature distributions [56, 57] that would be consistent with different atmosphere chemical composition as well as with different properties of the NSs.

5. CCOs as a “class”.

The picture of CCOs as weakly-magnetised, young NSs has to face two issues.

First issue: the highly-anisotropic, high contrast-ratio temperature surface distribution cannot be easily explained in this framework. The hot spots cannot be heated by rotation-powered particle bombardment (ruled out by the spin-down energetics); they cannot be accretion-powered (ruled out by timing [27] as well as by deep optical limits to any companion star, e.g., [58]). Hot spots could be due to localized crustal heating by magnetic field decay. This would require a magnetic field with strong non-dipolar components ($10^{14} - 10^{15}$ G to account for the observed luminosity [27]), coupled to a factor 10^4 less intense dipolar component. As an alternative possibility, hot spots could be powered by residual heat. This would require highly anisotropic heat transfer from the NS interior, pointing to a strong magnetic field in the crust. For instance, a toroidal field of order 10^{14} G could screen large part of the surface around the equator, channelling the heat flux towards the poles [28, 59]. A poloidal field as low as $10^{10} - 10^{11}$ G could be enough to make the magnetic field configuration stable [60].

Second issue: the region in the $P - \dot{P}$ parameter space where CCO pulsars are located is highly underpopulated, at odds with expectations [61]. Such region is well above the “death line” for radio pulsars. Indeed, rotation-powered radio emission is seen from several sources with similar spin parameters. Moreover, CCOs are a common outcome of core-collapse supernovae – they are found in SNR as frequently as other classes of NSs (more frequently than magnetars, see, e.g., [62]). Frequent formation, coupled with a very slow evolution of the spin period would lead to a large number of low B -field sources above the death line. As a possible explanation of this puzzle, CCOs could be intrinsically radio-quiet and become essentially invisible after their host SNR fade away. Alternatively, radio luminosity in rotation-powered emission could depend on spin-down power, as suggested by radio pulsar population synthesis (e.g., [63]). As a further possibility, CCOs could evolve and move to a different region of the $P - \dot{P}$ diagram. This would

ease a further related issue: the sum of the inferred birth rates for different classes of NSs exceeds the Galactic core-collapse supernova rate (see, e.g., [64], although these authors did not discuss the case of CCOs) – this points to evolutionary links among different NS families.

Different frameworks for CCOs as young INs with weak dipole field have been discussed. A first possible scenario [65, 66] links the origin of the weak field to a slow rotation of the collapsing progenitor star core, which would result in an inefficient dynamo mechanism [67]. Such a picture does not seem very appealing: the birth period for CCOs is not “long”, according to population synthesis models for radio pulsars (e.g., [63] point to a wide distribution of birth periods with a mean of 300 ms and a dispersion of 150 ms). Moreover, the origin of the peculiar thermal anisotropy would remain unexplained. As a second possibility, CCOs could be quiescent magnetars, with an extremely weak dipole field, but with a strong crustal magnetic field, emerging in local “sunspot” structures. Such a scenario, however, seems rather unlikely because of the general lack of variability in CCOs, at odds with the ubiquitous variability seen in magnetars (but we cannot exclude the picture to be correct for a fraction of the sample). A third picture is known as the “buried field” scenario: prompt accretion of supernova fallback material [18] could bury the magnetic field of the newborn NS beneath its surface; the field could then re-emerge by diffusion on a time scale depending on the amount of accreted matter [19, 68, 69]. Recent models suggest that spherical accretion of $\sim 10^{-4} - 10^{-2} M_{\odot}$ is required to screen typical fields in the $10^{12} - 10^{14}$ G range for $\sim 10^3 - 10^5$ yr [70, 71, 72, 73, 74, 75]. According to these studies, (i) the strong, hidden crustal field could explain peculiarities in the thermal map; (ii) a low dipole field could be common in young NSs (\leq few kyr); (iii) CCOs could turn into radio pulsars at an older age (see also [76]). It is difficult to test the buried field scenario on CCOs by X-ray timing: evidence for a re-emerging magnetic field could come from a measurement of their braking index $n = 2 - P\ddot{P}/\dot{P}^2$, expected to be equal to 3 in the case of pure magneto-dipole braking for a constant magnetic field, and smaller than 3 in case of a growing dipole field. However, the tiny \dot{P} of CCOs make any measurement of their n extremely challenging. Some supporting evidence for this picture is provided by timing investigation of young ($\tau_c \sim 10^3 - 10^4$ yr) rotation-powered pulsars, yielding low values for the braking index that can be interpreted as a hint of a growing magnetic field [77, 78]; indirect support is also provided by the absence of evidence for weakly magnetised ($\leq 10^{11}$ G) NSs in High-mass X-ray binaries, consistent with field re-emergence in these sources on a time scale of $\sim 10^4$ yr [79].

A different way to attack the problem is to search for CCO descendants. Such investigations assume that at least some CCO may be a radio pulsar; CCOs would be much younger than “standard” pulsars in a given $P - \dot{P}$ parameter space region and thus they should be distinguishable because of a high thermal luminosity. A search for CCO descendants was performed among “mildly-recycled” pulsars (NSs that have accreted gas from a companion star in a binary system for a short time before a second supernova explosion halted their spin-up) – a class of pulsars proposed to explain the sparse population of sources in the same $P - \dot{P}$ region where CCOs have been recently located. However, none was found [80]. A second attempt was carried out, considering seemingly old radio pulsars spatially coincident with known SNRs, with negative results [81]. A third investigation did not identify any plausible CCO among radio pulsars with weak field ($B < 10^{11}$ G), but energetics larger than mildly recycled pulsars [82]. These studies suggest that CCO descendants, if not radio-quiet, should hide among pulsars with larger B-field and energetics. Interestingly, the peculiar neutron star dubbed *Calvera* [83] has been proposed as a possible example of an evolved CCO, whose SNR is no more detectable in X-rays. It is a radio-quiet pulsar, with $P \sim 59$ ms [84] and $\dot{E}_{\text{rot}} \sim 6 \times 10^{35}$ erg s $^{-1}$ [85], only showing soft X-ray thermal-like emission in spite of its high spin-down luminosity. *Calvera* lies in the $P - \dot{P}$ plane along the path of growing B -field between CCOs and young radio pulsars; however, a lack of any constraint on its distance (hence on its luminosity) as well as on its true age does not allow us to draw a firm conclusion [86].

6. Conclusions.

CCOs, a handful of candidate young INs with elusive properties, proved to include extraordinarily interesting objects. Explanation of their phenomenology challenges standard models and points to a very complex scenario, in which accretion of supernova fallback material in different regimes, coupled to a variety of initial conditions for the magnetic field strength and configuration, plays an important role in shaping the properties of newborn NSs. This has very important implications towards a physical understanding of the different phenomenological classes of INs and of their evolution. There is still a large space for discoveries in CCO studies, especially in view of forecoming observing facilities such as *eROSITA*, *ATHENA*, SKA. Deep searches for periodicity and long-term flux monitoring in the X-rays will allow one to assess the nature of non-pulsating CCOs – proving them to be similar to the three CCO pulsars, or unveiling any (peculiar) magnetar among them – as a matter of fact, the spectrum of a quiescent magnetar and of a CCO are almost undistinguishable. Large surveys will possibly allow one to identify new candidate CCOs and/or candidate CCO descendants. (Targeted) radio pulsar searches will unveil if CCOs are intrinsically radio-quiet, or simply radio-faint. Using already available data, modeling the phase-resolved behaviour of CCO pulsar will allow one to constrain the physics of the spectral features, the surface thermal map and magnetic field topology, as well as to test expectations of the buried field scenario.

6.1. Acknowledgments

I warmly thank the SOC and the LOC for organizing a very interesting meeting and for the invitation to speak. I thank P Esposito for a critical reading of the manuscript and G F Bignami (who left us prematurely on May 24th, 2017), A Borghese, P A Caraveo, F Coti Zelati, G L Israel, S Mereghetti, N Rea, A Tiengo for many useful discussions.

References

- [1] Pavlov G G, Zavlin V E, Aschenbach B, Trümper J and Sanwal D 2000 *ApJ* **531** L53-6
- [2] Pavlov G G, Sanwal D and Teter M A 2004 *IAUS* **218** 239
- [3] De Luca A 2008 *AIPC* **983** 311-9
- [4] Gotthelf E V, Halpern J P and Alford J 2013 *ApJ* **765** 58
- [5] Tuohy I and Garmire G 1980 *ApJ* **239** L107-10
- [6] De Luca A, Caraveo P A, Mereghetti S, Tiengo A and Bignami G F 2006 *Science* **313** 814-7
- [7] De Luca A *et al* 2008 *ApJ* **682** 1185-94
- [8] Li X-D 2007 *ApJ* **666** L81-4
- [9] Pizzolato F, Colpi M, De Luca A, Mereghetti S and Tiengo A 2008 *ApJ* **681** 530-42
- [10] Bhadkamkar H and Ghosh P 2009 *A&A* **506** 1297-307
- [11] Esposito P, Turolla R, De Luca A, Israel G L, Possenti A and Burrows D N 2011 *MNRAS* **418** 170-5
- [12] Ikhsanov N R, Kim V Y, Beskrovnaya N G and Pustil'nik L A 2013 *Ap&SS* **346** 105-9
- [13] Liu X W, Xu R X, van den Heuvel E P J, Qiao G J, Han J L, Han Z W and Li X D 2015 *ApJ* **799** 233
- [14] Rea N, Borghese A, Esposito P, Coti Zelati F, Bachetti M, Israel G L and De Luca A 2016 *ApJ* **828** L13
- [15] D'Ai A *et al* 2016 *MNRAS* **463** 2394-404
- [16] Tendulkar S P, Kaspi V M, Archibald R F and Scholz P 2017 *ApJ* **841** 11
- [17] Thompson C and Duncan R C 1993 *ApJ* **408** 194-217
- [18] Colgate S A 1971 *ApJ* **163** 221
- [19] Chevalier R A 1989 *ApJ* **346** 847-59
- [20] Tong H, Wang W, Liu X W and Xu R X 2016 *ApJ* **833** 265
- [21] Ho W C G and Andersson N 2017 *MNRAS* **464** L65-9
- [22] Perna R, Duffell P, Cantiello M and MacFadyen A I 2014 *ApJ* **781** 119
- [23] Popov S B, Kaurav A A and Kaminker A D 2015 *PASA* **32** e018
- [24] Zavlin V E, Pavlov G G, Sanwal D and Trümper J 2000 *ApJ* **540** L25-8
- [25] Gotthelf E V, Halpern J P and Seward F D 2005 *ApJ* **627** 390-6
- [26] Gotthelf E V and Halpern J P 2009 *ApJ* **695** L35-9
- [27] Halpern J P and Gotthelf E V 2010 *ApJ* **709** 436-46
- [28] Shabaltas N and Lai D 2012 *ApJ* **748** 148

- [29] Bogdanov S 2014 *ApJ* **790** 94
- [30] Gotthelf E V, Perna R and Halpern J P 2010 *ApJ* **724** 1316-24
- [31] De Luca A *et al* 2012 *MNRAS* **421** L72-6
- [32] Sanwal D, Pavlov G G, Zavlin V E and Teter M A 2002 *ApJ* **574**, L61-4
- [33] Mereghetti S, De Luca A, Caraveo P A, Becker W, Mignani R and Bignami G F 2002 *ApJ* **581** 1280-5
- [34] Bignami G F, Caraveo P A, De Luca A and Mereghetti S 2003 *Nature* **423** 725-7
- [35] De Luca A, Mereghetti S, Caraveo P A, Moroni M, Mignani R P and Bignami G F 2004 *A&A* **418** 625-37
- [36] Suleimanov V F, Pavlov G G and Werner K 2010 *ApJ* **714** 630-5
- [37] Potekhin A Y 2010 *A&A* **518** A24
- [38] Suleimanov V F, Pavlov G G and Werner K 2012 *ApJ* **751** 15
- [39] Tananbaum H 1999 *IAUC* **7246** 1
- [40] Krause O *et al* 2005 *Sci* **308** 1604-6
- [41] Kim Y, Rieke G H, Krause O, Misselt K, Indebetouw R and Johnson K E 2008 *ApJ* **678** 287-96
- [42] Dwek E and Arendt R G 2008 *ApJ* **685** 976-87
- [43] Heinke C O and Ho W C G 2010 *ApJ* **719** L167-71
- [44] Shternin P S, Yakovlev D G, Heinke C O, Ho W C G and Patnaude D J 2011 *MNRAS* **412** L108-12
- [45] Yakovlev D G, Ho W C G, Shternin P S, Heinke C O and Potekhin A Y 2011 *MNRAS* **411** 1977-88
- [46] Page D, Prakash M, Lattimer J M and Steiner A W 2011 *PhRvL* **106** 081101
- [47] Yang S-H, Pi C-M and Zheng X-P 2011 *ApJ* **735** L29
- [48] Noda T, Hashimoto M, Yasutake N, Maruyama T, Tatsumi T and Fujimoto M 2013 *ApJ* **765** 1
- [49] Elshamouty K G *et al* 2013 *ApJ* **777** 22
- [50] Posselt B, Pavlov G G, Suleimanov V and Kargaltsev O 2013 *ApJ* **779** 186
- [51] Ho W C G and Heinke C O 2009 *Nature* **462** 71-3
- [52] Klochkov D, Puehlhofer G, Suleimanov V, Simon S, Werner K and Santangelo A 2013 *A&A* **556** A41
- [53] Klochkov D, Suleimanov V, Puehlhofer G, Yakovlev D G, Santangelo A and Werner K 2015 *A&A* **573** A53
- [54] Klochkov D, Suleimanov V, Sasaki M and Santangelo A 2016 *A&A* **592** L12
- [55] Ofengeim D D, Kaminker A D, Klochkov D, Suleimanov V and Yakovlev D G 2015 *MNRAS* **454** 2668-76
- [56] Pavlov G G and Luna G J M 2009 *ApJ* **703** 910-21
- [57] Suleimanov V F, Klochkov D, Poutanen J and Werner K 2017 *A&A* **600** A43
- [58] De Luca A *et al* 2011 *A&A* **525** A106
- [59] Pérez-Azorín J F, Miralles J A and Pons J A 2006 *A&A* **451** 1009-24
- [60] Braithwaite J 2009 *MNRAS* **397** 763-74
- [61] Kaspi V M 2010 *PNAS* **107** 7147-52
- [62] Halpern J P and Gotthelf E V 2010 *ApJ* **710** 941-7
- [63] Faucher-Giguère C-A and Kaspi V M 2006 *ApJ* **643** 332-55
- [64] Keane E F and Kramer M 2008 *MNRAS* **391** 2009-16
- [65] Gotthelf E V and Halpern J P 2007 *ApJ* **664** L35-8
- [66] Halpern J P, Gotthelf E V, Camilo F and Seward F D 2007 *ApJ* **665** 1304-10
- [67] Bonanno A, Urpin V and Belvedere G 2006 *A&A* **451** 1049-52
- [68] Muslimov A and Page D 1995 *ApJ* **440** L77-80
- [69] Geppert U, Page D and Zannias T 1999 *A&A* **345** 847-54
- [70] Ho W C G 2011 *MNRAS* **414** 2567-75
- [71] Viganò D and Pons J A 2012 *MNRAS* **425** 2487-92
- [72] Bernal C G, Page D and Lee W H 2013 *ApJ* **770** 106
- [73] Torres-Forné A, Cerdá-Durán P, Pons J A and Font J A 2016 *MNRAS* **456** 3813-26
- [74] Bernal C G and Fraija N 2016 *MNRAS* **462** 3646-59
- [75] Igoshev A P, Elfriz J G and Popov S B 2016 *MNRAS* **462** 3689-702
- [76] Rogers A and Safi-Harb S 2016 *MNRAS* **457** 1180-9
- [77] Ho W C G 2015 *MNRAS* **452** 845-51
- [78] Marshall F E, Guillemot L, Harding A K, Martin P and Smith D A 2016 *ApJ* **827** L39
- [79] Popov S B and Turolla R 2012 *ASPC* **466** 191
- [80] Gotthelf E V, Halpern J P, Allen B and Knispel B 2013 *ApJ* **773** 141
- [81] Bogdanov S, Ng C-Y and Kaspi V M 2014 *ApJ* **792** L36
- [82] Luo J, Ng C-Y, Ho W C G, Bogdanov S, Kaspi V M and He C 2015 *ApJ* **808** 130
- [83] Rutledge R E, Fox D B and Shevchuk A H 2008 *ApJ* **672** 1137-43
- [84] Zane S *et al* 2011 *MNRAS* **410** 2428-45
- [85] Halpern J P, Bogdanov S and Gotthelf E V 2013 *ApJ* **778** 120
- [86] Halpern J P and Gotthelf E V 2015 *ApJ* **812** 61