



Publication Year	2017
Acceptance in OA @INAF	2020-11-20T16:48:17Z
Title	Cosmic ray-driven winds in the Galactic environment and the cosmic ray spectrum
Authors	Recchia, S.; Blasi, P.; MORLINO, GIOVANNI
DOI	10.1093/mnras/stx1214
Handle	http://hdl.handle.net/20.500.12386/28496
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	470

Cosmic ray driven winds in the Galactic environment and the cosmic ray spectrum

S. Recchia ^{1*} P. Blasi, ^{2,1†} G. Morlino ^{1,2‡}

¹ *Gran Sasso Science Institute (INFN), Viale F. Crispi 7 - 67100 L' Aquila, Italy*

² *INAF/Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5 - 50125 Firenze, Italy*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Cosmic Rays escaping the Galaxy exert a force on the interstellar medium directed away from the Galactic disk. If this force is larger than the gravitational pull due to the mass embedded in the Galaxy, then galactic winds may be launched. Such outflows may have important implications for the history of star formation of the host galaxy, and in turn affect in a crucial way the transport of cosmic rays, both due to advection with the wind and to the excitation of waves by the same cosmic rays, through streaming instability. The possibility to launch cosmic ray induced winds and the properties of such winds depend on environmental conditions, such as the density and temperature of the plasma at the base of the wind and the gravitational potential, especially the one contributed by the dark matter halo. In this paper we make a critical assessment of the possibility to launch cosmic ray induced winds for a Milky-Way-like galaxy and how the properties of the wind depend upon the conditions at the base of the wind. Special attention is devoted to the implications of different conditions for wind launching on the spectrum of cosmic rays observed at different locations in the disc of the galaxy. We also comment on how cosmic ray induced winds compare with recent observations of Oxygen absorption lines in quasar spectra and emission lines from blank-sky, as measured by XMM-Newton/EPIC-MOS.

Key words: keyword1 – keyword2 – keyword3

1 INTRODUCTION

Galactic winds have been observed in many galaxies (see e.g. [Veilleux et al. \(2005\)](#); [Martin et al. \(2012\)](#); [King & Pounds \(2015\)](#)) and constitute an important feedback process to the evolution of galaxies. In fact, by removing gas from the galactic disk galactic winds regulate the star formation rate (see e.g. [Crain et al. \(2007\)](#)) and pollute galactic halos and intergalactic space with hot plasma and metals, thus affecting the chemical composition, the temperature and the degree of ionization of the interstellar medium (ISM) and of the intergalactic medium (IGM) (see e.g. [Miller & Bregman \(2015\)](#)). Moreover, the expelled material may contain an appreciable fraction of the number of baryons in the Universe (see e.g. [Kalberla & Dedes \(2008\)](#); [Miller & Bregman \(2015\)](#)) and may contribute to solving the problem of missing baryons in the local Universe (see e.g. [Anderson & Bregman \(2010\)](#)). Although observations of the Milky Way have not yet led to conclusive evidence of the existence of a

Galactic wind, hot dilute gas, possibly connected with such winds, has been detected through the continuous emission in the X-ray band ([Breitschwerdt & Schmutzler \(1994\)](#); [Breitschwerdt & Schmutzler \(1999\)](#); [Everett et al. \(2008\)](#)) and through Oxygen absorption lines (O_{VII} and O_{VIII} lines) in the spectrum of distant Quasars ([Miller & Bregman \(2015\)](#)). The features of such lines are compatible with a halo with a mass of $\sim 10^{10}M_{\odot}$, an extension of ~ 100 kpc, a temperature in the range $10^6 - 10^7$ K and an average metallicity of $Z \sim 0.2 - 0.3$. In particular the latter data imply that the gas in the halo comes from the Galaxy and not from intergalactic space ([Miller & Bregman \(2015\)](#)). Moreover, the recent observation of the so called Fermi Bubbles may be associated to recent bursts near the Galactic Center region (see [Cheng et al. \(2011\)](#); [Zubovas et al. \(2011\)](#)) or to past starburst activities (see [Lacki \(2014\)](#)).

Galactic winds may be 1) thermally-driven, namely generated by the heating of the ISM due to SN explosions (see e.g. [Chevalier & Clegg \(1985\)](#)) or by accretion onto the super-massive black holes in the center of AGN (see e.g. [King & Pounds \(2015\)](#)); 2) radiation pressure-driven (see e.g. [Scoville \(2003\)](#); [Murray et al. \(2005\)](#)); 3) CR-driven, namely due to the pressure exerted by escaping CRs on the ISM (see

* E-mail: sarah.recchia@gssi.infn.it

† E-mail: blasi@arcetri.astro.it

‡ E-mail: giovanni.morlino@gssi.infn.it

e.g. [Ipavich \(1975\)](#), [Breitschwerdt et al. \(1991\)](#); [Uhlig et al. \(2012\)](#)). The first two mechanisms are likely to take place in starburst galaxies and galaxies with active nuclei.

In the Milky Way the thermal and radiation pressure gradients are expected to be too small (except perhaps for the Galactic Center region) to drive outflows. However, CRs escaping from the Galaxy lead to a gradient in the CR density with several implications. First, the CR pressure gradient exerts a force, $-\nabla P_c$, on the background plasma directed away from the disk and opposite to the Galactic gravitational force (due to both baryonic and dark matter). If this force is larger than the gravitational pull it may launch a wind. Second, the CR density gradient can excite Alfvén waves, due to CR streaming instability ([Kulsrud & Cesarsky \(1971\)](#); [Skillington \(1975\)](#)), which move in the direction of the decreasing CR density. Such waves affect the scattering properties of CRs, namely their diffusive transport. Third, winds and the waves that propagate inside the outgoing plasma, advect CRs and contribute to change the spectrum of CRs as observed in the Galaxy.

CRs have been widely recognized as an appealing way to launch winds in our Galaxy and the hydrodynamics of CR-driven winds has been extensively studied in stationary one dimensional calculations (see e.g. [Ipavich \(1975\)](#); [Breitschwerdt et al. \(1991\)](#); [Everett et al. \(2008\)](#); [Zirakashvili et al. \(1996\)](#)), in time-dependent calculations ([Dorfi & Breitschwerdt \(2012\)](#)), in hydrodynamical simulations (see [Uhlig et al. \(2012\)](#); [Booth et al. \(2013\)](#); [Salem & Bryan \(2014\)](#)) and in MHD simulations (see [Girichidis et al. \(2016\)](#); [Ruszkowski et al. \(2016\)](#)). The paper by [Breitschwerdt et al. \(1991\)](#) represented a crucial step forward in the investigation of CR-driven Galactic winds: their computation techniques have been adopted by much of the subsequent work on the subject.

In all this bulk of work, no matter the level of refinement, only the hydrodynamics of CR-driven winds was addressed, while no information was retained on the CR spectrum in such winds. On the other hand, the CR spectrum represents a crucial observable that we can use in order to constraint wind models, hence its prediction is essential. Recently, our group ([Recchia et al. 2016](#), hereafter Paper I) presented an extensive study of the problem of the CR transport in a CR-driven wind. We developed a semi-analytical method which allows to compute self-consistently the hydrodynamical properties of CR-driven winds and the corresponding CR distribution function in the presence of self-generated Alfvén waves. In our approach, the hydrodynamics of the wind is calculated following the one-dimensional stationary model developed by [Breitschwerdt et al. \(1991\)](#) and the CR transport is due to scattering of CRs on the self-generated turbulence and to the CR advection with such waves and with the wind. Both the flow and the CR transport only occur along the magnetic field lines and the flow geometry is preassigned. The CR diffusion coefficient is calculated from the spectrum of waves excited by streaming instability and locally damped through non linear Landau damping (NLLD). The importance of CRs for wind launching was confirmed by our work, although we found that in general, the CR spectrum in the presence of winds launched in the vicinity of the Sun is not in agreement with observations. This is mainly due to the fact that CR advection in such winds is strong, leading to spectra at low energies

(below ~ 200 GeV) which are harder than the observed spectrum. At high energy (above ~ 200 GeV), instead, the wind expansion together with the steep energy dependence of the self-generated diffusion coefficient, leads to spectra that are steeper than the observed spectrum. However, these conclusions strongly depend, at low energy, on the wind launching parameters and, at high energy, on the physical conditions in the near-disk region, in particular on the amount of neutral gas in such region and on its volume filling factor. For instance, we showed that at high energy the CR spectrum is very sensitive to the diffusion properties of the near-disk region ($\lesssim 1$ kpc) where the presence of a pre-existing turbulence would lead to a spectral hardening in the high energy part of the spectrum (see also [Aloisio et al. \(2015\)](#)), in agreement with recent observations of the proton and helium spectrum (see [Adriani et al. \(2011\)](#); [Aguilar et al. \(2015\)](#)).

In this paper we use the calculations developed in [Paper I](#) and we study the possibility to launch winds, the dependence of the wind properties on environmental parameters and the implications of such dependences for the spectrum of CRs observed at the Sun position. Special attention is devoted to investigating the wind properties as functions of the ISM properties (gas density and temperature, Galactic magnetic field), of the CR pressure, of the flow geometry and on the Galactic gravitational potential.

The spatial dependence of these quantities in the Galaxy is constrained by observations (see e.g. [Ferrière \(2001\)](#); [Cox \(2005\)](#); [Wolfire et al. \(2003\)](#); [Miller & Bregman \(2015\)](#)). In particular, the observation of the Oxygen absorption lines in the spectrum of distant Quasars analyzed by [Miller & Bregman \(2015\)](#) suggests the presence of a large Galactic halo, presumably due to a Galactic wind, and provides us with indications on the gas mass distribution and temperature of such halo, and consequently on the flow geometry. Moreover, the baryonic and DM content of the Milky Way and the corresponding gravitational potential can be constrained by analyzing the dynamics of stars and gas (see e.g. [Bertone et al. \(2005\)](#); [Sofue \(2012\)](#); [Nesti & Salucci \(2013\)](#)). We discuss how our results change by adopting different distributions of Galactic dark matter and compare them with the halo structure deduced by [Miller & Bregman \(2015\)](#).

The paper is organized as follows: in [Section 2](#) we describe our wind model and we briefly summarize the method used to solve the hydrodynamic wind equations and the CR transport equation, as illustrated in [Paper I](#). In [Section 3](#) we describe the model used for the gravitational potential of the Galaxy, including several DM halo models. In [Section 4](#) we summarize the main properties of the ISM as deduced from observations and their relevance for the models of CR driven winds. In [Section 5](#), we discuss in detail the dependence of the wind properties on the parameters of the problem, as well as on the geometry of the outflow and the Galactic gravitational potential. Finally, in [Section 6](#), we illustrate the implications of CR driven winds launched at the position of the Sun on the spectrum of CRs.

2 THE WIND MODEL

The dynamics of a CR-driven wind in the stationary regime is described by the system of hydrodynamic equations for the wind (i.e the equations of mass, momentum and energy conservation) and by the kinetic transport equation for CRs. The hydrodynamic wind equations have been written down and discussed in detail by Breitschwerdt et al. (1991). A method to calculate the combined solution of the hydrodynamical equations for the wind and the transport equation of CRs has been recently developed in Paper I. This method allows one to determine at the same time the wind structure and the related CR spatial and momentum distribution. Here we briefly summarize the equations and the main features of the wind model.

Following Breitschwerdt et al. (1991), we assume that the wind is launched from a surface at distance z_0 above (and below) the Galactic disc. The geometry of the outflow induced by the CR pressure gradient is assumed to be pre-assigned and reflecting the geometry of magnetic field lines. In our calculation the CR diffusion coefficient is due to scattering off self-generated Alfvén waves, as due to the excitation of streaming instability, saturated through NLLD. Advection of CRs with self-generated waves and with the wind are taken into account. Wave damping is considered to occur on a timescale much shorter than any other considered process (definitely justified for NLLD) and contribute to the gas heating. Under these assumptions, as shown in Paper I, the hydrodynamic wind equations and the CR transport equation become effectively one-dimensional and only depend on the distance z from the Galactic disk. The hydrodynamic equations read

$$\rho u A = \text{const}, \quad (1)$$

$$A B = \text{const}, \quad (2)$$

$$\frac{du}{dz} = u \frac{c_*^2 \frac{1}{A} \frac{dA}{dz} - \frac{d\Phi}{dz}}{u^2 - c_*^2}, \quad (3)$$

$$\frac{dP_g}{dz} = \gamma_g \frac{P_g}{\rho} \frac{d\rho}{dz} - (\gamma_g - 1) \frac{v_A}{u} \frac{dP_c}{dz} \quad (4)$$

$$\frac{dP_c}{dz} = \gamma_{eff} \frac{P_c}{\rho} \frac{2u + v_A}{2(u + v_A)} \frac{d\rho}{dz}, \quad (5)$$

$$c_*^2 = \gamma_g \frac{P_g}{\rho} + \gamma_{eff} \frac{P_c}{\rho} \left[1 - (\gamma_g - 1) \frac{v_A}{u} \right] \frac{2u + v_A}{2(u + v_A)}, \quad (6)$$

$$\frac{\gamma_{eff}}{\gamma_{eff} - 1} = \frac{\gamma_c}{\gamma_c - 1} - \frac{\bar{D}}{(\gamma_c - 1)(u + v_A)P_c} \frac{dP_c}{dz}. \quad (7)$$

where $\rho(z)$, $u(z)$ and $P_g(z)$ are the gas density, velocity and pressure, $B(z)$ is the background magnetic field strength and $\gamma_g = 5/3$ is the adiabatic index of the gas. $P_c(z)$ is the CR pressure and $\gamma_c(z)$ is the adiabatic index of the CR gas, while the average CR diffusion coefficient \bar{D} is defined in equation 11. Notice that $\gamma_c(z)$ is actually calculated as a function of z from the distribution function $f(p, z)$ that solves the CR transport equation. The space-dependent Alfvén velocity is $v_A(z) = B(z)/\sqrt{4\pi\rho(z)}$, while $\Phi(z)$ is the gravitational potential of the Galaxy. The wave pressure does not appear explicitly since the assumption of fast damping results in a wave pressure much smaller than the gas and CR pressure. Two important quantities appear in these equations: the generalized sound speed c_* and the effective adiabatic index for CRs, γ_{eff} . Looking at the definition of c_*^2 , equation 6, one can recognize the sum of the sound speed term (square

root of $\gamma_g P_g/\rho$) and of a "CR sound speed" term (the term $\gamma_{eff} P_c/\rho$ is formally identical to the square root of the sound speed) which depends on the Alfvén Mach number u/v_A . Notice that the term $-(\gamma_g - 1)v_A/u$ is due to the non adiabatic gas heating induced by wave damping in equation 4. The importance of c_* resides in the fact that $u = c_*$ is the condition for the flow to become sonic but also is a critical point for the wind equation 3. The effective adiabatic index for CRs, γ_{eff} , takes into account the CR diffusivity in the hydrodynamic equations (see also Zirakashvili et al. (1996)). In general, γ_{eff} is a function of z , however, it has been verified that in general (including in the cases presented here) γ_{eff} shows a weak dependence on z and we can safely treat it as a constant. Typical values of $\gamma_{eff} \sim 1.1 - 1.2$, while $\gamma_c \sim 4/3$. Breitschwerdt et al. (1991) also introduced the generalized sound speed c_* but with two important differences compared with the calculation discussed here. First, they neglected the CR diffusion by assuming that $\bar{D} = 0$ so that in their calculation γ_c appears instead of γ_{eff} . Neglecting diffusion is not a good approximation, at least not at all distances z from the disc. Second, Breitschwerdt et al. (1991) did not take wave damping into account and retained the wave pressure in the hydrodynamic equations. Without damping, the growth of waves due to CR streaming can in fact lead to a wave pressure which is comparable with the CR pressure. For this reason, instead of having the term $-(\gamma_g - 1)v_A/u$ in the definition of c_*^2 , they have a "wave sound speed" term which takes into account the effect of the wave pressure.

The CR transport equation reads

$$\frac{\partial}{\partial z} \left[A D \frac{\partial f}{\partial z} \right] - A U \frac{\partial f}{\partial z} + \frac{d(AU)}{dz} \frac{1}{3} \frac{\partial f}{\partial \ln p} + A Q = 0, \quad (8)$$

where $f(z, p)$ and $D(z, p)$ are the CR distribution function and diffusion coefficient as functions of position z and momentum p and $U(z) = u(z) + v_A(z)$ is the advection velocity including the wind speed and the Alfvén speed.

The term $Q(z, p)$ represents the injection of CRs in the Galaxy, which we assume to be limited to the Galactic disc (considered as infinitely thin) and can be written as $Q(z, p) = Q_0(p)\delta(z)$. The function $Q_0(p)$ describes the injection spectrum,

$$Q_0(p) = \frac{\mathcal{N}_{SN}(p)\mathcal{R}_{SN}}{\pi R_d^2}, \quad (9)$$

where R_d is the Galactic disk and $\mathcal{N}_{SN}(p)$ is the spectrum contributed by individual supernovae (SN) occurring at a rate \mathcal{R}_{SN} and reads

$$\mathcal{N}_{SN}(p) = \frac{\xi_{CR} E_{SN}}{I(\gamma)c(mc)^4} \left(\frac{p}{mc} \right)^{-\gamma}. \quad (10)$$

In the last expression ξ_{CR} is the CR injection efficiency (typically $\sim 10\%$), E_{SN} is the energy released by a SN explosion ($\sim 10^{51}$ erg), \mathcal{R}_{SN} is the rate of SN explosions ($\sim 1/30 \text{ yr}^{-1}$), and $I(\gamma)$ is a normalization factor (see Paper I).

The average diffusion coefficient of equation 7 is defined as:

$$\bar{D}(z) = \frac{\int_0^\infty dp p^2 T(p) D(z, p) \nabla f}{\int_0^\infty dp p^2 T(p) \nabla f}, \quad (11)$$

where $T(p)$ is the kinetic energy of particles with momentum p . The diffusion coefficient $D(z, p)$ is determined by the local balance between the CR-driven growth of Alfvén waves

and their damping through NLLD. Since the self-generated perturbations in the magnetic field are relatively weak, one can use quasi-linear theory to write the diffusion coefficient as:

$$D(z, p) = \frac{1}{3} \frac{v(p)r_L(z, p)}{\mathcal{F}(z, k_{res})} \Big|_{k_{res}=1/r_L}, \quad (12)$$

where \mathcal{F} is the normalized energy density per unit logarithmic wavenumber k , calculated at the resonant wavenumber $k_{res} = 1/r_L(p)$. In the regions where the background gas is totally ionized, waves are damped through NLLD at a rate (see [Zhou & Matthaues \(1990\)](#); [Ptuskin & Zirakashvili \(2003\)](#)):

$$\Gamma_D = (2c_k)^{-3/2} k v_A \mathcal{F}^{1/2}, \quad (13)$$

where $c_k = 3.6$. On the other hand the CR-driven growth occurs at a rate which is given by [Skilling \(1971\)](#):

$$\Gamma_{CR} = \frac{16\pi^2}{3} \frac{v_A}{\mathcal{F}B^2} \left[p^4 v(p) \left| \frac{\partial f}{\partial z} \right| \right]_{p=p_{res}}. \quad (14)$$

Equating the two rates, $\Gamma_D = \Gamma_{CR}$, one obtains:

$$\mathcal{F}(z, p) = 2c_k \left[\frac{p^4 v(p) \left| \frac{\partial f}{\partial z} \right| \frac{16\pi^2}{3} r_L(z, p)}{B^2(z)} \right]^{2/3}, \quad (15)$$

where \mathcal{F} is expressed as a function of momentum by means of the resonant condition $p = p_{res}(k)$.

As for the flow geometry, we adopt one that is similar to the one introduced by [Breitschwerdt et al. \(1991\)](#) namely we assume that the wind is launched at the wind base z_0 and that it retains a cylindrical geometry out to a characteristic distance Z_b . At larger distances the flow opens up as $\sim z^\alpha$. The wind area transverse to z is then assumed to be in the form:

$$A(z) = A_0 \left[1 + \left(\frac{z}{Z_b} \right)^\alpha \right], \quad (16)$$

which is only function of the spatial coordinate z , namely of the distance from the Galactic disk. Notice that in [Breitschwerdt et al. \(1991\)](#) and in [Paper I](#) the authors assumed $Z_b \sim 15$ kpc and $\alpha = 2$, which corresponds to spherical opening. Here, instead, we explore the implications of changing these two parameters for the wind properties.

Following [Paper I](#), the hydrodynamic equations for the wind and the CR transport equation are solved together following an iteration scheme. For a given set of input parameters at the wind base z_0 (namely the gas density ρ_0 , the gas temperature T_0 , the magnetic field strength B_0 and the CR pressure P_{c0}), for each iteration we determine the velocity u_0 at the base of the wind, for which the flow experiences a smooth transition from the subsonic to the supersonic regime. Given u_0 , both the mass and energy flux of the wind are fixed, and it is possible to compute all the hydrodynamic quantities (gas density and pressure, wind velocity and CR pressure) as functions of the distance from the Galactic disk. For each iteration, the transport equation 8 is solved for the CR distribution function $f(z, p)$ and the diffusion coefficient $D(p, z)$. The iteration is then repeated until convergence is reached within a given accuracy.

3 THE GALACTIC GRAVITATIONAL POTENTIAL

The mass distribution of the Milky Way, and of other galaxies as well, is mainly inferred from analyses of the dynamics of stars and gas, namely by studying kinematic data such as rotation velocities, velocity dispersions and motion of satellite galaxies. Current models for the mass distribution of the Galaxy include a central bulge, a stellar and gas disk and a DM halo (see e.g [Irrgang et al. \(2013\)](#)).

Following [Breitschwerdt et al. \(1991\)](#) and [Irrgang et al. \(2013\)](#), the gravitational potential of the Galactic bulge and disk can be modelled using the parametrization of [Miyamoto & Nagai \(1975\)](#):

$$\Phi_{B,D}(R_0, z) = - \sum_{i=1}^2 \frac{GM_i}{\sqrt{R_0^2 + (a_i + \sqrt{z^2 + b_i^2})^2}}, \quad (17)$$

where z and R_0 are the distance from the Galactic disk and the Galactocentric distance respectively. For the parameters of this model we used the values proposed by [Sofue \(2012\)](#), namely $(M_1, a_1, b_1) = (1.652 \times 10^{10} M_\odot, 0.0$ kpc, 0.522 kpc) for the bulge and $(M_2, a_2, b_2) = (3.4 \times 10^{10} M_\odot, 3.19$ kpc, 0.289 kpc) for the disk.

For the dark matter mass distribution we considered three models, namely the Navarro-Frenk-White ([Navarro et al. 1996](#)), the Burkert ([Burkert 1995](#)) and the Einasto ([Retana-Montenegro et al. 2012](#)) profiles:

- Navarro-Frenk-White (NFW) ([Navarro et al. \(1996\)](#)):

the density profile is of the form

$$\rho_{NFW} = \frac{\rho_0}{x(1+x)^2}, \quad (18)$$

where $x = r/r_c$ and r_c is the scale radius of the distribution. For the two quantities ρ_0 and r_c we considered both the values given for the Milky Way by [Sofue \(2012\)](#) ($\rho_0 = 1.06 \times 10^7 M_\odot \text{ kpc}^{-3}$; $r_c = 12.0$ kpc) and by [Nesti & Salucci \(2013\)](#) ($\rho_0 = 1.3 \times 10^7 M_\odot \text{ kpc}^{-3}$; $r_c = 16.0$ kpc). We refer to the two sets of parameters as NFW-Sofue and NFW-Salucci respectively. The DM halo is assumed to extend out to a maximum distance that equals the virial radius, which corresponds to $r_{vir} \approx 240$ kpc for NFW-Sofue and $r_{vir} \approx 320$ kpc for NFW-Salucci.

- Burkert (BUR) ([Burkert \(1995\)](#)):

the density profile is of the form

$$\rho_{BUR} = \frac{\rho_0}{(1+x)(1+x^2)}, \quad (19)$$

where $x = r/r_c$ and r_c is the core radius. For the two quantities ρ_0 and r_c we considered the values given for the Milky Way by [Nesti & Salucci \(2013\)](#) ($\rho_0 = 4.13 \times 10^7 M_\odot \text{ kpc}^{-3}$; $r_c = 9.3$ kpc). The virial radius for this model is $r_{vir} \approx 300$ kpc.

- Einasto (EIN) ([Retana-Montenegro et al. \(2012\)](#)):

the density profile is of the form

$$\rho_{EIN} = \rho_0 \exp(-x^\alpha), \quad (20)$$




Figure 1. Galactic gravitational acceleration as function of the distance from the galactic plane, z , at the Sun position produced by different DM profiles. The contribution due to the disk and the bulge are also shown.

where $x = r/r_c$. For the three quantities ρ_0 , r_c and α we considered the values given for the Milky Way by Bernal & Palomares-Ruiz (2012) ($\rho_0 = 3.5 \times 10^{11} \text{ M}_\odot \text{ kpc}^{-3}$; $r_c = 6.7 \times 10^{-6} \text{ kpc}$, $\alpha = 0.17$).

These halo models predict a DM density at the Sun position in the range $0.2 - 0.4 \text{ GeV/cm}^3$, in agreement with most of the current literature (see Salucci et al. (2010) for a discussion on the local DM density).

A comparison between the gravitational acceleration produced by different DM models at the solar galactocentric distance $R = R_\odot$ and as a function of the Galactic altitude is shown in Fig. 1 where also the contribution due to the disk and the bulge are reported. One can see that the Burkert and the Einasto profiles are quite similar in magnitude below $z \sim 50 \text{ kpc}$ and that the Einasto profile becomes systematically larger than the Burkert profile above $z \sim 50 \text{ kpc}$. On the other hand, the NFW profile shows large differences in magnitude, compared to the other two profiles, below $z \sim 30 - 40 \text{ kpc}$. Since these profiles predict a similar DM density at the Sun, this is due to the different functional dependence on $x = r/r_c$ and to the different values of the scale radius r_c .

4 FIDUCIAL VALUES FOR THE ENVIRONMENTAL PARAMETERS

The ISM is a rich environment in which several components, distinguished by the level of ionization and temperature, co-exist and interact: molecular and atomic neutral gas, warm and hot ionized gas, interstellar magnetic field and CRs. In the cooler, denser regions of the ISM, matter is primarily in molecular form and can reach densities as high as 10^6 cm^{-3} . On the other hand, in the diffuse regions of the ISM, matter is mainly ionized and can reach densities as low as 10^{-4} cm^{-3} . In addition, the colder and denser phases of the ISM are mainly confined in the Galactic disk and have filling factor of order $\sim 1 - 5\%$, much smaller than the dilute hot gas filling factor, which is of order $30 - 70\%$. A detailed study of

Table 1. Components of the ISM and their properties in the vicinity of the Sun (see Ferrière (2001)).

Component	Temperature (K)	Density (cm^{-3})	Scale Height (kpc)
Molecular	10-20	$10^2 - 10^6$	~ 0.1
Cold Atomic	50-100	20-50	$\sim 0.1 - 0.4$
Warm Atomic	6000-10000	0.2-0.5	$\sim 0.1 - 0.4$
Warm Ionized	8000	0.2-0.5	~ 1
Hot Ionized	$\sim 10^6$	~ 0.006	> 1

the gas components of the ISM has been carried out by several authors (see e.g Cox (2005); Wolfire et al. (2003)) while a comprehensive review of the present knowledge of the ISM can be found in (Ferrière (2001)). In Table 1 we summarize the main properties of the gas components of the ISM in the vicinity of the Sun as reported by Ferrière (2001).

The Galactic magnetic field shows a rich structure in which at least a disk component, which follows the spiral arms, and an out of disk component can be found (see Jansson & Farrar (2012)). The regular field strength changes with the Galactocentric distance. An estimation of the magnetic field strength in the vicinity of the Sun is reported for instance in Jansson & Farrar (2012), which quote the value $\sim 1 - 2 \mu\text{G}$ for the disk component and $\sim 1 \mu\text{G}$ for the out of disk component. Ferrière (2001) quote a value $B_\odot \sim 1.5 \mu\text{G}$, while Cox (2005) suggest a value of $\sim 3 - 5 \mu\text{G}$ (which includes all field components). Here we are mainly interested in the out of disk component.

Recently Miller & Bregman (2015) analyzed the Oxygen absorption lines in quasar spectra and emission lines from blank-sky regions, measured by XMM-Newton/EPIC-MOS, and inferred the presence of a hot gaseous halo which could in fact be the result of a Galactic wind. Constraints on the structure of the hot halo can be found by fitting a radial model,

$$n(r) \approx n_0 \left(\frac{r_c}{r} \right)^{3\beta}, \quad (21)$$

for the halo density distribution, from which the expected emission, to compare with observations, is computed. Here $r = \sqrt{R^2 + z^2}$, where R is the Galactocentric distance and z the distance from the Galactic disk. The best fit parameters for the halo density given by Miller & Bregman (2015) are $\beta = 0.5 \pm 0.3$ and $n_0 r_c^{3\beta} = 1.35 \pm 0.24 \text{ cm}^{-3} \text{ kpc}^{3\beta}$. The authors also infer a nearly constant halo temperature of $\sim 2 \times 10^6 \text{ K}$ and a sub-solar gas metallicity that decreases with r , but that also must be $\gtrsim 0.3 Z_\odot$ to be consistent with the pulsar dispersion measure toward the Large Magellanic Cloud.

This halo model implies a gas density of $3 - 6 \times 10^{-3} \text{ cm}^{-3}$ in the vicinity of the Sun, a density scaling with z that reads $n(z) \sim z^{-1.5}$, and a halo mass of $M(< 50 \text{ kpc}) = (3.8 \pm 0.3) \times 10^9 \text{ M}_\odot$ and $M(< 250 \text{ kpc}) = (4.3 \pm 0.8) \times 10^{10} \text{ M}_\odot$. Notice that such halo mass would account for $\lesssim 50\%$ of the Milky Way

Table 2. Fiducial values for the wind input parameters at the Sun position.

parameter	fiducial range
area- α	1.5-2.0
area- Z_b	5-15 kpc
gas density	$3 - 6 \times 10^{-3} \text{ cm}^{-3}$
gas temperature	$1 - 3 \times 10^6 \text{ K}$
regular B	$1 - 2 \mu\text{G}$
CR pressure	$4 \times 10^{-13} \text{ erg/cm}^{-3}$

missing baryons.

All these pieces of information allow us to define reasonable fiducial values for the input parameters of our wind problem (see Table 2 for a summary): we are interested in the hot dilute phase of the ISM, for which we adopt a density $3 - 6 \times 10^{-3} \text{ cm}^{-3}$ and a temperature $1 - 3 \times 10^6 \text{ K}$, and in the out of disk regular magnetic field for which we retain a field strength $1 - 2 \mu\text{G}$.

The pressure in the form of CR protons measured at Earth is $\sim 4 \times 10^{-13} \text{ erg/cm}^{-3}$.

Finally, Breitschwerdt et al. (1991), Everett et al. (2008) and Paper I assumed that the flux-tube opens up spherically, namely $\alpha = 2$ in equation 16, while the results of Miller & Bregman (2015) suggest $\alpha \sim 1.5$. As for the area length scale Z_b , Breitschwerdt et al. (1991) and Paper I adopted the value 15 kpc, while Everett et al. (2008) treated Z_b as a fitting parameter and allowed it to vary around ~ 5 kpc. Here we consider $\alpha \sim 1.5 - 2$ and $Z_b \sim 5 - 15$ kpc.

5 DEPENDENCE OF THE WIND PROPERTIES ON THE GALACTIC ENVIRONMENT

In this section we present a purely hydrodynamical analysis of Galactic winds, launched at the Sun position, focusing on how the wind's properties are affected when changing the environmental parameters within the range allowed by observations, as summarized in Section 4. In Section 5.3 we briefly discuss the role of radiative cooling while we describe the implications for the CR spectrum in such winds in Section 6.

The topology of the solutions of the hydrodynamic equations 1-7 depends on the nature of the critical points of the wind equation 3 (see Breitschwerdt et al. (1991); Paper I), i.e. of the points in which the velocity derivative has zero numerator ($c_*^2 = \frac{d\Phi}{dz} / \frac{1}{A} \frac{dA}{dz}$) and/or zero denominator ($u^2 = c_*^2$). The point for which both the numerator and the denominator vanish is the critical (sonic) point, and it corresponds to the location where the flow velocity equals the compound sound speed, i.e. $u = c_*$. The solution relevant for our problem is the one with velocity that starts

subsonic at the wind base z_0 , increases with z , goes through the critical point where it becomes supersonic and keeps increasing (wind acceleration). The wind launching velocity u_0 for this solution is found by imposing crossing through the critical point. For launching velocities smaller than u_0 , the flow remains subsonic everywhere and there is a point in which the numerator of equation 3 vanishes. Such solutions are called “breezes”. For launching velocities larger than u_0 but still subsonic, there is a point in which the denominator of equation 3 vanishes and the corresponding solutions are unphysical.

The hydrodynamic equations, and in particular the mass and energy conservation equations and the wind equation,

$$\rho u A = \text{const} \quad (22)$$

$$\frac{u^2}{2} + \frac{\gamma_g}{\gamma_g - 1} \frac{P_g}{\rho} + \Phi + \frac{\gamma_{eff}}{\gamma_{eff} - 1} \frac{P_c}{\rho} \frac{u + v_A}{u} = \text{const} \quad (23)$$

$$\frac{du}{dz} = u \frac{c_*^2 a(z) - g(z)}{u^2 - c_*^2}, \quad (24)$$

(with reference to equation 3 we define $a(z) \equiv \frac{1}{A} \frac{dA}{dz}$ and $g(z) \equiv \frac{d\Phi}{dz}$), present many similarities with the Solar wind and the De Laval nozzle problem. In analogy with the Solar wind, we define a wind (“coronal”) base where the boundary conditions for the problem are assigned, and we look for a solution in which the flow experiences a smooth transition from subsonic to supersonic regime, as in the De Laval nozzle problem. As in the Parker model for the Solar wind (see Parker (1965)) the only possible transonic solution is the one which passes through the critical point, namely the point where both the numerator and the denominator of the wind equation are zero. The similarity with the De Laval nozzle is even more evident if we compare the formal expression for the wind equation reported above and the De Laval nozzle:

$$\frac{1}{u} \frac{du}{dz} = \frac{a(z) - \frac{g(z)}{c_*^2}}{M^2 - 1} \quad (\text{Wind equation}) \quad (25)$$

$$(26)$$

$$\frac{1}{u} \frac{du}{dz} = \frac{a(z)}{M^2 - 1} \quad (\text{De Laval nozzle equation})$$

where $M = u/c_*$ is the Mach number and $a(z) \equiv \frac{1}{A} \frac{dA}{dz}$ for the De Laval nozzle. In Fig. 5 the scheme of a De Laval nozzle is shown. Notice that the nozzle has a converging and a diverging section. If we look at equation 27, we see that this converging-diverging duct makes a smooth subsonic-supersonic transition possible: the flow starts subsonic in the converging duct ($a(z) < 0$ and $M < 1$), becomes sonic at the throat ($a(z) = 0$ and $M = 1$) and is accelerated to supersonic speed in the diverging duct ($a(z) > 0$ and $M > 1$). In such configuration it is possible to constantly accelerate the flow along the nozzle (du/dz remains positive). The sonic transition is only possible at the nozzle throat (choking), where $a = 0$.

The numerator of the CR-driven wind equation 25 presents an “effective area” term given by $a(z) - \frac{g(z)}{c_*^2}$, which plays the same role of $a(z)$ in the De Laval equation. Thus, in order to have a smooth subsonic-supersonic transition, the effective area must have the same convergent-divergent behavior of the De Laval nozzle:

- $u^2 < c_*^2$ (subsonic regime)




Figure 2. Scheme of a De Laval nozzle: the flow is smoothly accelerated ($du/dz > 0$ in all the duct) from the subsonic ($M < 1$) to the supersonic regime ($M > 1$). The flow is subsonic in the converging section ($dA/dz < 0$), becomes sonic at the throat ($dA/dz=0$, $M=1$) and continues to accelerate to supersonic speed in the diverging section ($dA/dz > 0$).

the gravitational term $g(z)$ must dominate in the numerator in order to have $du/dz > 0$. Thus, $g(z)$ is a "converging duct" term,

- $u^2 > c^{*2}$ (supersonic regime)


the area term $a(z)$ must dominate in the numerator in order to have $du/dz > 0$. Thus, $a(z)$ is a "diverging duct" term.

It is clear from this analysis that if $a(z) = 0$, i.e. if the flux-tube area is constant, a wind solution cannot be achieved. Similarly, if the $a(z) > 0$, namely the flow only expands (as in our wind model), the gravitational term $\frac{g(z)}{c^{*2}}$ allows for the presence of a stationary wind solution.


Finally, the passage through the critical point (choking condition for the De Laval nozzle), fixes the wind launching velocity u_0 . Because all other magnitudes in equation 22 and 23 are assigned at the wind base, this also fixes the mass and energy flux of the wind.

In order to carry out an analysis of the effects of the input parameters on the flow, it is helpful to define a reference model to which all other cases can be compared. Because we are considering winds launched at the Sun position, we define our reference case starting from observations and we choose: $z_0 = 100$ pc for the wind base (winds launched near the Galactic disk), $n_0 = 6 \times 10^{-3} \text{ cm}^{-3}$ for the gas density, $T_0 = 2 \times 10^6$ K for the gas temperature, $P_{c0} = 4 \times 10^{-13} \text{ erg/cm}^3$ for the CR pressure, $B_0 = 1 \mu\text{G}$ for the magnetic field, $Z_b = 15$ kpc and $\alpha = 2.0$ for the area parameters and NFW-Sofue for the DM profile. The latter choice is motivated by the fact that the NFW profile is one of the most commonly used models for dark matter halos.


In Fig. 3(a)-3(c) we reported the spatial profile of the wind related quantities for the reference case, while the various components of the Galactic gravitational acceleration have been previously shown in Fig. 1. Notice that, unlike the bulge and disk components, which die off at ~ 10 kpc, the



(a) Reference case: velocities.



(b) Reference case: gas density and temperature.



(c) Reference case: pressures.

Figure 3. Reference case: a) wind, Alfvén and compound sound speed; b) gas density and temperature; c) gas and CR pressures. The input parameters are: $z_0 = 100$ pc, $n_0 = 6 \times 10^{-3} \text{ cm}^{-3}$, $T_0 = 2 \times 10^6$ K, $P_{c0} = 4 \times 10^{-13} \text{ erg/cm}^3$, $B_0 = 1 \mu\text{G}$, $Z_b = 15$ kpc, $\alpha = 2.0$ and the NFW-Sofue dark matter profile.

Table 3. Dependence of the wind properties on the gas density. The gas temperature (2×10^6 K) and CR pressure (4×10^{-13} erg/cm³) are fixed. \dot{m} is the mass loss rate of the wind.

Gas density $n_0^R = 6 \times 10^{-3}$ (cm ⁻³)	u_0 (km/s)	z_c (kpc)	u_f (km/s)	\dot{m} 4.7×10^{-4} (M _⊙ kpc ⁻² yr ⁻¹)
0.2	39.7	18.9	595	2.5
0.5	10.7	30.9	428	1.7
1.0	3.1	40.6	403.7	1.0
3.0	0.8	72.2	296	0.73
6.0	0.3	96.5	269	0.53

DM contribution to the acceleration remains important out to the virial radius (~ 240 kpc for the NFW-Sofue).

For the reference case, at the wind base the flow is subsonic and sub-Alfvénic and most of the work on the gas is due to the thermal pressure, which is four times larger than the CR pressure. CRs contribute to push the gas, however a considerable fraction of their energy is used to generate Alfvén waves, which are quickly damped and heat the gas. This can be seen in Fig. 3(b), where it is shown that the gas temperature increases up to ~ 10 kpc and then decreases in z much slower than the gas density. Because of the CR-induced gas heating, also the gas pressure falls off in z slower than it would in the absence of wave damping. Nevertheless the gas pressure decreases with height much faster than the CR pressure, and at ~ 20 kpc the CR pressure starts to dominate over the gas pressure. This is due to the fact that CRs have a smaller adiabatic index compared to the gas ($\gamma_c \sim 4/3$; $\gamma_g = 5/3$), which has important consequences on the relative contribution of the thermal and CR pressure to the wind launching. For instance, while the thermal pressure is more efficient in accelerating the gas near the wind base, the CR pressure is able to push the gas also at large z , often making an outflow possible where the thermal pressure alone would have failed (see [Ipavich \(1975\)](#); [Breitschwerdt et al. \(1991\)](#); [Everett et al. \(2008\)](#)).

Having defined a reference set of parameters, below we study the dependence of the outflow properties on the input parameters at the wind base. In Section 5.1 we focus on the effect of changes of the gas density, gas temperature and CR pressure, while the analysis of the dependence on the area parameters is reported in Section 5.2.

5.1 Effect of the gas density and temperature and of the Cosmic Ray pressure

In order to emphasize the dependence of the results on the input parameters, here we also include in our analysis parameters' values that are not strictly compatible with observations at the position of the Sun (discussed in Section 4).

In Table 3-5 we report the launching velocity u_0 , the position of the sonic point z_c and the terminal velocity u_f , for different values of the gas density (Table 3), gas temperature

Table 4. Dependence of the wind properties on the gas temperature. The gas density (6×10^{-3} cm⁻³) and CR pressure (4×10^{-13} erg/cm³) are fixed.

Gas temperature $T_0^R = 2 \times 10^6$ (K)	u_0 (km/s)	z_c (kpc)	u_f (km/s)	\dot{m} 4.7×10^{-4} (M _⊙ kpc ⁻² yr ⁻¹)
0.5	3.09	34.3	419	0.98
1.0	3.1	40.6	403.7	1
2.0	6.9	104.9	243	2.2
3.0	17.5	367.6	152.4	5.6
4.0	182	13.32	263	58
5.0	280	7.8	441.3	89
6.0	362	4.8	573.5	116

Table 5. Dependence of the wind properties on the CR pressure. The gas density (6×10^{-3} cm⁻³) and temperature (2×10^6 K) are fixed.

CR pressure $P_{c0}^R = 4 \times 10^{-13}$ (erg/cm ³)	u_0 (km/s)	z_c (kpc)	u_f (km/s)	\dot{m} 4.7×10^{-4} (M _⊙ kpc ⁻² yr ⁻¹)
0.2	0.72	64.1	339	0.2
0.5	1.84	51.2	351	0.6
1.0	3.1	40.6	403.7	1
2.0	8.6	36.9	376.5	2.8
3.0	14.5	32.6	394	4.6
4.0	21.8	29.2	415	7.0
5.0	30.7	26.2	439.6	9.8
6.0	41.1	23.6	573.5	13

(Table 4) and CR pressure (Table 5), as compared with the reference case of Fig. 3 (indicated by the superscript "R").

Some basic considerations can be put forward based on purely energetic grounds. Consider for instance what happens at fixed gas density and CR pressure values as in Table 4: for temperatures below $\sim 0.4 T_0^R$ it is impossible to launch a wind simply because the energy available is too small. In the opposite limit, for instance for temperature $\gtrsim 7 T_0^R$, the gas is simply too hot and the outflow resembles more an evaporation process, which in reality would result in a non-stationary outflow that cannot be described using our formalism (see also [Everett et al. \(2008\)](#)).

At fixed gas temperature and CR pressure, an increase in the gas density leads to a decrease of both the launching velocity u_0 and the terminal velocity u_f (see Table 3). This

is not surprising, since a larger mass load makes more difficult for the pressure forces to launch a wind. Notice that the decrease of u_0 with increasing density is fast enough to drive down the mass loss rate of the wind (which is proportional to $n_0 u_0$) despite the increasing density (see Table 3).

As shown in Table 4 and Table 5, increasing the temperature or the CR pressure at fixed gas density leads to an increase in the launching velocity (and because n_0 is fixed, also in the mass loss rate). Notice however that multiplying by factor of 6 the temperature and the CR pressure results in an increase of factor ~ 100 and ~ 14 in u_0 respectively. This fact can be explained keeping in mind that momentum and energy depositions before or after the sonic point affect the wind in different ways (see e.g. Lamers & Cassinelli (1999); Everett et al. (2008)): momentum and energy added before the critical point is reached can increase both the mass loss and the terminal velocity, while momentum and energy input after crossing the critical point can only affect the terminal velocity, since the mass loss is determined by the passage through the critical point. The gas adiabatic index (5/3) is larger than the CR adiabatic index ($\sim 4/3$), implying a larger gas pressure gradient compared to the CR pressure gradient. Thus the thermal pressure is more efficient than the CR pressure at increasing the mass loss. In addition, notice that the z evolution of the CR pressure also depends on whether the flow is sub-Alfvénic or super-Alfvénic. In fact, with reference to equation 5, we have the following limits for the CR pressure gradient (see Everett et al. (2008)):

$$\lim_{u < v_A} \frac{dP_c}{dz} = \gamma_{eff} \frac{P_c}{2\rho} \frac{d\rho}{dz} \quad (27)$$

$$\lim_{u \gg v_A} \frac{dP_c}{dz} = \gamma_{eff} \frac{P_c}{\rho} \frac{d\rho}{dz}. \quad (28)$$

The effective CR adiabatic index is $\gamma_{eff}/2$ in the sub-Alfvénic regime and γ_{eff} in the super-Alfvénic regime. As a consequence, CRs are less efficient at driving the wind in the sub-Alfvénic regime than in the super-Alfvénic.

The behavior of the terminal velocity with the gas temperature and the CR pressure is a bit more complicated being non monotonic. In Table 4 and Table 5 we can see that, when increasing both P_{c0} and ρ_0 , the terminal velocity in general does not increase. Keeping the gas density and the CR pressure at the reference values and increasing the temperature from $0.5 T_0^R$ to $3 T_0^R$, u_f decreases by a factor ~ 3 , while from $3 T_0^R$ to $6 T_0^R$ it increases by factor of ~ 4 . Similarly, keeping the gas density and temperature at the reference values and increasing the CR pressure from $0.2 P_{c0}^R$ to P_{c0}^R , u_f increases. It decreases when the CR pressure goes from P_{c0}^R to $2 P_{c0}^R$, and finally increases for CR pressure from $2 P_{c0}^R$ to $6 P_{c0}^R$.

In order to understand this behavior, notice that the terminal velocity is computed from the energy conservation equation 23. At large z , where all other quantities drop to zero, all the energy density of the wind goes into kinetic energy of the gas, thus u_f is related to the quantities at the wind base through

$$\frac{u_f^2}{2} = \frac{u_0^2}{2} + \frac{\gamma_g}{\gamma_g - 1} \frac{P_{g0}}{\rho_0} + \Phi(z_0) + \frac{\gamma_{eff}}{\gamma_{eff} - 1} \frac{P_{c0}}{\rho_0} \frac{u_0 + v_{A0}}{u_0}. \quad (29)$$

Unlike the gas pressure term, which does not depend on u_0 , the CR term depends on the Alfvénic Mach number

$M_{A0} = u_0/v_{A0}$ through the factor $(1 + M_{A0})/M_{A0}$. This factor becomes large when M_{A0} is small ($\ll 1$) and approaches unity when M_{A0} gets large ($\gg 1$). Considering the results of Table 4, we have that, starting from $0.5 T_0^R$ and increasing the temperature, the launching velocity and the gas pressure increase, together with the kinetic and gas terms in equation 29. However, the CR term decreases, due to the factor $(1 + M_{A0})/M_{A0}$. In the range $0.5 T_0^R - 3 T_0^R$ the decrease of the CR term dominates over the increase of the kinetic and gas terms, thus leading to a decrease of the terminal velocity with increasing temperature. At larger gas temperature the gas term finally starts to dominate over the CR term and the terminal velocity increases with the gas temperature. Notice that at large temperatures the wind is launched super-Alfvénic ($v_{A0} = 28$ km/s) and the CR term becomes practically independent of M_{A0} . The results of Table 5 can be explained with similar considerations.

The location of the sonic point is closer to the Galactic disk for smaller gas density and for larger gas temperature and CR pressures. This is due to the fact that in all three cases the gas is accelerated more easily and in general reaches the sonic point at smaller z .

Is it also worth stressing the role played by wave damping. Breitschwerdt et al. (1991) did not study the effect of wave damping in detail and the paper mainly focused on a model where Alfvén waves, produced through CR streaming instability, are not damped and can grow indefinitely. In such a model, the parameter space for the wind launching was much larger than in the case where wave damping is included, resulting in the possibility to easily launch winds also at relatively low temperatures and high gas densities. In the presence of wave damping the parameter space is reduced, mainly due to two factors. First, if waves are not damped, their pressure contribution, which grows to reach more or less the same magnitude as the CR pressure, can significantly help in pushing the gas. Second when the waves are damped, the heat input from wave damping increases the temperature (and thus the gas pressure) along z , which results in a smaller gas pressure gradient, i.e a smaller force acting on to the wind. If the contribution of wave damping is too intense, the gas heating can be so important so as to make the gas pressure increase with z , thus creating a stall in the outflow. In this situation the wind formation is prevented (see Everett et al. (2008)).

We conclude this overview with a discussion of the mass loss rate of the Galaxy due to winds. In Tables 3, 4 and 5 we report the mass loss rate per unit area for different values of the launching parameters of the wind. If we assume that such mass loss rate per unit area is the same as in the whole Galactic disk and we consider input parameters compatible with the halo observations, we obtain a mass loss rate roughly in the range $\sim 0.5 - 1.5 M_\odot/\text{yr}$. Such values of the mass loss rate are of the same order of magnitude of the Galactic star formation rate ($\sim M_\odot/\text{yr}$; see e.g. Robitaille & Whitney (2010)), which means that CR-driven Galactic winds may play an important role in the evolution of the Milky Way.

5.2 Effect of the flux-tube geometry

Here we analyze the impact of the flux-tube geometry through the variation of the two parameters α and Z_b of

equation 16, keeping the gas density, temperature and CR pressure to the reference values of Fig. 3 and using the NFW-Sofue DM profile. The "mushroom" type geometry of equation 16 reflects what we would intuitively expect for outflows from disk galaxies: the flow proceeds in nearly cylindrical symmetry up to about a distance Z_b from the galactic disk and then opens up in a nearly spherical way. Of course, in a full treatment of the CR-driven wind problem, the flow geometry should be calculated self-consistently by accounting for the large-scale magnetic field. However, such study is beyond the scope of the present work, hence we retain the geometry described by equation 16 and we study how the wind properties get modified when Z_b and α are changed.

The flux-tube geometry directly affects the acceleration of the outflow, as well as the gas density (see equation 22 and 24) and consequently the pressure gradients (see equation 4 and 5). Thus, it is not surprising that the shape chosen for the function $A(z)$ has important implications for the wind properties. In what follows we first discuss the effect produced by changing the length scale, Z_b , and then the effect of changing the exponent α :

5.2.1 Effect of changing the length scale Z_b

We fix $\alpha = 2.0$ and we change Z_b in the range 5-20 kpc, stepped by 2.5 kpc. In Table 6 we show the wind launching velocity u_0 , the wind terminal velocity u_f and the location of the sonic point z_c as functions of Z_b , while in Fig. 4 we report the wind profile (relevant velocities, densities, temperature and pressures) for the two extremal cases $Z_b = 5$ kpc and $Z_b = 20$ kpc.

The plots in Fig. 4 show that for smaller values of Z_b the wind is launched with higher speed u_0 , reaching the critical point z_c farther away from the disc, but reaching a smaller final velocity u_f . These results can be explained as follows: for smaller values of Z_b , the adiabatic expansion of the gas associated to the geometric expansion of the outflow begins closer to the disc (smaller values of z), resulting in a larger density gradient, as shown in Fig. 4(b). Larger density gradients also correspond to larger pressure gradients (see equation 4 and 5) at smaller z . This results in an increase of the wind launching velocity (and, because the gas density is fixed, also in an enhanced mass loss rate).

Notice that, since $v_A = B/\sqrt{4\pi\rho}$ and $BA = \text{const}$ (see equation 2), a steeper density profile also corresponds to a faster fall-off of the Alfvén velocity with z (see Fig. 4(a)). For smaller values of Z_b this leads to a flow which is typically "more super-Alfvénic" along z , causing a more efficient driving of the CR wind (see discussion in Section 5.1), namely an extra contribution to the increase of u_0 .

Being all parameters at the wind base fixed, the increase of the wind launching velocity with decreasing Z_b results in a decrease of the CR term and in an increase in the kinetic term in the energy conservation equation 23. However, in the case considered, the first effect is dominant, and the terminal velocity decreases with decreasing Z_b .

In the subsonic regime, the numerator of the wind equation 24 is dominated by the gravitational term. A smaller value of Z_b makes the term $a(z)$ become important at smaller z (the area starts to open up at smaller z), while, as it can be seen in Fig. 4(a), c^* depends weakly on Z_b . This results in a flatter velocity profile for smaller values of Z_b . The larger

Table 6. Dependence of the wind properties on the area length scale Z_b . The gas density ($3 \times 10^{-3} \text{cm}^{-3}$), gas temperature (2×10^6 K) and CR pressure ($4 \times 10^{-13} \text{erg/cm}^3$) are fixed. $\alpha = 2.0$.

Z_b (kpc)	u_0 (km/s)	z_c (kpc)	u_f (km/s)
5.0	4.3	189.9	218.4
7.5	4.0	89.7	266.0
10.0	3.7	56.4	315.7
12.5	3.4	44.9	362.6
15.0	3.1	40.6	403.7
17.5	2.9	39.0	439.0
20.0	2.8	38.6	469.1

Table 7. Dependence of the wind properties on the expansion index α . The gas density ($3 \times 10^{-3} \text{cm}^{-3}$), gas temperature (2×10^6 K) and CR pressure ($4 \times 10^{-13} \text{erg/cm}^3$) are fixed. $Z_b = 15$ kpc.

α	u_0 (km/s)	z_c (kpc)	u_f (km/s)
1.2	2.5	73.0	513.6
1.4	2.7	61.2	482.0
1.6	2.9	52.6	453.5
1.8	3.0	45.9	427.6
2.0	3.1	40.6	403.7
2.2	3.3	36.2	382.0

launching velocity at smaller Z_b does not compensate the decrease in acceleration, leading to a sonic point located farther away from the disc.

5.2.2 Effect of changing the expansion index α

Here we fix $Z_b = 15$ kpc and we vary α in the range 1.2-2.2, stepped by 0.2. In Table 7 we show the wind launching velocity u_0 , the wind terminal velocity u_f and the location of the sonic point z_c as functions of α . In Fig. 5 we show the wind properties only for the two extreme cases $\alpha = 1.2$ and $\alpha = 2.2$.

As it can be inferred from these plots, a larger value of α corresponds to smaller z_c , larger u_0 and smaller u_f . This behavior can be explained with considerations analogous to those invoked above for the case of a changing scale height Z_b . At $z > Z_b$, an increase in α corresponds to steeper density profiles (see Fig. 5(b)), i.e. to larger pressure gradients, which act both in the subsonic and in the supersonic region (notice that in all cases $z_c > Z_b$). For this reason, when α increases, the launching velocity increases too (and consequently the mass loss rate increases), while the terminal velocity of the wind decreases.

Notice also that, as it can be seen in Fig. 5(a), the compound sound speed c^* does not change appreciably when changing α , especially in the subsonic regime. Moreover, for a given Z_b the area of the flux tube $A(z)$ does not change much for $z < Z_b$. This leads to velocity profiles which are practically parallel up to $\sim Z_b$ at all α (see equation 24 and Fig. 5(a)), and to a decrease of z_c when α increases. In fact, because the velocity profiles are nearly parallel within $\sim Z_b$ and c^* is weakly dependent on α in the subsonic regime, at larger u_0 correspond smaller z_c . Finally, notice the steeper c^* profile at $z > Z_b$ corresponding to larger values of α . This is due to a more rapid fall off in z of the density and pressures.

5.3 Radiative cooling

For gas temperature of order $\sim 10^6$ K radiative cooling may be important and affect in a potentially important manner the properties of CR driven winds or even the possibility to launch such winds.

As shown by Dalgarno & McCray (1972), in this range of temperature the ISM is cooled by the emission of forbidden lines and soft X-rays, at a rate

$$\frac{1}{T} \frac{dT}{dt} \sim \Lambda(T)n^2, \quad (30)$$

where the cooling function is $\Lambda(T \sim 10^6) \sim 2 \times 10^{-23}$ erg cm^3/s for Solar metallicity. The dependence of the cooling rate as the square of the gas density suggests that the importance of the cooling will be much more important in the cases of dense winds.

From the formal point of view, radiative cooling can be easily included in the wind hydrodynamic equations: equation 4 for the gas pressure becomes

$$\frac{dP_g}{dz} = \gamma_g \frac{P_g}{\rho} \frac{d\rho}{dz} - (\gamma_g - 1) \frac{v_A}{u} \frac{dP_c}{dz} - (\gamma_g - 1) \Lambda(T) \frac{n^2}{u}, \quad (31)$$


while the wind equation 3 gets modified as

$$\frac{1}{u} \frac{du}{dz} = \frac{c_*^2 a(z) - g(z) + (\gamma_g - 1) \Lambda(T) \frac{n^2}{u}}{u^2 - c_*^2}. \quad (32)$$


These expressions have been derived also by Dougherty et al. (1990), where it was also shown that the solution topology remains the same with the addition of radiative cooling. Notice that the cooling term appears in the numerator of the wind equation 32 with the same sign of the area term, namely it behaves as a "diverging duct" term in the De Laval nozzle picture discussed above.

Comparing the radiative cooling time and the wave heating time for the reference case of Fig. 3 ($n_0 = 6 \times 10^{-3} \text{cm}^{-3}$), we have $\tau_{cool} \sim 10^7$ yr and $\tau_{heat} \sim 10^8$ yr. This example shows that, in the absence of additional heating terms, the radiative cooling can be a quite fast process near the Galactic disk (where the gas density is larger) and can in principle prevent the wind launching. From the formal point of view, a large cooling term would result in a positive numerator at the wind base in equation 32, making a wind solution impossible.


On the other hand, the ISM is also heated by SN explosions through the injection of hot gas and magnetic turbulence that will eventually dissipate into thermal energy. Recently it was suggested that also Coulomb losses of CRs



(a) Changing Z_b : velocities.



(b) Changing Z_b : gas density and temperature.



(c) Changing Z_b : pressures.

Figure 4. Comparison of wind profiles in case of changing the flux tube geometry. Two cases are shown, $Z_b = 5$ kpc and $Z_b = 20$ kpc, while the value of α is fixed to 2. The different plots are: a) wind, Alfvén and compound sound speed; b) gas density and temperature; c) gas and CR pressure.




(a) Changing α : velocities.(b) Changing α : gas density and temperature.(c) Changing α : pressures.

Figure 5. Comparison of wind profiles in case of changing the flux tube geometry. Two cases are shown, $\alpha = 1.2$ and $\alpha = 2.2$, while the value of Z_b is fixed to 15 kpc. The different plots are: a) wind, Alfvén and compound sound speed; b) gas density and temperature; c) gas and CR pressures.

themselves might be a substantial source of heating (see Walker (2016)). The relevance of cooling was already recognized by Breitschwerdt et al. (1991) which, nevertheless, assumed that some sort of energy balance must exist between heating and cooling processes. It is currently unclear which mechanism may be responsible for sufficient heating to compensate cooling, but observations reveal that the temperature in the halo is indeed in the range $\sim 10^5 - 10^6$ K (see for instance Miller & Bregman (2015)), hence supporting the idea that either cooling is negligible or it is balanced by one or more heating processes.

It is worth remarking that the cooling function in equation 30 has been estimated assuming that the halo has the same metallicity as the ISM in the solar vicinity, which might not be strictly true. In case of smaller metallicity, as one might expect for the gas in the halo, the cooling rate is somewhat reduced.

Given the lack of knowledge of detailed physical processes that may heat the base of the wind, here we adopted the same pragmatic attitude as Breitschwerdt et al. (1991) and assume that radiative cooling is balanced by some heating process, so as to not include an explicit heating and cooling term in the wind equations (see also (Paper I)).

6 COSMIC RAY SPECTRUM

As discussed in the sections above, the possibility to launch a CR driven wind and the properties of such winds depend upon the local conditions at the base of the wind, and the gravitational potential along the trajectory of the outflow. Of course, one of the important parameters of the problem is the CR pressure at the base of the wind. If we are interested in solving the problem at the position of the Sun we should recall that not only the CR pressure but also the CR spectrum is measured, so that the problem of launching a wind is much more constrained. In a non-linear theory, such as the one we adopt in this paper, the two quantities are not independent: the number of CRs at a given momentum determines the energy density of the self-produced waves and hence the diffusion coefficient, which in turn affects the spectrum and energy density of CRs. Moreover the same quantities have a direct effect on the wind properties, which also have a feedback on the CR spectrum through advection. Notice that in our iterative approach to solving the combined set of hydrodynamical equations for the wind and kinetic equation for CRs, the CR pressure is taken from observations ($P_{CR\odot} \sim 4 \times 10^{-13}$ erg/cm³), but the spectrum is an output of the kinetic equation. Imposing that the predicted spectrum matches the observed flux at a given energy does not ensure that the spectrum also fits observations. In fact, it is in general the case that the spectrum corresponding to wind solutions is quite unlike the CR spectrum observed at the Earth (see also the discussion in Paper I).

This will be best discussed in Section 6.1, where we present some selected cases in which we solve the wind problem for input parameters suitable for the Sun's position in the Galaxy and we check the computed spectrum versus observations. We will show that typically the CR spectrum associated to wind solutions is harder than the observed one, as a result of a dominant role of advection of CRs with the wind. In some cases, the advection velocity can be so

large as to dominate the CR transport up to energies of hundreds of GeV. A smaller Alfvén speed (and possibly a smaller wind speed) is obtained by launching the wind with larger gas density (and/or at smaller magnetic field). On the other hand, it may happen that increasing the gas density at the base of the wind, so as to reduce the Alfvén speed and the wind speed, at some point the wind can no longer be launched. In some borderline cases, whether the wind is in fact launched or not may depend on the spatial distribution of dark matter, namely the contribution of dark matter to the gravitational potential.

At CR energies $E \gtrsim 200$ GeV) the opening of the flux-tube area together with the energy dependence of the self-generated diffusion coefficient (see equation 12) makes the CR spectrum typically steeper than the observed one. However, as discussed in Paper I, the high energy behaviour of the CR flux may be considerably affected by the presence of non self-generated turbulence in the near disc region (within ~ 1 kpc from the base of the wind). In fact such turbulence should be postulated in this class of models if to account for the spectral hardening measured by the PAMELA (Adriani et al. 2011) and AMS-02 (Aguilar et al. 2015) experiments.

In Section 6.2, we discuss the impact of the DM halo profile on the CR spectrum, and we show that it is possible to find a wind model which is consistent with both the recent observations illustrated by Miller & Bregman (2015) and with the observed CR spectrum.

6.1 Reference models

Here we discuss three scenarios in which a wind is launched at the position of the Sun (Galactocentric distance $R_\odot = 8.5$ kpc, $z_0 = 100$ pc) with parameters that are compatible with observations. Following the discussion reported in Section 4, we fix the gas temperature as $T_0 = 2 \times 10^6$ K and the parameters of the flux-tube geometry as $Z_b = 15$ kpc and $\alpha = 1.5$. The calculations are carried out assuming that dark matter is distributed according to a NFW-Sofue profile and we fix the slope of the injection spectrum as $\gamma = 4.3$. The three scenarios differ in the values of the gas density n_0 and magnetic field B_0 at the wind base and in the normalization of the CR injection spectrum (which is normalized either to reproduce the observed CR pressure or to match the computed and observed spectra at 50 GeV): in the first scenario (Model A) we assume $n_0 = 0.003$ cm $^{-3}$ and $B_0 = 2$ μ G at the wind launching point z_0 . In addition, we normalize the injection spectrum in order to get the observed CR pressure $P_{c0} = 4 \times 10^{-13}$ erg/cm 3 at z_0 . This implies the following condition to be fulfilled:

$$\frac{\xi_{CR}}{0.1} \frac{\mathcal{R}_{SN}}{1/30 \text{ yr}^{-1}} \approx 1.1$$

for the injection term (equation 9). The results for Model A are summarized in Fig. 6: Fig. 6(a) shows the wind velocity, the Alfvén speed and the compound sound speed, while wind density and temperature are shown in Fig. 6(b). The wind is launched sub-Alfvénic with $u_0 = 14$ km/s and becomes sonic at $z_c = 35$ kpc, while the Alfvén speed at z_0 is 79 km/s. The CR pressure is shown in Fig. 6(c) together with the gas pressure and the wave pressure. The latter is computed from the equilibrium wave spectrum, obtained by equating the growth rate of waves due to CR streaming instability and

the damping rate due to NLLD (see equation 15). Notice that the wave pressure is much smaller than the gas and CR pressure, thereby justifying a posteriori the fact that it is neglected in the hydrodynamic equations.

In the second model (Model B) we assume again a gas density $n_0 = 0.003$ cm $^{-3}$ and a magnetic field $B_0 = 2$ μ G at the base of the wind. However, we normalize the injection spectrum in a way that the computed flux of CR protons at the location of the Sun is the same as measured at an energy of 50 GeV. This constraint implies the following condition on the injection parameters:

$$\frac{\xi_{CR}}{0.1} \frac{\mathcal{R}_{SN}}{1/30 \text{ yr}^{-1}} \approx 0.75.$$

The corresponding wind velocity, the Alfvén speed and the compound sound speed are shown in Fig. 7(a) while the wind density and temperature are shown in Fig. 7(b). The wind is launched sub-Alfvénic with $u_0 = 9$ km/s and becomes sonic at $z_c = 39$ kpc, while the Alfvén speed at z_0 is 79 km/s. The CR pressure is shown in Fig. 7(c), together with the gas pressure and wave pressure. Notice that, because we are normalizing the spectrum to the observed proton flux at 50 GeV rather than on the CR pressure at z_0 , the latter is an output of the calculation, with value $P_{c0} = 2.6 \times 10^{-13}$ erg/cm 3 . This explains the difference in wind velocity between Model A and Model B.

In the third scenario (Model C), we assume a gas density $n_0 = 0.006$ cm $^{-3}$ and a magnetic field $B_0 = 1$ μ G at the base of the wind. As for Model B, we normalize the injection spectrum in order to reproduce the observed CR flux at 50 GeV, which implies:

$$\frac{\xi_{CR}}{0.1} \frac{\mathcal{R}_{SN}}{1/30 \text{ yr}^{-1}} \approx 0.32.$$

The wind velocity, the Alfvén speed and the compound sound speed for Model C are shown in Fig. 8(a), while Fig. 8(b) illustrates the evolution of the wind density and temperature. The wind is launched sub-Alfvénic with $u_0 = 2$ km/s and becomes sonic at $z_c = 60$ kpc, while the Alfvén speed at z_0 is 28 km/s. The CR pressure is shown in Fig. 8(c), together with the gas and the wave pressure. As for Model B, the CR pressure at z_0 is an output of the calculation, $P_{c0} = 3.2 \times 10^{-13}$ erg/cm 3 .

The spectrum of CR protons at the position of the Sun in the three scenarios discussed above is shown in Figs. 9(a), 9(b) and 9(c) compared with the proton flux as measured by the VOYAGER-I (Stone et al. 2013) at low energies and by AMS-02 (Aguilar et al. 2015) at higher energies. The red curve illustrates the effect of solar modulation, with the simple recipe of Bernardo et al. (2010). One should not overestimate the importance of the comparison of our predicted CR spectrum with observations at energies below ~ 1 GeV, because in such energy region energy losses become important and such effects have not been included in the calculations presented in this paper.

In all the three cases introduced above, the CR advection velocity near the wind base is dominated by the Alfvén speed. The advection velocity at the wind base is $u_0 + v_{A0} \sim 93$ km/s, ~ 88 km/s and ~ 30 km/s for model A, B and C respectively. An important feature of the spectra in the disk, shown for the three models in Fig. 9, is the presence of hard spectra which extend to higher energies for




Figure 6. Results for Model-A: at the wind base, z_0 , the gas density is $n_0 = 3 \times 10^{-3} \text{ cm}^{-3}$ and the magnetic field is $B_0 = 2 \mu\text{G}$. The CR injection spectrum has been normalized in order to get the observed CR pressure $P_{c0} = 4 \times 10^{-13} \text{ erg/cm}^3$ at z_0 .

increasing advection velocity. This is due to the fact that, the larger the CR advection velocity, the higher the CR energy at which advection dominates upon diffusion in the CR transport equation, thus producing a spectrum with spectral index close to the injection spectral index.


From Fig. 9(c), one can see that below $\sim 200 \text{ GeV}$ Model-C resemble the observed CR spectrum, while the harder spectra in Model A and B appear to be quite unlike the observed one, even qualitatively. This is a rather general conclusion: even using values for the environmental parameters that are compatible with observations (see Section 5) and that lead to wind launching, the corresponding CR spectrum at the Sun’s position is, in general, qualitatively different from the observed one, even if the total CR pressure may be close to the measured one.

Making use of the points discussed in Section 5 concerning the dependence of the wind properties on the launching parameters, we can infer what would happen by changing the gas density and temperature and the magnetic field at the wind base (always within the observational constraints). By increasing the magnetic field and/or decreasing the gas density the Alfvén speed would increase and, consequently, harder spectra would be inferred. In addition, a decrease of the gas density, as well as an increase of the gas temperature, leads to an increase of the wind launching velocity, with similar implications for the CR spectrum. On the other hand, an increase in the gas density would lead to smaller advection velocities (decreasing both the Alfvén and the wind speed), hence a generally steeper spectrum. However, when the density becomes too high it may become impossible to launch a wind. The same considerations hold when decreasing the gas temperature: the Alfvén speed is not affected but the wind velocity decreases. This situation is particularly important for winds in which the launching velocity is comparable or larger than the Alfvén speed. Also in this case, for smaller temperatures the wind launching may be inhibited.


6.2 Models versus observations: the effect of the Dark Matter halo

Here we present a wind model whose parameters are compatible with properties of the Galactic halo derived in Miller & Bregman (2015), i.e Model-C of Section 6.1 (see discussion below), and we study how the CR spectrum depends on the DM halo profile. In particular, we show that it is possible to have a good agreement with the observed CR proton spectrum, depending on the choice of the DM halo potential.




In Fig. 11(a) we show the wind density and temperature obtained with the input parameters of Model-C and for the three DM halos NFW-Sofue, BUR and EIN. The NFW-Salucci model is missing since with such halo no wind was launched. This is a typical example of how, fixed any other parameter, an increase of the gravitational pull (the NFW-Salucci has the largest force among the proposed models) could prevent the wind launching. In Fig. 10(a) and 10(b) we show the corresponding wind speed, sound speed and Alfvén velocity, while in Fig. 11(b) we show the gas and CR pressures. For all the three DM models, the wind is launched as sub-Alfvénic with: ($u_0 = 2 \text{ km/s}$, $z_c = 60 \text{ kpc}$, $u_f = 450 \text{ km/s}$) for the NFW-Sofue profile, ($u_0 = 3 \text{ km/s}$, $z_c = 285 \text{ kpc}$, $u_f = 273 \text{ km/s}$) for the BUR profile




(a) Wind velocity, Alfvén velocity and sound speed.



(a) Wind velocity, Alfvén velocity and sound speed.

(b) Density in units of 10^{-3}cm^{-3} and temperature in units of 10^6 K.(b) Density in units of 10^{-3}cm^{-3} and temperature in units of 10^6 K.


(c) Gas pressure, CR pressure obtained from the hydrodynamic and kinetic calculations, wave pressure obtained from the CR transport equation.




(c) Gas pressure, CR pressure obtained from the hydrodynamic and kinetic calculations, wave pressure obtained from the CR transport equation.

Figure 7. Results for Model-B: at the wind base, z_0 , the gas density is $n_0 = 3 \times 10^{-3}\text{cm}^{-3}$ and the magnetic field is $B_0 = 2\mu\text{G}$. The CR injection spectrum has been normalized in order to reproduce the observed CR spectrum at 50 GeV.


Figure 8. Results for Model-C: at the wind base, z_0 , the gas density is $n_0 = 6 \times 10^{-3}\text{cm}^{-3}$ and the magnetic field is $B_0 = 1\mu\text{G}$. The CR injection spectrum has been normalized in order to reproduce the observed CR spectrum at 50 GeV.



(a) Model-A: CR spectrum.




(b) Model-B: CR spectrum.




(c) Model-C: CR spectrum.

Figure 9. CR spectrum compared to the VOYAGER and AMS-02 data for Model A, B and C. Solar modulation has been applied with $\Phi = 500$ MV.



(a) Wind and compound sound speed.




(b) Wind and Alfvén speed.

Figure 10. Dependence of the CR spectrum on the DM halo: a) wind and compound sound speed; b) wind and Alfvén speed.


and ($u_0 = 2.7$ km/s, $z_c = 590$ kpc, $u_f = 253$ km/s) for the EIN profile.

The mass loss rate and the halo mass of the wind is ($\dot{M} = 0.46 M_\odot/\text{yr}$, $M(< 50\text{kpc}) \sim 1.2 \times 10^9 M_\odot$, $M(< 250\text{kpc}) \sim 1.9 \times 10^9 M_\odot$) for the NFW-Sofue profile, ($\dot{M} = 0.7 M_\odot/\text{yr}$, $M(< 50\text{kpc}) \sim 3.2 \times 10^9 M_\odot$, $M(< 250\text{kpc}) \sim 6.4 \times 10^9 M_\odot$) for the BUR profile and ($\dot{M} = 0.63 M_\odot/\text{yr}$, $M(< 50\text{kpc}) \sim 3.1 \times 10^9 M_\odot$, $M(< 250\text{kpc}) \sim 8 \times 10^9 M_\odot$) for the EIN profile. Such values of the halo mass are of the same order of magnitude of the Galactic halo mass estimated by Miller & Bregman (2015) ($M(< 50\text{kpc}) \sim 3.8 \times 10^9 M_\odot$, $M(< 250\text{kpc}) \sim 4.3 \times 10^{10} M_\odot$).

Notice that at larger gravitational acceleration, i.e for the NFW-Sofue profile, we get a smaller launching velocity u_0 and a larger terminal velocity u_f . Moreover, we get a smaller z_c . This fact can be explained as follows: in the subsonic regime the gravitational term dominates the numerator of the wind equation 3, while c^* is weakly dependent on the DM profile (see Fig. 10). Thus a larger gravitational force leads to a larger gas acceleration in the subsonic region, large enough to provide a smaller z_c despite the smaller launching velocity, as it can be seen in Fig. 10. The rapid increase of the wind velocity, which starts at smaller z and with a



(a) Gas density and temperature.




(b) Gas and CR pressure.


Figure 11. Dependence of the CR spectrum on the DM halo: a) gas density and temperature; b) gas and CR pressure.

steeper profile for the NFW-Sofue, corresponds to the rapid decrease of the gas density and of the CR and gas pressure (see Fig. 11(a) and Fig. 11(b)). In the range $z \sim 1 - 100$ kpc, the NFW-Sofue profile shows the largest Alfvén speed (due to the smaller gas density), and the smallest CR pressure. However the first effect is still dominant, leading to a larger heating due to wave damping (recall that the heating term is $\sim v_A \nabla P_c$), and hence to a slightly larger temperature (see Fig. 11(a)) compared to the BUR and EIN halos. Notice that the effect of wave damping is in fact to keep the gas temperature around 2×10^6 K up to ~ 100 kpc (this is especially true for the NFW-Sofue profile), in line with the results of [Miller & Bregman \(2015\)](#) on the Galactic halo temperature.


In Fig. 12, we show the CR spectrum for the NFW-Sofue, BUR and EIN profiles. Below ~ 200 GeV, the best (qualitative) match to the observed proton spectrum is achieved with the NFW-Sofue model. The larger wind launching velocity of the BUR and EIN models leads to a low energy spectrum harder than the observed one.



(a) NFW-Sofue halo.



(b) Burkert halo.



(c) Einasto halo.

Figure 12. Dependence of the CR spectrum on the DM halo: a) CR spectrum with the NFW halo; b) CR spectrum with the BUR halo; c) CR spectrum with the EIN halo.

7 CONCLUSIONS

In this paper we studied the possible presence of CR-driven Galactic winds launched at the Sun location and the dependence of their properties and of the related CR spectrum on the conditions of the Galactic environment in the vicinity of the Sun. We used the semi-analytical calculation developed in [Paper I](#), which allows to compute at the same time both the hydrodynamical structure of the wind, the CR distribution function and CR self-generated diffusion coefficient. The CR transport is considered to be due to the diffusion off Alfvén waves generated by CRs through streaming instability, saturated through NLLD, and to the advection with such waves and with the wind. The wind launching and the CR spectrum depend on the properties of the ISM (gas density and temperature, Galactic magnetic field), on the CR injection, on the flow geometry and on the Galactic gravitational potential. All these quantities depend on the position in the Galaxy and are constrained by observations.

We found that, in agreement with previous hydrodynamic calculations of CR-driven winds, the combined action of the thermal and CR pressures can drive winds for a variety of environmental conditions. In particular, it is possible to launch winds with input parameters compatible with observations of the ISM in the vicinity of the Sun. We also found that in many cases, winds launched with environmental parameters compatible with observations lead to a CR spectrum which is not in agreement with the observed one. As pointed out in [Paper I](#), this is mainly due to the fact that CR advection in such winds is strong, leading to spectra at low energies (below ~ 200 GeV) which are harder than the observed spectrum. At high energy (above ~ 200 GeV), instead, the wind expansion, together with the steep energy dependence of the self-generated diffusion coefficient, leads to spectra which are steeper than the observed spectrum. However, the high energy spectrum may be affected by other factors, such as the presence of turbulence non generated by CRs ([Aloisio et al. \(2015\)](#); [Paper I](#)).

As for the low energy part of the spectrum, we found that it is possible to find specific cases in which the wind launching parameters and the resulting wind properties are compatible with observations, in particular with the Galactic halo properties deduced by [Miller & Bregman \(2015\)](#), and the resulting CR spectrum is also in agreement with observations. In this analysis we also found that an important role is played by the choice of the dark matter halo model. In fact, we showed that, for given values of all parameter, the DM gravitational potential can determine whether the wind is launched and the quality of the agreement with the observed CR spectrum.

This analysis highlights that the low energy part of the CR spectrum provides a strong constraint for wind models and that an accurate modeling of the Galactic environment and of the Galactic gravitational potential (in particular of the DM halo) are of crucial importance in the understanding of Galactic winds.

The study presented here can be extended to any other locations in the Galaxy different from the Sun's position and

can be used to analyze the possible presence of winds and the related CR spectrum in the whole Galaxy. Such investigation would allow us to calculate the rate of mass loss and the mass of the baryonic halo induced by the wind (both are crucial ingredients in models of galaxy formation and evolution), as well as the expected intensity in emission and absorption lines from the halo (for instance in the X-ray band) and the CR density gradient in the Galaxy (which can be tracked by the γ -ray emission resulting from CR interactions with the background plasma).

ACKNOWLEDGEMENTS

The authors are glad to acknowledge a continuous ongoing collaboration with the rest of the Arcetri high energy Group. G.M. want also to acknowledge the hospitality of ISSI for Teamwork 351 on "The origin and composition of Galactic cosmic rays".

REFERENCES

- Adriani O., et al., 2011, [Science](#), **332**, 69
Aguilar M., et al., 2015, [Physical Review Letters](#), **114**, 171103
Aloisio R., Blasi P., Serpico P. D., 2015, [Astronomy & Astrophysics](#), **583**, A95
Anderson M. E., Bregman J. N., 2010, [The Astrophysical Journal](#), **714**, 320
Bernal N., Palomares-Ruiz S., 2012, [J. Cosmology Astropart. Phys.](#), **1**, 006
Bernardo G. D., Evoli C., Gaggero D., Grasso D., Maccione L., 2010, [Astroparticle Physics](#), **34**, 274
Bertone G., Hooper D., Silk J., 2005, [Phys. Rep.](#), **405**, 279
Booth C. M., Agertz O., Kravtsov A. V., Gnedin N. Y., 2013, [ApJ](#), **777**, L16
Breitschwerdt D., Schmutzler T., 1994, [Nature](#), **371**, 774
Breitschwerdt D., Schmutzler T., 1999, [A&A](#), **347**, 650
Breitschwerdt D., McKenzie J. F., Voelk H. J., 1991, [A&A](#), **245**, 79
Burkert A., 1995, [ApJ](#), **447**, L25
Cheng K.-S., Chernyshov D. O., Dogiel V. A., Ko C.-M., Ip W.-H., 2011, [ApJ](#), **731**, L17
Chevalier R. A., Clegg A. W., 1985, [Nature](#), **317**, 44
Cox D. P., 2005, [Annual Review of Astronomy and Astrophysics](#), **43**, 337
Crain R. A., Eke V. R., Frenk C. S., Jenkins A., McCarthy I. G., Navarro J. F., Pearce F. R., 2007, [MNRAS](#), **377**, 41
Dalgarno A., McCray R. A., 1972, [ARA&A](#), **10**, 375
Dorfi E. A., Breitschwerdt D., 2012, [A&A](#), **540**, A77
Dougherty K. M., Breitschwerdt D., Völk J. H., 1990, [International Cosmic Ray Conference](#), **3**, 319
Everett J. E., Zweibel E. G., Benjamin R. A., McCammon D., Rocks L., Gallagher J. S., 2008, [The Astrophysical Journal](#), **674**, 258
Ferrière K. M., 2001, [Reviews of Modern Physics](#), **73**, 1031
Girichidis P., et al., 2016, [ApJ](#), **816**, L19
Ipavich F. M., 1975, [Astrophysical Journal](#), **196**, 107
Irrgang A., Wilcox B., Tucker E., Schiefelbein L., 2013, [A&A](#), **549**, A137
Jansson R., Farrar G. R., 2012, [ApJ](#), **757**, L4
Kalberla P. M. W., Dedes L., 2008, [A&A](#), **487**, 951
King A., Pounds K., 2015, [ARA&A](#), **53**, 115
Kulsrud R. M., Cesarsky C. J., 1971, [Astrophys. Lett.](#), **8**, 189
Lacki B. C., 2014, [MNRAS](#), **444**, L39

- Lamers H. J. G. L. M., Cassinelli J. P., 1999, Introduction to Stellar Winds: Cambridge University Press, Cambridge, doi:10.1017/CBO9781139175012, <https://www.cambridge.org/core/books/introduction-to-stellar-winds/6E24600EDAD99F2B7839BBA190C115C0>
- Martin C. L., Shapley A. E., Coil A. L., Kornei K. A., Bundy K., Weiner B. J., Noeske K. G., Schiminovich D., 2012, *The Astrophysical Journal*, 760, 127
- Miller M. J., Bregman J. N., 2015, *The Astrophysical Journal*, 800, 14
- Miyamoto M., Nagai R., 1975, *PASJ*, 27, 533
- Murray N., Quataert E., Thompson T. A., 2005, *ApJ*, 618, 569
- Navarro J. F., Frenk C. S., White S. D. M., 1996, *ApJ*, 462, 563
- Nesti F., Salucci P., 2013, *J. Cosmology Astropart. Phys.*, 7, 016
- Parker E. N., 1965, *Space Sci. Rev.*, 4, 666
- Ptuskin V. S., Zirakashvili V. N., 2003, *A&A*, 403, 1
- Recchia S., Blasi P., Morlino G., 2016, *Monthly Notices of the Royal Astronomical Society*
- Retana-Montenegro E., van Hese E., Gentile G., Baes M., Frutos-Alfaro F., 2012, *A&A*, 540, A70
- Robitaille T. P., Whitney B. A., 2010, *ApJ*, 710, L11
- Ruszkowski M., Yang H.-Y. K., Zweibel E., 2016, preprint, (arXiv:1602.04856)
- Salem M., Bryan G. L., 2014, *MNRAS*, 437, 3312
- Salucci P., Nesti F., Gentile G., Frigerio Martins C., 2010, *A&A*, 523, A83
- Scoville N., 2003, *Journal of Korean Astronomical Society*, 36, 167
- Skilling J., 1971, *Astrophysical Journal*, 170, 265
- Skilling J., 1975, *MNRAS*, 173, 255
- Sofue Y., 2012, *PASJ*, 64
- Stone E. C., Cummings A. C., McDonald F. B., Heikkila B. C., Lal N., Webber W. R., 2013, *Science*, 341, 150
- Uhlig M., Pfrommer C., Sharma M., Nath B. B., Enflin T. A., Springel V., 2012, *MNRAS*, 423, 2374
- Veilleux S., Cecil G., Bland-Hawthorn J., 2005, *ARA&A*, 43, 769
- Walker M. A., 2016, *ApJ*, 818, 23
- Wolfire M. G., McKee C. F., Hollenbach D., Tielens A. G. G. M., 2003, *ApJ*, 587, 278
- Zhou Y., Matthes W. H., 1990, *J. Geophys. Res.*, 95, 10291
- Zirakashvili V. N., Breitschwerdt D., Ptuskin V. S., Voelk H. J., 1996, *A&A*, 311, 113
- Zubovas K., King A. R., Nayakshin S., 2011, *MNRAS*, 415, L21

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.