




Publication Year	2017
Acceptance in OA @INAF	2020-09-09T11:02:41Z
Title	Massive ionized outflows in quasars
Authors	MARZIANI, Paola; Negrete, Alenka; Dultzin, Deborah; Martinez-Aldama, Mary Loli; Del Olmo, Ascension; et al.
DOI	10.5281/zenodo.571299
Handle	http://hdl.handle.net/20.500.12386/27241

Massive ionized outflows in quasars


P. Marziani¹, C. A. Negrete², D. Dultzin², M. L. Martinez-Aldama³, A. Del Olmo³,
M. D'Onofrio⁴, G. M. Stirpe⁵

¹INAF, Osservatorio Astronomico di Padova, Italy; ²Instituto de Astronomía, UNAM, Mexico; ³Instituto de Astrofísica de Andalucía (IAA-CSIC), Granada, Spain; ⁴Università di Padova, Padova, Italia; ⁵INAF, Osservatorio Astronomico di Bologna, Italia; ⁶Belgrade Observatory, Serbia

The most luminous quasars in the Hamburg-ESO (HE) survey show, at a high prevalence, CIV $\lambda 1549$ and [OIII] $\lambda \lambda 4959, 5007$ emission line profiles with high-amplitude blueshifts which indicate outflows occurring over a wide range of spatial scales. We found evidence in favor of the nuclear origin of the outflows diagnosed by [OIII] $\lambda \lambda 4959, 5007$. The derived ionized gas mass, kinetic power, and radiation thrust are extremely high, and suggest widespread feedback on the host galaxies of very luminous quasars, at cosmic epochs between 2 and 6 Gyr from the Big Bang.




The left panel shows the almost fully blueshifted (with respect to rest frame) CIV $\lambda 1549$ emission to which the H β profile is superimposed (from Sulentic et al. 2017). The right panel shows the H β spectral range with an enlargement on [OIII] $\lambda \lambda 4959, 5007$, semi-broad, boxy and blueshifted (not a unique case at high L: e.g., Marziani et al. 2016, Bischetti et al. 2017, and references therein).




Demographics of ionized outflows

“Large” scale (< few kpc) outflows are traced by the **blue outliers** ([OIII] $\lambda 5007$ peak shift < -250 km s⁻¹). BOs are rare in low z samples (they are real statistical outliers) but much more frequent in the high z and L HE sample. Their [OIII] shift and FWHM distributions are remarkably different from those of low- z , low L samples (Marziani et al. 2016; histograms aside).




Our samples confirm the [OIII] Baldwin effect for the general quasar population: $W \propto L_{5100}^{-0.26 \pm 0.03}$, steeper than the “classical” CIV Baldwin effect (which may be entirely due to selection effects). **The absence of a Baldwin effect for the BOs** is consistent with [OIII] emission from gas photoionized by the nucleus. The very high luminosity implies that the [OIII] outflowing nuclear component may “overswamp” narrow-line emission whose dynamics is dominated by the inner bulge of the host galaxy.

The green panel aside explains the high prevalence of blue outliers as a luminosity effect.




The relation between the CIV and [OIII] blueshifts is such that **all blue outliers show large CIV blueshifts** in our sample. The converse is not true, as expected if [OIII] emission on larger spatial scales is affecting the unresolved [OIII] profile. This is apparently valid also at low- z (the grey spots are for two low- z NLSy1s).



The samples

52 high luminosity HE with H β and [OIII] ($47 \leq \log L \leq 48$ [erg s⁻¹] and intermediate redshift ($1 \leq z \leq 2.6$) coverage from VLT/ISAAC

28 sources at $z > 1.4$ with CIV 1549, H β and [OIII] coverage from VLT/FORS





High luminosity Hamburg ESO (HE) quasar sample + lower z comparison sample



(Sulentic et al. 2017)

They almost uniformly cover 4 dex in quasar luminosity, and include 80 radio-quiet quasars with both H β , CIV $\lambda 1549$, and [OIII] $\lambda \lambda 4959, 5007$



A significant feedback effect

The absence of a Baldwin effect and the relation to the CIV blueshifts suggest that the [OIII] $\lambda \lambda 4959, 5007$ “blue outliers” trace an outflows of nuclear origin, also revealed by CIV $\lambda 1549$ blueshifted profiles (Marziani et al. 2016; 2016a).

Computing the kinetic power and the thrust is possible under several caveats and assumptions. For [OIII], since there is no spatial resolution, derived values lower limits (blue spots). For CIV, parameters are model dependent. We estimated the black hole mass and Eddington ratio of each quasar, and assumed an emitting region radius $r_{\text{CIV}} \propto L^b$ (Kaspi et al. 2007) from which we derived the local escape velocity. We then computed the ionized gas terminal velocity above escape velocity considering that the gas at r_{CIV} is still being accelerated by radiation forces (following Netzer & Marziani 2010).

The ionized gas mass, kinetic power, and thrust derived from the [OIII] $\lambda 5007$ and CIV $\lambda 1549$ shift and luminosity are extremely high because they ultimately depend on line luminosity. The total energy provided by the outflow over $\sim 10^8$ yr is comparable to the internal gravitational energy of a massive spheroid.

References

Bischetti M., et al. 2017, arXiv 161203728B
Carniani et al., 2016, AAp, 591, 28
Harrison, C. M., et al. 2014, MNRAS, 441, 3306
Kaspi, S. Et al. 2007, ApJ, 659, 997
Marziani P., et al. 2016, ApSS, 361, 3
Marziani P., et al. 2016a, ApSS, 361, 29
Netzer, H., et al. 2004, ApJ, 614, 558
Netzer, H., Marziani, P. 2010, ApJ, 724, 318
Sulentic J. W., et al. 2017, submitted