



<b>Publication Year</b>	2017
<b>Acceptance in OA @INAF</b>	2020-08-20T10:46:59Z
<b>Title</b>	The evolution of S0s with UV bright rings A SWIFT-UVOT study
<b>Authors</b>	RAMPAZZO, Roberto; MAZZEI, Paola; Marino, A.; TRINCHIERI, Ginevra; USLENGHI, MICHELA; et al.
<b>DOI</b>	10.1017/S1743921316009315
<b>Handle</b>	<a href="http://hdl.handle.net/20.500.12386/26745">http://hdl.handle.net/20.500.12386/26745</a>
<b>Series</b>	PROCEEDINGS OF THE INTERNATIONAL ASTRONOMICAL UNION
<b>Number</b>	vol. 11, S321

# The evolution of S0s with UV bright rings A SWIFT-UVOT study

R. Rampazzo<sup>1</sup>, P. Mazzei<sup>1</sup>, A. Marino<sup>1</sup>, G. Trinchieri<sup>2</sup>, M. Uslenghi<sup>3</sup>  
and A. Wolter<sup>2</sup>

<sup>1</sup>INAF Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy

<sup>2</sup>INAF Osservatorio Astronomico di Brera, Via Brera 28, 20121 Milano, Italy

<sup>3</sup>INAF-IASF, Via Bassini 15, 20133 Milano, Italy

**Abstract.** We report about an ongoing study of the evolution of 24 S0s with UV-bright ring/arm-like structures observed with SWIFT-UVOT and interpreted via SPH simulations with chemo-photometric implementation.

**Keywords.** galaxies: elliptical and lenticular, cD, galaxies: photometry, galaxies: evolution

---

## 1. Introduction

A significant fraction of S0s is far from being passively evolving. Signatures of activity (e.g. star formation/AGN) maximize their presence in their centre (e.g. Rampazzo *et al.* 2013). However, spectacular signatures of activity have been extensively mapped by GALEX. FUV bright rings, pseudo-rings and arm-like structures in the disk, shells and tails in the galaxy outskirts are revealed (Thilker *et al.* 2007; Salim & Rich 2010; Marino *et al.* 2011). Sometimes UV-structures are barely visible in the optical bands.

## 2. The project and few preliminary results

We select from the ARRAKIS compilation of galaxies with rings (Comerón *et al.* 2014) 24 S0s observed with SWIFT-UVOT. We obtained their luminosity profiles and structure and their SED. These information single out, among a grid of SPH simulations with chemo-photometric implementation, the evolution of these S0s providing the history of the development of their ring/arm-like structures (see Mazzei *et al.* 2014a; Mazzei *et al.* 2014b). These are found in UV images of S0s with different family classification (see e.g. Laurikainen *et al.* 2010). UV bright ring/arm-like structures have different shapes, from very diffuse (IC 2006) to filamentary (NGC 1543), to irregular (NGC 1533). SPH simulations suggest that mergers and interactions drive the evolution of ring/arm-like structures which mark transient phases during the galaxy assembly history.

## References

- Comerón, S., Salo, H., Laurikainen, E., *et al.* 2014, *A&A*, 562, A121  
Laurikainen, E., Salo, H., Buta, R., *et al.* 2010, *MNRAS*, 405, 1089  
Marino, A., Bianchi, L., Rampazzo, R., *et al.* 2011, *ApJ*, 736, 154  
Mazzei, P., Marino, A., & Rampazzo, R. 2014a, *ApJ*, 782, 53  
Mazzei, P., Marino, A., Rampazzo, R., *et al.* 2014b, *Adv. Sp. Res.*, 53, 950  
Salim, S. & Rich, R. M. 2010, *ApJ*, 714, L290  
Rampazzo, R., Panuzzo, P., Vega, O., *et al.* 2013, *MNRAS*, 432, 374  
Thilker, D. A., Bianchi, L., Meurer, G., *et al.* 2007, *ApJS*, 173, 538