



Publication Year	2016
Acceptance in OA @INAF	2020-09-16T11:20:22Z
Title	Baryon Content of Massive Galaxy Clusters ($0.57 < z < 1.33$)
Authors	I. Chiu; J. Mohr; M. Mcdonald; S. Bocquet; M. L. Ashby; et al.
DOI	10.1093/mnras/stv2303
Handle	http://hdl.handle.net/20.500.12386/27414
Journal	MONTHLY NOTICES OF THE ROYAL ASTRONOMICAL SOCIETY
Number	455

Baryon content of massive galaxy clusters at $0.57 < z < 1.33$

I. Chiu,^{1,2★} J. Mohr,^{1,2,3} M. McDonald,⁴ S. Bocquet,^{1,2} M. L. N. Ashby,⁵
M. Bayliss,^{5,6} B. A. Benson,^{7,8,9} L. E. Bleem,^{8,10,11} M. Brodwin,¹² S. Desai,^{1,2}
J. P. Dietrich,^{1,2} W. R. Forman,⁵ C. Gangkofner,^{1,2} A. H. Gonzalez,¹³ C. Hennig,^{1,2}
J. Liu,^{1,2} C. L. Reichardt,¹⁴ A. Saro,^{1,2} B. Stalder,^{5,15} S. A. Stanford,^{16,17} J. Song,¹⁸
T. Schrabback,¹⁹ R. Šuhada,¹ V. Strazzullo¹ and A. Zenteno^{1,20}

Affiliations are listed at the end of the paper

Accepted 2015 October 2. Received 2015 October 1; in original form 2014 December 24

ABSTRACT

We study the stellar, brightest cluster galaxy (BCG) and intracluster medium (ICM) masses of 14 South Pole Telescope (SPT) selected galaxy clusters with median redshift $z = 0.9$ and mass $M_{500} = 6 \times 10^{14} M_{\odot}$. We estimate stellar masses for each cluster and BCG using six photometric bands, the ICM mass using X-ray observations and the virial masses using the SPT Sunyaev–Zel’dovich effect signature. At $z = 0.9$, the BCG mass M_{\star}^{BCG} constitutes 0.12 ± 0.01 per cent of the halo mass for a $6 \times 10^{14} M_{\odot}$ cluster, and this fraction falls as $M_{500}^{-0.58 \pm 0.07}$. The cluster stellar mass function has a characteristic mass $M_0 = 10^{11.0 \pm 0.1} M_{\odot}$, and the number of galaxies per unit mass in clusters is larger than in the field by a factor of 1.65 ± 0.20 . We combine our SPT sample with previously published samples at low redshift and correct to a common initial mass function and for systematic virial mass differences. We then explore mass and redshift trends in the stellar fraction f_{\star} , the ICM fraction f_{ICM} , the collapsed baryon fraction f_c and the baryon fraction f_b . At a pivot mass of $6 \times 10^{14} M_{\odot}$ and redshift $z = 0.9$, the characteristic values are $f_{\star} = 1.1 \pm 0.1$ per cent, $f_{\text{ICM}} = 9.6 \pm 0.5$ per cent, $f_c = 10.7 \pm 1.1$ per cent and $f_b = 10.7 \pm 0.6$ per cent. These fractions all vary with cluster mass at high significance, with higher mass clusters having lower f_{\star} and f_c and higher f_{ICM} and f_b . When accounting for a 15 per cent systematic virial mass uncertainty, there is no statistically significant redshift trend at fixed mass. Our results support the scenario where clusters grow through accretion from subclusters (higher f_{\star} , lower f_{ICM}) and the field (lower f_{\star} , higher f_{ICM}), balancing to keep f_{\star} and f_{ICM} approximately constant since $z \sim 0.9$.

Key words: Galaxy: evolution – galaxies: clusters: general – large-scale structure of Universe – X-rays: galaxies: clusters.

1 INTRODUCTION

The utility of galaxy clusters for cosmological parameter studies was recognized quite early (Frenk et al. 1990; Henry & Arnaud 1991; Lilje 1992; White, Efstathiou & Frenk 1993a; White et al. 1993b), but the overwhelming evidence of widespread merging in the cluster population (Forman & Jones 1982; Geller & Beers 1982; Dressler & Shectman 1988; Mohr et al. 1995) together with the high scatter in the X-ray luminosity–temperature relation (e.g. Fabian 1994) left many with the impression that clusters were too complex and varied to ever be useful for cosmological studies. It was some time later that the first evidence that clusters exhibit significant regularity in their intracluster medium (ICM) properties

appeared (Mohr & Evrard 1997; Arnaud & Evrard 1999; Cavaliere, Menci & Tozzi 1999; Mohr, Mathiesen & Evrard 1999); X-ray observations showed that clusters as a population exhibit a size–temperature scaling relation with ≈ 10 per cent scatter, a level of regularity comparable to that known in elliptical galaxies (i.e. Djorgovski & Davis 1987). This regularity together with the emergence of evidence for cosmic acceleration (Riess et al. 1998; Perlmutter et al. 1999) focused renewed interest in the use of galaxy clusters for precise cosmological studies (e.g. Haiman, Mohr & Holder 2001). Moreover, the existence of low-scatter, power-law relations among cluster observables provided a useful tool to study the variation in cluster structure with mass and redshift.

Soon thereafter, the regularity seen in the X-ray properties of clusters was shown to exist also in the optical properties of clusters (Lin, Mohr & Stanford 2003, hereafter L03). L03 carried out an X-ray and near-infrared (NIR) 2MASS *K*-band study of an ensemble

* E-mail: inonchiu@usm.lmu.de

of 27 nearby clusters, measuring the mass fraction of the stellar component inside the galaxies (f_*), the ICM mass fraction (f_{ICM}), the total baryon fraction (f_b), the cold baryon fraction (f_c , hereafter we refer to this as the collapsed baryon fraction) and the metal enrichment of the ICM. This study showed an increasing f_b and decreasing f_* and f_c in the more massive haloes, suggesting that the star formation efficiency is higher in the low-mass haloes as well as that feedback associated with this enhanced star formation was having a larger structural impact in low-mass than in high-mass haloes. Over the last decade, additional studies using larger samples and better data have largely confirmed this result (e.g. Gonzalez, Zaritsky & Zabludoff 2007; Giodini et al. 2009; Andreon 2010; Zhang et al. 2011, hereafter Zha11; Gonzalez et al. 2013, hereafter GZ13).

Understanding how the cluster and group baryon components evolve with redshift is a key question today. While there have been many detailed studies of intermediate- and high-redshift galaxy clusters, most previous observational studies of large cluster samples have focused on nearby systems due to the difficulty of defining high-redshift samples and of following them up in the X-ray and with adequately deep optical or NIR imaging. That is changing now with the recent analyses of Sunyaev–Zel’dovich effect (hereafter SZE; Sunyaev & Zel’dovich 1970, 1972) selected clusters and groups at intermediate and high redshift. The SZE results from inverse Compton interactions of the hot ionized ICM with cosmic microwave background (CMB) photons; because it is a CMB spectral distortion rather than a source of emission, it does not suffer from cosmological dimming. Since the first SZE selected clusters were discovered in the SPT-SZ survey (Staniszewski et al. 2009), this method has been demonstrated to be a useful tool for discovering and studying galaxy cluster populations out to high redshift (Zenteno et al. 2011; Hilton et al. 2013, hereafter H13; Bayliss et al. 2014). In addition, NIR selected clusters and groups at high redshift are now also being used to study the evolution of galaxy populations (e.g. van der Burg et al. 2014, hereafter vdB14). In the present work, we focus on an SZE selected cluster sample at redshift higher than 0.6 that originates from the first 720 deg² of the South Pole Telescope (Carlstrom et al. 2011) SZE (SPT-SZ) survey (Song et al. 2012b; Reichardt et al. 2013).

To study the evolution of f_* , one needs robust stellar and virial mass estimates. Stellar masses are typically estimated by converting the observed galaxy luminosity into the stellar mass using the mean mass-to-light ratio constructed from theoretical models. This approach is sensitive to the galaxy spectral templates and needs to be modelled carefully to reduce possible biases (vdB14). For accurate stellar mass measurements with less model dependence, one requires deep multiwavelength observations that allow the spectral energy distribution (SED) to be measured on a galaxy-by-galaxy basis. For clusters at $z \approx 1$, this typically requires photometry using 8 m telescopes like the Very Large Telescope (VLT) together with space-based NIR data from the *Spitzer Space Telescope*.

The cluster virial mass measurements typically have come from X-ray mass proxies such as the emission weighted mean temperature or from galaxy velocity dispersions. The calibration of the X-ray mass proxies has often been based on the assumption of hydrostatic equilibrium, which in some circumstances can underestimate the mass by 20–40 per cent due to the non-thermal pressure components in these young structures (see Molnar et al. 2010; Chiu & Molnar 2012, and references therein). Velocity dispersion mass estimates, although likely less biased than hydrostatic mass estimates, have been shown to have quite high scatter on a single cluster basis (e.g. White, Cohn & Smit 2010; Saro et al. 2013; Ruel et al.

2014). Therefore, a study of the redshift variation of f_* would benefit from a low-scatter mass proxy from the X-ray or SZE that has been calibrated to mass using low-bias measurements such as weak lensing or velocity dispersions together with a method that accounts for selection effects and cosmological sensitivity. The masses we use in this analysis are based on the SZE signal-to-noise for each cluster as observed in the SPT-SZ survey and are calibrated in just such a manner (Bocquet et al. 2015).

In addition to robust, low-scatter mass estimates, one should use a uniformly selected cluster sample whose selection is not directly affected by variations in f_* . ICM-based observables such as the X-ray luminosity or the SZE signature enable this, although connections between the physics of star formation and the structure of the ICM remain a concern. Also, if one wishes to probe the regime beyond the group scale at high redshift, one must survey enough volume to find significant numbers of the rare, massive clusters. Large solid angle SZE surveys like those from SPT, the Atacama Cosmology Telescope (ACT; Fowler et al. 2007) and *Planck* (Tauber 2000) provide a clean way to discover clusters. Indeed, because the SZE signature for a cluster of a given mass evolves only weakly with redshift in an arcminute resolution SZE survey, the SPT-SZ survey provides a cluster sample that is well approximated as a mass-limited sample above redshift $z \approx 0.3$ (e.g. Vanderlinde et al. 2010).

In this paper, we seek to study the baryon content, including the ICM and the stellar mass components, of massive high-redshift clusters discovered within the SPT-SZ survey. We attempt also to constrain the evolution of the baryon content of these clusters by combining our high-redshift, massive clusters with other samples, primarily studied at low redshift. The paper is organized as follows. We describe the cluster sample and the data in Section 2. In Section 3, we provide detailed descriptions of the ICM, the stellar mass and the total mass measurements for the clusters. We present the stellar mass function (SMF) in Section 4 and present results on the mass and redshift trends of the baryon composition in Section 5. We discuss these results in Section 6 and summarize our conclusions in Section 7.

We adopt the concordance cold dark matter cosmological model with the cosmological parameters measured in Bocquet et al. (2015) throughout this paper: $\Omega_M = 0.299$, $\Omega_\Lambda = 0.701$ and $H_0 = 68.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$. These constraints are derived from a combination of the SPT-SZ cluster sample, the *Planck* temperature anisotropy, *Wilkinson Microwave Anisotropy Probe* (WMAP) polarization anisotropy and baryon acoustic oscillation (BAO) and SN Ia distances. Unless otherwise stated, all uncertainties are indicated as 1σ , the quantities are estimated at the overdensity of 500 with respect to the critical density (ρ_{crit}) at the cluster’s redshift, all celestial coordinates are quoted in the epoch J2000 and all photometry is in the AB magnitude system.

2 CLUSTER SAMPLES AND DATA

In this section, we briefly summarize the SPT cluster sample and the follow-up data acquisition, reduction, calibration as well as the literature cluster sample we compare to. The deep optical observations from the VLT and the *Hubble Space Telescope* (HST), together with the NIR observations from the *Spitzer*, enable us to measure the integrated stellar masses of our clusters accurately. The ICM masses are extracted from *Chandra* and *XMM-Newton* X-ray observations. Cluster total masses are derived from the SPT SZE observable ξ as calibrated using the external data sets (see Section 3.1). The literature sample we compare with in this study is described in Section 2.2.




Figure 1. VLT pseudo-colour images of SPT-CL J2331–5051 constructed from b_H , I_B and z_G . The left- and right-hand panels, respectively, show cluster R_{500} and $R_{500}/3$ regions centred on the X-ray peak. The SZE signal-to-noise contours from 0 to 10 with steps of 2 are white, the R_{500} region is the green circle and the BCG is marked by the yellow circle. The VLT pseudo-colour images for the other 13 clusters are available online.

2.1 SPT cluster sample

The 14 clusters we analyse are drawn from early SPT-SZ cluster catalogues, which covered the full 2500 deg^2 with shallower data (Williamson et al. 2011) or included the first 720 deg^2 of the full depth SPT-SZ survey (Reichardt et al. 2013). The full 2500 deg^2 catalogue has meanwhile been released (Bleem et al. 2015). These 14 systems have high detection significance ($\xi > 4.8$) and were selected for further study using *HST* and the VLT. All 14 have measured spectroscopic redshifts (Song et al. 2012b).

We study the virial region defined by R_{500} in each cluster, where R_{500} is extracted from a virial mass estimate (M_{500}) that is derived from the SPT SZE observable (see Section 3.1). We adopt the X-ray centroid as the cluster centre, because the SZE cluster centre measurement uncertainties are larger. A previous analysis of the offset between the SPT measured cluster centre and the brightest cluster galaxy (BCG) positions in a large ensemble of the SPT clusters (Song et al. 2012b) indicated that once the SPT positional measurement uncertainties are accounted for, this offset distribution is consistent with that seen in local samples where the X-ray centre is used (e.g. Lin & Mohr 2004). In our sample, the BCG positions, X-ray centres and SZE centres are all in reasonably good agreement (see Fig. 1). Importantly, these offsets have a negligible impact on our analysis, because we are comparing average properties determined within the radius R_{500} .

We present the names, redshifts and the sky positions in J2000 (α , δ) of the X-ray centre and BCG of our SPT sample in Table 1. The virial mass M_{500} and the virial radius R_{500} for each cluster are listed in Table B1.

2.1.1 Optical and infrared photometry

VLT/FOcal Reducer and low dispersion Spectrograph 2 (FORs2) imaging in the bands b_{High} (b_H), I_{Bessel} (I_B) and z_{Gunn} (z_G) was obtained for the 14 clusters under programmes 088.A-0889 and 089.A-0824 (PI: Mohr). Observations were carried out in queue

mode under clear conditions. The nominal exposure times for the different bands are 480 s (b_H), 2100 s (I_B) and 3600 s (z_G). These exposure times are achieved by co-adding dithered exposures with 160 s (b_H), 175 s (I_B) and 120 s (z_G). Deviations from the nominal exposure times are present for some fields due to repeated observations when conditions violated specified constraints or observing sequences that could not be completed during the original planned semester. The pseudo-colour images of the 14 SPT clusters constructed from VLT bands b_H , I_B and z_G are shown in Fig. 1. Each image shows also SZE contours (white), the R_{500} virial region (green circle) and the BCG (yellow circle).

Data reduction is performed with the THELI pipeline (Erben et al. 2005; Schirmer 2013). Twilight flats are used for flat-fielding. The I_B - and z_G -band data are defringed using fringe maps extracted from night sky flats constructed from the data themselves. To avoid oversubtracting the sky background, the background subtraction is modified from the pipeline standard as described by Applegate et al. (2014).

The FORs2 field of view is so small that only a few astrometric standards are found in the common astrometric reference catalogues. Many of them are saturated in our exposures. While we use the overlapping exposures from all passbands to map them to a common astrometric grid, the absolute astrometric calibration is adopted from mosaics of $F606W$ images centred on our clusters from the complementary Advanced Camera for Surveys (ACS)/*HST* programmes C18-12246 (PI: Stubbs) and C19-12447 (PI: High). Each cluster is observed in the well-dithered mode through $F606W$ and $F814W$ filters. For $F606W$ imaging, the cluster is imaged by four pointings with minimal overlap to remove the chip gap; these mosaics span a field of view of $6.7 \times 6.7 \text{ arcmin}^2$ centred on the cluster core. For $F814W$ imaging, only one pointing centred on cluster core is acquired. In this work, we use only the $F606W$ observations for deriving the stellar masses.

Cataloguing of the VLT images is carried out using SExtractor (Bertin & Arnouts 1996) in dual image mode. The detection image is created through the combination of I_B and z_G . Cataloguing of

Table 1. Cluster properties and photometric depths: The columns contain the cluster name, redshift and coordinates of the X-ray centre and BCG followed by the 10σ depths in each band.

Cluster	Redshift	α_X (deg)	δ_X (deg)	α_{BCG} (deg)	δ_{BCG} (deg)	$m_{b_H}^{10\sigma}$	$m_{F606W}^{10\sigma}$	$m_{I_B}^{10\sigma}$	$m_{z_G}^{10\sigma}$	$m_{[3.6]}^{10\sigma}$	$m_{[4.5]}^{10\sigma}$
SPT-CL J0000–5748	0.702	0.2500	–57.8093	0.2502	–57.8093	23.61	26.36	24.94	24.19	22.04	20.79
SPT-CL J0102–4915	0.870	15.7340	–49.2656	15.7407	–49.2720	24.34	26.31	24.51	24.14	22.21	21.86
SPT-CL J0205–5829	1.320	31.4437	–58.4856	31.4511	–58.4801	24.51	26.44	24.54	23.74	22.21	20.76
SPT-CL J0533–5005	0.881	83.4060	–50.0965	83.4144	–50.0845	24.64	26.84	24.66	23.99	22.01	20.56
SPT-CL J0546–5345	1.067	86.6548	–53.7590	86.6569	–53.7586	24.64	26.56	24.51	23.71	21.86	20.89
SPT-CL J0559–5249	0.609	89.9329	–52.8266	89.9300	–52.8242	24.49	26.46	24.31	24.06	22.09	20.71
SPT-CL J0615–5746	0.972	93.9570	–57.7780	93.9656	–57.7802	24.49	26.26	24.11	23.86	21.99	20.76
SPT-CL J2040–5726	0.930	310.0631	–57.4287	310.0552	–57.4208	24.51	26.34	24.69	24.21	22.24	20.76
SPT-CL J2106–5844	1.132	316.5179	–58.7426	316.5192	–58.7411	24.84	26.24	24.61	23.71	22.31	20.49
SPT-CL J2331–5051	0.576	352.9634	–50.8649	352.9631	–50.8650	24.04	26.41	24.94	23.51	22.29	20.71
SPT-CL J2337–5942	0.775	354.3523	–59.7056	354.3650	–59.7013	24.66	26.36	24.59	23.86	22.24	20.86
SPT-CL J2341–5119	1.003	355.3000	–51.3287	355.3014	–51.3291	24.59	26.24	24.81	23.89	22.26	20.49
SPT-CL J2342–5411	1.075	355.6916	–54.1849	355.6913	–54.1847	24.46	26.31	24.31	23.91	22.26	20.64
SPT-CL J2359–5009	0.775	359.9327	–50.1697	359.9324	–50.1722	24.84	26.19	24.66	23.91	21.74	20.66

the *HST* images is carried out separately, also using `SEXTRACTOR`. Galaxy photometry is extracted using `MAG_AUTO`. The VLT and *HST* photometry is matched at the catalogue level with a 1 arcsec match radius.

Because VLT data are generally not taken in photometric conditions, the photometric calibration is also carried out using data from the *HST* programmes. We derive a relation between $F814W$ magnitudes and the FORS2 I_B filter

$$m_{I_B} - m_{F814W} = -0.052 + 0.0095(m_{F606W} - m_{F814W})$$

from the Pickles (1998) stellar library, which is valid for stars with $(m_{F606W} - m_{F814W}) < 1.7$ mag. After deriving the absolute photometric calibration of the FORS2 I_B passband from this relation, the relative photometric calibrations of the other bands are fixed using a stellar locus regression (e.g. High et al. 2009; Desai et al. 2012) in the $(m_{b_H}, m_{F606W}, m_{I_B}, m_{z_G})$ colour space. The inclusion of $F606W$ data in this process is necessary because the stellar locus in $(m_{b_H}, m_{I_B}, m_{z_G})$ colours has no features.

All our clusters were observed with the *Spitzer* Infrared Array Camera (IRAC; Fazio et al. 2004) at both 3.6 and 4.5 μm under programmes PID 60099, 70053 and 80012 (PI: Brodwin). The images are acquired in dithered mode with exposure times of 8×100 and 6×30 s for 3.6 and 4.5 μm , respectively. We follow standard data reduction procedures to reduce the IRAC observations (Ashby et al. 2009). For each field, we generate a pair of spatially registered infrared mosaics: a relatively deep 3.6 μm image and a shallower 4.5 μm image. These images are catalogued with `SEXTRACTOR` in dual image mode, using the 3.6 μm mosaic as the detection image. We use the `SEXTRACTOR` `MAG_AUTO` and its associated uncertainty. We verify our detections by visually inspecting the `SEXTRACTOR` object check image. Because the IRAC point spread function is significantly larger than in either the *HST* or VLT imaging, we match our two-band IRAC photometry ([3.6] and [4.5]) to the nearest optical counterpart at the catalogue level, using a 1 arcsec match radius. If an object has multiple matches within the *Spitzer* point spread function, we then deblend the IRAC fluxes into the counterparts as described below.

For the objects in the *Spitzer*/IRAC catalogue with multiple optical counterparts, we deblend the [3.6] and [4.5] fluxes using the properties of the optical counterparts in z_G . Specifically, we deblend the *Spitzer*/IRAC fluxes assuming that the flux ratios of the neigh-

bouring objects in the IRAC band are the same as in the reddest optical band:

$$R_{(i,j)}^{[3.6],[4.5]} \equiv \frac{f_i}{f_j} \Bigg|_{[3.6],[4.5]} = \frac{f_i}{f_j} \Bigg|_{z_G}, \quad (1)$$

where f_i is the flux of object i .

We test the relationship between the flux ratios in z_G and the two IRAC bands by estimating the flux ratios of matched objects without close optical neighbours. We find that the intrinsic scatter of $R_{(i,j)}^{[3.6]}$ and $R_{(i,j)}^{[4.5]}$ are of the order of 0.6 and 0.8 dex, respectively. We add this scatter into the flux uncertainties in [3.6] and [4.5] of deblended objects.

Although the uncertainties in the deblended fluxes are large, we find that adding these two IRAC bands – deblended using our method – reduces the uncertainties of the stellar mass estimates by a mean value of 20 per cent and reduces the lognormal scatter of the reduced χ^2 (Section 3.3.1) by 29 per cent. Moreover, through studying an ensemble of pairs of unblended sources that we first artificially blend and then deblend, we find that our method does not introduce biases in the resulting mass estimates.

The fraction of blended IRAC sources lying projected within R_{500} for the 14 clusters varies from 11 to 20 per cent with a mean of 16 per cent and a standard deviation of 2.3 per cent. From 25 to 55 per cent of the sources are blended within $0.5R_{500}$. Thus, the majority (>80 per cent) of sources used in our analysis are not affected by flux blending.

We derive 10σ depth $m_{\text{filter}}^{10\sigma}$ for six passbands (filter = $b_H, F606W, I_B, z_G, [3.6], [4.5]$) of each cluster in the catalogue stage by estimating the magnitude where the median of the `MAG_AUTO` error distribution is equal to 0.11. These values are listed in Table 1. The $m_{\text{filter}}^{10\sigma}$ depths show good consistency to the 10σ depths estimated by measuring the sky variance in 2 arcsec apertures within the VLT images. The $m_{[3.6]}^{10\sigma}$ depths are about 2 mag deeper than our estimated m^* for each cluster, and hence the cluster galaxies should be detected without significant incompleteness.

We estimate the m^* of each passband for each cluster using a composite stellar population (CSP) model (Bruzual & Charlot 2003). This model has a burst at $z = 3$ that decays exponentially with e-folding time-scale of $\tau = 0.4$ Gyr. The tilt of the red sequence is modelled by using six CSPs with different metallicities and by calibrating those models to reproduce the Coma red sequence (for more details, see Song et al. 2012a). This model has been shown to

be adequate to derive accurate red sequence redshifts within SPT selected clusters to $z > 1$ with the root mean square of the cluster's photo- z error $\Delta z/(1+z)$, calibrated with spectroscopic clusters, of 0.02 (Song et al. 2012b; Hennig, in preparation). This model provides a good representation of the colour and tilt of the red sequence and the evolution of m^* in SPT selected galaxy clusters extending to $z \approx 1.2$ (Hennig, in preparation).

2.1.2 X-ray data

Out of the 14 clusters in our sample, 11 have been targeted by the *Chandra* X-ray telescope with programme nos. 12800071, 12800088 and 13800883. The remaining three clusters, SPT-CL J0205–5829 ($z = 1.32$; see Stalder et al. 2013), SPT-CL J0615–5746 ($z = 0.972$) and SPT-CL J2040–5726 ($z = 0.93$), have been observed with *XMM-Newton* with programme 067501 (PI: Andersson). The X-ray follow-up observations are designed to observe the SPT clusters uniformly with the goal of obtaining between 1500 and 2000 source photons within R_{500} . These photons enable us to measure the ICM projected temperature, the density profile and the mass proxy Y_X (the product of the ICM mass and X-ray temperature) with ~ 15 per cent accuracy.

The *Chandra* data reduction is fully described in previous publications (Andersson et al. 2011; Benson et al. 2013; McDonald et al. 2013). We include an additional cluster with *Chandra* data (ObsID 12258), the massive merging cluster SPT-CL J0102–4915 (Menanteau et al. 2012; Jee et al. 2014) at $z = 0.87$, which we analyse in an identical way to those previous works (Benson et al., in preparation). For the *XMM-Newton* data, we use *SAS* 12.0.1 to reduce and reprocess the data. All three cameras (MOS1, MOS2 and pn) are used in our analysis. The background flare periods are removed in both hard and soft bands using 3σ clipping after point source removal. We describe the ICM mass measurements in Section 3.2.

2.2 Comparison samples for this study

To place our results in context and to have a more complete view of the possible redshift variation of the baryon content in galaxy clusters, we compare our measurements with the published results from the local universe at $z \leq 0.1$. We include L03, Zha11 and GZ13 because they all provide estimates of f_* , f_{ICM} and f_b for large samples over a broader mass range than we are able to sample with the SPT selected clusters. L03 study 27 nearby galaxy clusters selected by optical/X-ray with masses ranging from 10^{14} to $10^{15} M_\odot$; 13 of these have available ICM mass measurements (Mohr et al. 1999). There are 19 clusters in Zha11, in which M_{500} is estimated using velocity dispersions. We discard two clusters, A2029 and A2065, from Zha11 because they argue that the virial mass estimates are biased due to the substructures. GZ13 estimate mass fractions for 15 nearby clusters, 12 of those have stellar mass measurements. In addition, we include the clusters and groups from H13 and vdB14 that extend to $z \geq 0.8$, allowing a more complete study at high redshifts. H13 study the stellar composition of 10 SZE selected clusters from ACT, and vdB14 study the Gemini CLuster Astrophysics Spectroscopic Survey (GCLASS) sample, consisting of 10 low-mass clusters selected by *Spitzer*/IRAC imaging. We restrict the cluster sample to those with virial masses above $3 \times 10^{14} M_\odot$, which is the mass regime probed by the SPT-SZ sample. This results in a total of 34 clusters in the comparison sample. We note that the majority of the vdB14 sample is in the low-mass regime and therefore falls below our mass threshold; our results should not be extrapolated into this lower mass regime.

There are several important differences between these studies and ours. We note that the groups or the clusters in these samples, with the exception of those in H13, are either selected from X-ray or optical/NIR surveys. Thus, these differences in selection method could potentially lead to observable differences in the samples. In addition to these selection differences, there are differences in the stellar mass and virial mass estimates. We describe below the corrections we apply to the comparison sample to address these differences, thereby enabling a meaningful combination with the SPT sample.

2.2.1 Correcting to a common IMF

The most important systematic factor for estimating stellar mass is the choice of the initial mass function (IMF) for the stellar population models that are then employed when converting from galaxy light to galaxy stellar mass. Different assumed IMFs introduce systematic shifts in the mass-to-light ratios of the resulting stellar populations (Cappellari et al. 2006). For instance, the conventional Salpeter (1955) IMF with a power-law index of -2.35 would predict a mass-to-light ratio higher by a factor of 2 than the one using the Kroupa (2001) IMF (Chabrier 2003; Kauffmann et al. 2003; Cappellari et al. 2006). For this analysis, we adopt the Chabrier (2003) IMF (see more detailed discussion in Section 3.3) and apply a correction to the literature results so all measured stellar masses are appropriate for this IMF. Specifically, we reduce the stellar mass measurements of L03 and Zha11 by 24 per cent (Lin et al. 2012; Zhang et al. 2012), the measurements of GZ13 by 24 per cent (or 0.12 dex) and the measurements of H13 by 42 per cent (or 0.24 dex). Because vdB14 use the same Chabrier IMF as in this work, no IMF correction is needed.

2.2.2 Correcting for virial mass systematics

To enable a meaningful comparison of the baryon content across samples, it is crucial to use a consistent virial mass estimate for all samples. Zha11, H13 and vdB14 estimate M_{500} using velocity dispersions, while the other analyses all use X-ray mass proxies (ICM temperature) to estimate virial masses. Our SPT masses arise from a recent analysis (Bocquet et al. 2015) that includes corrections for selection effects, marginalization over cosmological parameters and systematic uncertainties and combination with external cosmological data sets (see discussion in Section 3.1).

The Bocquet et al. (2015) analysis quantifies the systematic mass shifts that result for SPT clusters when using only X-ray data, only velocity dispersion data or the full combined data set of X-ray, velocity dispersions and external cosmological constraints from CMB, BAO and SNe. Namely, when compared to our cluster mass estimates obtained using the full combined data set, the SPT cluster masses inferred from the X-ray mass proxy Y_X alone have a systematically lower mass by 44 per cent, and masses inferred from velocity dispersions alone have systematically lower masses by 23 per cent. As explained in more detail in Section 3.1, we adopt the full combined data set masses for the analysis of our SPT cluster sample.

For the comparison here, it is not crucial to know which virial mass estimate is most accurate. What we must do is adopt one mass calibration method for our SPT sample and then correct the virial mass estimates in the other samples to a consistent mass definition before making comparisons of the baryon content. To make these corrections, we use the results from the recent SPT mass analysis

(Bocquet et al. 2015) to apply a correction to the virial mass scale in each literature sample to bring it into better consistency with our SPT sample.

Specifically, we estimate the M_{500} of the clusters in L03 by using the same T_X – M_{500} relation (Vikhlinin et al. 2009) used in GZ13; then we increase the L03 and GZ13 masses by 44 per cent, assuming that the systematic offset of Y_X derived SPT virial masses is the same for these clusters whose masses were derived using the T_X – M_{500} relation. Similarly, we increase the masses in Zha11, H13 and vdB14 by 23 per cent, because their masses are derived from velocity dispersion measurements.

Increasing M_{500} increases the virial radius and therefore also increases the stellar and ICM masses. Specifically, a 44 per cent (23 per cent) increase in virial mass leads to a 13.2 per cent (7.4 per cent) and 12.9 per cent (7.1 per cent) increment in M_* and M_{ICM} , respectively, assuming that the cluster galaxies are distributed as an Navarro–Frenk–White (NFW) model with concentration $c_{500} = 1.9$ and the ICM near the virial radius falls off as a β -model (Cavaliere & Fusco-Femiano 1978) with $\beta = 2/3$. In correcting the literature results for comparison to the SPT sample, we apply a correction that accounts for the shifts in all the different masses.

Correcting previously published masses to account for different data sets and analysis methods allows us to more accurately compare the results, but this correction procedure has inherent uncertainties. It is challenging to quantify these remaining uncertainties, but for this analysis we adopt a systematic virial mass uncertainty of 15 per cent (1σ) when constraining the redshift variation with the combined sample. We return to this discussion in Section 5 where we present our fitting procedure in detail. Also, in the conclusions we comment on the impact of adopting other systematic uncertainties.

We note in passing that if we had adopted the SPT masses calibrated only using the X-ray mass proxy Y_X , the SPT cluster virial masses M_{500} would decrease on average by a factor of $1/1.44$. The new values for the SPT sample quantities M_* , M_{ICM} , f_* , f_{ICM} , f_c and f_b can be approximated by applying the scale factors 0.87, 0.88, 1.26, 1.27, 0.99 and 1.27, respectively, to the measurements presented in Table B1.

3 MASS MEASUREMENT METHODS

In this section, we describe the method for estimating the virial, the ICM and the stellar masses.

3.1 SPT cluster virial mass M_{500} measurements

The virial masses (M_{500}) that we use come from the mass calibration and cosmological analysis of Bocquet et al. (2015). They are derived using the SPT SZE observable ξ , the cluster redshift, and a combination of internal and external calibration data. These data include direct mass information from 63 measured cluster velocity dispersions (observed using Gemini South, the VLT, and the Magellan Baade and Clay telescopes; see Ruel et al. 2014) and 16 Y_X measurements (Andersson et al. 2011; Foley et al. 2011; Benson et al. 2013). In addition, mass information derives from the 100 cluster candidates extracted from the first 720 deg² of the SPT-SZ survey. These SPT data are then jointly analysed in combination with *Planck* temperature anisotropy, *WMAP9* polarization anisotropy, BAO and SNIa constraints.

As explained in Bocquet et al. (2015, see fig. 2), adopting such strong external cosmological constraints has a dramatic impact on

the cluster masses, pushing them higher to better match the masses expected within the preferred cosmological model, given the ξ and redshift distribution of the cluster sample. In contrast, the Y_X constraints prefer lower masses, and the velocity dispersions prefer masses in the middle. By combining all the constraints, one ends with a mass calibration that prefers higher masses than the masses one would obtain when using solely the Y_X 's or velocity dispersions as calibrators (see also further discussion in Section 2.2.2). We adopt these masses that arise from a combination of internal and external calibration data for the analysis below.

Our SPT masses are corrected for Eddington bias that arises from the scatter between the mass and the selection variable ξ and the steep cluster mass distribution. The intrinsic scatter in mass at fixed ξ is approximately 16 per cent, and there is an additional measurement scatter that reaches ≈ 14 per cent at $\xi = 5$. Final mass uncertainties include marginalization over all cosmological and scaling relation parameters. Thus, our masses and mass uncertainties include a combination of the systematic and statistical uncertainties. Typical final mass uncertainties are ~ 20 per cent. The masses are then used to calculate R_{500} , which has a characteristic uncertainty of ≈ 7 per cent. We refer the reader to Bocquet et al. (2015) for additional details. The virial mass systematics correction for the comparison sample is described in Section 2.2.2.

3.2 ICM mass measurements

In this work, we adopt the X-ray ICM mass M_{ICM} measurements extracted within R_{500} . We determine the centre of the cluster (α_X, δ_X) iteratively as the centroid of X-ray emission in the 0.7–2.0 keV energy band within a 250–500 kpc annulus (see Table 1). The final centroid is visually verified on the smoothed X-ray emission map and is adjusted to match the centre of the most circularly symmetric isophote if it deviates significantly from the peak. The ICM density profile is estimated by fitting the X-ray surface brightness profile extracted in the energy range 0.7–2.0 keV assuming spherical symmetry and centred on the derived centroid. For *Chandra* observations, we fit the modified single β -model (equations 1 and 2 in Vikhlinin et al. 2006) to the X-ray surface brightness profile. The details of the X-ray analysis are given elsewhere (Andersson et al. 2011; McDonald et al. 2013).

Because we cannot simultaneously constrain all the parameters in the modified single β -model for the *XMM-Newton* observations, we instead fit a single β -model for SPT-CL J0205–5829 ($z = 1.32$), SPT-CL J0615–5746 ($z = 0.972$) and SPT-CL J2040–5726 ($z = 0.93$). For these clusters, the single β -model provides a good fit to the *XMM-Newton* X-ray surface brightness profile. The best-fitting X-ray surface brightness profile then provides the radial distribution of the ICM, and we use the flux of the cluster within the 0.15–1.0 R_{500} annulus to determine the central density (e.g. Mohr et al. 1999). We assume the metal abundance of the ICM is 0.3 solar, resulting in $n_e/n_p = 1.199$ and $\mu \equiv \rho_{\text{ICM}}/(m_p n_e) = 1.16$, where the subscripts p and e denote proton and electron, respectively.

To estimate M_{ICM} , we integrate the measured ICM profile to R_{500} obtained from the SZE derived M_{500} . The uncertainty of M_{ICM} is estimated by propagating the uncertainties of the best-fitting parameters. Deriving the X-ray temperature of the ICM free from the instrumental calibration bias can be challenging; however, the ICM mass and density profile is insensitive to the temperature (Mohr et al. 1999) and to instrumental systematics (Donahue et al. 2014; Martino et al. 2014; Schellenberger et al. 2015). Thus, we do not expect significant systematics in the ICM masses.

3.3 Stellar mass measurements

In the sections below, we describe the SED fitting to determine galaxy stellar masses and our method of making a statistical background correction.

3.3.1 SED fitting

We use the multiband photometry to constrain the SED of each galaxy and to estimate its stellar mass. The photometry of the six bands (b_H , $F606W$, I_B , z_G , [3.6], [4.5]) is used for each galaxy. We use the LE PHARE SED fitting routine (Arnouts et al. 1999; Ilbert et al. 2006) together with a template library that consists of stellar templates (Pickles 1998) and galaxy templates from CSP models (Bruzual & Charlot 2003) derived using a Chabrier (2003) IMF. The systematics correction for the different IMF used in the comparison sample is described in Section 2.2.1. The stellar templates include all normal stellar spectra together with the spectra of metal-weak F through K dwarfs and G through K giants. The galaxy library includes templates that cover (1) a wide range in metallicity $Z = 0.004, 0.008, 0.02$, (2) an e-folding exponentially decaying star formation rate with characteristic time-scale $\tau = 0.1, 0.3, 1.0, 2.0, 3.0, 5.0, 10.0, 15.0, 30.0$ Gyr, (3) a broad redshift range between 0.0 and 3.0 with steps of 0.05, and (4) the Calzetti et al. (2000) extinction law evaluated at $E(B - V) = 0.0, 0.2, 0.4, 0.6, 0.8, 1.0$. Our galaxy library contains no templates with emission lines.

We run the LE PHARE routine with this template library on every object that lies projected within R_{500} and is brighter than $m^* + 2.0$ within the z_G passband (except that we use [3.6] for the two clusters at $z > 1.1$). This ensures that we are selecting the galaxy population in a consistent manner over the full redshift range. For each galaxy, we adopt a uniform prior on the extinction law $E(B - V)$ between 0.0 and 1.0 and a weak, flat prior on the stellar mass between 10^8 and $10^{13} M_\odot$.

For the SED fit, we increase the MAG_AUTO flux uncertainties for all six passbands by a factor of 2. We estimate this correction factor by examining the *photometric repeatability* of the galaxies that appear in multiple individual VLT exposures (Desai et al. 2012; Liu et al. 2015). With this correction, the resulting magnitude uncertainties correctly describe the scatter in the repeated photometric measurements of the same galaxies. Rescaling the uncertainties has no significant impact on the final result but increases the uncertainty of the stellar mass estimate for each galaxy by 25 per cent.

For each cluster, we first estimate the stellar mass of the BCG, M_*^{BCG} , fixing the redshift to the cluster redshift. The BCG is chosen to be the brightest cluster galaxy projected within R_{500} ; we select this galaxy visually using the NIR and optical imaging and then confirm in the catalogue (z_G and [3.6]) that it is the brightest galaxy. We find that the BCGs in our cluster sample all prefer the templates with the characteristic e-folding time-scale for the star formation rate to be $\tau \leq 1$ Gyr. This indicates that the rapid star-forming activity seen in the SPT selected Phoenix Cluster BCG (McDonald et al. 2014) is not present in our cluster sample. This result is consistent with the view that the evolution of the typical BCG is well described by a CSP model with $\tau \approx 0.9$ Gyr out to redshift 1.5 (Lidman et al. 2012). For the final M_*^{BCG} estimates, we restrict the template library to $\tau \leq 1$ Gyr, which results in an ≈ 6 per cent reduction in the stellar mass uncertainties as compared to fitting across the full range of τ . This small change in uncertainty has no impact on our final result. We then estimate the stellar mass for the remaining galaxies using the same configuration except that we allow the redshift to float and fit the templates without restricting τ .

We adjust the LE PHARE routine to output the best-fitting mass M_*^{best} , the median mass M_*^{med} , the mass at the lower (higher) 68 per cent confidence level M_*^{lo} (M_*^{hi}) and the best-fitting χ^2 extracted over the full template library. We discard the objects from the analysis where the best-fitting χ^2 arises for a stellar template. This stellar removal works well; testing on the COSMOS field (Cappak et al. 2007; Sanders et al. 2007; Ilbert et al. 2009) indicates that we have a residual stellar contamination and a false identification rate for galaxies under 1.5 per cent and ~ 0.15 per cent, respectively. The mass-to-light ratios Υ and their rms variations in the observed frame [3.6] band for all clusters are provided in Table B1. These are extracted from the subset of galaxies projected within the virial region that have photo- z 's that are within $\Delta z = 0.1$ of the cluster spectroscopic redshift.

We examine those galaxies with $M_*^{\text{best}} > M_*^{\text{BCG}}$ closely, because we expect no galaxy to be more massive than the BCG. We find that most of these galaxies can be excluded because they have redshifts far higher than the cluster. In total, there are 37 out of 2640 galaxies with $M_*^{\text{best}} > M_*^{\text{BCG}}$ within R_{500} of the 14 clusters. That is, about 1.5 per cent of the objects are discarded through this process. However, one must take special care in cases of merging clusters, which could host one or more galaxies with masses similar to the most massive one. In a few cases (three galaxies to be exact), we find through photo- z and visual inspection that these galaxies likely lie in the cluster and have mass estimates slightly larger than the mass of our selected BCG. In these cases, we include those galaxies in the stellar mass estimate. We provide further details of our SED fit performance in Appendix A.

The stellar mass estimate for the region within R_{500} , including the foreground and background galaxies, is the sum of M_*^{best} . The uncertainty includes the uncertainties on the single galaxy masses (using M_*^{lo} and M_*^{hi} and assuming that the probability distribution for the stellar mass is Gaussian).

We also calculate the fraction of objects f_{cor} with unphysical mass outputs (i.e. $\log(M_*^{\text{med}}) = -99.0$), which occur mostly due to data corruption. We correct for these missing galaxies by assuming that they share the same distribution of stellar masses as the uncorrupted galaxies. We note that this fraction varies between 3 and 10 per cent. A correction for the masking of the bright stars is also applied. Thus, for each cluster, we estimate the total stellar mass M_*^{field} projected within R_{500} as

$$M_*^{\text{field}} = \frac{\sum_i M_{*,i}^{\text{best}}}{(1 - f_{\text{mask}})(1 - f_{\text{cor}})}, \quad (2)$$

where f_{mask} is the fraction of area within R_{500} that is masked and $M_{*,i}^{\text{best}}$ is the best stellar mass estimate for galaxy i in the cluster.

3.3.2 Background correction

We correct the stellar mass from the cluster field M_*^{field} for the contribution from the foreground and background galaxies M_*^{bkg} using a statistical correction. Because the FORS2 field of view is small, the background regions outside R_{500} are contaminated by cluster galaxies. Thus, we use the COSMOS survey to estimate the background correction.

The COSMOS survey has 30-band photometry with wavelength coverage from UV to mid-infrared. To minimize systematics, we take two steps to make the COSMOS data set as similar as possible to our SPT data set. First, we choose the passbands which are most similar to ours (from Subaru Suprime-Cam and *Spitzer*) and apply colour corrections where needed to convert the COSMOS

photometry into our passbands. MAG_AUTO photometry is used in the COSMOS field. Secondly, we degrade the COSMOS photometry to have the same measurement noise as in our data set.

We then measure the stellar mass for each galaxy in the COSMOS field using the converted photometry, the same *Spitzer* object detection, the same matching algorithm and the same fitting strategy as we applied to our own data. We correct this background estimate for the fraction of corrupted galaxies as described for the cluster fields in equation (2).

Then, correcting the COSMOS background estimates to the area of each cluster field, we subtract the background estimate M_*^{bkg} , obtaining our estimate of the cluster stellar mass projected within R_{500} . We then apply a geometric factor f_{geo} to correct this projected quantity to the stellar mass within the virial volume M_* using a typical radial galaxy profile with concentration $c_{500}^{\text{gal}} = 1.9$ (Lin, Mohr & Stanford 2004; Hennig, in preparation), which corresponds to a normalization correction of $f_{\text{geo}} = 0.71$,

$$M_* = M_*^{\text{BCG}} + f_{\text{geo}} (M_*^{\text{field}} - M_*^{\text{bkg}}), \quad (3)$$

where we have defined M_* to include M_*^{BCG} , the BCG stellar mass.

In Appendix B, we compare the COSMOS background to the background estimated in the non-cluster portions of the VLT imaging where a correction for cluster contamination has been applied. We find that the two backgrounds agree at the 10 per cent level, leading to background-corrected cluster stellar mass estimates M_* that are consistent at the 4 per cent level. Thus, we adopt this difference as the amplitude of the systematic uncertainty associated with our statistical background correction.

4 BCG AND CLUSTER SMF

In this section, we present the components of the cluster SMF. These include the BCGs (Section 4.1), which we discuss first, followed by the full SMF and the luminosity function (LF) of the satellite galaxies (Section 4.2).

4.1 BCG stellar mass

We present M_*^{BCG} for the 14 SPT clusters and compare them with the measurements of H13 and vdB14, in which groups and clusters at $z \geq 0.3$ are studied. H13 estimated M_*^{BCG} based on the mass-to-light technique assuming a passive evolution model with the [3.6] magnitude MAG_AUTO. vdB14 applied the same technique using the K_s luminosity together with the Sérsic model fitting to the light profile. As noted in vdB14, the magnitude inferred by the Sérsic profile could differ from MAG_AUTO by up to 0.2 mag, depending on the shape of the light profile. In this work, we estimate M_*^{BCG} using an SED fit to the six bands available in our survey. No special attempt is made to include or deblend the extended halo or intracluster light (ICL) in any of these studies. We have three clusters in common with H13: SPT-CL J0102–4915, SPT-CL J0546–5345 and SPT-CL J0559–5249. For SPT-CL J0546–5345, our M_*^{BCG} is about a factor of 2 higher, but in the other two clusters M_*^{BCG} agrees at better than 10 per cent. We compare the [3.6] photometry of SPT-CL J0546–5345, and find that the magnitude reported by H13 is about 0.5 mag fainter, suggesting that a more sophisticated deblending algorithm is needed for the crowded core of SPT-CL J0546–5345 in this work. We adopt the SED mass estimates for the BCGs in the analyses that follow. The BCGs are marked by yellow circles in the optical images presented in Fig. 1.

As is clear in Fig. 2, neither the SPT nor the H13 sample provides strong evidence for a correlation between the BCG mass and the




Figure 2. The BCG stellar mass (M_*^{BCG}) versus cluster virial mass M_{500} for the SPT sample (blue), H13 (cyan) and vdB14 (red). The H13 sample is corrected to Chabrier IMF. The blue dashed line is the best fit for the SPT sample alone and the black solid line is the best fit for the combined sample (see equation 4).

cluster halo mass. However, in combination with the vdB14 sample that extends to much lower mass, we find a best-fitting $M_*^{\text{BCG}}-M_{500}$ relation of

$$M_*^{\text{BCG}} = (5.30 \pm 0.39) \times 10^{11} \left(\frac{M_{500}}{3 \times 10^{14} M_{\odot}} \right)^{0.42 \pm 0.07}, \quad (4)$$

for the combined sample, and this relation is plotted in Fig. 2 (black dashed line). Thus, the BCG stellar mass constitutes about 0.12 per cent of the cluster halo mass at $M_{500} = 6 \times 10^{14} M_{\odot}$. Because M_*^{BCG} scales with cluster halo mass with a power-law index less than 1, the fraction of the cluster mass made up by the BCG falls as $M_*^{\text{BCG}}/M_{500} \propto M_{500}^{-0.58 \pm 0.07}$.

The SPT sample scatters significantly about this relation, providing evidence of intrinsic scatter in M_*^{BCG} at fixed cluster halo mass of $\sigma_{\text{int}} = 0.17 \pm 0.034$ dex. The full sample exhibits a consistent value $\sigma_{\text{int}} = 0.15 \pm 0.021$ dex. Thus, the characteristic scatter of the BCG masses at a fixed cluster halo mass is 41 per cent.

4.2 Cluster LF and SMF

We extract the [3.6] LF and the SMF using a statistical background subtraction with the COSMOS field as the source of the background (see Section 3.3.2). We apply a correction from the virial cylinder to the virial volume in the same manner as in Section 3.3.2. The measured LF and SMF are in physical density units of Mpc^{-3} . The uncertainty of each bin is estimated by the Poisson error associated with the galaxy counts in the case of the LF and this error combined with the galaxy stellar mass measurement uncertainties for the SMF.

We stack the LF and SMF from 14 SPT clusters using inverse-variance weighting within each bin. The number densities are corrected to the median redshift of the SPT clusters, $z = 0.9$, by multiplying by the ratio of the critical densities, $\left(\frac{E(0.9)}{E(z)} \right)^2$, where $E(z)^2 \equiv \Omega_{\Lambda} + \Omega_{\text{M}}(1+z)^3$ and z is the redshift of the cluster. We stack the LF within the space of $m - m^*$ with magnitude bins of width 0.5, where




Figure 3. The stacked LF of 14 SPT clusters extracted from the [3.6] photometry (black points). The grey point is fainter than $m^* + 2$ and is not included in the fit. The line marks the best-fitting Schechter function. The LF is plotted versus $m - m^*$, where m^* is obtained from the passive evolution model described in the text (Section 2.1.1). The stacked number densities are corrected for evolution of the critical density ($\rho_{\text{crit}} \propto E(z)^2$) and normalized to median redshift $z = 0.9$.

m^* comes from the CSP model described in Section 2.1.1. Given that the galaxy population in SPT selected clusters has been shown to be well described by the CSP model (Song et al. 2012b; Hennig et al., in preparation), stacking LFs with respect to the m^* predicted at the redshift of each cluster provides a simple way to extract the information for the normalization and shape of the composite LF. We stack the SMF in the stellar mass range from 10^{10} to $10^{12} M_{\odot}$ with bin width of 0.2 dex. Finally, we characterize the stacked LF and SMF with the standard Schechter function (Schechter 1976). Specifically, we fit the stacked LF directly in log space to

$$\Phi_L(m) = 0.4 \ln(10.0) \phi^* \times 10.0^{-0.4(\alpha_L+1)(m-m_0)} \times \exp(-10.0^{-0.4(m-m_0)}), \quad (5)$$

where m is the magnitude, m_0 is the characteristic magnitude predicted by the passively evolving model (see Section 2.1.1), ϕ^* is the characteristic density and α_L is the faint-end slope. We fit the stacked SMF directly in log space to

$$\Phi_M(M_*) = \ln(10) \phi_M \times 10^{(\alpha_M+1)(m_*-M_0)} \times \exp(-10^{(m_*-M_0)}), \quad (6)$$

where m_* is the stellar mass in units of $\log_{10}(m_*/M_{\odot})$, M_0 is the characteristic mass, ϕ_M is the characteristic density and α_M is the faint-end slope. We restrict our fit to those galaxies brighter than $m^* + 2$ in the LF analysis. Because the stellar mass is not a linearly rescaled version of the magnitude, we choose the conservative depth limit used in the SMF analysis, which is based on the mass-to-light-inferred mass at brighter magnitude, $m^* + 1.5$, assuming the passively evolving model for SMF analysis.

The stacked LF and SMF are shown in Figs 3 and 4, respectively. The best-fitting parameters are given in Table 2. We convert the SMF and LF from physical number density to the abundance per mass of $10^{15} M_{\odot}$ (total baryon and dark matter mass) by using the mean density within the virial region at $z = 0.9$, which is




Figure 4. The measured SMF obtained by stacking 14 SPT clusters. The black line is the best-fitting Schechter function (see Table 2). The grey points are measurements beyond the depth limit and are not used in the fit. For comparison, we show the field SMF from vdB13.

Table 2. LF and SMF parameters. The LF (top) characteristic density, characteristic magnitude, faint-end slope and reduced χ^2 are shown followed (below) by the equivalent SMF parameters.

ϕ^* ($\text{Mpc}^{-3} \text{mag}^{-1}$)	m_0 (mag)	α_L	χ_{red}^2
14.90 ± 1.0	-0.18 ± 0.1	-0.19 ± 0.1	0.6
ϕ_M ($\text{Mpc}^{-3} \text{dex}^{-1}$)	M_0 (dex)	α_M	χ_{red}^2
12.30 ± 1.5	11.06 ± 0.1	-0.32 ± 0.2	1.4

$500\rho_{\text{crit}}(z = 0.9)$. This value is shown on the right y-axis. Similarly, to compare to a field LF or SMF, one would convert from Mpc^{-3} to per unit mass by using the mean density of the universe at that redshift ($\rho(z) = \Omega_M(z) \times \rho_{\text{crit}}(z)$).

The best-fitting m_0 indicates that the LF deviates from the predicted characteristic $m_{[3.6]}^*$ for the passive evolution model (Section 2.1.1) by -0.18 ± 0.1 , suggesting the mild evidence (about 1.8σ) of the blue population at the high-redshift clusters. The best-fitting SMF and LF are consistent with one another; the characteristic $m_{[3.6]}^*$ at median redshift $z = 0.9$ predicted by the passively evolving model corresponds to the stellar mass of $10^{10.96} M_{\odot}$, while the measured characteristic mass is $10^{11.0 \pm 0.1} M_{\odot}$. The faint-end slopes and characteristic densities are also in good agreement.

In a recent paper, van der Burg et al. (2013, hereafter vdB13) compare the SMFs of the GCLASS low-mass clusters to the field at redshift $z = 0.85$ – 1.2 and find that the number density of galaxies per unit mass (dark matter plus baryons) in the field SMF is lower than that in the groups over the mass range 10^{10} – $10^{11.5} M_{\odot}$. This suggests that the galaxy formation rate has been lower over time in the field than in the dense group and cluster environments. A similar picture had previously emerged in the local Universe ($z < 0.1$; Lin et al. 2004, 2006), where the LFs of K -band selected galaxies and of radio sources within clusters are also significantly higher than the field after corrections for the mean matter density differences in the two environments. As seen in Fig. 4, the normalization of the SMF

for the SPT clusters on the massive end ($\log_{10}(m_*/M_\odot) \approx 11.2$ – 11.5) is significantly higher than the field (open triangle) measured by [vdB13](#). By integrating the best-fitting SMF of SPT above our single galaxy stellar mass threshold of $2.5 \times 10^{10} M_\odot$, we estimate that the number of galaxies per unit total mass for SPT clusters is $\approx 1.65 \pm 0.20$ times higher than the field at $z = 0.85$ – 1.2 . Our result reinforces this picture that the cluster environment contains a more biased galaxy population than the field.

5 BARYON COMPOSITION

In this section, we present our measurements for the stellar mass fraction, ICM mass fraction, collapsed baryon fraction and baryon fraction:

$$f_* \equiv \frac{M_*}{M_{500}} \quad (7)$$

$$f_{\text{ICM}} \equiv \frac{M_{\text{ICM}}}{M_{500}} \quad (8)$$

$$f_c \equiv \frac{M_*}{M_b} \quad (9)$$

$$f_b \equiv \frac{M_b}{M_{500}}, \quad (10)$$

where M_* is the stellar mass (see equation 3), M_{ICM} is the ICM mass (see Section 3.2) and $M_b \equiv M_* + M_{\text{ICM}}$ is the total mass in baryons. M_{500} is the halo virial mass, estimated using the SZE observable (see Section 3.1).

In addition, we study mass and redshift trends in our SPT clusters and in the combined sample that includes the clusters studied in the literature (discussed in Section 2.2). Note that we are not probing the evolution of the baryon content by directly tracing the progenitors, because our SPT sample lacks low-mass groups at all redshifts. We instead estimate the baryon content of the massive clusters with respect to the characteristic mass at the different epochs statistically by fitting the scaling relation simultaneously in mass and redshift space (see Section 5.1). We also compare our cluster results with more general results coming from external, non-cluster data sets. We use the universal baryon fraction f_b estimated using the *Planck* CMB anisotropy observations (*Planck* Collaboration XVI 2014), and we estimate the universal stellar density parameter Ω_* , where the mean stellar density at $z = 0$ is extracted from the local *K*-band galaxy LF (Kochanek et al. 2001) and the mean stellar density at $z = 1$ is extracted from the [vdB13](#) analysis. These values have been corrected to our fiducial cosmology and are appropriate for a Chabrier IMF, enabling comparison to our cluster measurements.

5.1 Fitting procedure

We fit these measurements from our cluster ensemble and also from the combined sample to a power-law relation in both mass and redshift:

$$f_{\text{obs}}(M_{500}, z) = \alpha_{\text{obs}} \left(\frac{M_{500}}{M_{\text{piv}}} \right)^{\beta_{\text{obs}}} \left(\frac{1+z}{1+z_{\text{piv}}} \right)^{\gamma_{\text{obs}}}, \quad (11)$$

where M_{piv} and z_{piv} are the mass and redshift pivot points, obs corresponds to the different observables, and α_{obs} , β_{obs} and γ_{obs} correspond to normalization of the best-fitting relation, the power-law index of the mass dependence and the power-law index of the redshift dependence, respectively. We perform χ^2 fitting directly in log space using the measurement uncertainties and accounting for

intrinsic scatter. For the SPT and combined samples, we choose the pivot points to be the median mass $M_{500} = 6 \times 10^{14} M_\odot$. For the SPT sample, we adopt the redshift pivot $z_{\text{piv}} = 0.9$, consistent with the median redshift of the sample, but for the combined sample we adopt a redshift pivot of $z_{\text{piv}} = 0$.

The parameters for the best-fitting relations for the SPT sample and for the combined sample are listed in Table 3, while the measured cluster virial masses, ICM masses, stellar masses and the derived quantities above are listed in Table B1. These results are summarized in Figs 5 and 6, where the first figure focuses on the mass trends and the second focuses on the redshift trends. In the subsections below, we discuss each derived quantity in turn.

5.1.1 Accounting for M_{500} systematic uncertainties

We account for systematic differences in M_{500} estimation between the low-redshift comparison sample ([L03](#), [Zha11](#) and [GZ13](#)) and the high-redshift sample (SPT with two additional samples of [H13](#) and [vdB14](#) added when comparing f_*) by adopting a 15 per cent (1σ) systematic virial mass uncertainty (see discussion in Section 2.2). These virial mass uncertainties imply corresponding R_{500} uncertainties that lead also to systematic uncertainties in the stellar mass and ICM mass for each cluster. We estimate the systematic uncertainties in the redshift variation parameter γ_{obs} (Table 3) by perturbing the virial masses of the high-redshift sample by ± 15 per cent and extracting the best-fitting parameters in each case. The 1σ systematic uncertainty is estimated as half the difference between the two sets of parameters. This virial mass systematic is only important for the measured redshift trends.

5.1.2 Accounting for differences in measurement uncertainties

We also account for systematic differences in the measurement uncertainties among the different samples by solving for a best-fitting intrinsic scatter separately for each sample. For the SPT sample, where mass uncertainties include both measurement and systematic uncertainties (Section 3.1), we find no need for an additional intrinsic scatter. The best-fitting estimates of the intrinsic scatter for the other samples are 9 per cent for f_{ICM} and f_b in [L03](#), 14 per cent for f_* , f_{ICM} and f_b in [GZ13](#), 18 per cent for f_* in [H13](#) and 20 per cent in [vdB14](#), and 20–22 per cent for the fractions in [Zha11](#), [H13](#) and [vdB14](#). Three of the samples with the largest intrinsic scatter ([Zha11](#), [H13](#) and [vdB14](#)) employ velocity dispersions for single cluster mass estimation as opposed to X-ray or SZE mass indicators. This is not surprising, because it has been shown that cluster velocity dispersions provide high scatter single cluster mass estimates (see [Saro et al. 2013](#), and references therein). Velocity dispersions can be effectively used in ensemble to calibrate ICM-based single cluster mass estimates ([Bocquet et al. 2015](#)).

5.2 Stellar mass fraction f_*

The stellar mass fraction we estimate here is the mass in stars within cluster galaxies. We make no attempt to account for the ICL component. Fig. 5 contains a plot of our results (blue). The mean f_* of our 14 clusters is 0.011 ± 0.001 , and the characteristic value at $z = 0.9$ and $M_{500} = 6 \times 10^{14} M_\odot$ is 0.011 ± 0.001 . The SPT sample provides no evidence for a mass or redshift trend, but the large mass trend uncertainty ($f_* \propto M_{500}^{-0.09 \pm 0.27}$) means the sample is statistically consistent with the trend for more massive clusters to have lower f_* ([L03](#)). In the combined sample, there is 3.7σ evidence

Table 3. Mass and redshift trends of baryon composition with $M_{\text{piv}} \equiv 6 \times 10^{14} M_{\odot}$. The columns contain the quantity of interested, the normalization at the pivot mass and redshift, mass dependence and redshift dependence (see equation 11) for the SPT sample alone (above) and for the SPT sample together with the literature sample (below).

f_{obs}	α_{obs}	β_{obs}	γ_{obs}^a
SPT sample results with $z_{\text{piv}} \equiv 0.9$			
f_{\star}	0.011 ± 0.001	-0.09 ± 0.27	1.07 ± 1.08
f_{ICM}	0.096 ± 0.005	0.43 ± 0.13	0.20 ± 0.49
f_{c}	0.107 ± 0.011	-0.55 ± 0.22	0.81 ± 0.93
f_{b}	0.107 ± 0.006	0.39 ± 0.13	0.32 ± 0.50
Combined sample results with $z_{\text{piv}} \equiv 0$			
f_{\star}	0.0099 ± 0.0005	-0.37 ± 0.09	$0.26 \pm 0.16 \pm 0.08$
f_{ICM}	0.1120 ± 0.0032	0.22 ± 0.06	$-0.20 \pm 0.11 \pm 0.22$
f_{c}	0.0859 ± 0.0049	-0.65 ± 0.10	$0.39 \pm 0.15 \pm 0.16$
f_{b}	0.1227 ± 0.0035	0.22 ± 0.06	$-0.17 \pm 0.11 \pm 0.22$

Notes. ^aThe second γ_{obs} uncertainty arises from the 15 per cent M_{500} systematic uncertainty.




Figure 5. The baryonic fractions f_{\star} , f_{ICM} , f_{c} and f_{b} are shown as a function of cluster virial mass M_{500} for the combined sample. In all cases, the measurements have been corrected to $z = 0$ using the best-fitting redshift trend. The best-fitting mass trend is shown in green (Table 3). The colour coding and point styles are defined in the upper panel and is the same throughout. The red shaded region indicates the universal baryon composition from combining the best-fitting cosmological parameters from Bocquet et al. (2015) together with the local K -band LF (Kochanek et al. 2001).

for a mass trend $f_{\star} \propto M_{500}^{-0.37 \pm 0.09}$, which is also consistent with the L03 result. The combined sample exhibits no significant redshift variation ($f_{\star} \propto (1+z)^{0.26 \pm 0.16 \pm 0.08}$), where the second uncertainty reflects the 15 per cent (1σ) systematic virial mass uncertainty. The

characteristic value at $z = 0$ is $f_{\star} = 0.010 \pm 0.0005$ (statistical), which is in good agreement with the SPT value at $z_{\text{piv}} = 0.9$.

Also shown in the shaded region is the f_{\star} constraint emerging from a combination of the stellar mass density from the K -band local LF (Kochanek et al. 2001), $\Omega_{\star} h = 3.4 \pm 0.4 \times 10^{-3}$ with $h = 0.683$, with the most recent combined results (*Planck* + *WMAP* polarization+SNe+BAO+SPT clusters) on the cosmological matter density $\Omega_{\text{M}} = 0.299 \pm 0.009$ (Bocquet et al. 2015). The cluster f_{\star} is in good agreement with this estimate of the universal average field value $f_{\star} = (0.95 \pm 0.12)$ per cent at $z = 0$. However, the average field $f_{\star} = 0.22 \pm 0.003$ per cent (see Fig. 6) inferred from the SMF measurements at $z = 0.85$ – 1.2 (vdB13) is significantly lower than the cluster f_{\star} . The cluster or group f_{\star} may be altered over time through either the accretion of lower mass clusters or groups (higher f_{\star}) or through infall from the field (lower f_{\star}). Presumably, these influences must combine to produce the transformation in f_{\star} from a lower mass cluster at $z = 1$ to a higher mass cluster at $z = 0$. We return to this discussion in Section 6.

We compare the high-redshift SPT results to two other samples at high redshift: vdB14 and H13. The virial masses for the majority of the vdB14 systems are below $3 \times 10^{14} M_{\odot}$ and therefore lower mass than our SPT clusters. The one remaining system in this mass range falls near the bottom of our distribution of f_{\star} . The H13 sample shows stellar mass fractions that are in good agreement with ours. We have three clusters in common; combining these measurements we determine that the differences are 1.11σ , 0.69σ and 0.52σ for M_{\star} , M_{500} and f_{\star} measurements, respectively. vdB14 express concern that the M_{\star} estimated by the mass-to-light technique in H13 could possibly be overestimated by as much as a factor of 2. While the largest difference with our sample is indeed with M_{\star} , the level of agreement between the H13 results and our SED fitting results would suggest that the bias is likely smaller (about 10 ± 23 per cent).

5.3 ICM mass fraction f_{ICM}

The majority of the baryonic mass within clusters lies within the hot ICM. The arithmetic mean of f_{ICM} for the 14 SPT clusters is 0.1027 ± 0.0073 , and the characteristic value at $z = 0.9$ and $M_{500} = 6 \times 10^{14} M_{\odot}$ is 0.096 ± 0.005 . A clear mass trend ($f_{\text{ICM}} \propto M_{500}^{0.43 \pm 0.13}$), significant at the 3.3σ level, is seen for SPT clusters. This trend is steeper than (but statistically consistent with) the trends, $\beta_{\text{ICM}} = 0.30 \pm 0.07$ and 0.26 ± 0.03 , presented by Zha11 and GZ13, but steeper at 2.1σ than the result $\beta_{\text{ICM}} = 0.15 \pm 0.03$




Figure 6. The redshift trends of f_* , f_{ICM} , f_c and f_b for the combined cluster sample. The colour coding of the points and the red bands are the same as in Fig. 5. For f_* and f_c , we show the universal value at $z = 0$ (red) and at $z = 0.85\text{--}1.2$ (blue). Measurements have been corrected using the best-fitting mass trends to the mass $6 \times 10^{14} M_\odot$, and the best-fitting redshift trend is shown in green (Table 3).

presented by Andreon (2010). The combined sample has a preferred mass trend $\beta_{\text{ICM}} = 0.22 \pm 0.06$, which is 1.5σ shallower than the SPT sample.

These results, extending to redshift $z = 1.32$, show the clear tendency for f_{ICM} to be suppressed in lower mass clusters – first shown in studies of individual low-redshift clusters (David et al. 1993) and later with a uniform analysis of a large sample of low-redshift clusters (Mohr et al. 1999).

The characteristic value of the combined sample at $z_{\text{piv}} = 0$ and $M_{\text{piv}} = 6 \times 10^{14} M_\odot$ is $f_{\text{ICM}} = 0.112 \pm 0.0032$ (statistical only), which is higher than the $z = 0.9$ SPT sample $f_{\text{ICM}} = 0.096 \pm 0.005$. However, neither the SPT high-redshift sample nor the combined sample exhibits evidence for redshift variation in f_{ICM} with $\gamma_{\text{ICM}} = -0.20 \pm 0.11 \pm 0.22$. The impact of the halo mass 15 per cent systematic uncertainty on the redshift trend introduces an additional systematic uncertainty of $\sigma_{\gamma_{\text{ICM}}} = 0.22$ which is larger than the statistical uncertainty. This underscores the importance of using a homogeneous sample with consistently derived virial mass measurements.

5.4 Collapsed baryon fraction f_c

The collapsed baryon fraction f_c is the fraction of baryonic mass that has cooled to form stars that lie in galaxies and thereby reflects

an integral of the star formation efficiency within the halo averaged over cosmic time (e.g. David & Blumenthal 1992). As already noted, we make no attempt to include an estimate of the ICL contribution here. The arithmetic mean f_c of our 14 clusters is 0.099 ± 0.007 , and the characteristic value at $z = 0.9$ and $M_{500} = 6 \times 10^{14} M_\odot$ is 0.107 ± 0.011 (statistical). The SPT sample exhibits no evidence for either a mass or redshift trend. The combined sample exhibits a 6.5σ significant mass trend $f_c \propto M^{-0.65 \pm 0.10}$ with the collapsed baryon fraction falling to lower values in high-mass clusters and a characteristic value of 0.0859 ± 0.0049 (statistical) at $z = 0$. The redshift trend from the combined sample $f_c \propto (1+z)^{0.39 \pm 0.15 \pm 0.16}$ is significant at 1.8σ if we add the statistical and systematic uncertainties in quadrature.

Note that in the case of f_c , the 15 per cent systematic uncertainty in M_{500} has no impact on the f_c measurement, because a shift in R_{500} has approximately the same impact on M_* and M_{ICM} . However, because of the steep mass trend for f_c ($\beta_c \approx -0.6$), a shift in the virial mass of the high-redshift sample impacts the best-fitting redshift trend, because this shift masquerades as a shift in f_c of $\delta f_c \sim -0.6\delta M_{500}/M_{500}$. This impacts the estimate of the systematic uncertainty in γ_c . This process is also at work for the other fractions, but because their mass dependences are weaker, the impact is smaller.

In Fig. 5, we see that f_c decreases with cluster mass, and the scatter about this trend (especially in the Zha11 sample) is less than in the case of f_* . This measure is interesting because the M_{500} measurements come in only through defining the virial radius, and if the radial variation in f_{ICM} and f_* are mild, then f_c has only a weak dependence on the virial mass estimates. Thus, in the cases where the M_{500} estimates exhibit large uncertainties, the f_c can be an effective way of exploring trends in the mix of stars and ICM within clusters.

Our study indicates that over cosmic time the collapsed baryon fraction f_c at fixed cluster halo mass falls. This redshift trend is driven by the slight rise in the ICM mass fraction f_{ICM} and slight fall in stellar mass fraction f_* .

5.5 Baryon fraction f_b

The arithmetic mean of the baryon fraction for our SPT clusters is 0.114 ± 0.008 (statistical only), and the characteristic value at $z = 0.9$ and $M_{500} = 6 \times 10^{14} M_\odot$ is 0.107 ± 0.006 (statistical). This is lower than the characteristic values of the combined sample at $z = 0$ of 0.1227 ± 0.0035 (statistical). However, neither the SPT sample nor the combined sample ($f_b \propto (1+z)^{-0.17 \pm 0.11 \pm 0.22}$) provides clear evidence for a redshift trend. The mass systematics between the low- and high-redshift samples introduce an uncertainty in the redshift trend parameter of $\sigma_{\gamma_b} = 0.22$, which is larger than the statistical uncertainty, implying that controlling mass systematics among the different samples is crucial. The SPT sample exhibits a mass trend $f_b \propto M_{500}^{0.39 \pm 0.13}$ that has 3σ significance. The combined sample exhibits a mass trend $f_b \propto M_{500}^{0.22 \pm 0.06}$, which is somewhat shallower and is significant at 3.6σ .

6 DISCUSSION

Because our sample includes the highest redshift massive ($M_{500} > 3 \times 10^{14} M_\odot$) clusters studied to date, our analysis is useful for constraining the redshift variation of the ICM and stellar mass components on cluster mass scales. While we do consider intrinsic scatter in fitting the observed properties within our sample, the sample does not provide meaningful constraints on this scatter; thus, our

results shed no light on assembly bias, which would link the baryon properties of individual clusters to the properties of the large-scale environment within which they formed. A joint analysis of the SPT sample and a comparison sample indicates that the cluster collapsed baryon fraction (accounting only for stars in galaxies) within R_{500} is decreasing from 10.7 to 8.6 per cent on the characteristic mass scale $M_{\text{div}} = 6 \times 10^{14} M_{\odot}$ since $z \approx 0.9$; the redshift trend is significant at the 1.8σ confidence level when accounting for a 15 per cent virial mass systematic uncertainty between the literature and SPT samples. Moreover, our analysis indicates that this change is driven by a weak increase in the ICM fraction (f_{ICM} changes from 9.6 to 11.2 per cent) and a weak decrease in the stellar fraction from 1.1 to 1.0 per cent over the same redshift range. The same trends in f_{ICM} and f_{\star} lead to a weak trend in the baryon fraction (from 10.7 to 12.3 per cent) from $z = 0.9$ to the present, a change that is only 0.7σ significant given the systematic mass uncertainties between the high-redshift and local comparison samples.

To build a physical picture, it is important to take note of the mass trends in the stellar mass fraction $f_{\star} \propto M^{-0.37 \pm 0.09}$ that indicate that high-mass clusters have f_{\star} values that lie below those of groups and that are comparable to or even higher than the field f_{\star} at $z = 0$ (see also L03 and vdB13). The ICM mass fractions f_{ICM} behave oppositely $f_{\text{ICM}} \propto M^{0.22 \pm 0.06}$ (see also Mohr et al. 1999), with groups having lower values than massive clusters, whose f_{ICM} values are approaching but still lower than the universal baryon fraction. These mass trends then give rise to the trend of falling collapsed baryon fraction f_c with mass $f_c \propto M^{-0.65 \pm 0.10}$.

Because of the clear mass trends and weak redshift trends in f_{\star} and f_{ICM} , a simple merging scenario for halo formation, where the accretion of low-mass (group-scale) haloes is dominating the mass assembly of massive clusters, does not provide an adequate explanation of the observations. In general, such a scenario would lead to f_{\star} that is approximately independent of cluster halo mass (Balogh et al. 2008). The massive haloes of today form from haloes that were lower mass at higher redshift, so if these low-mass subclusters had lower f_{\star} or higher f_{ICM} at higher redshift, then the simple merger scenario could in principle be consistent with the data. However, the weak redshift variation in these fractions at fixed halo mass that we estimate here for massive haloes does not help to resolve the situation, because it indicates that f_{\star} and f_{ICM} at fixed halo mass have changed only weakly over time; if trends on the massive end are coupled with similar trends on the lower mass end, then the simple merging scenario must be flawed. The conclusion that infall from the field and/or the inclusion of stripping processes that modify the apparent stellar fraction during the process of the growth of massive, cluster-scale haloes is inescapable.

Infall from the field likely plays a critical role in the growth of massive clusters. Studies of the standard hierarchical structure formation scenario on the mass scales of interest here indicate that ~ 40 per cent of the cluster galaxies have previously been in lower mass group or cluster haloes (McGee et al. 2009) and that the rest infall from the field. In the case of f_{\star} , we have shown that the field has lower f_{\star} in comparison to massive clusters at redshift $z = 0.9$ (Section 4.2) and that it has comparable values of f_{\star} at $z = 0$ (see Fig. 6). Through an appropriate mix of field and group accretion, the f_{\star} values in massive clusters could in principle either increase or decrease with cosmic time. Our results indicate that this mix of field and group accretion to build up the haloes of the most massive clusters must produce haloes with f_{\star} that are similar (at ~ 10 per cent level) up to (or weakly decreasing since) redshift $z \sim 1$.

Add to this the likely stripping of stellar material from infalling galaxies during the accretion and relaxation process, and one has an

additional mechanism to reduce the observed f_{\star} over cosmic time, because the ICL from these stripped stars is not included in the f_{\star} measurements here. Lin & Mohr (2004) suggested just such a mechanism to reconcile the falling f_{\star} with halo mass they observed in the local Universe. They presented a toy model that suggested such a mechanism would have to lead to an ICL mass fraction that increases with halo mass and reaches high values of ≈ 40 per cent of the stars in the central galaxy of the clusters. Neither this trend nor ICL fractions at this high redshift have been observed in recent observational studies (Zibetti et al. 2005; GZ13). Presumably, as massive clusters grow, a reduction of f_{\star} through both accretion from the field and stripping of stars from cluster galaxies is counterbalancing the increase of f_{\star} due to accretion of lower mass subclusters. Together these processes must transform high- f_{\star} , low-mass clusters into lower f_{\star} , high-mass clusters. Moreover, these processes must maintain a roughly constant f_{\star} at fixed halo mass over cosmic time on cluster mass scales.

A similar scenario of infall from the field and accretion of subclusters could explain the trends in f_{ICM} as a function of halo mass and redshift. In the case of f_{ICM} , the field value, which is inferred by the *Planck* measurement, is higher than that in the clusters at $z = 0.9$ and remains so to $z = 0$ (see Fig. 6). Thus, given our observed weakly increasing cluster f_{ICM} at fixed halo mass since $z \sim 1$, the increases in f_{ICM} during cluster growth from infall from the field are compensating for the decreases in f_{ICM} from accretion of subclusters. These constraints, when coupled to a detailed hydrodynamical study, would presumably enable one to constrain processes such as early preheating as well as entropy injection from AGN residing in groups and clusters.

7 CONCLUSIONS

In this work, we study the SMF and baryon composition of 14 high-redshift SZE selected clusters between redshifts 0.572 and 1.32 that have a median mass M_{500} of $6 \times 10^{14} M_{\odot}$. We estimate f_{\star} , f_{ICM} , f_c and f_b within R_{500} (Table B1). Our sample provides the highest redshift, uniformly selected sample to date for the study of the baryon content in massive clusters; our measurements together with low-redshift measurements in the literature enable us to constrain the redshift variation of these quantities. We summarize our results here.

(i) We examine the $M_{\star}^{\text{BCG}}-M_{500}$ relation by combining our sample with the sample of H13 and vdB14 (Section 4.1, equation 4). On the cluster mass scale of $6 \times 10^{14} M_{\odot}$, the BCG stellar mass constitutes 0.12 ± 0.01 per cent of the halo mass. That fraction falls with cluster mass as $M_{500}^{-0.58 \pm 0.07}$. BCG stellar masses scatter about the best-fitting $M_{\star}^{\text{BCG}}-M_{500}$ relation with a characteristic value of 41 per cent, a measure of the considerable variation in the BCG population.

(ii) We measure the stacked SMF of these clusters and fit it to a Schechter function (Table 2; Section 4.2). The characteristic mass is $M_0 = 10^{11.0 \pm 0.1} M_{\odot}$, consistent with values derived in low-mass clusters at high redshift (vdB14) and at low redshift (Vulcani et al. 2013). Moreover, through comparison to constraints on the field SMF in the same redshift range (vdB13), we show that the number of galaxies with stellar mass above our threshold ($2.5 \times 10^{10} M_{\odot}$) per unit total mass is higher in clusters than in the field by a factor of 1.65 ± 0.20 .

We take the measurements of the baryon composition in each of our clusters and fit to power-law relations in redshift and mass (equation 11). We present best-fitting trends for the SPT sample and for a

combined sample that includes several samples from the literature (Table 3). In combining with external samples, we homogenize the stellar mass measurements to the Chabrier IMF (Section 2.2.1), we apply corrections for the differences in the virial mass estimates (Section 2.2.2), we adopt a 15 per cent (1σ) systematic virial mass uncertainty (Section 5.1.1) and we account for differences in the estimates of measurement uncertainties by solving for independent intrinsic scatter estimates for each subsample (Section 5.1.2). The key results are described below.

(iii) The stellar mass fraction has a characteristic value 1.1 ± 0.1 per cent (statistical) for clusters with mass $M_{500} = 6 \times 10^{14} M_{\odot}$ at $z = 0.9$ and 1.0 ± 0.05 per cent (statistical) at $z = 0$. It falls with cluster halo mass $f_{\star} \propto M^{-0.37 \pm 0.09}$ and mildly decreases with cosmic time $f_{\star} \propto (1+z)^{0.26 \pm 0.16 \pm 0.08}$ with 1.45σ significance, where the second component of the uncertainty represents the impact of the 15 per cent systematic mass uncertainty between the low- and high-redshift samples. A similar result for the mass trend $\beta_{\star} \approx -0.26$ is also seen for low-mass clusters and groups at $0.8 \leq z \leq 1.0$ (Balogh et al. 2014). The mass trend and mild redshift trend indicate that the infall from subclusters (which would tend to increase f_{\star}) and infall from the field and stripping of stars from cluster galaxies (which would both tend to decrease the observed f_{\star}) must combine to enable the transformation of f_{\star} from low-mass clusters into that of higher mass clusters having similar f_{\star} over the redshift range $0 < z < 1$. Numerical simulations suggest that approximately 40 per cent of cluster galaxies have been accreted as members of subclusters, and the remainder from the field (McGee et al. 2009), but additional study is warranted to test whether the observed trends in f_{\star} (now constrained both as a function of mass and of redshift) can be reproduced by current structure formation scenarios.

(iv) The ICM mass fraction has a characteristic value in clusters with mass $M_{500} = 6 \times 10^{14} M_{\odot}$ of 9.6 ± 0.5 per cent (statistical) at $z = 0.9$ and 11.2 ± 0.32 per cent (statistical) at $z = 0$. It rises with cluster halo mass $f_{\text{ICM}} \propto M^{0.22 \pm 0.06}$ and evolves weakly with redshift at fixed halo mass as $f_{\text{ICM}} \propto (1+z)^{-0.20 \pm 0.11 \pm 0.22}$, where the 0.22 is due to the 15 per cent systematic mass uncertainty between the low- and high-redshift samples. The trend of increasing f_{ICM} with mass has been previously observed (Mohr et al. 1999) and can be explained through entropy injection through early preheating or from cluster AGN. A weakly varying f_{ICM} with cosmic time could be explained by infall from the field, where f_{ICM} is larger than that in clusters at $z = 0.9$ and 0 (see Fig. 6). Hydrodynamical studies of this scenario are needed.

(v) The collapsed baryon fraction determines the fraction of the baryonic component that has cooled to form stars. It is the ratio of the stellar mass to the ICM plus stellar mass. The characteristic value at cluster masses $M_{500} = 6 \times 10^{14} M_{\odot}$ is 10.7 ± 0.1 per cent (statistical) at $z = 0.9$ and 8.6 ± 0.5 per cent (statistical) at $z = 0$. It falls with halo mass as $f_c \propto M^{-0.65 \pm 0.10}$, indicating with 6.5σ significance that a smaller fraction of halo baryons is in the form of stars in the most massive haloes. The redshift trend is $f_c \propto (1+z)^{0.39 \pm 0.15 \pm 0.16}$, where the second uncertainty is due to the 15 per cent systematic mass uncertainty between the low- and high-redshift samples. Thus, there is $\approx 1.8\sigma$ evidence that the collapsed baryon fraction is falling with cosmic time, and this is driven by the weak trends of rising f_{ICM} and falling f_{\star} presented above.

(vi) The baryon fraction f_b is the fraction of the halo mass that is in ICM and stars. The characteristic value at cluster mass $M_{500} = 6 \times 10^{14} M_{\odot}$ is 10.7 ± 0.6 per cent (statistical) at $z = 0.9$ and 12.3 ± 0.4 per cent (statistical) at $z = 0$. It rises with halo mass as $f_b \propto M^{0.22 \pm 0.06}$, and this 3.7σ mass trend is affected both

by the increase in f_{ICM} and the decrease in f_{\star} with cluster mass. The evidence for redshift variation at fixed halo mass is weak $f_b \propto (1+z)^{-0.17 \pm 0.11 \pm 0.22}$, where the second uncertainty is due to the 15 per cent systematic mass uncertainty between the low- and high-redshift samples. If the two uncertainties are added in quadrature, there is no significant evidence (0.7σ) that the baryon fraction is evolving.

As already discussed in Section 6, these mass and redshift trends in baryon quantities are not consistent with a simple hierarchical structure formation merger model where massive clusters form solely through the accretion of lower mass clusters and groups. Significant accretion of galaxies and ICM from the field must also occur, and this accretion together with infall of subclusters can likely explain the weak variation (at fixed cluster halo mass) in f_{\star} and in f_{ICM} over cosmic time. Additionally, the loss of stellar mass from galaxies through stripping is an additional mechanism that would allow for f_{\star} to fall as low-mass clusters grow to higher mass.

This analysis of the first homogeneously selected high-mass cluster sample extending to high redshift allows for interesting initial constraints on the redshift trends in the baryon content; however, these trends are dependent to some extent on the adopted systematic virial mass uncertainty between the low- and high-redshift samples. If the systematic virial mass uncertainty is 15 per cent, there are no statistically significant redshift trends. Higher virial mass systematic uncertainties would further reduce the significance of trends. A reduction of the 15 per cent systematic virial mass uncertainty to 10 or 5 per cent would result in a fractional reduction (to 2/3 or 1/3, respectively) for the redshift trend systematic uncertainties $\gamma_{\text{obs}}^{\text{sys}}$. In the case of a 5 per cent systematic virial mass uncertainty, the significance of the redshift trends for f_{\star} (f_{ICM} , f_c , f_b) would increase from 1.12σ (0.81σ , 1.6σ , 0.69σ) to 1.27σ (1.53σ , 2.2σ , 1.28σ). It is clear that what is needed is a systematic study of a large, homogeneously selected cluster sample with high-quality mass estimates that spans a broad redshift range.

ACKNOWLEDGEMENTS

We acknowledge the support by the DFG Cluster of Excellence ‘Origin and Structure of the Universe’ and the Transregio programme TR33 ‘The Dark Universe’. The calculations have been carried out on the computing facilities of the Computational Center for Particle and Astrophysics (C2PAP) and of the Leibniz Supercomputer Center (LRZ). BB is supported by the Fermi Research Alliance, LLC under Contract No. De-AC02-07CH11359 with the United States Department of Energy. BS acknowledges the support of the NSF grants at Harvard and SAO (AST-1009012, AST-1009649 and MRI-0723073). TS acknowledges the support from the German Federal Ministry of Economics and Technology (BMWi) provided through DLR under project 50 OR 1210. The South Pole Telescope is supported by the National Science Foundation through grant ANT-0638937. Partial support is also provided by the NSF Physics Frontier Center grant PHY-0114422 to the Kavli Institute of Cosmological Physics at the University of Chicago, the Kavli Foundation and the Gordon and Betty Moore Foundation.

Optical imaging data from the VLT programmes 088.A-0889 and 089.A-0824, *HST* imaging data from programmes C18-12246 and C19-12447, and *Spitzer Space Telescope* imaging from programmes 60099, 70053 and 80012 enable the SED fitting in this analysis. X-ray data obtained with *Chandra* X-ray Observatory programmes and *XMM-Newton* Observatory programme 067501 enable the ICM mass measurements. The SPT survey programme SPT-SZ enabled

the discovery of these high-redshift clusters and subsequent analyses have enabled virial mass estimates of these systems. Optical spectroscopic data from VLT programmes 086.A-0741 and 286.A-5021 and Gemini programme GS-2009B-Q-16, GS-2011A-C-3 and GS-2011B-C-6 were included in this work. Additional spectroscopic data were obtained with the 6.5 m Magellan Telescopes.

Facilities: South Pole Telescope, *Spitzer/IRAC*, VLT: Antu (FORIS2), *HST/ACS*, *Chandra*, *XMM-Newton*, Magellan

REFERENCES

- Andersson K. et al., 2011, *ApJ*, 738, 48
 Andreon S., 2010, *MNRAS*, 407, 263
 Applegate D. E. et al., 2014, *MNRAS*, 439, 48
 Arnaud M., Evrard A. E., 1999, *MNRAS*, 305, 631
 Arnouts S., Cristiani S., Moscardini L., Matarrese S., Lucchin F., Fontana A., Giallongo E., 1999, *MNRAS*, 310, 540
 Ashby M. L. N. et al., 2009, *ApJ*, 701, 428
 Balogh M. L., McCarthy I. G., Bower R. G., Eke V. R., 2008, *MNRAS*, 385, 1003
 Balogh M. L. et al., 2014, *MNRAS*, 443, 2679
 Bayliss M. B. et al., 2014, *ApJ*, 794, 12
 Benson B. A. et al., 2013, *ApJ*, 763, 147
 Bertin E., Arnouts S., 1996, *A&AS*, 117, 393
 Bleem L. E. et al., 2015, *ApJS*, 216, 27
 Bocquet S. et al., 2015, *ApJ*, 799, 214
 Bruzual G., Charlot S., 2003, *MNRAS*, 344, 1000
 Calzetti D., Armus L., Bohlin R. C., Kinney A. L., Koornneef J., Storchi-Bergmann T., 2000, *ApJ*, 533, 682
 Capak P. et al., 2007, *ApJS*, 172, 99
 Cappellari M. et al., 2006, *MNRAS*, 366, 1126
 Carlstrom J. E. et al., 2011, *PASP*, 123, 568
 Cavaliere A., Fusco-Femiano R., 1978, *A&A*, 70, 677
 Cavaliere A., Menci N., Tozzi P., 1999, *MNRAS*, 308, 599
 Chabrier G., 2003, *PASP*, 115, 763
 Chiu I.-N. T., Molnar S. M., 2012, *ApJ*, 756, 1
 David L. P., Blumenthal G. R., 1992, *ApJ*, 389, 510
 David L. P., Slyz A., Jones C., Forman W., Vrtilik S. D., Arnaud K. A., 1993, *ApJ*, 412, 479
 Desai S. et al., 2012, *ApJ*, 757, 83
 Djorgovski S., Davis M., 1987, *ApJ*, 313, 59
 Donahue M. et al., 2014, *ApJ*, 794, 136
 Dressler A., Shectman S. A., 1988, *AJ*, 95, 985
 Erben T. et al., 2005, *Astron. Nachr.*, 326, 432
 Fabian A. C., 1994, *ARA&A*, 32, 277
 Fazio G. G. et al., 2004, *ApJS*, 154, 10
 Foley R. J. et al., 2011, *ApJ*, 731, 86
 Forman W., Jones C., 1982, *ARA&A*, 20, 547
 Fowler J. W. et al., 2007, *Appl. Opt.*, 46, 3444
 Frenk C. S., White S. D. M., Efstathiou G., Davis M., 1990, *ApJ*, 351, 10
 Geller M. J., Beers T. C., 1982, *PASP*, 94, 421
 Giodini S. et al., 2009, *ApJ*, 703, 982
 Gonzalez A. H., Zaritsky D., Zabludoff A. I., 2007, *ApJ*, 666, 147
 Gonzalez A. H., Sivanandam S., Zabludoff A. I., Zaritsky D., 2013, *ApJ*, 778, 14 (GZ13)
 Haiman Z., Mohr J. J., Holder G. P., 2001, *ApJ*, 553, 545
 Henry J. P., Arnaud K. A., 1991, *ApJ*, 372, 410
 High F. W., Stubbs C. W., Rest A., Stalder B., Challis P., 2009, *AJ*, 138, 110
 Hilton M. et al., 2013, *MNRAS*, 435, 3469 (H13)
 Ilbert O. et al., 2006, *A&A*, 457, 841
 Ilbert O. et al., 2009, *ApJ*, 690, 1236
 Jee M. J., Hughes J. P., Menanteau F., Sifón C., Mandelbaum R., Barrientos L. F., Infante L., Ng K. Y., 2014, *ApJ*, 785, 20
 Kauffmann G. et al., 2003, *MNRAS*, 341, 33
 Kochanek C. S. et al., 2001, *ApJ*, 560, 566
 Kroupa P., 2001, *MNRAS*, 322, 231
 Lidman C. et al., 2012, *MNRAS*, 427, 550
 Lilje P. B., 1992, *ApJ*, 386, L33
 Lin Y.-T., Mohr J. J., 2004, *ApJ*, 617, 879
 Lin Y., Mohr J. J., Stanford S. A., 2003, *ApJ*, 591, 749 (L03)
 Lin Y., Mohr J. J., Stanford S. A., 2004, *ApJ*, 610, 745
 Lin Y., Mohr J. J., Gonzalez A. H., Stanford S. A., 2006, *ApJ*, 650, L99
 Lin Y.-T., Stanford S. A., Eisenhardt P. R. M., Vikhlinin A., Maughan B. J., Kravtsov A., 2012, *ApJ*, 745, L3
 Liu J. et al., 2015, *MNRAS*, 449, 3370
 McDonald M. et al., 2013, *ApJ*, 774, 23
 McDonald M. et al., 2014, *ApJ*, 784, 18
 McGee S. L., Balogh M. L., Bower R. G., Font A. S., McCarthy I. G., 2009, *MNRAS*, 400, 937
 Martino R., Mazzotta P., Bourdin H., Smith G. P., Bartalucci I., Marrone D. P., Finoguenov A., Okabe N., 2014, *MNRAS*, 443, 2342
 Menanteau F. et al., 2012, *ApJ*, 748, 7
 Mohr J., Evrard A., 1997, *ApJ*, 491, 38
 Mohr J. J., Evrard A. E., Fabricant D. G., Geller M. J., 1995, *ApJ*, 447, 8
 Mohr J. J., Mathiesen B., Evrard A. E., 1999, *ApJ*, 517, 627
 Molnar S. M., Chiu I.-N., Umetsu K., Chen P., Hearn N., Broadhurst T., Bryan G., Shang C., 2010, *ApJ*, 724, L1
 Perlmutter S. et al., 1999, *ApJ*, 517, 565
 Pickles A. J., 1998, *PASP*, 110, 863
 Planck Collaboration XVI, 2014, *A&A*, 571, A16
 Reichardt C. L. et al., 2013, *ApJ*, 763, 127
 Riess A. G. et al., 1998, *AJ*, 116, 1009
 Ruel J. et al., 2014, *ApJ*, 792, 45
 Salpeter E. E., 1955, *ApJ*, 121, 161
 Sanders D. B. et al., 2007, *ApJS*, 172, 86
 Saro A., Mohr J. J., Bazin G., Dolag K., 2013, *ApJ*, 772, 47
 Schechter P., 1976, *ApJ*, 203, 297
 Schellenberger G., Reiprich T. H., Lovisari L., Nevalainen J., David L., 2015, *A&A*, 575, A30
 Schirmer M., 2013, *ApJS*, 209, 21
 Sifón C. et al., 2013, *ApJ*, 772, 25
 Song J., Mohr J. J., Barkhouse W. A., Warren M. S., Rude C., 2012a, *ApJ*, 747, 58
 Song J. et al., 2012b, *ApJ*, 761, 22
 Stalder B. et al., 2013, *ApJ*, 763, 93
 Staniszewski Z. et al., 2009, *ApJ*, 701, 32
 Sunyaev R. A., Zel'dovich Y. B., 1970, *Comments Astrophys. Space Phys.*, 2, 66
 Sunyaev R. A., Zel'dovich Y. B., 1972, *Comments Astrophys. Space Phys.*, 4, 173
 Tauber J. A., 2000, in Lasenby A., Wilkinson A., eds, *Proc. IAU Symp. 201, New Cosmological Data and the Values of the Fundamental Parameters*. Astron. Soc. Pac., San Francisco, 145
 van der Burg R. F. J. et al., 2013, *A&A*, 557, A15 (vdB13)
 van der Burg R. F. J., Muzzin A., Hoekstra H., Wilson G., Lidman C., Yee H. K. C., 2014, *A&A*, 561, A79 (vdB14)
 Vanderlinde K. et al., 2010, *ApJ*, 722, 1180
 Vikhlinin A., Kravtsov A., Forman W., Jones C., Markevitch M., Murray S. S., Van Speybroeck L., 2006, *ApJ*, 640, 691
 Vikhlinin A. et al., 2009, *ApJ*, 692, 1033
 Vulcani B. et al., 2013, *A&A*, 550, A58
 White S. D. M., Efstathiou G., Frenk C. S., 1993a, *MNRAS*, 262, 1023
 White S., Navarro J., Evrard A., Frenk C., 1993b, *Nature*, 366, 429
 White M., Cohn J. D., Smit R., 2010, *MNRAS*, 408, 1818
 Williamson R. et al., 2011, *ApJ*, 738, 139
 Zenteno A. et al., 2011, *ApJ*, 734, 3
 Zhang Y.-Y., Laganá T. F., Pierini D., Puchwein E., Schneider P., Reiprich T. H., 2011, *A&A*, 535, A78 (Zha11)
 Zhang Y.-Y., Laganá T. F., Pierini D., Puchwein E., Schneider P., Reiprich T. H., 2012, *A&A*, 544, C3
 Zibetti S., White S. D. M., Schneider D. P., Brinkmann J., 2005, *MNRAS*, 358, 949




Figure A1. A test of SED fitting using galaxies with spectroscopic redshifts. The x -axis is the normalized difference of photo- z and spec- z , and the y -axis is the resulting stellar mass difference averaged on a per-cluster basis. The colour code from blue to red indicates the clusters from the lowest to the highest redshift. SPT-CL J0205–5829 at $z = 1.32$ is marked as it has the largest mass difference. The black dot indicates the mean of $(\log_{10} M_{\star}^{\text{photoz}} - \log_{10} M_{\star}^{\text{specz}})$ and $\Delta z/(1+z)$ of the ensemble of clusters.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

(<http://mnras.oxfordjournals.org/lookup/suppl/doi:10.1093/mnras/stv2303/-/DC1>).

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: PERFORMANCE OF SED FITTING

With the published spectroscopic sample for SPT clusters (Sifón et al. 2013; Ruel et al. 2014), we are able to quantify how the uncertainty of the photo- z impacts on the stellar mass estimates based on the SED fit using the six band photometry (b_H , $F606W$, I_B , z_G , $[3.6]$, $[4.5]$). We cross-match our photometry identified sample with the galaxy sample in Ruel et al. (2014) and repeat the whole SED fit analysis with the redshift fixed to the measured spectroscopic redshift. We show the comparison in Fig. A1. The photo- z performance is estimated as the mean $\Delta z/(1+z) \equiv (z_{\text{photo}} - z_{\text{spec}})/(1+z_{\text{spec}})$ to be 0.037 ± 0.0083 . The difference of the stellar mass estimates ($\log_{10} M_{\star}^{\text{photoz}} - \log_{10} M_{\star}^{\text{specz}}$) when using z_{photo} and z_{spec} is at the level of $\lesssim 0.2$ with a mean ≈ 0.03 . Except for the highest redshift cluster (SPT-CL J0205–5829 at $z = 1.32$), which has only 5 spectroscopic redshifts available for the cluster members, the SED fitting using our six-band photometry returns unbiased estimates of the stellar masses.




Figure B1. The magnitude distribution (MD, left) and stellar mass function (SMF, right) for the full population of galaxies in the SPT clusters. We show the cluster + background estimates from within R_{500} (green), the uncorrected local background (grey), the corrected local background (black) and the background estimated from COSMOS (magenta). The SMFs are derived using SED fitting of six-band photometry. The dashed lines indicate the results for individual clusters and the heavy-solid lines are the averages over all clusters. The COSMOS and local, contamination-corrected background estimates are in good agreement. We adopt the COSMOS background correction in this work.

APPENDIX B: TESTS OF STATISTICAL BACKGROUND CORRECTION

To test the COSMOS background, we extract the local background information from our SPT data set, applying a correction for the cluster galaxy contamination. We extract the corrected local background between $1.2R_{500}$ and $2.5R_{500}$ for each cluster. We correct for cluster contamination by assuming that the cluster galaxies are distributed as an NFW model with concentration of $c_{500}^{\text{gal}} = 1.9$ (Lin et al. 2004; Hennig, in preparation), and the SMF and the magnitude distribution (MD) are the same for the region within the cluster R_{500} and for the cluster population that is contaminating the background region. Together with the area extracted for the region within R_{500} and the local background, we solve for the surface number densities of the SMF and MD using the corrected local background for each cluster. The SMF and MD derived using the corrected local background are noisy for each individual cluster, especially for the lower redshift clusters where the area available for the local background is typically less than 5 arcmin^2 . We combine 9 of the 14 independent estimates (those with background area larger than 8 arcmin^2) to create an average local background estimate. In averaging, we use the area weighted average of the individual background estimates so that clusters with greater area (but not necessarily higher number density) receive higher weight.

Fig. B1 contains a comparison of the COSMOS and local background estimates for the SMF (right-hand panels) and MD (left-hand panels). The corrected local background estimates (black) for the SMF and MD are in a good agreement with the COSMOS backgrounds (magenta). There is poorer agreement on the bright (massive) end with the tendency that the local background is slightly higher than COSMOS. The cluster plus background SMF and MDs extracted from within R_{500} (green) show significant overdensities with respect to the background estimates. In both the case of the local background estimates (black) and the cluster plus

Table B1. Measured quantities for the SPT cluster sample. Columns contain the cluster name, spectroscopic redshift, M_{500} estimated from the SZE signature, R_{500} inferred from the given M_{500} and redshift, ICM mass M_{ICM} , the BCG mass M_{BCG}^* , the total stellar mass M_* , the stellar mass fraction f_* , the collapsed baryon fraction f_c , the baryon fraction f_b , the ICM mass fractions f_{ICM} and the stellar-mass-to-light ratios Υ (rms in the parentheses) in the [3.6] band in the observed frame.

Cluster	Redshift	M_{500} ($10^{14} M_{\odot}$)	R_{500} (Mpc)	M_{ICM} ($10^{13} M_{\odot}$)	M_{BCG}^* ($10^{11} M_{\odot}$)	M_* ($10^{12} M_{\odot}$)	f_* (per cent)	f_c (per cent)	f_b (per cent)	f_{ICM} (per cent)	Υ ($\frac{M_{\odot}}{L_{\odot}}$)
SPT-CL J0000–5748	0.702	4.35 ± 1.16	0.90	4.33 ± 0.73	$13.49^{+1.18}_{-1.06}$	6.34 ± 2.22	1.46 ± 0.64	12.76 ± 4.33	11.42 ± 3.51	9.96 ± 3.14	$0.37(0.146)$
SPT-CL J0102–4915	0.870	15.75 ± 3.22	1.30	25.51 ± 0.98	$9.23^{+0.82}_{-0.73}$	10.29 ± 1.92	0.65 ± 0.18	3.88 ± 0.71	16.85 ± 3.50	16.19 ± 3.37	$0.32(0.196)$
SPT-CL J0205–5829	1.320	5.65 ± 1.14	0.78	5.26 ± 1.13	$3.95^{+0.92}_{-1.02}$	5.01 ± 0.96	0.89 ± 0.25	8.70 ± 2.28	10.20 ± 2.87	9.31 ± 2.74	$0.42(0.158)$
SPT-CL J0533–5005	0.881	4.24 ± 1.13	0.83	2.70 ± 0.51	$5.89^{+0.56}_{-0.47}$	2.95 ± 0.98	0.70 ± 0.30	9.85 ± 3.39	7.06 ± 2.24	6.36 ± 2.07	$0.33(0.221)$
SPT-CL J0546–5345	1.067	5.48 ± 1.16	0.85	7.05 ± 1.23	$17.41^{+1.86}_{-4.57}$	14.07 ± 2.19	2.57 ± 0.67	16.63 ± 3.24	15.44 ± 3.99	12.87 ± 3.53	$0.38(0.208)$
SPT-CL J0559–5249	0.609	7.16 ± 1.44	1.11	8.36 ± 0.55	$4.83^{+0.41}_{-0.38}$	8.10 ± 1.64	1.13 ± 0.32	8.83 ± 1.71	12.80 ± 2.70	11.67 ± 2.47	$0.30(0.171)$
SPT-CL J0615–5746	0.972	11.75 ± 2.35	1.13	13.60 ± 2.25	$14.39^{+1.75}_{-4.61}$	14.70 ± 2.45	1.25 ± 0.33	9.75 ± 2.07	12.83 ± 3.21	11.57 ± 3.00	$0.36(0.186)$
SPT-CL J2040–5726	0.930	4.10 ± 0.97	0.81	4.25 ± 0.95	$5.44^{+0.49}_{-0.45}$	3.44 ± 0.93	0.84 ± 0.30	7.50 ± 2.43	11.22 ± 3.53	10.38 ± 3.38	$0.28(0.146)$
SPT-CL J2106–5844	1.132	9.35 ± 1.84	0.99	11.68 ± 1.43	$9.96^{+0.87}_{-0.79}$	14.06 ± 2.01	1.50 ± 0.37	10.75 ± 1.80	13.99 ± 3.16	12.49 ± 2.89	$0.42(0.171)$
SPT-CL J2331–5051	0.576	6.45 ± 1.34	1.08	6.07 ± 0.83	$3.03^{+2.67}_{-0.66}$	2.78 ± 1.31	0.43 ± 0.22	4.38 ± 2.06	9.84 ± 2.42	9.41 ± 2.34	$0.37(0.197)$
SPT-CL J2337–5942	0.775	9.44 ± 1.83	1.14	8.52 ± 0.79	$10.33^{+1.00}_{-0.80}$	11.24 ± 2.14	1.19 ± 0.32	11.66 ± 2.18	10.21 ± 2.16	9.02 ± 1.94	$0.40(0.156)$
SPT-CL J2341–5119	1.003	5.9 ± 1.31	0.92	6.85 ± 1.00	$9.41^{+1.62}_{-1.84}$	7.12 ± 1.60	1.08 ± 0.32	9.42 ± 2.29	11.47 ± 2.75	10.39 ± 2.57	$0.34(0.183)$
SPT-CL J2342–5411	1.075	4.43 ± 1.07	0.79	3.15 ± 0.64	$8.25^{+0.74}_{-0.67}$	5.03 ± 1.17	1.14 ± 0.38	13.77 ± 3.66	8.25 ± 2.47	7.11 ± 2.24	$0.39(0.167)$
SPT-CL J2359–5009	0.775	4.98 ± 1.16	0.92	3.50 ± 0.34	$6.92^{+0.88}_{-0.53}$	4.80 ± 1.31	0.96 ± 0.35	12.06 ± 3.07	8.00 ± 2.00	7.03 ± 1.77	$0.43(0.182)$

background estimates (green), the individual cluster results are shown with dashed lines and the thick solid lines represent the ensemble average.

On the other hand, the corrected local background for the SMF and MD for the red population is generally lower than the COSMOS estimates. This suggests that we are overcorrecting the local backgrounds for cluster contamination in the case of the red population, and this is to be expected given that we do not have the right filter combinations (blue band containing 4000 Å break and one band redwards of the break) for the half of our sample that lies at $z > 0.9$. For these reasons, we do not present any analyses of the red sequence selected subpopulation in this paper.

We compare the differences between the cumulative stellar mass estimates for the full population when using the two different background corrections. We fit a simple linear relation $M_{\text{local}} = 10^x \times M_{\text{COSMOS}}$, allowing the normalization 10^x to float, where M_{local} and M_{COSMOS} are the mass estimations for using the local and COSMOS backgrounds, respectively. The resulting best-fitting x is -0.018 ± 0.005 (0.045 ± 0.012) for the cluster (background) stellar mass estimation. That is, using the COSMOS background results in ~ 4 per cent higher stellar mass estimates for the cluster and ~ 10 per cent lower mass estimates in the background as compared to those using the corrected local background.

¹Department of Physics, Ludwig-Maximilians-Universität, Scheinerstr. 1, D-81679 München, Germany

²Excellence Cluster Universe, Boltzmannstr. 2, D-85748 Garching, Germany

³Max-Planck-Institut für extraterrestrische Physik, Giessenbachstr., D-85748 Garching, Germany

⁴Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

⁵Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁶Department of Physics, Harvard University, 17 Oxford Street, Cambridge, MA 02138, USA

⁷Fermi National Accelerator Laboratory, Batavia, IL 60510-0500, USA

⁸Kavli Institute for Cosmological Physics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637, USA

⁹Department of Astronomy and Astrophysics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637, USA

¹⁰Department of Physics, University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637, USA

¹¹Argonne National Laboratory, 9700 S. Cass Avenue, Argonne, IL 60439, USA

¹²Department of Physics and Astronomy, University of Missouri, 5110 Rockhill Road, Kansas City, MO 64110, USA

¹³Department of Astronomy, University of Florida, Gainesville, FL 32611, USA

¹⁴School of Physics, University of Melbourne, Parkville, VIC 3010, Australia

¹⁵Institute for Astronomy, University of Hawaii at Manoa, Honolulu, HI 96822, USA

¹⁶Department of Physics, University of California, One Shields Avenue, Davis, CA 95616, USA

¹⁷Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

¹⁸Korea Astronomy and Space Science Institute, 776 Daedeokdae-ro, Yuseong-gu, Daejeon 305-348, Republic of Korea

¹⁹Argelander-Institut für Astronomie, Auf dem Hügel 71, D-53121 Bonn, Germany

²⁰Cerro Tololo Inter-American Observatory, Casilla 603, La Serena, Chile

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.