



Publication Year	2019
Acceptance in OA @INAF	2021-01-05T16:20:59Z
Title	High-Resolution X-ray Imaging Studies of Neutron Stars, Pulsar Wind Nebulae and Supernova Remnants
Authors	Safi-Harb, Samar; AMATO, Elena; Gotthelf, Eric V.; Katsuda, Satoru; Sasaki, Manami; et al.
Handle	http://hdl.handle.net/20.500.12386/29499
Journal	BULLETIN OF THE AMERICAN ASTRONOMICAL SOCIETY
Number	51

Astro2020 Science White Paper

High-Resolution X-ray Imaging Studies of Neutron Stars, Pulsar Wind Nebulae and Supernova Remnants

Thematic Areas: Planetary Systems Star and Planet Formation
 Formation and Evolution of Compact Objects Cosmology and Fundamental Physics
 Stars and Stellar Evolution Resolved Stellar Populations and their Environments
 Galaxy Evolution Multi-Messenger Astronomy and Astrophysics

Principal Author:

Name: Samar Safi-Harb

Institution: University of Manitoba (Canada)

Email: samar.safi-harb@umanitoba.ca – Phone: (204) 474-7104

Co-authors: Elena Amato (INAF-Arcetri, Italy), Eric V. Gotthelf (Columbia, USA), Satoru Katsuda (Saitama, Japan), Manami Sasaki (Erlangen-Nürnberg, Germany), Yasunobu Uchiyama, Naomi Tsuji (Rikkyo, Japan), Benson Guest (University of Manitoba, Canada)

Abstract: Supernova remnants serve as nearby laboratories relevant to many areas in Astrophysics, from stellar and galaxy evolution to extreme astrophysics and the formation of the heavy elements in the Universe. The *Chandra* X-ray mission has enabled a giant leap forward in studying both SNRs and their compact stellar remnants on sub-arcsecond scale. However, such high-resolution imaging studies have been mostly limited to the nearby and/or relatively bright objects. There is no question that we are missing a large population, especially in external galaxies. Within our own Galaxy, we are presented with new fundamental questions related to neutron stars' diversity, kicks, relativistic winds and the way these objects interact with, and impact, their host environments. In this white paper, we highlight some of the breakthroughs to be achieved with future X-ray missions (such as the proposed *AXIS* probe) equipped with sub-arcsecond imaging resolution and an order of magnitude improvement in sensitivity.

Endorsed by: Katie Auchettl (University of Copenhagen), Aya Bamba (Univ. of Tokyo), Harsha Blumer (West Virginia Univ.), Niccoló Bucciantini (INAF-Arcetri), Yang Chen (Nanjing Univ.), Roger Chevalier (Univ. of Virginia), Emma de Ona Wilhelmi (IEEC-CSIC and DESY), Gloria Dubner (Universidad de Buenos Aires), Gilles Ferrand (RIKEN), Chris Fryer (LANL), Joseph Gelfand (NYU Abu Dhabi), Oleg Kargaltsev (George Washington Univ.), Noel Klingler (Penn State Univ.), Roland Kothes (NRC Herzberg), Denis Leahy (Univ. of Calgary), Laura Lopez (Ohio State Univ.), Marco Miceli (Palermo), Shigehiro Nagasaki (RIKEN), Barbara Olmi (Università di Firenze), Salvatore Orlando (INAF-Palermo), Sangwook Park (Univ. of Texas at Arlington), George Pavlov (Penn State Univ.), Paul Plucinsky (Harvard-Smithsonian CfA), Adam Rogers (Brandon Univ.), Ashley Ruitter & Ivo Sietenzahl (University of New South Wales), Patrick Slane (CfA, Harvard & Smithsonian), Tuguldur Sukhbold (Ohio State Univ.), Takaaki Tanaka (Kyoto Univ.), Tea Temim (STSci), Diego Torres (IEEC-CSIC, Barcelona), Jacco Vink (Univ. of Amsterdam), Frédéric Vogt (ESO, Chile), Jennifer West (Dunlap Institute), Brian Williams (NASA GSFC), George Younes (George Washington Univ.), Ping Zhou (Univ. of Amsterdam)

1 Motivation

Supernova Remnants (SNRs) are among the most fascinating astrophysical objects in the Universe. They impact the chemical enrichment and evolution of galaxies, accelerate cosmic rays to extremely high energies, and those resulting from core-collapse explosions make the most magnetic and compact objects in the Universe: neutron stars (NSs). NSs are the best laboratories to study extreme physical conditions that can not be achieved even in the most advanced laboratories on Earth, as well as relativistic outflows and jets that are ubiquitous in Astrophysics. These objects have not only driven scientific breakthroughs, technology development and interdisciplinary connections, but they also fascinate the public and young people.

In this white paper, we focus on SNRs and associated isolated NSs. We highlight outstanding science breakthroughs that can be achieved *only with sub-arcsecond resolution and high sensitivity* in the X-ray band – at least comparable to, or better than, *Chandra’s* imaging resolution, and with an order of magnitude improvement in sensitivity. Such capabilities will be met by the proposed probe *AXIS* [23].

We note that the future missions *Athena* and *Lynx* are expected to achieve high-spectral resolution that will significantly benefit SNR science; however the high-resolution spectroscopy aspect is discussed in separate white papers (B. Williams et al.; L. Lopez et al.).

We here specifically aim to address the following questions: (i) How do pulsars’ relativistic winds communicate with, and energize, their surrounding medium?; (ii) How do NSs evolve and what drives their kicks; (iii) How do SNRs impact cosmic magnetism and galaxies’ evolution?

2 Neutron Star Winds: How do they impact their surroundings?




Figure 1: Simulated *AXIS* images of PWNe, using *Chandra* images as input, illustrating the fine structures that can be observed with modest exposure times ranging from 25 ks to 100 ks. The high-resolution over a larger FoV and low background will allow the detection of faint, thin structures out to large distances from the powering pulsar.

Magnetized relativistic plasmas phenomena are ubiquitous among many classes of astrophysical objects. Of these, Pulsar Wind Nebulae (PWNe) [10] are some of the best particle accelerators in the Universe, with efficiencies close to 30%, generating particle with energy up to 1 PeV, the “knee” in the cosmic ray spectrum (cf. Crab Nebula). With its arcsec resolution, *Chandra* opened a new window to resolve torii and jet structures originating from particles accelerated in shocked regions [16]. Some of these features (resolved also in the optical with HST) are observed to move on sub-arcsecond scales with relativistic velocities. The location of

the termination shock, where the ram pressure of the NS’s relativistic wind is balanced by the nebular pressure, often lies $\lesssim 0.1$ pc from the pulsar. Deep, high resolution *AXIS* observations offer the possibility of resolving these shocks and providing new insight on the origin, internal dynamics,

and evolution of PWNe, and the magnetic field in which they are embedded.

Fig. 1 shows examples of PWNe simulated with *AXIS* to illustrate the high-resolution structures that can be imaged from young and evolved nebulae using exposure times 5-10 times shorter than *Chandra*'s. Such high-resolution imaging studies have been mostly limited to the nearby, brightest, or youngest objects; or else very deep exposures are needed for resolving faint structures. *AXIS* will make a leap in probing a much larger sample of objects in our Galaxy and beyond.

PWNe in the Magellanic clouds are rare but will be finally within reach. An *AXIS*-like resolution of $0.3''$ corresponds to 0.1 pc at the LMC distance, exactly the scale of termination shocks in PWNe. Increasing the population of spatially resolved PWNe will shed light on star formation as a function of metallicity, but will also answer open fundamental questions about PWNe, pulsars and high-energy astrophysics processes; some of which are listed here for their broad implications:

- Pulsar (PSR) wind magnetization and anisotropy: These properties of the PSR wind show directly in the appearance of the nebulae, when imaged at high energy with sufficiently high spatial resolution: high speed jets can only appear if the wind is sufficiently magnetized and anisotropic [7]. On the other hand, these properties of the wind tell us about the inner workings of the PSR magnetosphere and about magnetic dissipation in relativistic plasmas, a topic which is relevant for many high energy sources and phenomena.
- Particle acceleration at a transverse relativistic shock: High spatial resolution X-ray imaging and the study of time variability of small-scale features in the inner part of PWNe (X-ray rings and wisps) can constrain where and in what physical conditions particle acceleration occurs [24] and assess whether it is the shock or some other form of dissipation, like magnetic reconnection, that can provide such efficient acceleration. In addition, the process behind particle acceleration bears information on the pair multiplicity of the PSR magnetosphere, namely the number of electron-positron pairs produced by each electron extracted from the star surface. NSs and PWNe are in fact likely to be the primary contributors of the so-called positron excess [1], that in recent years has attracted much attention both in the cosmic rays and dark matter communities.
- Pulsar contribution to leptonic cosmic-rays: In bow-shock PWNe (BSPWNe), after a high-speed PSR has left its parent SNR, the wind is only confined by the ISM ram-pressure so that electrons and positrons are free to leave the system and be released in the ISM in the back of the shock, hence contributing to the cosmic ray flux [2]. Details of the particle release are important to determine the PWN contribution to the above-mentioned positron excess. Mapping the spectral index close to the NS and out to large distances is essential to assess the particle acceleration process and particle aging effects. Furthermore, multi-TeV particles are sometimes released from the head of the system, through very long and thin X-ray bright channels [17, 3]. *AXIS* will be essential to establish the particle release from BSPWNe and properly synthesize the spectrum of cosmic ray leptons.
- Magnetar Wind Nebulae: There is growing evidence of compact and faint PWNe associated with magnetars, or magnetars-in disguise, whose powering mechanism remains a puzzle. Their analysis is complicated by their faintness, compactness and contamination from a dust scattering halo [26], [31], [5]. It is not clear if their X-ray emission is powered by rotation, magnetism or both (e.g., [27]). *AXIS* will play an important role in studying the nature and origin of nebulae around systems that display a magnetar like activity, thus also requiring a rapid response to bursting sources.

3 Neutron Stars Diversity: Nature or Nurture?

The combination of high spatial resolution, low background and high effective area is ideal for detecting a new generation of fainter young pulsars (PSRs) in SNRs. This can help address important questions on the birth and evolution of PSRs, and explain their seemingly diverse properties.

PSRs were originally detected by their radio emission (e.g., Crab, Vela) but X-ray observations over the past two decades have discovered young, isolated NSs with spectral and timing properties markedly different from those of the typical rotation-powered radio PSRs. The radio-quiet PSRs are best described by their implied magnetic field that range from $10^{10} - 10^{14}$ G. The largest of these are associated with the magnetars, slow rotators ($\sim 2-12$ s), that display a variety of temporal phenomena, such as short and long scale transient outbursts, random episodes of short (~ 1 s) burst of hard X-rays, and erratic spin-down. Most notably, their X-ray emission far exceeds that predicted for a rotation-powered PSR, based on their spin-down luminosity. X-ray emission of magnetars is believed to arise from magnetic losses from a strongly magnetized ($B > 4.4 \times 10^{13}$ G) isolated NS [8]. However, the detection of magnetar-like activity from seemingly classical rotation-powered PSRs [11, 21, 13] complicates this picture. Similarly, pulsations detected from the central compact objects (CCOs), PSRs in SNRs with extremely small magnetic dipole fields [12], are a puzzle since the only mechanism thought to be capable of creating a non-uniform surface temperature is anisotropic heat conduction in a strong magnetic field.

The latter problem can be addressed with *AXIS* spectral-temporal observations of young NSs to model their surface emission using phase-resolved spectroscopy. This will allow a better understanding of their magnetic field configurations. A leading explanation for the generation of the observed hot spots on CCOs requires crustal toroidal fields to insulate the magnetic equator from heat conduction. This toroidal component is expected to be generated by differential rotation in the proto-neutron star dynamo. To have a significant effect on the heat transport, the crustal toroidal field required in all models is $> 10^{14}$ G, far greater than the poloidal field, if the latter is measured by the spin-down. Do all CCOs harbor an inner magnetar buried in their crust without (currently) contributing to its external dipole responsible for its slow spin-down? Most importantly, how do the CCOs and the magnetars relate to each other and to the classical rotation-powered radio PSRs?

The proposed very high spatial resolution of *AXIS* coupled with its high time resolution CCD imaging modes can provide breakthrough science for faint CCOs in SNRs. Of great interest is detecting pulsations from the 300 year-old CCO in SNR Cassiopeia A, the long-sought compact object discovered with the first light *Chandra* observation [25]. This will provide the critical energetics and magnetic field estimates for a PSR close to its birth values. Alternatively, a strong upper limit on any pulsations would help advance theories of atmospheric physics of NSs. Lastly, the order of magnitude increase in sensitivity will be needed to discover the missing CCO descendants, and in turn address the population of core-collapse SNRs in our Galaxy.

4 Neutron Star Velocities: What drives their kicks?

The origin of high velocities in NSs is a long-standing mystery in astrophysics. There are two main competing mechanisms to kick NSs: (a) anisotropic ejection of the stellar debris (“hydrodynamic kick”, [15]) and (b) asymmetric-neutrino emission (“neutrino-induced kick”, [4]). Fortunately, the two scenarios predict a clear difference in NS kick velocities and SN asymmetries. The hydrodynamic kick mechanism predicts that NS velocities are directed opposite to the stronger explosion

where explosive nucleosynthesis elements from Si to Fe are preferentially expelled, whereas the neutrino-induced kick mechanism either suggest no correlation between NS velocities and SN asymmetries, or predict the strongest mass ejection in the direction of NS motion.

Recent X-ray observations of SNRs revealed that NSs preferentially move opposite to the bulk of either X-ray emission [14] or intermediate-mass elements [18], supporting the hydrodynamic kick scenario. However, in many cases, NS velocities are indirectly inferred from displacements between NS positions and geometric centers. The number of robust samples is still quite small. An instrument like *AXIS* will significantly increase the observational sample, as described below. To distinguish between the two NS kick scenarios, it is critically important to measure both NS proper motions and detailed distributions of SN ejecta. NS proper motions are important not only because they allow us to estimate NS velocities, but also because they help to check if a NS is really associated with a SNR and to infer an explosion site by tracing back the proper motion. The long time baseline, which will be available with the combination of *Chandra* and *AXIS*, will be the key to reduce systematic uncertainties on NS proper motions. In addition, *AXIS*'s superior throughput and wide field of view, together with its moderate spectral resolution, will allow us to map detailed SN ejecta distributions for faint and large SNRs.

So far, we have only three SNRs (Puppis A, Cas A, G292.0+1.8) for which both NS kick velocities and ejecta distributions are robustly estimated. All three systems show an anti-correlation between NS kick velocities and ejecta distributions, favoring the hydrodynamic origin for the NS kicks. *AXIS* will increase the number of such samples substantially, and will reveal if the hydrodynamic kick scenario is the only process that can accelerate the NS or if other mechanisms, such as the neutrino-induced kick scenario, can play a role. Increasing the samples is also important to search for correlations between the degrees of explosion asymmetries and NS kick velocities, and between the NS surface magnetic fields and NS kick velocities.

5 SNR shock impact on cosmic magnetism




Figure 2: Simulated radial profiles of shocked emission from a young, 20 pc diameter SNR, at a distance of 1 kpc. The 100 ks *AXIS* simulations, for an amplified magnetic fields of $100\mu\text{G}$ and 1 mG, illustrate that the shock narrows in width for stronger fields.

variability of the synchrotron X-rays can be seen, which is another evidence in favor of MFA [28].

Turbulent magnetic fields at young SNR shocks are expected to be significantly amplified by a cosmic-ray current driven instability that develops in the shock precursor. Magnetic field amplification (MFA) is thought as the key element in non-linear Diffusive Shock Acceleration theory [6]. X-ray observations with *Chandra* have revealed the presence of narrow synchrotron X-ray filaments at the outer edge of young SNRs, demonstrating that the strong shocks at young SNRs are indeed capable of amplifying the interstellar magnetic field by large factors [29]. The narrowness of synchrotron X-ray filaments could be due to rapid synchrotron cooling of high-energy electrons in the postshock flow if the magnetic field reaches ~ 0.1 mG. In some cases, time-

The turbulent magnetic fields, likely amplified by CR current driven instabilities, can be imprinted in the spatial structures of the synchrotron X-ray filaments. Testing the theoretical predictions is best achieved with high-angular resolution measurements of the energy dependence on the width of the filamentary structures (see e.g., [32]). This is illustrated in Fig. 2 showing the radial profiles of the synchrotron X-ray filaments expected to be observed by *AXIS*.

6 Population Studies: From our Galaxy to the Nearby Universe

SNRs radiate copiously at energies of 0.5 – 2 keV, a range that is difficult to study globally in the Milky Way because of absorption by matter in the Galactic disk, with an absorbing column density that amounts to a few 10^{21} cm^{-2} within less than a kpc. In order to study the population of SNRs in a galaxy as a whole we have to look beyond the Milky Way.

By studying SNRs in nearby galaxies, we particularly want to address following questions:

- What is the fraction and spatial distribution of core-collapse SNRs vs. type Ia SNRs? These can be identified based on their morphology combined with spectral properties.
- What is the X-ray luminosity function (XLF) of SNRs? How are the XLFs of different galaxies related to the underlying stellar population, ISM, metallicity and SNR evolution?
- What is the distribution of SNRs in comparison to that of the cold ISM? Are SNRs correlated with large structures in the ISM or with star-forming regions?
- How many of the SNRs show correlations with molecular clouds? Can the SNR population explain the cosmic ray density in galaxies?

First X-ray surveys of the larger galaxies in the Local Group, the Magellanic Clouds, M31 and M33 were performed with *Einstein* and *ROSAT*, yielding catalogs of SNRs and candidates in these galaxies. A detailed list of X-ray SNRs in M31 was created using an XMM-Newton survey. While SNRs in the Magellanic Clouds can be generally well resolved spatially and studied in detail [22], so far, only a few SNRs in M31 have been resolved with *Chandra* [19, 20, 30].

The Magellanic Clouds, M31 and M33 have very different ISM densities, metallicities, and star formation rates, making differences in their ensemble of X-ray SNRs of great interest for testing theories for the dominant SNR and ISM characteristics that cause the X-ray emission of SNRs. *AXIS* will allow us to extend the study of SNR populations to nearby galaxies outside the Local Group including large spiral galaxies like M81, M83, or NGC 300, in which candidates of X-ray SNRs have been detected using *XMM-Newton* or *Chandra*, but more detailed studies have not been possible. In addition, $0.3''$ corresponds to ~ 1.1 pc at the distance of M31 which will allow us to detect and resolve all mature SNRs in M31 and M33, allowing for the first time detailed X-ray population studies of SNRs in galaxies beyond the Magellanic Clouds.

Last but not least, the combined sub-arcsecond resolution and high sensitivity will be needed to resolve small/young remnants, including SN 1987A which will should enter the ejecta-dominated phase in the 2030's [9]. This is also crucial for comparing the X-ray emission with that at other wavelengths, where high-resolution images are or will be available in the future.

In summary: Following up on the legacy of *Chandra*, an X-ray telescope with sub-arcsecond resolution combined with high sensitivity and ToO capabilities will further revolutionize the field of SNRs, PWNe and NSs in our Galaxy and the nearby Universe. These capabilities will be an absolute requirement to advance the field, especially in synergy with upcoming high-resolution facilities across the electromagnetic spectrum.

References

- [1] M. Aguilar et al. First Result from the Alpha Magnetic Spectrometer on the International Space Station: Precision Measurement of the Positron Fraction in Primary Cosmic Rays of 0.5–350 GeV. *PRL*, 110:141102, Apr 2013.
- [2] Elena Amato and Pasquale Blasi. Cosmic ray transport in the Galaxy: A review. *Advances in Space Research*, 62:2731–2749, Nov 2018.
- [3] Maxim V. Barkov, Maxim Lyutikov, Noel Klingler, and Pol Bordas. Kinetic ‘jets’ from fast-moving pulsars. *MNRAS*, 485:2041–2053, May 2019.
- [4] G. S. Bisnovatyi-Kogan. Asymmetric neutrino emission and formation of rapidly moving pulsars. *Astronomical and Astrophysical Transactions*, 3:287–294, Jan 1993.
- [5] Harsha Blumer, Samar Safi-Harb, and Maura A. McLaughlin. PSR J1119-6127 and Its Pulsar Wind Nebula Following the Magnetar-like Bursts. *ApJ*, 850:L18, Nov 2017.
- [6] Andrei M. Bykov, Donald C. Ellison, Sergei M. Osipov, and Andrey E. Vladimirov. Magnetic Field Amplification in Nonlinear Diffusive Shock Acceleration Including Resonant and Non-resonant Cosmic-Ray Driven Instabilities. *ApJ*, 789:137, Jul 2014.
- [7] L. Del Zanna, E. Amato, and N. Bucciantini. Axially symmetric relativistic MHD simulations of Pulsar Wind Nebulae in Supernova Remnants. On the origin of torus and jet-like features. *A&A*, 421:1063–1073, Jul 2004.
- [8] Robert C. Duncan and Christopher Thompson. Formation of Very Strongly Magnetized Neutron Stars: Implications for Gamma-Ray Bursts. *ApJ*, 392:L9, Jun 1992.
- [9] Kari A. Frank, Svetozar A. Zhekov, Sangwook Park, Richard McCray, Eli Dwek, and David N. Burrows. Chandra Observes the End of an Era in SN 1987A. *ApJ*, 829:40, Sep 2016.
- [10] Bryan M. Gaensler and Patrick O. Slane. The Evolution and Structure of Pulsar Wind Nebulae. *Annual Review of Astronomy and Astrophysics*, 44:17–47, Sep 2006.
- [11] F. P. Gavriil, M. E. Gonzalez, E. V. Gotthelf, V. M. Kaspi, M. A. Livingstone, and P. M. Woods. Magnetar-Like Emission from the Young Pulsar in Kes 75. *Science*, 319:1802, Mar 2008.
- [12] E. V. Gotthelf, J. P. Halpern, and J. Alford. The Spin-down of PSR J0821-4300 and PSR J1210-5226: Confirmation of Central Compact Objects as Anti-magnetars. *ApJ*, 765:58, Mar 2013.
- [13] Ersin Göğüş, Lin Lin, Yuki Kaneko, Chryssa Kouveliotou, Anna L. Watts, Manoneeta Chakraborty, M. Ali Alpar, Daniela Huppenkothen, Oliver J. Roberts, George Younes, and Alexander J. van der Horst. Magnetar-like X-Ray Bursts from a Rotation-powered Pulsar, PSR J1119-6127. *ApJ*, 829:L25, Oct 2016.

- [14] Tyler Holland-Ashford, Laura A. Lopez, Katie Auchettl, Tea Temim, and Enrico Ramirez-Ruiz. Comparing Neutron Star Kicks to Supernova Remnant Asymmetries. *ApJ*, 844:84, Jul 2017.
- [15] H. T. Janka and E. Mueller. Neutron star recoils from anisotropic supernovae. *A&A*, 290:496–502, Oct 1994.
- [16] O. Kargaltsev and G. G. Pavlov. Pulsar Wind Nebulae in the Chandra Era. In C. Bassa, Z. Wang, A. Cumming, and V. M. Kaspi, editors, *40 Years of Pulsars: Millisecond Pulsars, Magnetars and More*, volume 983 of *American Institute of Physics Conference Series*, pages 171–185, Feb 2008.
- [17] O. Kargaltsev, G. G. Pavlov, N. Klingler, and B. Rangelov. Pulsar wind nebulae created by fast-moving pulsars. *Journal of Plasma Physics*, 83:635830501, Oct 2017.
- [18] Satoru Katsuda, Mikio Morii, Hans-Thomas Janka, Annap Wongwathanarat, Ko Nakamura, Kei Kotake, Koji Mori, Ewald Müller, Tomoya Takiwaki, Masaomi Tanaka, Nozomu Tomonaga, and Hiroshi Tsunemi. Intermediate-mass Elements in Young Supernova Remnants Reveal Neutron Star Kicks by Asymmetric Explosions. *ApJ*, 856:18, Mar 2018.
- [19] A. K. H. Kong, M. R. Garcia, F. A. Primini, and S. S. Murray. The Discovery of a Spatially Resolved Supernova Remnant in M31 with Chandra. *ApJL*, 580:L125–L128, December 2002.
- [20] A. K. H. Kong, L. O. Sjouwerman, B. F. Williams, M. R. Garcia, and J. R. Dickel. Discovery of Radio/X-Ray/Optical-resolved Supernova Remnants in the Center of the Andromeda Galaxy. *ApJL*, 590:L21–L24, June 2003.
- [21] Harsha Sanjeev Kumar and Samar Safi-Harb. Variability of the High Magnetic Field X-Ray Pulsar PSR J1846-0258 Associated with the Supernova Remnant Kes 75 as Revealed by the Chandra X-Ray Observatory. *ApJ*, 678:L43–L46, May 2008.
- [22] P. Maggi, F. Haberl, P. J. Kavanagh, M. Sasaki, L. M. Bozzetto, M. D. Filipović, G. Vasilopoulos, W. Pietsch, S. D. Points, Y.-H. Chu, J. Dickel, M. Ehle, R. Williams, and J. Greiner. The population of X-ray supernova remnants in the Large Magellanic Cloud. *A&A*, 585:A162, January 2016.
- [23] R. Mushotzky. AXIS: a probe class next generation high angular resolution x-ray imaging satellite. In *Space Telescopes and Instrumentation 2018: Ultraviolet to Gamma Ray*, volume 10699 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, page 1069929, Jul 2018.
- [24] B. Olmi, L. Del Zanna, E. Amato, R. Bandiera, and N. Bucciantini. On the magnetohydrodynamic modelling of the Crab nebula radio emission. *MNRAS*, 438:1518–1525, Feb 2014.
- [25] G. G. Pavlov, V. E. Zavlin, B. Aschenbach, J. Trümper, and D. Sanwal. The Compact Central Object in Cassiopeia A: A Neutron Star with Hot Polar Caps or a Black Hole? *ApJ*, 531:L53–L56, Mar 2000.

- [26] Samar Safi-Harb. Pulsar Wind Nebulae: On their growing diversity and association with highly magnetized neutron stars. In Joeri van Leeuwen, editor, *Neutron Stars and Pulsars: Challenges and Opportunities after 80 years*, volume 291 of *IAU Symposium*, pages 251–256, Mar 2013.
- [27] Diego F. Torres. Rotationally Powered Magnetar Nebula around Swift J1834.9-0846. *ApJ*, 835:54, Jan 2017.
- [28] Yasunobu Uchiyama, Felix A. Aharonian, Takaaki Tanaka, Tadayuki Takahashi, and Yoshitomo Maeda. Extremely fast acceleration of cosmic rays in a supernova remnant. *Nature*, 449:576–578, Oct 2007.
- [29] Jacco Vink and J. Martin Laming. On the Magnetic Fields and Particle Acceleration in Cassiopeia A. *ApJ*, 584:758–769, Feb 2003.
- [30] B. F. Williams, L. O. Sjouwerman, A. K. H. Kong, J. D. Gelfand, M. R. Garcia, and S. S. Murray. Two New X-Ray/Optical/Radio Supernova Remnants in M31. *ApJL*, 615:720–726, November 2004.
- [31] G. Younes, C. Kouveliotou, O. Kargaltsev, R. Gill, J. Granot, A. L. Watts, J. Gelfand, M. G. Baring, A. Harding, G. G. Pavlov, A. J. van der Horst, D. Huppenkothen, E. Göğüş, L. Lin, and O. J. Roberts. The Wind Nebula around Magnetar Swift J1834.9-0846. *ApJ*, 824:138, Jun 2016.
- [32] V. N. Zirakashvili and F. Aharonian. Analytical solutions for energy spectra of electrons accelerated by nonrelativistic shock-waves in shell type supernova remnants. *A&A*, 465:695–702, Apr 2007.