

Palaeo-geoecological significance of Pleistocene trees in the Lluta Valley, Atacama Desert

Heinz Veit,^{1*} Jan-Hendrik May,² Andrea Madella,³ Romain Delunel,³ Fritz Schlunegger,³ Sönke Szidat⁴ and José M. Capriles^{5,6}

¹ Institute of Geography, University of Bern, Switzerland

² Institute of Earth and Environmental Sciences – Geology, Albert-Ludwigs-Universität Freiburg, Germany

³ Geological Institute, University of Bern, Switzerland

⁴ Department of Chemistry and Biochemistry & Oeschger Centre for Climate Change Research, University of Bern, Switzerland

⁵ Departamento de Antropología, Universidad de Tarapacá, Arica, Chile

⁶ Department of Anthropology, The Pennsylvania State University, PA 16802, USA

*Correspondence: Heinz Veit, as above.

E-mail: veit@giub.unibe.ch

Running head: PLEISTOCENE TREES IN THE LLUTA VALLEY

Graphical abstract **JQS 2857 Graphics sizing and other information**

Fig. 1. Full page depth with caption below

Fig. 2. Full page depth with caption below

Fig. 3. Full page width (17.6 cm)

Fig. 4. $\frac{3}{4}$ of full page width (13.2 cm)

Fig. 5. $\frac{2}{3}$ page width (11.7 cm), caption at the side

Number of Tables: 2.

Supporting information. None.

Please send the proofs to submitting Editor (Dr Joe Licciardi (Email: joe.licciardi@unh.edu) and to the E-in-C (Geoff Duller: ggd@aber.ac.uk)

Abstract

In the Lluta Valley, northern Chile, climate is hyperarid and vegetation is restricted to the valley floors and lowermost footslopes. Fossil tree trunks and leaves of predominantly *Escallonia angustifolia*, however, are abundant up to ~15 m above the present valley floor, where they are intercalated with slope deposits, reflecting higher water levels in the past. A total of 17 samples have been radiocarbon dated, yielding ages between 38 and 15k cal a BP. The youngest ages of 15.4k cal a BP are interpreted as reflecting the beginning of river incision and lowering of the valley floor, impeding the further growth of trees at higher parts of the slopes. The most plausible scenario for this observation is intensified river incision after 15.4k cal a BP due to increased stream power and runoff from the Rio Lluta headwaters in the Western Cordillera and Altiplano corresponding to the highstand of the Tauca and CAPE wet phase.

KEYWORDS: Atacama desert; climate change; fluvial dynamics; late Pleistocene; northern Chile.

Introduction

The Atacama Desert in northern Chile is one of the driest deserts on earth. Rainfall along the coast between Arica and Iquique is $\sim 1 \text{ mm a}^{-1}$ (Schulz *et al.*, 2011). Cosmogenic ^{21}Ne exposure dates on surfaces in Quebrada Tiliviche near Pisagua ($19^{\circ}35'\text{S}$) indicate the onset of extreme hyper-aridity probably as early as 25–22 Ma (Dunai *et al.*, 2005). Hyperarid conditions might have persisted through most of the Neogene as the Andes reached their current elevation around 15 Ma, creating the pronounced rain-shadow effect still in place today (Alpers and Brimhall, 1988; Houston and Hartley, 2003), even though ages around 10 Ma have also been proposed (Jordan *et al.*, 2014). As a consequence, the landscape in the Atacama Desert is characterized by very smooth slopes and interfluves which do not sustain noteworthy plant growth, and are characterized by extremely low erosion rates (Kober *et al.*, 2007). In contrast, the main rivers in northern Chile, such as the Río Lluta, reach the Pacific via canyon-like valleys (Figs 1 and 2a). Downcutting via headward migration of erosional fronts (i.e. knickzones) probably commenced at $\sim 8\text{--}11$ Ma in response to surface uplift (García and Hérail, 2005; Schlunegger *et al.*, 2006; Schildgen *et al.*, 2007, 2010; Thouret *et al.*, 2007; García *et al.*, 2011), and had lowered the valley floor to near-modern elevations by ~ 2.7 Ma in proximity to the Pacific coast, as evidenced by the Lauca Ignimbrite deposits (Schröder and Wörner, 1996; Uhlig, 1999; Schlunegger *et al.*, 2006; García *et al.*, 2011). This implies relative long-term stability of the valley floor elevation along the graded river reach beneath the knickzone (Fig. 1e), and dominant sediment bypass in wide braided river channels and floodplains. In this setting, vegetation is restricted to the lowermost footslopes and marginal valley floors, representing modern ‘treelines’ (Fig. 2a). Fossil tree trunks and leaves, however, are abundant up to $\sim 15\text{--}20$ m above the present valley floor of the Lluta Valley in northern Chile, where they are intercalated with slope deposits, suggesting past changes in water availability and humidity. Therefore, these deposits may provide new information on changes in discharge and/or sediment supplies over late Quaternary timescales. Here we present a comprehensive radiocarbon-based chronology for these stratified slope deposits in the Lluta Valley, consider potential mechanisms of their formation, and discuss their palaeoclimatic and tectonic significance.

Study area

Geologically, northern Chile is characterized (from west to east) by a Coastal Cordillera, Central Depression, Precordillera and the Western Cordillera (Fig. 1b). Directly north of Arica, the Coastal Cordillera is completely absent, whereas south of Arica it is characterized by a steep escarpment and up to 1000-m-high cliffs, deeply incised by some of the major rivers in northern Chile (Fig. 1b,c). With their headwaters in the Western Cordillera of the Andes, these rivers reach the Pacific via canyon-like valleys such as the Quebrada de Camarones, Quebrada de Vitor, Quebrada de Azapa or the Valle de Lluta which have incised up to 1000 m into the surrounding Tertiary pediments and peneplains of the Precordillera and Central Depression (Fig. 1). Further south, between $\sim 19^{\circ}30'S$ and $26^{\circ}30'S$, most rivers are endorheic and terminate in the hyperarid Atacama Desert. Climatic conditions in northern Chile and the Atacama region are generally hyperarid, and are characterized by a strong meridional gradient in annual precipitation with 0.4 mm at the coast to 237 mm at Putre (3650 m a.s.l.) and values of >350 mm in the highest parts of the catchment (DGA, 2004). The region's hyperaridity owes its existence to a combination of: (i) the extreme rainshadow effect of the high Andes, blocking advection of tropical/subtropical moisture from the east; (ii) limited influence of winter storm tracks from the south due to the presence of the semi-permanent South Pacific Anticyclone; and (iii) the generation of a temperature inversion at ~ 1000 m by the cold, north-flowing Humboldt Current that limits the inland (upslope) penetration of Pacific moisture. Precipitation variability in both summer and winter is modulated primarily by Pacific sea surface temperature gradients and the associated upper-air circulation anomalies (Vuille *et al.*, 2000; Garreaud *et al.*, 2003). A significant fraction of the inter-annual variability of summer precipitation is currently related to the El Niño Southern Oscillation (Vuille, 1999). Thus, wet summers on the Andean Altiplano are associated with an El Niño Southern Oscillation-related cooling of the tropical Pacific (i.e. La Niña phase). Rainfall and discharge maxima occur mainly during the 'Invierno Boliviano' (December–March; DGA, 2004; Houston, 2006) in relation to the South American Summer Monsoon system (Lenters and Cook, 1997; Schulz *et al.*, 2011), which transports moisture from the Amazon lowlands over the Andes down to ~ 2800 m a.s.l. on the western Andean slopes, but does not cause significant rainfall in the central Atacama (Rech *et al.*, 2006).

The catchment of the Río Lluta comprises an area of ~ 3400 km², with a river length of approximately 150 km (Fig. 1c). The highest peaks in the upper catchment are the 'Nevados de Putre' (5861 m a.s.l.), 'Volcán Tacora' (5988 m a.s.l.) and 'Volcán Tarapacá' (5860 m a.s.l.). The Lluta River originates in the Western Cordillera at an altitude of about 3900 m a.s.l. and reaches the Pacific Ocean ~ 4 km north of Arica at $18^{\circ}24'55''S$, $70^{\circ}19'35''W$ (Fig. 1c). Whereas a pediplain developed on top of the Coastal Cordillera during the Paleogene ('Tarapacá Pediplain'; Mortimer and Sarič, 1975), river incision in northern Chile commenced at ~ 8 – 11 Ma in response to surface uplift (García and Hérail, 2005; Schildgen *et al.*, 2007, 2010; Thouret *et al.*, 2007; García *et al.*, 2011). In the Lluta Valley incision had lowered the valley floor nearly to modern elevations by 2.7 Ma, as evidenced by the Lauca Ignimbrite deposits (Schröder and Wörner, 1996; Uhlig, 1999; García *et al.*, 2011). Today, the Lluta Valley long-profile exhibits a ~ 30 - km-long, steep and narrow knickzone of mixed bedrock–alluvial channel morphology, which divides the Lluta River into an upstream and downstream segment of wider valley floors (Fig. 1e). Beneath the knickzone, the valley floor is up to 1200 m wide, encompassing a gravelly braided river channel and variably wide floodplains. Only very few remnants of older terraces are visible, but they have not been studied up to now and their age and genesis are unknown. Today, the floodplain is subject to intensive irrigation and agricultural use (Fig. 2a). The Lluta River has incised 1–3 m into this valley bottom, which seems to have been the active floodplain for most of the Holocene. Apart from agriculture, *Escallonia angustifolia* and *Schinus molle* are the dominating trees of the riparian plant communities near the current valley floor. *Schinus molle* can live from groundwater, with root systems of 10–20 m in length, whereas *Escallonia angustifolia* needs surficial running water and humidity. Surface discharge along the Río Lluta is highly seasonal and can reach mean monthly flows of ~ 16 m³ s⁻¹ in February at the Pacific coast (DGA, 2004), and sediment loads up to 1700 tonnes per day have been measured during peak flow episodes (Campos Ortega *et al.*, 2007). The annual distribution of discharge is strongly controlled by precipitation in the Andean headwaters of the Lluta River in the Western Cordillera (Fig. 1c), which is

mainly built up by volcanic and sedimentary sequences (García *et al.*, 2011). In the Central Depression and the Precordillera, the geology is characterized by Oligo-Miocene sediments and ignimbrites (Azapa, Oxaya, El Diablo and Huaylas formations, Tobar *et al.*, 1968; Seyfried *et al.*, 1995; Uhlig, 1999; García *et al.*, 2011). Uplift of the Precordillera has induced major landslides, and parts of the Oxaya Formation collapsed >2.5 Myr ago building up the large Lluta Collapse deposit (Seyfried *et al.*, 1998; Wörner *et al.*, 2002; Strasser and Schlunegger, 2005; García *et al.*, 2011). All profiles presented here are situated in the area of the Lluta Collapse (Fig. 1c,d), east of Poconchile and along Chile's National Road 11 (18°27'6"S, 70°4'1"W; ±750 m a.s.l.). Along this segment of the valley, profiles can be studied easily along road cuts. However, the fossil trunks and leaves exist in a far broader area, from the coast and the mouth of the Lluta River up to > 1000 m a.s.l. along the valley floor. Further upstream of our study area the main road soon leaves the valley and crosses to the southern Azapa Valley. The smaller road which continues in the Lluta catchment runs on the present valley floor. We concentrated our investigation on the easily accessible part with open profiles.

Methods

Fieldwork consisted of detailed investigation of road cut exposures in the lower slopes along the southern side of the Lluta Valley east of Poconchile (Fig. 1c,d). An exhaustive sampling campaign was conducted on the organic horizons situated at 30–50 km from the coast (Figs 2–4). As the up to 50-cm-thick leaf deposits could be identified as almost completely consisting of leaves of *Escallonia angustifolia*, no further in-depth study of the palaeovegetation was undertaken. Trunks, which are completely imbedded within the leaf deposits, were not identified botanically. Profiles of these slope sedimentary sequences were described in the field. Profile location as well as an approximate elevation were documented with a hand-held GPS. Organic layers, mostly leaves and trunks, were stratigraphically sampled at five locations for ¹⁴C dating (Profiles km 36.2, km 36.5, km 37.0, km 37.5 and km 37.7; Table 1; ten samples), and were analysed at the radiocarbon laboratory of the Institute of Physics at the University of Bern. To check for spatial validity and chronological accuracy of these results, a set of additional samples was collected from exposures situated in a wider area (Profiles LVW-1, LVW-2, LVW-3, LVW-4, LVW-5, LVW-6 and LVW-7; Table 2; seven samples) and analysed with the accelerator mass spectrometer MICADAS at the LARA laboratory at the University of Bern (Szidat *et al.*, 2014). All results were calibrated with CALIB version 7.0 (Stuiver and Reimer, 1993, 2011) and the SHcal13 calibration curve (Hogg *et al.*, 2013). These dates are referred to as 'cal a BP' or 'k cal a BP'.

Results

The slopes along the Lluta Valley are very smooth and undissected without any sign of linear erosion (Fig. 2a). In the hyperarid setting of northern Chile, these slopes are the combined product of the slow salt- and gravity-related surficial displacement of material (salt-creep), granular flows, and the lack of linear fluvial erosion (Mortensen, 1927; Abele, 1987, 1990). These hillslope deposits are made up of well-sorted, horizontally stratified and cross-bedded sandy and coarser detritus, lacking a sedimentary matrix, with a thickening of the layers downslope. Stratification is on the order of millimetres, dipping parallel to the surface. These slope sediments are frequently intercalated with layers of fossil organic material such as trunks and leaves (Figs 2b–d and 3). Trunks and branches do not show signs of transport and are well preserved (Figs 2d and 3d). Therefore, they are more or less in an *in situ* position. It is particularly in the surroundings of Poconchile where this organic material is nicely recognizable along road cuts. In one case the slope sediments have buried

the Ah-horizon of a palaeosol (Fig. 2e). The fossil organic material reaches up to ~9 m above the present distribution of living trees at the intersection of the modern valley floor and slopes (Fig. 3a), which is generally up to 11–15 m (exception: profile LVW-2 at 17.9 m and Profile LVW-7 at 20.3 m) above the valley bottom. Further upslope, no trunks or leaves have been found. Most documented profiles contain several layers of organic material, reflecting an alternation of tree growth and sedimentation (Figs 2 and 4). All radiocarbon ages from these organic layers and a bAh horizon are summarized in Tables 1 and 2. The profiles documented in the photographs (Fig. 2) and sketches (Fig. 4) can be characterized as follows:

- At km 36.2 (Figs 2c and 4), leaves from a single organic layer at ~80 cm in the profile are dated to 23.9 ± 0.1 k cal a BP.
- The profile at km 36.5 (Figs 2b and 4) shows at least five organic layers and a layer of whitish volcanic ash. A wood fragment from the lowermost exposed organic layer at 250 cm below the surface yielded an age of 37.1 ± 0.4 k cal a BP. Two samples from leaf-dominated layers at 120 and 50 cm have ages of 28.9 ± 0.1 and 18.3 ± 0.1 k cal a BP, respectively. The uppermost 30 cm of the profile contains reeds, faecal pellets and abundant ceramics. The reed sample was dated to 0.5 ± 0.01 k cal a BP (~AD 1406–1439), thus reflecting the locally strong influence of pre-Colombian cultures in the Lluta Valley.
- At km 37.0 (Figs 2d and 4), a thick leaf deposit at a depth of 80 cm is dated to 19.4 ± 0.1 k cal a BP. Wood from the same layer yielded an age of 19.5 ± 0.1 k cal a BP.
- At km 37.5 (Fig. 4) leaves at a depth of 100–120 cm below the surface are dated to 15.6 ± 0.1 k cal a BP.
- At km 37.7 (Figs 2e and 4), the profile consists of a volcanic ash layer and an organic layer at the bottom intercalated with stratified slope deposits. Approximately 110 cm below the surface, a dark, greyish palaeosol (bAh-horizon) indicates a period of relative surface stability at 15.4 ± 0.4 k cal a BP. It is directly overlain by leaves with an age of 16.1 ± 0.1 k cal a BP. Both the palaeosol horizon and the leaf layer are truncated by the palaeochannel of a gully or small creek. The palaeochannel is filled by stratified slope deposits without organic layers and is no longer visible at the surface.

In summary, all dated wood and leaf samples belong stratigraphically to the Upper Pleistocene between ~15 and 38 k cal a BP. In all profiles, the lowermost layers are the oldest ones, becoming younger towards the surface. ^{14}C ages from leaves and wood from the same layer are in good accordance with this. No age inversions between layers occur. The results from ^{14}C dating of seven additional samples (Table 2), which were processed using two different methodologies, are internally consistent, and range between ~16 and 35 k cal a BP. Hence, they corroborate the chronological results obtained from the other profiles.

Discussion

The modern upper limit of living trees ('treeline', Figs 3a and 5) is characterized by a zone of non-climatic struggle between valley floor vegetation that depends on the presence of perennial surficial water and groundwater in the Rio Lluta floodplain deposits, and the vegetation-bare hillslopes with extreme aridity and active gravitational processes, mainly surficial creep, induced by wetting and drying of salt (Abele, 1987, 1990). This seems to be the present-day process, reflecting the dry environmental conditions on the slopes. Small gullies, as in profile km 37.7 (Fig. 2e), may have formed in a very short period, indicating linear erosion on the slope after 16.1–15.4 k cal a BP. They are filled with stratified slope deposits, which formed under dry conditions, comparable to those of today. Thus, in the marginal floodplain and on the lowermost slopes, accumulating organic matter, leaves and wood debris are successively covered by sediments of salt-creep, ultimately forming stratified deposits of alternating organic material and slope debris. The distribution of these deposits including the fossil trees and leaves is restricted to the lowermost slopes ~15–20 m above the modern floodplain in relative proximity to the modern treeline, indicating that the presence of fossilized vegetation remains does not reflect wetter climatic conditions at a local scale and extensively vegetated hillslopes,

for example as the result from substantially increased precipitation. A wider distribution of organic material and related palaeosols could be expected if more humid climatic conditions had prevailed at low elevations in the Atacama Desert, but has not been observed in the numerous hillslope quarries. In combination with the clear dominance of tree taxa in the organic layers, and the lack of hillslope taxa in the fossil record, this implies a higher palaeo-‘treeline’ (treeline A, Figs 3a and 5). In our profiles, the available radiocarbon dates from organic remains within the stratified slope deposits range between ~38 and 15.4k cal a BP and thus reflect tree growth up to ~9 m above the modern ‘treeline’ for this timeframe. In contrast, the youngest and stratigraphically uppermost organic layers consistently date to ~15.4k cal a BP, and no younger radiocarbon ages are available. Based on our model of quasi-continuous formation of stratified slope deposits along the marginal floodplain, this suggests a ~9-m downslope shift of the ‘treeline’ (from treeline A to B, Fig. 5) after ~15.4k cal a BP in relation to a major drop of the valley floor at that time.

Several scenarios could explain this drop on the order of ~ 9 m at our study site along the Rio Lluta: (i) the re-organization of the valley floor and river long-profile as the result of tectonics, (ii) the influence of global sea-level oscillations through the last glacial cycle, (iii) blocking of the valley by mass wasting (landslides, alluvial fans from tributaries, etc.) or dunes, (iv) reduced runoff and/or groundwater recharge in the Rio Lluta catchment due to drier climatic conditions in the Western Cordillera of the Andes, or – vice versa – (v) river incision induced by intensified runoff from the Western Cordillera and increased stream power.

Neotectonic activity is common throughout the Andean forearc and the Coastal Cordillera, even though the exact mechanisms and magnitudes of deformation related to stress accumulation and release during the seismic cycle are not yet fully understood (Allmendinger and González, 2010; A. Madella *et al.*, unpubl. data). Marine terraces dating to 140 ka are found up to 250 m a.s.l. in northern Chile and up to 300 m a.s.l. in southern Peru (Regard *et al.*, 2010), indicating variable uplift rates of up to 2.1 m ka⁻¹. In the area immediately north of Arica at the mouth of the Rio Lluta, however, the Coastal Cordillera is absent (Fig. 1a). In contrast, the broad Rio Lluta–Tacna plain seems to indicate relative subsidence rather than uplift. However, the discovery of ca. 10 ka raised fluvial terraces directly at the mouth of the Rio Lluta does suggest the occurrence of uplift, at least at the coast (A. Madella *et al.*, unpubl. data). Nevertheless, this uplift is unlikely to explain the 9-m lowering of groundwater table in the study reach because it occurred ca. 5000 years later. In addition, the study area is characterized by the lack of active structures (A. Madella *et al.*, unpubl. data), which implies that the smooth slopes and related deposits that embed the fossil wood are not affected by any tectonic deformation. This observation indicates that a tectonic disturbance alone could not be the major control for the onset of incision in this area.

During the Last Glacial Maximum (LGM), global sea level was ca. 125–135 m lower than today (Sidall *et al.*, 2006; Lambeck *et al.*, 2014). Today, the valley floor at our study site is ~750 m a.s.l. and up to 30–50 km inland from the coastline. The gradient at Poconchile is ~25 m km⁻¹ and gradually decreases to ~15 m km⁻¹ along the coast. During the LGM sea-level lowstand, the coastline would have been ~20 km west of the modern coastline (Becker *et al.*, 2009), indicating even lower gradients of around ~6 m km⁻¹. Additionally, sea level approached the LGM lowstand gradually over the last glacial cycle, in combination probably limiting the potential for significant upstream migration of a knickpoint. In contrast, the postglacial rise of global sea level was rapid between ~20 and 8 ka with a particularly rapid increase around 15 ka (Lambeck *et al.*, 2014). As a consequence, upstream valley floor aggradation and a tendency of increasing water levels would be expected. However, our results clearly indicate a decrease in water levels and/or valley floor elevations, suggesting that sea-level changes were probably not a major control on the observed drop in treeline after 15k cal a BP, particularly considering that our study site is located at altitudes >750 m a.s.l. The same holds true for a possible incursion of salinized groundwater and a corresponding lethal effect on the vegetation, following sea-level rise. This scenario cannot explain dying trees at >750 m a.s.l.

Landslides or alluvial input from tributary rivers might have caused local blocking of the Lluta River. So far, no age control on visible landslides in the valley is available. Because fossil trunks and leaves exist from the coast, a few metres above sea level to far into the catchment up to >1000 m a.s.l., and due

to the complete lack of limnic sediments indicating former lake conditions, this catastrophic explanation does not seem realistic. Additionally, the vertical sequence of ^{14}C ages in the profiles would have to be interpreted as several blocking periods, with each (younger) one resulting in higher lake levels, which also has no indication in the sediments.

Pleistocene dunes might have blocked the Lluta River during periods of low sea level such as the LGM. The rising sea level after 15 ka might have hindered new dune formation, allowing the Lluta River to incise into the valley bottom. However, as no major dune system has yet been identified and datings are not available, these considerations are speculative. In addition and similar to the potential effect of landslide blocking, the dated profiles are situated at an altitude of >750 m a.s.l., where a direct effect of blocking dunes in the lower part of the valley does not seem to be very likely.

The temporal coincidence between a well-known wet phase on the Altiplano and the initiation of valley lowering along the Río Lluta suggests an environmental driver. In this context, some authors report wet LGM conditions in the Western Cordillera/Altiplano (e.g. Baker *et al.*, 2001a,b; Bobs *et al.*, 2001; Rigsby *et al.*, 2005; Steffen *et al.*, 2009). In turn, a drop in treeline after ~15.4k cal a BP would consequently point to drier conditions. Most published proxies, however, point to wet climatic conditions on the Bolivian Altiplano, commonly referred to as the 'Tauca Period' (Servant and Fontes, 1978). Palaeolake Tauca was a freshwater lake, indicating enhanced precipitation between 18.1 and 14.1k cal a BP, with highest lake levels from 16.4 to 14.1k cal a BP (Servant and Fontes, 1978; Sylvestre *et al.*, 1999; Placzek *et al.*, 2006, 2013). Lake levels dropped after 14.1k cal a BP with a minor, secondary highstand during the minor Coipasa phase, between 13 and 11k cal a BP (Placzek *et al.*, 2006). Nester *et al.* (2007) and Gayo *et al.* (2012) report perennial stream discharge in the endorheic basins of the Atacama (Pampa del Tamarugal, 19°30' to 22°S) between 16.4 and 13.7k cal a BP and consider this the most important groundwater recharge event of the past 18 ka. It might appear as a contradiction that the alluvial fans in the Pampa del Tamarugal reacted with aggradation, whereas the Río Lluta eroded and incised at the same time. However, this can be easily explained by the different geomorphological systems; the fans in the endorheic basins of the Atacama are generally inactive during dry periods. There, increased precipitation leads to erosion in the catchment and corresponding accumulation on the fans. In a perennial river system with a much larger catchment area like the Río Lluta, it depends on the balance between water energy and material transported to react with accumulation or incision. After 15.4k cal a BP, there was probably a more consistent and/or elevated discharge available to allow for incision. At the Río Salado catchment (22°S), rodent middens from 3000 m a.s.l. have been interpreted as a two-fold precipitation increase at 17.5–16.3k cal a BP (Latorre *et al.*, 2006). A number of palaeowetland deposits additionally point to a major wet phase between >15 and 9k cal a BP (Betancourt *et al.*, 2000; Latorre *et al.*, 2002, 2003; Rech *et al.*, 2002, 2003; Quade *et al.*, 2008). The regional climatic pattern reflected in all these records has been summarized as the 'Central Andean Pluvial Event' ('CAPE'; Quade *et al.*, 2008). The onset of the CAPE (ca. 18k cal a BP) occurred 2000–2500 years before the lowering of the valley floor in the Lluta valley (15.4k cal a BP), as reflected in the youngest ages of the fossil leaves and trunks, but coincides well with the highstand of Lake Tauca and the wet period in the Pampa del Tamarugal at about 16.4–14k cal a BP. It is unlikely that fluvial incision at our study sites started at exactly the time when precipitation started to increase on the Altiplano. Instead, environmental changes in the catchment are progressively propagated through the fluvial system. This leads to a time lag (i.e. response time) which varies from catchment to catchment (compare e.g. Romans *et al.*, 2016) and was probably ~2000 years in duration after the onset of the CAPE in the Lluta Valley. Glacial ice modelling and dating of glacial advances also indicate rainfall increases of up to 100% in the Central Andes during the Tauca period (Kull *et al.*, 2002, 2008; Zech *et al.*, 2007, 2008, 2010). In contrast, the preceding period before and around the LGM (44–17 ka) was probably cold and dry in this area, as inferred from the small number of middens and the widespread lack of wetland deposits of this age, and palaeobotanical studies on fossil plants contained in them (Latorre *et al.*, 2002; Quade *et al.*, 2008).

In summary, the most plausible scenario for a ~9-m drop in treeline is river incision and lowering of the valley floor after 15.4k cal a BP due to increased stream power and runoff from the Rio Lluta headwaters in the Western Cordillera and Altiplano, and corresponding to the highstand of the CAPE and Tauca wet phase.

Conclusions

Along the large valley systems in the hyperarid Atacama Desert of northern Chile, vegetation is markedly restricted to the floodplains and valley floors. As such, the treeline is non-climatic and is only indirectly controlled by precipitation via groundwater availability and humidity at the valley floor. Despite the notion of long-term climatic stability along the lower reaches of the Lluta River, the observation of fossil tree remains, leaves and organic material intercalated with slope deposits significantly above the modern treeline indicates a drop in treeline elevation. Our radiocarbon-based chronology for these organic deposits suggests that this drop occurred at ~15.4k cal a BP, and was probably related to ~9 m of incision triggered by a pulse of increased discharge during the CAPE, which was characterized by significantly increased tropical moisture and precipitation totals in much of the Central Andes.

Acknowledgments. We very much appreciate the intense review and the valuable comments made by two reviewers.

Abbreviations. CAPE, Central Andean Pluvial Event; LGM, Last Glacial Maximum.

References

- Abbott MB, Wolfe BB, Wolfe AP *et al.* 2003. Holocene paleohydrology and glacial history of the central Andes using multiproxy lake sediment studies. *Palaeogeography, Palaeoclimatology, Palaeoecology* **194**: 123–138 [DOI: 10.1016/S0031-0182(03)00274-8].
- Abele G. 1987. Das Relief der Andenwestflanke bei Antofagasta (Nordchile) unter dem Einfluß langfristiger und extremer Trockenheit. *Erdkunde* **41**: 299–310 [DOI: 10.3112/erdkunde.1987.04.04].
- Abele G. 1990. Salzkrusten, salzbedingte Solifluktion und Steinsalzkarst in der nordchilenisch-peruanischen Wüste. *Mainzer Geographische Studien* **34**: 23–46.
- Allmendinger RW, González G. 2010. Invited review paper: Neogene to Quaternary tectonics of the coastal Cordillera, northern Chile. *Tectonophysics* **495**: 93–110 [DOI: 10.1016/j.tecto.2009.04.019].

- Alpers CN, Brimhall GH. 1988. Middle Miocene climatic change in the Atacama Desert, northern Chile: Evidence from supergene mineralization at La Escondida. *Geological Society of America Bulletin* **100**: 1640–1656 [DOI: 10.1130/0016-7606(1988)100<1640:MMCCIT>2.3.CO;2].
- Ariazza B. 1995. Chile's Chinchorro mummies. *National Geographic* **187**: 68–89.
- Baker PA, Rigsby CA, Seltzer GO *et al.* 2001b. Tropical climate changes at millennial and orbital timescales on the Bolivian Altiplano. *Nature* **409**: 698–701 [DOI: 10.1038/35055524] [PubMed: 11217855],
- Becker JJ, Sandwell DT, Smith WHF *et al.* 2009. Global bathymetry and elevation data at 30 arc seconds resolution: SRTM30_PLUS. *Marine Geodesy* **32**: 355–371 [DOI: 10.1080/01490410903297766].
- Baker PA, Seltzer GO, Fritz SC *et al.* 2001a. The history of South American tropical precipitation for the past 25,000 years. *Science* **291**: 640–643 [DOI: 10.1126/science.291.5504.640] [PubMed: 11158674].
- Betancourt JL, Latorre C, Rech JA *et al.* 2000. A 22,000-year record of monsoonal precipitation from northern Chile's Atacama Desert. *Science* **289**: 1542–1546 [DOI: 10.1126/science.289.5484.1542] [PubMed: 10968788].
- Bobs AL, Lowenstein TK, Jordan TE *et al.* 2001. A 106ka paleoclimate record from drill core of the Salar de Atacama, northern Chile. *Palaeogeography, Palaeoclimatology, Palaeoecology* **173**: 21–42 [DOI: 10.1016/S0031-0182(01)00308-X].
- Campos Ortega H, Díaz Muñoz G, Campos Ortega C. 2007. Aportes sedimentarios de los ríos Lluta y San José en la zona costera de la rada de Arica, Chile. *Idesia* **25**: 37–48.
- DGA. 2004. *Cuenca del Río Lluta*. Dirección General de Aguas, Gobierno de Chile.
- Dunai TJ, González López GA, Juez-Larré J. 2005. Oligocene–Miocene age of aridity in the Atacama Desert revealed by exposure dating of erosion-sensitive landforms. *Geology* **33**: 321–324 [DOI: 10.1130/G21184.1].
- Durán-Gonzales JA, Cabrera-Fajardo G. 2001. Field determination of stream–aquifer interrelations caused by intensive groundwater pumping for water right purposes: integrated water resources management. *IAHS Publication* **272**: 205–212.
- Eitel B, Hecht S, Mächtle B *et al.* 2005. Geoarcheological evidence from desert loess in the Nazca-Palpa region, southern Peru: palaeoenvironmental changes and their impact on pre-Columbian cultures. *Archaeometry* **47**: 137–158 [DOI: 10.1111/j.1475-4754.2005.00193.x].
- Fritz SC, Baker PA, Lowenstein TK *et al.* 2004. Hydrologic variation during the last 170'000 years in the southern hemisphere tropics of South America. *Quaternary Research* **6**: 95–104.

- García M, Hérail G. 2005. Fault-related folding, drainage network evolution and valley incision during the Neogene in the Andean Precordillera of Northern Chile. *Geomorphology* **65**: 279–300 [DOI: 10.1016/j.geomorph.2004.09.007].
- García M, Riquelme R, Farías M. 2011. Late Miocene–Holocene canyon incision in the western Altiplano, northern Chile: tectonic or climatic forcing? *Journal of the Geological Society of London* **168**: 1047–1060.
- Garreaud R, Vuille M, Clement AC. 2003. The climate of the Altiplano: observed current conditions and mechanisms of past changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* **194**: 5–22 [DOI: 10.1016/S0031-0182(03)00269-4].
- Garreaud RD, Vuille M, Compagnucci R *et al.* 2009. Present-day South American climate. *Palaeogeography, Palaeoclimatology, Palaeoecology* **281**: 180–195 [DOI: 10.1016/j.palaeo.2007.10.032].
- Gayo EM, Latorre C, Jordan TE *et al.* 2012. Late Quaternary hydrological and ecological changes in the hyperarid core of the northern Atacama Desert (~21°S). *Earth-Science Reviews* **113**: 120–140 [DOI: 10.1016/j.earscirev.2012.04.003].
- Hogg AG, Hua Q, Blackwell PG *et al.* 2013. SHCal13 southern hemisphere calibration, 0–50,000 cal yr BP. *Radiocarbon* **55**: 1889–1903.
- Houston J. 2006. Variability of precipitation in the Atacama Desert: its causes and hydrological impact. *International Journal of Climatology* **26**: 2181–2198 [DOI: 10.1002/joc.1359].
- Houston J, Hartley AJ. 2003. The Central Andean west-slope rainshadow and its potential contribution to the origin of hyper-aridity in the Atacama Desert. *International Journal of Climatology* **23**: 1453–1464 [DOI: 10.1002/joc.938].
- Jordan TE, Kirk-Lawlor NE, Blanco NP *et al.* 2014. Landscape modification in response to repeated onset of hyperarid paleoclimate states since 14 Ma, Atacama Desert, Chile. *Geological Society of America Bulletin* **126**: 1016–1046 [DOI: 10.1130/B30978.1].
- Kober F, Ivy-Ochs S, Schlunegger F *et al.* 2007. Denudation rates and a topography-driven rainfall threshold in northern Chile: multiple cosmogenic nuclide data and sediment yield budgets. *Geomorphology* **83**: 97–120 [DOI: 10.1016/j.geomorph.2006.06.029].
- Kull C, Grosjean M, Veit H. 2002. Modelling Modern and Late Pleistocene glacio-climatological conditions in the North Chilean Andes (29°S –30°S). *Climate Change* **53**: 359–381.
- Kull C, Imhof S, Grosjean M *et al.* 2008. Late Pleistocene glaciation in the Central Andes: temperature versus humidity control – a case study from the eastern Bolivian Andes (17°S) and regional synthesis. *Global and Planetary Change* **60**: 148–164 [DOI: 10.1016/j.gloplacha.2007.03.011].

- Lambeck K, Rouby H, Purcell A *et al.* 2014. Sea level and global ice volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National Academy of Sciences of the United States of America* **111**: 15296–15303 [DOI: 10.1073/pnas.1411762111] [PubMed: 25313072].
- Latorre C, Betancourt JL, Arroyo MTK. 2006. Late Quaternary vegetation and climate history of a perennial river canyon in the Río Salado basin (22°S) of Northern Chile. *Quaternary Research* **65**: 450–466 [DOI: 10.1016/j.yqres.2006.02.002].
- Latorre C, Betancourt JL, Rylander KA *et al.* 2002. Vegetation invasions into absolute desert: A 45;th000 yr rodent midden record from the Calama–Salar de Atacama basins, northern Chile (lat 22°–24°S). *Geological Society of America Bulletin* **114**: 349–366 [DOI: 10.1130/0016-7606(2002)114<0349:VIIADA>2.0.CO;2].
- Latorre C, Betancourt JL, Rylander KA *et al.* 2003. A vegetation history from the arid prepuna of northern Chile (22–23°S) over the last 13 500 years. *Palaeogeography, Palaeoclimatology, Palaeoecology* **194**: 223–246 [DOI: 10.1016/S0031-0182(03)00279-7].
- Lenters JD, Cook KH. 1997. On the origin of the Bolivian High and related circulation features of the South American climate. *Journal of the Atmospheric Sciences* **54**: 656–678 [DOI: 10.1175/1520-0469(1997)054<0656:OTOOTB>2.0.CO;2].
- Lowenstein TK, Hein MC, Bobst AL *et al.* 2003. An assessment of stratigraphic completeness in climate-sensitive closed-basin lake sediments: Salar de Atacama, Chile. *Journal of Sedimentary Research* **73**: 91–104 [DOI: 10.1306/061002730091].
- Mortensen H. 1927. Der Formenschatz der nordchilenischen Wüste. *Abhandlungen der Akademie der Wissenschaften in Göttingen, Math.-Physik. Kl., N.F.* **12**: 1–191.
- Mortimer C, Sarić N. 1975. Cenozoic studies in northernmost Chile. *Geologische Rundschau* **64**: 395–420.
- Nester PL, Gayó E, Latorre C *et al.* 2007. Perennial stream discharge in the hyperarid Atacama Desert of northern Chile during the latest Pleistocene. *Proceedings of the National Academy of Sciences of the United States of America* **104**: 19724–19729 [DOI: 10.1073/pnas.0705373104] [PubMed: 18056645].
- Placzek C, Quade J, Patchett PJ. 2006. Geochronology and stratigraphy of late Pleistocene lake cycles on the southern Bolivian Altiplano: Implications for causes of tropical climate change. *Geological Society of America Bulletin* **118**: 515–532 [DOI: 10.1130/B25770.1].
- Placzek CJ, Quade J, Patchett PJ. 2013. A 130 ka reconstruction of rainfall on the Bolivian Altiplano. *Earth and Planetary Science Letters* **363**: 97–108 [DOI: 10.1016/j.epsl.2012.12.017].

- Quade J, Rech JA, Betancourt JL *et al.* 2008. Paleowetlands and regional climate change in the central Atacama Desert, northern Chile. *Quaternary Research* **69**: 343–360 [DOI: 10.1016/j.yqres.2008.01.003].
- Rech JA, Currie BS, Michalski G *et al.* 2006. Neogene climate change and uplift in the Atacama Desert, Chile. *Geology* **34**: 761–764 [DOI: 10.1130/G22444.1].
- Rech JA, Pigati JS, Quade J *et al.* 2003. Re-evaluation of mid-Holocene deposits at Quebrada Puripica, northern Chile. *Palaeogeography, Palaeoclimatology, Palaeoecology* **194**: 207–222 [DOI: 10.1016/S0031-0182(03)00278-5].
- Rech JA, Quade J, Betancourt JL. 2002. Late Quaternary paleohydrology of the central Atacama Desert (lat 22°–24°S), Chile. *Geological Society of America Bulletin* **114**: 334–348 [DOI: 10.1130/0016-7606(2002)114<0334:LQPOTC>2.0.CO;2].
- Regard V, Saillard M, Martinod J *et al.* 2010. Renewed uplift of the central Andes forearc revealed by coastal evolution during the Quaternary. *Earth and Planetary Science Letters* **297**: 199–210 [DOI: 10.1016/j.epsl.2010.06.020].
- Rigsby CA, Bradbury JP, Baker PA *et al.* 2005. Late Quaternary palaeolakes, rivers, and wetlands on the Bolivian Altiplano and their palaeoclimatic implications. *Journal of Quaternary Science* **20**: 671–691 [DOI: 10.1002/jqs.986].
- Romans BW, Castellort S, Covault JA *et al.* 2016. Environmental signal propagation in sedimentary systems across timescales. *Earth-Science Reviews* **153**: 7–29 [DOI: 10.1016/j.earscirev.2015.07.012].
- Saillard M, Hall SR, Audin L *et al.* 2011. Andean coastal uplift and active tectonics in southern Peru: ¹⁰Be surface exposure dating of differentially uplifted marine terrace sequences (San Juan de Marcona, ~15.4°S). *Geomorphology* **128**: 178–190 [DOI: 10.1016/j.geomorph.2011.01.004].
- Schildgen TF, Balco G, Shuster DL. 2010. Canyon incision and knickpoint propagation recorded by apatite ⁴He/³He thermochronometry. *Earth and Planetary Science Letters* **293**: 377–387 [DOI: 10.1016/j.epsl.2010.03.009].
- Schildgen TF, Hodges KV, Whipple KX *et al.* 2007. Uplift of the western margin of the Andean plateau revealed from canyon incision history, southern Peru. *Geology* **35**: 523–526 [DOI: 10.1130/G23532A.1].
- Schlunegger F, Zeilinger G, Kounov A *et al.* 2006. Scale of relief growth in the forearc of the Andes of Northern Chile (Arica latitude, 18oS). *Terra Nova* **18**: 217–223 [DOI: 10.1111/j.1365-3121.2006.00682.x].
- Schröder W, Wörner G. 1996. Widespread Cenozoic ignimbrites in N-Chile, W-Bolivia and S-Peru (17°–29°S/71°–68°E): stratigraphy, extension, correlation and origin. *3me symposium international sur la Géodynamique andine, Satin-Malo*. ORSTOM Collection Colloques et Séminaires: 645–648.

- Schulz N, Boisier JP, Aceituno P. 2012. Climate change along the arid coast of northern Chile. *International Journal of Climatology* **32**: 1803–1814 [DOI: 10.1002/joc.2395].
- Servant M, Fontes JC. 1978. Les lacs quaternaires des hauts plateaux des Andes boliviennes. Premières interprétations paléoclimatiques. *Cahiers Orstom, Serie Geologie* **10**: 9–23.
- Seyfried H, Wörner G, Uhlig D. 1995. Eine kleine Landschaftsgeschichte der Anden in Nordchile. Wechselwirkungen; Jahrbuch 1994: 60–72. University of Stuttgart.
- Seyfried H, Wörner G, Uhlig D *et al.* 1998. Introducción a la geología y morfología de los Andes en el norte de Chile. *Chungara* **30**: 7–39.
- Sidall M, Chappell J, Potter EK. 2006. Eustatic sea-level during past interglacials. In *The Climate of Past Interglacials*, Sirocko F, Litt T, Claussen M, Sanchez-Goni MF (eds). Elsevier: Amsterdam; 75–92.
- Steffen D, Schlunegger F, Preusser F. 2009. Drainage basin response to climate change in the Pisco valley, Peru. *Geology* **37**: 491–494 [DOI: 10.1130/G25475A.1].
- Strasser M, Schlunegger F. 2005. Erosional processes, topographic length-scales and geomorphic evolution in arid climatic environments: the ‘Lluta collapse’, northern Chile. *International Journal of Earth Sciences* **94**: 433–446 [DOI: 10.1007/s00531-005-0491-2].
- Stuiver M, Reimer PJ. 1993. Extended ¹⁴C database and revised CALIB radiocarbon calibration program. *Radiocarbon* **35**: 215–230.
- Sylvestre F, Servant M, Servant-Vildary S *et al.* 1999. Lake-level chronology on the Southern Bolivian Altiplano (18°–23°S) during Late-Glacial Time and the Early Holocene. *Quaternary Research* **51**: 54–66 [DOI: 10.1006/qres.1998.2017].
- Szidat S, Salazar GA, Vogel E, Battaglia M, Wacker L, Synal HA, Türlér A. 2014. ¹⁴C analysis and sample preparation at the New Bern Laboratory for the analysis of radiocarbon with AMS (LARA). *Radiocarbon* **56**: 561–566 [DOI: 10.2458/56.17457].
- Thouret J-C, Wörner G, Gunnell Y *et al.* 2007. Geochronologic and stratigraphic constraints on canyon incision and Miocene uplift of the Central Andes in Peru. *Earth and Planetary Science Letters* **263**: 151–166 [DOI: 10.1016/j.epsl.2007.07.023].
- Tobar A, Salas Y, Kast RF. 1968. *Cuadrángulos Camarca y Azapa, Provincia de Tarapacá*. Carta Geologica de Chile 19/20, 1:50 000, Inst. de Inv. Geológicas: Santiago.
- Uhlig D. 1999. Die Westabdachung der Zentralanden in den Provinzen Arica und Parinacota, Nordchile: Landschaftsentwicklung und Geologie. *Profil* **17**: 167–124.

- Vuille M. 1999. Atmospheric circulation over the Bolivian Altiplano during dry and wet periods and extreme phases of the Southern Oscillation. *International Journal of Climatology* **19**: 1579–1600 [DOI: 10.1002/(SICI)1097-0088(19991130)19:14<1579::AID-JOC441>3.0.CO;2-N].
- Vuille M, Bradley RS, Keimig F. 2000. Interannual climate variability in the Central Andes and its relation to tropical Pacific and Atlantic forcing. *Journal of Geophysical Research: Atmospheres* **105**: 12447–12460 [DOI: 10.1029/2000JD900134].
- Wörner G, Uhlig D, Kohler I *et al.* 2002. Evolution of the West Andean Escarpment at 18°S (N. Chile) during the last 25 Ma: uplift, erosion and collapse through time. *Tectonophysics* **345**: 183–198 [DOI: 10.1016/S0040-1951(01)00212-8].
- Zech J, Zech R, May JH *et al.* 2010. Lateglacial and Early Holocene glaciation in the tropical Andes caused by la Niña-like conditions. *Palaeogeography, Palaeoclimatology, Palaeoecology* **293**: 248–254 [DOI: 10.1016/j.palaeo.2010.05.026].
- Zech R, May J, Kull C *et al.* 2008. Timing of the late Quaternary glaciation in the Andes from ~15 to 40° S. *Journal of Quaternary Science* **23**: 635–647 [DOI: 10.1002/jqs.1200].
- Zech R, Kull C, Kubik PW *et al.* 2007. Exposure dating of Late Glacial and pre-LGM moraines in the Cordon de Doña Rosa, Northern/Central Chile (~31° S). *Climate of the Past* **3**: 1–14 [DOI: 10.5194/cp-3-1-2007].

Figure 1. Study area. (a) Overview; (b) northern Chile with morphotectonical units; (c) Lluta Valley with location of the study area; (d) location of studied profiles; (e) longitudinal profile of the Río Lluta. Triangle in (c) marks the location of dated wood at the coast, according to A. Madella (pers. comm.).

Figure 2. Field impressions from the Lluta Valley. (a) Overview, agriculture on the valley floor and dry slopes; (b–d) stratified slope deposits with intercalated wood and leaves; (e) gully, filled with slope deposits and older, buried organic soil horizons (note: green triangles mark organic layers with fossil trees, logs and leaves; white triangle marks volcanic ash).

Figure 3. Field impressions from the Lluta Valley. (a) Overview with palaeo-‘treeline’; (b) organic layers in surface parallel position; (c) detail of leaf deposit; (d) leaves of dominating *Escallonia angustifolia*.

Figure 4. Sketch of profiles with ¹⁴C dates. All dates are given as k cal a BP (Table 1).

Figure 5. Sketch, showing the evolution of the slopes and the valley bottom, the incision of the Río Lluta and the changes in ‘treeline’ due to lowering of the valley bottom.

Table 1. ¹⁴C results from samples VEC 06/.

Sample name	Latitude	Longitude	Profile	Height	Depth	Material	Lab code	δ ¹³ C (‰)	¹⁴ C age (a BP)	Age (cal a BP)	Age (k cal a BP)
-------------	----------	-----------	---------	--------	-------	----------	----------	-----------------------	----------------------------	----------------	------------------

	(°S)	(°W)		above valley floor (m)	(cm)										
VEC 06/35	-18.40221	-70.01646	36.2	12	80	Leaves	B-9016	-25.0	± 0.2	19 870	± 70	23 746	23 993	23.9	± 0.1
VEC 06/39	-18.40357	-70.01371	36.5	11.5	15	Reed	B-9012	-10.2	± 0.2	510	± 20	503	520	0.5	± 0.0
VEC 06/36	-18.40357	-70.01371	36.5	11.5	50	Leaves	B-9013	-23.6	± 0.2	15 120	± 90	18 192	18 459	18.3	± 0.1
VEC 06/37B	-18.40357	-70.01371	36.5	11.5	120	Leaves	B-9014	-23.2	± 0.2	24 900	± 100	28 732	29 001	28.9	± 0.1
VEC 06/38	-18.40357	-70.01371	36.5	11.5	250	Wood	B-9015	-24.4	± 0.2	33 090	± 180	36 733	37 514	37.1	± 0.4
VEC 06/34A	-18.40553	-70.00944	37.0	12.5	80	Leaves	B-9010	-23.4	± 0.2	16 120	± 60	19 298	19 511	19.4	± 0.1
VEC 06/34B	-18.40553	-70.00944	37.0	12.5	80	Wood	B-9011	-20.7	± 0.2	16 180	± 60	19 378	19 583	19.5	± 0.1
VEC 06/33	-18.40787	-70.00493	37.5	13.5	110	Leaves	B-9009	-24.2	± 0.2	13 080	± 70	15 457	15 757	15.6	± 0.2
VEC 06/30	-18.40865	-70.00325	37.7	13.5	115	TOC fAh	B-9007	-21.1	± 0.2	12 930	± 220	15 073	15 782	15.4	± 0.4
VEC 06/32	-18.40865	-70.00325	37.7	14	110	Leaves	B-9008	-23.4	± 0.2	13 420	± 80	15 973	16 226	16.1	± 0.1

Table 2. ^{14}C results from samples LVW-, presented with 95% confidence limits.

Sample name	Latitude (°S)	Longitude (°W)	Height above valley floor (m)	Depth (cm)	Material	Lab code	$\delta^{13}\text{C}$ (‰)	^{14}C age (a BP)	Age (cal a BP)	Age (k cal a BP)			
LVW-1	-18.45164	-70.05891	8.3	30	Wood	BE-1754.1.1	-18.5	30 593	± 140	34 300	34 897	35	± 0.3
						BE-1754.1.2	-19.8	30 842	± 178				
						BE-1754.2.1*	-20.5	30 936	± 285				
LVW-2	-18.40407	-70.03452	17.9	10	Wood	BE-1755.2.1*	-19.0	13 826	± 212	16 094	17 330	16.7	± 0.6
LVW-3	-18.40036	-70.02080	14.5	20	Wood	BE-1756.1.1	-20.1	18 641	± 41	21 999	22 588	22	± 0.3
						BE-1756.1.2	-18.9	18 415	± 51				
						BE-1756.2.1*	-22.5	18 891	± 216				
LVW-4	-18.40273	-70.01611	11.6	60	Wood	BE-1757.1.1	-19.7	25 547	± 81	29 294	29 842	30	± 0.3
						BE-1757.1.2	-18.4	25 495	± 99				
						BE-1757.2.1*	-20.4	25 326	± 234				
LVW-5	-18.40552	-70.00953	11.6	80	Wood	BE-1758.2.1*	-18.9	23 704	± 227	27 427	28 234	27.8	± 0.4
LVW-6	-18.40552	-70.00953	11.3	60	Wood	BE-1759.2.1*	-21.1	16 132	± 214	18 919	19 948	19.4	± 0.5
LVW-7	-18.39513	-69.96443	20.0	40	Wood	BE-1760.1.1	-18.0	14 576	± 30	17 538	17 880	18	± 0.2

* Due to advanced degradation, these samples were prepared using the simplified acid–base–acid (ABA) method and corrected for an average age offset of -454 ± 209 years by comparison with the base–acid–base–acid–bleaching (BABAB) method, i.e. the standard method for wood analysis (Szidat *et al.*, 2014).



Fig 3 .

Author Manuscript




Fig1 .

Author Manuscript







Fig2 .

Author Manuscript


Profile 'km 36.2'




Profile 'km 37.0'




Profile 'km 36.5'



Profile 'km 37.7'



Profile 'km 37.5'




Legend

- | | | | | |
|--|----------------------------------|--------------|-----------------------|-----------------------------|
| Slope Deposit / Salt Creep (Late Holocene) | Plant Remains (Wood, Leaves) | Paleosol | Ceramics | Radiocarbon Age (cal ka BP) |
| Slope Deposit / Salt Creep (Pleistocene) | Plant Remains (Reed, Corn/Maize) | Volcanic Ash | Fecal Pellets (Llama) | |

Fig4 .


Author Manuscript

a) >38-15.4 cal kyr BP



1) Burial and Formation of Stratified Slope Deposits

b) <15.4 cal kyr BP



2) Incision

3) Renewed Burial and Formation of Stratified Slope Deposits

Fig5 .

Author Manuscript



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Veit, H; May, J-H; Madella, A; Delunel, R; Schlunegger, F; Szidat, S; Capriles, JM

Title:

Palaeo-geoecological significance of Pleistocene trees in the Lluta Valley, Atacama Desert

Date:

2016-04-01

Citation:

Veit, H., May, J. -H., Madella, A., Delunel, R., Schlunegger, F., Szidat, S. & Capriles, J. M. (2016). Palaeo-geoecological significance of Pleistocene trees in the Lluta Valley, Atacama Desert. JOURNAL OF QUATERNARY SCIENCE, 31 (3), pp.203-213.
<https://doi.org/10.1002/jqs.2857>.

Persistent Link:

<http://hdl.handle.net/11343/291136>

File Description:

Accepted version