



eCOMMONS

Loyola University Chicago  
Loyola eCommonsAnthropology: Faculty Publications and Other  
Works

Faculty Publications

2008

## Technical note: dental microwear textures of "Phase I" and "Phase II" facets

Kristin L. Krueger

*kkruieger4, kkrueger4@luc.edu*

Jessica R. Scott

*University of Arkansas at Little Rock*

Richard F. Kay

*Duke University*

Peter S. Ungar

*University of Arkansas, Fayetteville*Follow this and additional works at: [https://ecommons.luc.edu/anthropology\\_facpubs](https://ecommons.luc.edu/anthropology_facpubs)Part of the [Anthropology Commons](#)

### Recommended Citation

Krueger, Kristin L.; Scott, Jessica R.; Kay, Richard F.; and Ungar, Peter S.. Technical note: dental microwear textures of "Phase I" and "Phase II" facets. *American Journal of Physical Anthropology*, 137, 4: 485-490, 2008. Retrieved from Loyola eCommons, Anthropology: Faculty Publications and Other Works, <http://dx.doi.org/10.1002/ajpa.20928>

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Anthropology: Faculty Publications and Other Works by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 License](#).  
© 2008 Wiley.



American Journal of  
Physical Anthropology

**Technical Note: Dental microwear textures of “Phase I” and “Phase II” facets**

Journal:	<i>American Journal of Physical Anthropology</i>
Manuscript ID:	draft
Wiley - Manuscript type:	Technical Note
Date Submitted by the Author:	n/a
Complete List of Authors:	Krueger, Kristin; University of Arkansas, Department of Anthropology Scott, Jessica; University of Arkansas, Department of Anthropology; University of Arkansas, Environmental Dynamics Kay, Richard; Duke University, Biological Anthropology and Anatomy Ungar, Peter; University of Arkansas, Anthropology
Key Words:	Microwear, Mastication, Primate, Diet



1  
2  
3  
4  
5  
6  
7  
8  
9 Technical Note: Dental microwear textures of “Phase I” and “Phase II” facets  
10  
11  
12  
13  
14  
15  
16

17 Kristin L. Krueger<sup>1</sup>, Jessica R. Scott<sup>2</sup>, Richard F. Kay<sup>3</sup>, and Peter S. Ungar<sup>1,2</sup>  
18  
19  
20  
21  
22  
23

24 <sup>1</sup>Department of Anthropology, University of Arkansas, Fayetteville, AR 72701  
25

26 <sup>2</sup>Environmental Dynamics Program, University of Arkansas, Fayetteville, AR 72701  
27

28 <sup>3</sup>Department of Biological Anthropology and Anatomy, Duke University  
29 Durham, NC 27708  
30  
31  
32

33 Address correspondence to:  
34  
35

36 Peter Ungar  
37 Department of Anthropology  
38 Old Main 330  
39 University of Arkansas  
40 Fayetteville, AR 72701 USA  
41 Phone: 479-575-6361  
42 FAX: 479-575-6595  
43 Email: [pungar@uark.edu](mailto:pungar@uark.edu)  
44  
45  
46  
47  
48  
49

50 Number of text pages: 12

51 Number of figures: 2

52 Number of tables: 4  
53  
54  
55

56 Grant sponsor: US National Science Foundation. Grant number: BCS 0315157  
57  
58  
59  
60

**ABSTRACT**

1  
2  
3  
4  
5  
6  
7  
8 The power stroke of mastication has been traditionally divided into two parts, one  
9  
10 which precedes centric occlusion, and the other which follows it -- “Phase I” and “Phase  
11  
12 II” respectively. Recent studies of primate mastication have called into question the role  
13  
14 of “Phase II” in food processing, as they have found little muscle activity or  
15  
16 accompanying bone strain following centric occlusion. That said, many researchers  
17  
18 today look to “Phase II” facets to relate diet to patterns of dental microwear. This  
19  
20 suggests the need to reevaluate microwear patterns on “Phase I” facets. Here we use  
21  
22 texture analysis to compare and contrast microwear on facets representing both phases in  
23  
24 three primate species with differing diets (*Alouatta palliata*, *Cebus apella*, and  
25  
26 *Lophocebus albigena*). Results reaffirm that microwear patterns on “Phase II” facets  
27  
28 better distinguish taxa by diet than do those on “Phase I” facets. Further, differences in  
29  
30 microwear textures between facet types for a given taxon may themselves reflect diet.  
31  
32 Some possible explanations for differences in microwear textures between facet types are  
33  
34 proposed.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## INTRODUCTION

The power stroke of mastication involves the application of forces to food particles between the teeth. It has traditionally been divided into two intervals: “Phase I” and “Phase II”, which immediately precede and follow centric occlusion respectively (Butler, 1952; Butler and Mills, 1959; Hiiemae and Kay, 1972; Kay and Hiiemae, 1974). Wear facets produced along the surfaces leading away from the crests as the teeth move upwards and lingually into centric occlusion were identified with “Phase I” of mastication. Wear facets produced in the tooth basins as the teeth come into full centric occlusion and then are moved parallel to the occluding surfaces were associated with “Phase II”.

However, the importance of “Phase II” to chewing has since been called into question as, at least for some primates, minimal muscle activity and jaw bone strain occur following centric occlusion. This suggests that little food processing is done once the lower teeth begin to separate from the uppers (Hylander and Crompton, 1980; Wall et al., 2006). Nevertheless, it is precisely the “Phase II” facets that are usually used in primate dental microwear analysis. This stems from research conducted in the early 1980s when microwear was first compared between facet types. These studies demonstrated that “Phase I” and “Phase II” facets could have differing patterns of microwear, even on an individual tooth. Thus, control over sampling area on an occlusal surface was deemed key to interpreting results. These analyses also showed that the sizes and shapes of microwear features that dominated “Phase II” facets separated primates with differing diets in expected, predictable ways (Gordon, 1982; Teaford and Walker, 1984).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Observations that patterns of microwear features on “Phase II” facets distinguish primates with differing diets, but that minimal bone strain and muscle activity occur during “Phase II” of the power stroke present a conundrum. If the work of mastication is not being done during “Phase II”, why are the facets associated with it considered to be the best for distinguishing microwear patterns caused by different diets? This paper reexamines the use of specific facets for primate microwear research in light of new understandings of the biomechanics of chewing and new approaches to the quantitative characterization and comparison of dental microwear. Results confirm that “Phase II” facets generally distinguish primates with differing diets better than “Phase I” facets. They also suggest that differences between facet types for a given taxon can themselves be of value for inferring diet from microwear textures.

## MATERIALS AND METHODS

We examined microwear on “Phase I” and “Phase II” facets of the M<sub>1S</sub> and M<sub>2S</sub> of 36 individuals, consisting of *Alouatta palliata* ( $n = 10$ ), *Cebus apella* ( $n = 13$ ) and *Lophocebus albigena* ( $n = 13$ ). *Alouatta palliata* is usually considered a tough food eater, consuming mostly leaves, stems, and fruit seeds and flesh. *Cebus apella* and *Lophocebus albigena* are soft-fruit eaters that fallback on hard, brittle foods such as nuts and palm fronds (Chalmers, 1968; Estrada, 1984; Estrada and Coates-Estrada, 1986; Lambert et al., 2004; Teaford, 1985). All specimens were wild-caught, with originals housed at the US National Museum of Natural History (*A. palliata* and *C. apella*) and in the Tappen Collection at the University of Minnesota Department of Anthropology (*L. albigena*).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

High-resolution replicas were produced following standard microwear molding and casting procedures (Grine, 1986), and surfaces were analyzed using published dental microwear texture analysis protocols (see Scott et al., 2006; Ungar et al., 2008). “Phase I” (5 or 7n) and “Phase II” (9, 10n or x) facets were scanned using a Sensofar Pl $\mu$  white-light scanning confocal profiler (Solarius Development Inc., Sunnyvale, CA) with a 100x objective lens. Point clouds were generated for each surface examined with a lateral sampling interval of 0.18  $\mu\text{m}$  and a vertical resolution of 0.005  $\mu\text{m}$ . Four adjoining fields of view were scanned, for a total work area of 276 x 204  $\mu\text{m}$  (Fig. 1). Scan data were then normalized and leveled using Solarmap Universal software (Solarius Development Inc., Sunnyvale, CA), and any identifiable defects, such as dust particles, were deleted from the dataset prior to analysis.

The four adjoining scans of each specimen were analyzed using Toothfrax (Surfract, [www.surfract.com](http://www.surfract.com)) and SFrax scale-sensitive fractal analysis (SSFA) software packages. Scale-sensitive fractal analysis operates on the principle that the apparent length of a profile from a rough surface, the apparent area of that surface, and its apparent volume change with scale of observation. Surface textures that appear smooth at a coarse scale can be rough at a finer scale. Scale-sensitive fractal analysis is valuable as it can be applied to three-dimensional surfaces, allowing for length, area, and volume scale analyses.

Six texture variables were analyzed for this study following previous analyses: complexity ( $Asfc$ ), anisotropy ( $epLsar$ ), scale of maximum complexity ( $Smc$ ), textural fill volume ( $Tfv$ ), and heterogeneity dividing each scan into 3x3 ( $HA_{sfc_9}$ ) and 9x9 ( $HA_{sfc_{81}}$ )

1  
2  
3 rows and columns. These attributes are described in detail in Scott et al. (2006) and  
4  
5 Ungar et al. (2008) and summarized in Table 1.  
6  
7

8 Data were collected for individual fields, and median values for each variable  
9  
10 were calculated for each surface analyzed. These were rank-transformed to mitigate  
11  
12 violation of assumptions inherent in parametric statistical analyses (Conover and Iman,  
13  
14 1981). Data for the variables were compared among species and facet type using a two-  
15  
16 factor multivariate analysis of variance model, with taxon and facet type as the factors,  
17  
18 the individual texture attributes (*Asfc*, *Smc*, *epLsar*, *Tfv*, *HAsfc<sub>9</sub>*, and *HAsfc<sub>8I</sub>*) as the  
19  
20 dependent variables, and values for each individual as the replicates. This model assesses  
21  
22 significance of variation among taxa, between facet types, and interactions between the  
23  
24 two factors in overall microwear surface textures. Factorial ANOVAs for each variable  
25  
26 (again, with taxon and facet type as the factors) were used to identify those attributes that  
27  
28 evinced significant variation between taxa, facet type, or an interaction between the two.  
29  
30 Multiple comparisons tests were then used to determine the sources of significant  
31  
32 variation. Given that these groups were chosen for the dietary (and expected microwear)  
33  
34 differences, Fisher's LSD *a priori* tests were used to compare species. Tukey's HSD *post*  
35  
36 *hoc* tests were also run to balance the risks of Type I and Type II errors (Cook and  
37  
38 Farewell, 1996).  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 48 RESULTS & DISCUSSION

49 The results are presented in Figures 1-2 and Tables 2-4. We found significant  
50  
51 variation between taxa and phases in overall microwear surface texture, as well as an  
52  
53 interaction between the two factors. The sources of these variations are identified in  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 individual ANOVAs for each variable. First, there was significant variation between  
4  
5 taxa, but no interaction between taxon and facet type for complexity (*Asfc*) and scale of  
6  
7 maximum complexity (*Smc*). In other words, the three taxa differed from one another in  
8  
9 *Asfc* and *Smc*, and those differences were independent of whether “Phase I” or “Phase II”  
10  
11 facets were examined. Significant variation was also found between taxa for textural fill  
12  
13 volume (*Tfv*) and heterogeneity 9x9 (*HAsfc<sub>81</sub>*) though a significant interaction between  
14  
15 taxon and facet type for these two texture variables required separate consideration of  
16  
17 taxa by facet type.  
18  
19  
20  
21

22 The facet types differed significantly from one another in complexity (*Asfc*) and  
23  
24 anisotropy (*epLsar*) in similar ways for the three taxa considered. “Phase II” facets were  
25  
26 both more complex and more anisotropic than were “Phase I” facets for the samples  
27  
28 examined. Fill volume (*Tfv*) also showed significant variation between facet types;  
29  
30 however, as with taxon, a significant interaction required further statistical analyses.  
31  
32  
33

34 Significant interactions between factors necessitated separate comparisons of  
35  
36 facet types by each taxon for *Tfv*, *HAsfc<sub>9</sub>*, and *HAsfc<sub>81</sub>*. Individual results indicated that  
37  
38 *C. apella* and *L. albigena* both had higher *Tfv* values on their “Phase II” facets, and that  
39  
40 *C. apella* had more heterogeneous “Phase II” microwear surface textures. Further, *A.*  
41  
42 *palliata* demonstrated no significant differences in these three variables between facets.  
43  
44 These differences between the facet types in complexity and anisotropy presumably  
45  
46 reflect differences in tooth-food interactions before and after centric occlusion, and  
47  
48 support earlier suggestions that facet types should not be mixed in microwear analyses  
49  
50  
51  
52  
53 (Gordon, 1982; Gordon & Walker, 1983).  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The interactions are especially interesting because they suggest that different taxa  
4 have different patterns of variation between “Phase I” and “Phase II” microwear textures.  
5 Specifically, while the more folivorous *A. palliata* had no significant difference between  
6 “Phase I” and “Phase II” facets in *Tfv*, the two “hard-object” fallback species had higher  
7 *Tfv* values on Phase II facets. Further, *C. apella* had more heterogeneous “Phase II”  
8 texture features than did any of the other taxa.  
9

10  
11 Further statistical analysis of the significant interactions of taxon and phase  
12 suggests that “Phase II” facets better distinguish taxa by diet than do “Phase I” facets.  
13 “Phase I” facets do not distinguish taxa for *Tfv*, *HAsfc<sub>9</sub>*, or *HAsfc<sub>81</sub>*. Conversely, all three  
14 of these texture attributes did differ significantly between species on “Phase II” facets.  
15 Although this offers additional support for the use of “Phase II” facets in microwear  
16 studies, it still does not provide a biomechanical explanation for this trend. It begs the  
17 question “what is occurring during ‘Phase II’ of the power stroke that accounts for its  
18 reliability in dental microwear research?”  
19

20  
21 Several possible reasons can be suggested to explain why wear patterns on “Phase  
22 I” and “Phase II” surfaces differ. These differences may relate to the volume of food and  
23 associated abrasives interposed between surfaces as they contact one another.  
24 Alternatively, differences may relate to the angle of masticatory forces relative to the  
25 planes of the wear facets themselves. In the first case, the most sharply delineated facets  
26 of “Phase I” are most likely formed at a stage in the masticatory process when few food  
27 particles are interposed between the teeth. The prevalence of tooth-tooth contact  
28 occurring later in a masticatory cycle might obliterate or obscure wear features produced  
29 more specifically by the food or the grit adhering to it (*i.e.*, in the earlier puncture-  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 crushing stages of mastication). In contrast, “Phase II” facets occur in the trigonid and  
4 talonid basins and are produced, at least in part, at the end of a “Phase I” movement as  
5 food is compressed between relatively flat opposing surfaces.  
6  
7  
8  
9

10 Even in the later stages of a masticatory cycle, one might expect to find food  
11 particles and abrasives trapped in the trigonid and talonid basins to still have a role in  
12 producing microwear features. In the second case, “Phase I” facets are produced largely  
13 at a time when the anatomy of the teeth dictates when the most precise occlusion should  
14 occur. Because of the way the “Phase I” occlusal surfaces are arranged, the masticatory  
15 forces are concentrated at the leading edges of the surfaces and likely oriented at an acute  
16 angle to the surfaces. “Phase II” movement and the accompanying facets it produces are  
17 less rigidly constrained by the anatomy of the teeth. It would not be surprising to find  
18 more complexity and anisotropy on facets produced by a less precise part of the occlusal  
19 cycle (occlusion produced by forces more-or-less normal to the plane of the facets).  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 In summary, the use of dental microwear texture analysis in this study confirms  
35 that “Phase II” facets better distinguish primates by diet than “Phase I” facets. This study  
36 also suggests that distinct biomechanical events leading to differences in dental  
37 microwear occur during “Phase I” and “Phase II”. Moreover, the variation among  
38 species in degree of difference between textures on the two facets suggests that  
39 comparisons of patterns of microwear across a tooth may yield even more information  
40 about diet than we can obtain using “Phase II” facets alone. Further studies including  
41 larger samples of primates with other diets will help us gain a better understanding of this  
42 phenomenon.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ACKNOWLEDGEMENTS

We are grateful to Mark Teaford for help collecting dental impressions used in this study, and Robert Scott, for generating the “Phase II” microwear data, which was first published in Scott et al. (2006) and Ungar et al. (2008). We also thank curators at the US National Museum of Natural History, and Martha Tappen for their permission to study specimens in their care. This study was funded by the US National Science Foundation (BCS 0315157).

## LITERATURE CITED

1  
2  
3  
4  
5  
6  
7  
8 Butler PM. 1952. The milk molars of the Perissodactyla, with remarks on molar  
9 occlusion. Proc Zool Soc Lond 121:777-817.  
10

11  
12  
13  
14  
15 Butler PM, and Mills JRE. 1959. A contribution to the odontology of *Oreopithecus*. Bull.  
16 Brit. Mus. Nat. Hist. Geol. 4:1-26.  
17

18  
19  
20  
21  
22 Conover WJ, and Iman RL. 1981. Rank transformations as a bridge between parametric  
23 and nonparametric statistics. Am Stat 35:124-129.  
24

25  
26  
27  
28  
29 Cook RJ, and Farewell VT. 1996. Multiplicity considerations in the design and analysis  
30 of clinical trials. Journal of the Royal Statistical Society Series A 159:93-110.  
31

32  
33  
34  
35  
36 Chalmers NR. 1968. Group composition, ecology and daily activities of free-living  
37 mangabeys in Uganda. Folia Primatol 8:247-262.  
38

39  
40  
41  
42  
43 Estrada A. 1984. Resource use by howler monkeys (*Alouatta palliata*) in the rain forest  
44 of Los Tuxtlas, Veracruz, Mexico. Int J Primatol 5:105-131.  
45

46  
47  
48  
49  
50 Estrada A, and Coates-Estrada RA. 1986. Use of leaf resources by howling monkeys  
51 (*Alouatta palliata*) and leaf-cutting ants *Atta cephalotes*) in the tropical rain forest of Los  
52 Tuxtlas, Mexico. Am J Primatol 10:51-66.  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 Gordon KD. 1982. A study of microwear on chimpanzee molars: implications of dental  
7  
8 microwear analysis. *Am J Phys Anthropol* 59:195-215.  
9

10  
11  
12 Gordon KD, and Walker AC. 1983. Playing ‘possum: a microwear experiment. *Am J*  
13  
14 *Phys Anthropol* 60:109-112.  
15  
16

17  
18  
19  
20 Grine FE. 1986. Dental evidence for dietary differences in *Australopithecus* and  
21  
22 *Paranthropus*. *J Hum Evol* 15:783-822.  
23  
24

25  
26  
27 Hiiemae K, and Kay RF. 1972. Trends in the evolution of primate mastication. *Nature*  
28  
29 240:486-487.  
30  
31

32  
33  
34 Hylander WL, and Crompton AW. 1980. Loading patterns and jaw movement during the  
35  
36 masticatory power stroke in macaques. *Am J Phys Anthropol* 52:239.  
37  
38

39  
40  
41 Kay RF, and Hiiemae K. 1974. Jaw movement and tooth use in recent and fossil  
42  
43 primates: *J Hum Evol* 40:227–256.  
44  
45

46  
47  
48 Kay RF, and Covert HH. 1983. True grit: a microwear experiment. *Am J Phys Anthropol*  
49  
50 61:33-38.  
51  
52

1  
2  
3 Lambert JE, Chapman CA, Wrangham RW, and Conklin-Brittain NL. 2004. The  
4  
5 hardness of Cercopithecine foods: implications for the critical function of enamel  
6  
7 thickness in exploiting fallback foods. *Am J Phys Anthropol* 214:363-368.  
8  
9

10  
11  
12 Scott RS, Ungar PS, Bergstrom TS, Brown CA, Childs BE, Teaford MF, and Walker A.  
13  
14 2006. Dental microwear texture analysis: technical considerations. *J Hum Evol* 51:339-  
15  
16 349.  
17  
18  
19

20  
21  
22 Teaford MF, and Walker AC. 1984. Quantitative differences in dental microwear  
23  
24 between primate species with different diets and a comment on the presumed diet of  
25  
26 *Sivapithecus*. *Am J Phys Anthropol* 64:191-200.  
27  
28  
29

30  
31  
32 Teaford MF. 1985. Molar microwear and diet in the genus *Cebus*. *Am J Phys Anthropol*  
33  
34 66:363-370.  
35  
36  
37

38  
39 Ungar PS, Scott RS, Scott JR, and Teaford MF. 2008. Dental microwear analysis:  
40  
41 historical perspectives and new approaches. In: Irish JD, Nelson GC, editors. *Dental*  
42  
43 *anthropology*. Cambridge: Cambridge University Press. p 389-425.  
44  
45  
46

47  
48 Wall CE, Vinyard CJ, Johnson KR, Williams SH, and Hylander WL. 2006. Phase II jaw  
49  
50 movements and masseter muscle activity during chewing in *Papio anubis*. *Am J Phys*  
51  
52 *Anthropol* 129:215-224.  
53  
54  
55  
56  
57  
58  
59  
60

## FIGURE LEGENDS

Figure 1. Photosimulations of “Phase I” and “Phase II” surfaces generated from elevation data: A) *Alouatta palliata*, B) *Cebus apella*, C) *Lophocebus albigena*. Each image represents a surface 276  $\mu\text{m}$  x 204  $\mu\text{m}$ .

Figure 2. Dental microwear texture attributes by species and facet type. Species and variables are as indicated on x- and y-axes. Black and white bars indicate “Phase I” and “Phase II” facets respectively.



Table 1: Dental microwear texture variables, definitions, and examples.

<b>Texture Variable</b>	<b>Definition</b>	<b>Example</b>
<i>Asfc</i>	Complexity	Pits and scratches of different sizes overlaying one another would present a complex surface.
<i>Smc</i>	Scale of Maximum Complexity	A surface dominated by large pits with an absence of fine scratches might have a high <i>Smc</i> .
<i>epLsar</i>	Anisotropy	A surface dominated by scratches all running in the same direction would have a high <i>epLsar</i> .
<i>Tfv</i>	Textural Fill Volume	A surface dominated by deep features would have a high <i>Tfv</i> .
<i>HAsfc</i>	Heterogeneity	High heterogeneity indicates differing patterns of scratching and pitting across a microwear surface.

Table 2. Statistical Analyses. Overall comparisons.

- A. MANOVA results. Variables considered include *Asfc*, *epLsar*, *Smc*, *Tfv*, *HAsfc<sub>9</sub>*, and *HAsfc<sub>81</sub>*. This analysis was conducted on ranked data.

Effect	Wilk's $\lambda$	<i>df</i>	<i>F</i>	<i>p</i>
Taxon	0.340	12, 122	7.276	0.000
Facet type	0.562	6, 61	7.912	0.000
Interaction	0.697	12, 122	2.009	0.029

- B. Univariate test results.

Variable	Taxon		Phase		Interaction	
	<i>df</i>	<i>F</i>	<i>df</i>	<i>F</i>	<i>df</i>	<i>F</i>
<i>Asfc</i>	2	31.701*	1	7.338*	2	3.003
<i>epLsar</i>	2	2.263	1	4.333*	2	1.329
<i>Smc</i>	2	3.286*	1	1.095	2	0.744
<i>Tfv</i>	2	8.025*	1	22.447*	2	5.279*
<i>HAsfc<sub>9</sub></i>	2	0.820	1	2.679	2	4.363*
<i>HAsfc<sub>81</sub></i>	2	3.578*	1	3.182	2	4.242*

---

\**p* < 0.05

Table 3. Tests for differences between facet types by taxon, and between taxa by facet types.

## A. ANOVA results for differences between facet types by taxon.

Variable	<i>A. palliata</i>		<i>C. apella</i>		<i>L. albigena</i>	
	<i>df</i>	<i>F</i>	<i>df</i>	<i>F</i>	<i>df</i>	<i>F</i>
<i>Tfv</i>	1	0.021	1	7.526*	1	45.117*
<i>HAsfc<sub>9</sub></i>	1	0.092	1	15.522*	1	0.386
<i>HAsfc<sub>8I</sub></i>	1	0.217	1	16.536*	1	0.384

## B. ANOVA results for differences between taxa by facet type.

Variable	Phase I		Phase II	
	<i>df</i>	<i>F</i>	<i>df</i>	<i>F</i>
<i>Tfv</i>	2	1.025	2	15.047*
<i>HAsfc<sub>9</sub></i>	2	0.828	2	3.966*
<i>HAsfc<sub>8I</sub></i>	2	0.115	2	7.339*

---

\* $p < 0.05$

Table 4. Pairwise comparisons

A. <i>Asfc</i> (Independent of Phase)			
	<i>A. palliata</i>	<i>C. apella</i>	<i>L. albigena</i>
<i>A. palliata</i>	-		
<i>C. apella</i>	33.325 <sup>a,b</sup>	-	
<i>L. albigena</i>	27.113 <sup>a,b</sup>	-6.212	-
B. <i>Smc</i> (Independent of Phase)			
	<i>A. palliata</i>	<i>C. apella</i>	<i>L. albigena</i>
<i>A. palliata</i>	-		
<i>C. apella</i>	9.671	-	
<i>L. albigena</i>	15.460 <sup>a,b</sup>	5.788	-
C. <i>Tfv</i> (Phase II)			
	<i>A. palliata</i>	<i>C. apella</i>	<i>L. albigena</i>
<i>A. palliata</i>	-		
<i>C. apella</i>	28.504 <sup>a,b</sup>	-	
<i>L. albigena</i>	29.696 <sup>a,b</sup>	1.192	-
D. <i>HAsfc<sub>9</sub></i> (Phase II)			
	<i>A. palliata</i>	<i>C. apella</i>	<i>L. albigena</i>
<i>A. palliata</i>	-		
<i>C. apella</i>	17.523	-	
<i>L. albigena</i>	-4.631	-22.154 <sup>a,b</sup>	-
E. <i>HAsfc<sub>81</sub></i> (Phase II)			
	<i>A. palliata</i>	<i>C. apella</i>	<i>L. albigena</i>
<i>A. palliata</i>	-		
<i>C. apella</i>	17.227 <sup>b</sup>	-	
<i>L. albigena</i>	-12.081	-29.308 <sup>a,b</sup>	-

<sup>a</sup>Tukey's HSD test  $p \leq 0.05$ <sup>b</sup>Fisher's LSD test  $p \leq 0.05$