



2014

# Peripubertal Binge Ethanol, Age and Sex Modulate MicroRNA Expression in the Ventral and Dorsal Hippocampus of the Adolescent Rat

Sarah Arianne Prins  
*Loyola University Chicago*

## Recommended Citation

Prins, Sarah Arianne, "Peripubertal Binge Ethanol, Age and Sex Modulate MicroRNA Expression in the Ventral and Dorsal Hippocampus of the Adolescent Rat" (2014). *Dissertations*. Paper 1297.  
[http://ecommons.luc.edu/luc\\_diss/1297](http://ecommons.luc.edu/luc_diss/1297)

This Dissertation is brought to you for free and open access by the Theses and Dissertations at Loyola eCommons. It has been accepted for inclusion in Dissertations by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 3.0 License](https://creativecommons.org/licenses/by-nc-nd/3.0/).  
Copyright © 2014 Sarah Arianne Prins

LOYOLA UNIVERSITY CHICAGO

PERIPUBERTAL BINGE ETHANOL, AGE AND SEX MODULATE  
MICRORNA EXPRESSION IN THE VENTRAL AND DORSAL HIPPOCAMPUS  
OF THE ADOLESCENT RAT

A DISSERTATION SUBMITTED TO  
THE FACULTY OF THE GRADUATE SCHOOL  
IN CANDIDACY FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY

PROGRAM IN CELL BIOLOGY, NEUROBIOLOGY AND ANATOMY

BY  
SARAH A. PRINS  
CHICAGO, IL  
DECEMBER 2014

Copyright by Sarah Prins, 2014  
All rights reserved.

## ACKNOWLEDGEMENTS

I would like to acknowledge first, Dr. Toni Pak, for her encouragement to join the Cell Biology, Neurobiology and Anatomy (CBNA) PhD program at Loyola University Chicago, for accepting me in her laboratory as a graduate student and for supporting me through completion of my dissertation. I would also like to acknowledge my dissertation committee for their support and guidance, which includes Dr. Pieter de Tombe, Dr. Gwendolyn Kartje, Dr. Evan Stubbs, and Dr. Sakthivel Sadayappan. In addition, many professors have provided invaluable mentorship to me including Dr. Phong Le, Dr. Elizabeth Kovacs and Dr. John Callaci. Lastly, I would like to collectively acknowledge the following elements of Loyola University Chicago for contributing to my graduate education: the CBNA graduate program, the Alcohol Research Program, the Neuroscience Research Institute, the Department of Cell and Molecular Physiology and all professors which instructed my courses.

I have particular gratitude for the sources which contributed funds during my training which includes Dr. Toni Pak, Dr. Elizabeth Kovacs, Dr. Pieter de Tombe, Loyola University Chicago, the National Institute on Alcohol Abuse and Alcoholism, and Loyola's Neuroscience Research Institute.

Acknowledging my friends and family for their support goes without saying, yet, a special acknowledgment is owing to my partner, Jonathan Kulpit, for continuously and entertainingly teaching and motivating me.

To Peter, Stephen and Matthew  
For value, curiosity and balance

## TABLE OF CONTENTS

<b>Acknowledgements</b>	i
<b>List of Tables</b>	vii
<b>List of Figures</b>	viii
<b>Abstract</b>	x
<b>CHAPTER ONE: STATEMENT OF THE PROBLEM</b>	1
Introduction	1
Hypothesis and Specific Aims	2
<b>CHAPTER TWO: LITERATURE REVIEW</b>	11
Adolescence	11
Propensity for Binge Drinking	11
Alcohol Alters Neurodevelopment	12
Dysregulation of the Hypothalamic-Pituitary-Adrenal Axis	12
Hippocampus Malfunctioning: Ventral and Dorsal Perspectives	13
Hippocampal Brain-Derived Neurotrophic Factor Role in Development	15
Hippocampal Sirtuin 1 Role in Development	16
microRNAs	17
What are microRNAs?	17
microRNA Biogenesis	18
microRNAs in Neuronal Development	19
Alcohol effects on microRNA	21
Gonadal hormone effects on microRNA	23
Summary	24
<b>CHAPTER THREE: LONG-TERM EFFECTS OF PERIPUBERTAL BINGE ETOH EXPOSURE ON HIPPOCAMPAL MICRORNA EXPRESSION IN THE RAT</b>	25
Introduction	25
Approach	28
Results	29
Discussion	54
<b>CHAPTER FOUR: SEXUALLY DIMORPHIC GENE EXPRESSION OF HIPPOCAMPUS ETOH-SENSITIVE MICRORNA, MICRORNA TARGET GENES AND MICRORNA PROCESSING ENZYMES DURING PUBERTAL DEVELOPMENT</b>	61
Introduction	61
Approach	63
Results	64
Discussion	75

<b>CHAPTER FIVE: SEX DIFFERENCES IN HIPPOCAMPUS 17<math>\beta</math>-ESTRADIOL-RESPONSIVE MICRORNA EXPRESSION DURING PUBERTAL DEVELOPMENT OF THE WISTAR RAT</b>	84
Introduction	84
Approach	86
Results	87
Discussion	92
<b>CHAPTER SIX: GENERAL DISCUSSION</b>	99
Summary	99
Key Findings	100
Final Remarks	105
Selection of Model Organism	105
microRNA Functions in the Brain: Relevance to Neuronal Development	106
The Role of microRNAs in Ethanol Neurotoxicity During Development	108
EtOH-Induced Changes in Hippocampal microRNA Correlate with Altered BDNF and SIRT1 mRNA: Proposed Role of Ethanol Metabolism	113
Possible Role for Gonadal Hormones in Mediating Sexually Dimorphic Pubertal microRNA Expression	116
Take Home Message	125
Future Directions	126
<b>CHAPTER SEVEN: GENERAL METHODS</b>	127
Animals	127
Ethics Statement	128
Binge Exposure Paradigm and Treatment Design	128
Tissue Collection	129
RNA Isolation	130
Rat RT <sup>2</sup> microRNA PCR Array	130
Quantitative Reverse Transcription PCR (qRT-PCR)	131
Hormone Measurements	133
Blood Alcohol Concentration Assay	133
Western Blot	134
Statistical Analysis	135
<b>Reference List</b>	137
<b>Vita</b>	170

## LIST OF TABLES

Table	Page
1. Statistical Analysis of Gene Expression Levels in the Dorsal Hippocampus	36
2. Statistical Analysis of Gene Expression Levels in the Ventral Hippocampus	40
3. Summary of Peripubertal EtOH Exposure on microRNA and mRNA Expression	43
4. Statistical Analysis of Dorsal Hippocampus Gene Expression	65
5. Statistical Analysis of Ventral Hippocampus Gene Expression	69
6. Statistical Analysis of microRNA Expression Levels in the Dorsal Hippocampus	89
7. Statistical Analysis of microRNA Expression Levels in the Ventral Hippocampus	93



## LIST OF FIGURES

Figure	Page
1. Peripubertal repeated binge alcohol paradigm alters blood alcohol concentration	30
2. Peripubertal binge EtOH increases miR-10a-5p and miR-423 expression in the male rat whole hippocampus	32
3. Peripubertal binge-pattern EtOH exposure alters microRNA and target gene expression in male rat hippocampus	33
4. Diagram of experimental paradigm	35
5. Peripubertal binge EtOH exposure alters microRNA expression during pubertal development in the dorsal hippocampus	37
6. Peripubertal binge EtOH alters microRNA expression during pubertal development in the ventral hippocampus	41
7. Peripubertal binge EtOH exposure differentially alters microRNA biosynthetic processing enzymes in the dorsal and ventral hippocampus	45
8. Diagram depicting predicted microRNA binding sites for BDNF and SIRT1	48
9. Peripubertal binge EtOH exposure differentially alters microRNA target genes, BDNF and SIRT1, in the dorsal and ventral hippocampus	49
10. Assessment BDNF and SIRT1 protein expression levels following peripubertal binge EtOH	52
11. Peri-pubertal binge EtOH exposure did not affect circulating testosterone levels	53
12. Sexually dimorphic microRNA expression is evident during pubertal development in the dorsal hippocampus	66
13. Sexually dimorphic microRNA expression is evident during pubertal development in the ventral hippocampus	70

14. Sexually dimorphic microRNA processing enzyme expression is evident during pubertal development in the dorsal and ventral hippocampus	72
15. Sexually dimorphic microRNA target gene expression is evident during pubertal development in the dorsal and ventral hippocampus	74
16. Diagram of experimental paradigm. Diagram depicting the age of sacrifice for each group of male and female Wistar rats	88
17. Sexually dimorphic microRNA expression is evident during pubertal development in the dorsal hippocampus	91
18. Sexually dimorphic microRNA expression is evident during pubertal development in the ventral hippocampus	94

## ABSTRACT

Adolescent binge ethanol (EtOH) abuse induces long-term changes in gene expression, resulting in an increased risk for the development of adult mood disorders. microRNAs (microRNAs) are small, noncoding RNAs that regulate gene expression by translational repression. microRNAs are implicated as important contributors to the neural mechanisms underlying alcohol addiction and are also altered in response to EtOH in the developing brain and can be expressed in sexually dimorphic manners. The biogenesis of mature 22-24 nucleotide (nt), single-stranded microRNAs involves 1) transcription of a 100-1000 nt primary-microRNA (pri-microRNA) product, 2) cropping of the transcript by the enzyme Drosha into the preliminary-microRNA (pre-microRNA), 3) cleaving of the pre-microRNA by the enzyme Dicer and 4) loading of the mature microRNA onto the RNA induced silencing complex (RISC). The microRNA/RISC complex (miRISC) is guided to the microRNA target gene and induces either cleavage or translational repression of the messenger RNA (mRNA) transcript. Our lab has found that repeated binge EtOH exposure alters gene expression in the hypothalamus and dysregulates the Hypothalamic-Pituitary-Adrenal (HPA) Axis in a sexually dimorphic, long-term and gonadal hormone-dependent manner in Wistar rats, and that a subset of microRNAs targeting brain-derived neurotrophic factor (BDNF) and sirtuin 1 (SIRT1) are differentially expressed in the ventral hippocampus dependent on age and sex. We examine whether microRNA expression, microRNA biogenesis enzymes Drosha and Dicer, and microRNA target genes BDNF and SIRT1 are altered by peripubertal binge

EtOH in the ventral and dorsal hippocampus. We also document sex differences in the expression of microRNAs sensitive to EtOH and 17 $\beta$ -estradiol during pubertal development. Overall, we demonstrate : 1) peripubertal binge EtOH exposure induces long-term alterations in mature microRNA expression levels in the male rat hippocampus, and has the potential to modulate the expression of their downstream target genes, 2) expression profiles of EtOH-sensitive microRNAs, miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495, and their target genes, are dependent on sex and age in the pubertal rat hippocampus and therefore may contribute to sexually dimorphic hippocampus neurodevelopment, 3) expression profiles of E<sub>2</sub>-responsive microRNAs, miR-7a, miR-9, miR-125a and miR-181a, are differentially dependent on sex and age throughout pubertal development, suggesting that they play distinct developmental roles during puberty and 4) EtOH-sensitive and E<sub>2</sub>-responsive microRNAs have distinct expression profiles in the dorsal and ventral hippocampus throughout pubertal development, suggesting that their respective functions are region-specific. This research increases our understanding of how pubertal binge EtOH exposure affects microRNA expression, provides evidence that microRNA are expressed in sexually dimorphic patterns throughout pubertal development, and suggest that microRNAs play a role in normal pubertal hippocampus development as well as hippocampus dysfunction following adolescent alcohol abuse.

## CHAPTER ONE

### STATEMENT OF THE PROBLEM

#### **Introduction**

A division of the Department of Health and Human Services, the Substance Abuse and Mental Health Services Administration (SAMHSA) has documented that alcohol is the most highly abused drug amongst adolescents, that binge drinking begins and peaks during puberty and that underage drinkers have the highest rates of alcohol dependence (SAMHSA 2000, 2002, 2003, 2008, 2009). Such reports led the U.S. Department of Health and Human Services Office of the Surgeon General to announce a call to action aiming to prevent and reduce underage drinking. Despite these efforts, more current reports from the Centers for Disease Control and Prevention (CDC) indicate that binge drinking is an even bigger problem than previously thought and that it is particularly serious and yet under-recognized among women and girls (CDC 2012, 2013). Indeed, binge drinking during puberty puts adolescents at an increased risk for neurological complications later in life [1,2,3,4,5,6,7]. For example, adolescent alcohol abuse has long-term consequences on many neurological systems responsible for learning, memory and the regulation of the stress response [8,9,10,11,12]. Indeed, research demonstrates that the pubertal brain continues to develop well into the twenties [13]. Adolescent brain development is characterized by a remodeling of neural networks established in the perinatal period [14,15,16] and the generation of new connections,

which together, refine the neural control of behavior [17]. Yet, despite our increased understanding of how the adolescent brain develops and how alcohol abuse alters this development, many questions remain about what regulates alcohol abuse-induced brain damage, particularly with respect to gene expression regulation and possible sex differences.

### **Hypothesis and Specific Aims**

Pubertal brain development is characterized, in part, by sexually dimorphic alterations in physiology and behavior and is predominantly thought to be regulated in a sex biased manner, yet, precise sex-specific mechanisms of gene regulation remain largely elusive. A major regulator of gene expression includes microRNAs (microRNAs), which are small, noncoding RNA molecules that target and bind, via imperfect base pairing, to protein-coding messenger RNA (mRNA) transcripts and lead to mRNA degradation or translational repression. Despite the relatively short time during which microRNAs have been researched (~12 years), they are currently recognized as regulators of at least 70% of fundamental biological processes [18,19]. Development of the pubertal brain is a dynamic biological process that involves sexually dimorphic anatomical changes in grey and white matter and cellular strengthening of relevant neuronal networks. Therefore, as master regulators of gene expression, microRNAs may play a role in orchestrating the complex molecular adaptations taking place during adolescent brain development. In this way, microRNAs may regulate the cellular, anatomical and behavioral transformations manifested throughout this significant stage of brain development.

Typical adolescent behaviors are often “high-risk,” such as binge drinking, and moreover, can manifest in a sex-specific manner. Alcohol abuse induces particular morphological, cellular and molecular damage to the hippocampus, which plays a pivotal role in brain development, as it is the primary source of adult neurogenesis, spatial and emotional memory, and is a major contributor to the regulation of mood. Importantly, microRNAs have been linked to alcohol-induced neurological afflictions including addiction and fetal alcohol spectrum disorder (FASD) and in addition, demonstrate age, cell type and sex-specific expression profiles in multiple species. However, the normal developmental expression profiles of microRNAs during puberty and how peri-pubertal alcohol abuse alters these profiles, remains undocumented. Using multiple target prediction algorithms, four microRNAs (miR-10a-5p, miR-26a, miR-103 and miR-495) were identified to target the 3'UTR of two genes important for hippocampus function and development, *BDNF* and *SIRT1*. Importantly, miR-26a and miR-103 are in the top 15 most highly expressed microRNAs in the rodent hippocampus, miR-10a is important for developmental processes as it regulates homeobox (Hox) gene expression [20],[21] and miR-495 regulates the neuroprotective effects of mood stabilizing drugs [22]. Therefore, I generated the following hypothesis that *adolescent hippocampal microRNA expression throughout pubertal development is sex biased, dependent on age, and altered by peri-pubertal binge EtOH exposure.* The following specific aims were developed to test this hypothesis:

**AIM 1:** Determine whether peripubertal binge EtOH alters expression of microRNA, microRNA processing enzymes and microRNA target genes throughout pubertal development.

Pubertal binge drinking induces long-term changes in the expression of genes that regulate the stress response. An altered stress response often underlies depression and anxiety-related disorders. Importantly, these conditions are commonly experienced by over 50% of alcohol-dependent patients and present in a sexually-dimorphic fashion. Sex biased mood and memory impairments are often present in tandem with alcohol abuse and neuropsychiatric illnesses that emerge post-puberty. Disruption of mature microRNA expression and/or function has been linked to alcohol-induced neurological afflictions including addiction and FASD.

The following questions were answered in this aim:

- 1) Is the normal expression profile of miR-10a-5p, miR-26a, miR-103 and miR-495 during pubertal development altered in response to peri-pubertal binge EtOH in an immediate manner?
- 2) What is the long-term effect of peri-pubertal binge EtOH exposure on the expression profile of miR-10a-5p, miR-26a, miR-103 and miR-495?
- 3) What are the expression profiles of microRNA biochemical processing enzymes and the downstream target genes of miR-10a-5p, miR-26a, miR-103 and miR-495 (BDNF and SIRT1) throughout pubertal development immediately following mid/peri-pubertal binge EtOH exposure as well as 30 days post EtOH?

In order to answer these questions, males were chosen for the binge studies in order to avoid potential experimental design and data analysis complications due to the effects that cycling hormone levels in females would have on the results. Male Wistar rats were distributed into 3 groups: early pubertal (PND 30), peripubertal (PND 37), and



late pubertal (PND 73). The early, peripubertal and late pubertal age animals each had an untreated group (N=10/age group) which were sacrificed at PND 30, PND 44 and PND 73, respectively. In addition to the untreated groups at each age, there were 2 groups administered treatments at peripubertal age (N= 20/treatment group; total of 40 animals): peripubertal water (control) and peripubertal binge EtOH treated. The water and EtOH groups were handled for 5 minutes once/day beginning at PND 30 to eliminate non-specific effects of handling stress. Peripubertal water or EtOH treatments were treated with water or the following repeated binge alcohol (EtOH) paradigm via oral gavage beginning at PND 37 (3g/kg; 1x/day/3days EtOH, +1x/day/2days water + 1x/day/3days EtOH). Half of the peripubertal water/EtOH treated animals were sacrificed 60 minutes following the last EtOH/water treatment at PND 44 (N = 10 water + N = 10 EtOH). The remaining animals (N = 10 water + N = 10 EtOH) were left undisturbed following the last EtOH treatment in their home cage until sacrificed at late puberty (PND 73). Brains were sectioned at 200  $\mu$ m on a freezing microtome and ventral and dorsal hippocampi were microdissected using a 0.75 mm Palkovit's brainpunch tool. Total RNA was isolated and cDNA was made using Invitrogen's NCode microRNA First-strand cDNA synthesis kit.

In summary, the data from this aim demonstrate that the expression of each microRNA tested (miR-10a-5p, miR-26a, miR-103 and miR-495) is dynamic across pubertal development and that the developmental profiles for each microRNA are distinct between the dorsal and ventral hippocampus. Moreover, peripubertal binge EtOH exposure altered normal pubertal development expression patterns of miR-10a-5p, miR-26a, miR-103, miR-495, Dicer, Drosha, BDNF and SIRT1 in an age- and brain region-

dependent manner. Most striking, our results showed that peripubertal binge EtOH exposure had significant long lasting effects on several microRNAs studied, as well as their processing enzymes and target genes. These effects were evident for as long as one-month following the last EtOH exposure, suggesting that EtOH could have lasting consequences on gene expression profiles in the male rat hippocampus through long-term regulation of microRNA expression patterns.

**AIM 2: Determine whether there are sex and brain region differences in EtOH-sensitive microRNAs, microRNA processing enzymes and microRNA target gene expression throughout pubertal development**

Pubertal binge EtOH changes in expression of genes regulating the stress response in a sexually dimorphic manner. microRNAs regulate gene expression by targeting RNA transcripts and have recently been recognized as critical mediators of nearly all basic cellular processes. microRNAs are small, non-protein coding RNAs which are sequentially processed to their mature form by the enzymes Drosha and Dicer, which allows them to bind their complementary mRNA sequences and lead to the prevention of gene translation. microRNAs regulate neuronal development during embryogenesis, postnatal neuronal maintenance and survival, and hippocampal neurogenesis throughout life. Preliminary data generated using an RT-microRNA-PCR array identified EtOH-sensitive hippocampal microRNAs and five microRNAs were chosen for further analysis: miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495. Importantly, these microRNAs target mutual genes—BDNF and SIRT1—which regulate synaptic plasticity [23,24,25,26,27] an essential element of hippocampus-dependent memory and mood processing. Moreover, pubertal EtOH abuse can alter the expression

of genes important for mood regulation in a sexually dimorphic manner, and some brain microRNAs are expressed differentially between males and females, but the fundamental sex differences in microRNA expression throughout puberty for EtOH-sensitive microRNAs is unknown.

The following questions were answered in this aim:

- 1) Are miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495 expressed in sex, age and brain region-dependent manners?
- 2) In the dorsal and ventral hippocampus, are Drosha and Dicer expressed in sex, age, and brain region-dependent manners and do their expression profiles suggest that they regulate miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495 expression profiles throughout pubertal development?
- 3) Are BDNF and SIRT1 expressed in sex, age, and brain region-dependent manners and do their expression profiles suggest that miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495 regulate their expression profiles throughout pubertal development?

In order to answer these questions male and female Wistar rats at PND 30, PND 44 and PND 73 (N=10/group) were sacrificed by decapitation, brains were rapidly frozen and sectioned at 200  $\mu\text{m}$ , and the ventral and dorsal hippocampus were microdissected using a 0.75 mm Palkovit's brainpunch tool. RNA was isolated using TriZol according to manufacturers' instructions, and used to reverse transcribe microRNA (NCode™ microRNA First-Strand) and mRNA (SuperScript® VILO™) into cDNA. microRNA and mRNA expression levels were quantified using qRT-PCR with Fast Start Universal

SYBR Green Master Mix (Roche) and primers designed for the respective genes of interest.

In summary, data from this aim demonstrate that miR-10a-5p and miR-103 expression in females and miR-26a and miR-495 expression in males was dynamic across pubertal development in the dorsal hippocampus. In the dorsal hippocampus, the normal expression profile throughout pubertal development was significantly different between males and females at early puberty for miR-10a-5p and at each time point for miR-26a. miR-10a-5p, miR-26a, miR-103 and miR-495 expression profiles were dynamic in males as well as sexually dimorphic throughout pubertal development in the ventral hippocampus. Expression of BDNF and SIRT1 in males and SIRT1 in females was dynamic across pubertal development. Expression of BDNF and SIRT1 is significantly increased at peripuberty in males as well as sexually dimorphic in the ventral hippocampus.

**AIM 3: Determine whether there are sex and brain region differences in expression of estrogen-responsive microRNAs throughout pubertal development.**

Circulating levels of gonadal hormones increase rapidly throughout pubertal development, particularly estradiol ( $E_2$ ) in females and play a role in the development of sexually dimorphic behaviors. Interestingly, microRNA expression has been demonstrated to be sensitive to  $E_2$  in many systems. For example, Rao et al. demonstrated that the expression of five microRNAs, including let-7i, miR-9, miR-125a and miR-181a, are dependent on age and  $E_2$  in female rats [28]. Furthermore, Morgan et al. demonstrated that blocking  $E_2$  synthesis can alter microRNA expression in neonates [29] and suggest

that microRNAs may play an important role in initiating brain sex differences during fetal development. Overall, E<sub>2</sub> may modulate neuronal target genes using microRNAs as a fine tuning mechanism and it is important to determine pubertal sex differences in the expression of microRNAs sensitive to E<sub>2</sub>, as such information may yield insight into whether microRNAs play a role in brain sex differences arising throughout pubertal development.

The following questions were answered in this aim:

- 1) Are let-7i, miR-7a, miR-9, miR-125a and miR-181a expressed in sex, age and brain region-dependent manners?
- 2) In the ventral hippocampus, specifically, how are let-7i, miR-7a, miR-9, miR-125a, miR-181a expressed in males and females throughout and is the expression of each microRNA dynamic throughout pubertal development?
- 3) In the dorsal hippocampus, specifically, how are let-7i, miR-7a, miR-9, miR-125a, miR-181a expressed in males and females throughout puberty and is the expression of each microRNA dynamic throughout pubertal development?

In order to answer these questions, male and female Wistar rats at PND 30, PND 44 and PND 73 (N=10/group) were sacrificed by decapitation, brains were rapidly frozen and sectioned at 200  $\mu$ m, and the ventral and dorsal hippocampus were microdissected using a 0.75 mm Palkovit's brainpunch tool. RNA was isolated using Trizol according to manufacturers' instructions, and used to reverse transcribe microRNA (NCode™ microRNA First-Strand) into cDNA. qRT-PCR was performed with Fast Start Universal

SYBR Green Master Mix (Roche), using primers designed for microRNAs of interest, to quantify microRNA expression levels.

In summary, data from this aim demonstrate that miR-7a, miR-9, miR-125a and miR-181a are expressed in sex, age and brain region-dependent manners. In the ventral hippocampus, expression of miR-7a, miR-9, miR-125a, miR-181a are sexually dimorphic throughout puberty, and miR-9 expression is dynamic throughout pubertal development. In the dorsal hippocampus, expression of miR-7a, miR-9, miR-125a, miR-181a are sexually dimorphic throughout puberty and expression of miR-7a, miR-9, miR-125a and miR-181a in female and miR-7a in males is dynamic throughout pubertal development.

## CHAPTER TWO

### LITERATURE REVIEW

#### **Adolescence**

##### *Propensity for Binge Drinking*

Adolescence in humans comprises, on average, the years between 10-25 years of age [30,31]. During this period, the brain is in an exceptionally plastic state, such that morphological changes occur in gray matter, white matter and ventricle volume [32,33] as well as in neural networks as due to highly active molecular events taking place involving neurogenesis [15] and synaptogenesis [34,35] as well as apoptosis [36] and synapse elimination [37,38]. This combination of complex molecular events gives rise to the adolescents' brain being extremely sensitive to environmental stimuli, yet, impulse control and inhibitory drive are some of the last behaviors to develop [39]. Together, these characteristics lend adolescents to engage in increased risky behaviors such as binge drinking [17]. Binge drinking is defined as raising the blood alcohol concentration (BAC) greater than the legal limit ( $>0.08$  g/100 g) within a two-hour period, typically 3-4 servings of alcohol for a woman and 4-5 drinks for a man [40]. In 2007, SAMSA documented that on average, binge drinking begins around age 13 and peaks between ages 18-22 and that adolescents' drink 2 times more EtOH per occasion than adults (SAMSA, 2007). Using rat models of adolescent binge drinking, studies from our lab and others have demonstrated that peripubertal binge drinking induces long-lasting alterations

in hypothalamus and hippocampus gene expression, suggesting that this behavior impairs normal brain development [8,10,11,12,41].

## **Alcohol Alters Neurodevelopment**

### *Dysregulation of the Hypothalamic-Pituitary-Adrenal Axis*

Alcohol is a potent physiological stressor, as it activates the neuroendocrine stress response controlled by the hypothalamic-pituitary-adrenal (HPA) axis [42]. HPA axis activation can be induced by acute psychological or physical stressors and entails hypothalamic release of corticotrophin-releasing hormone (CRH), which stimulates the release of adrenocorticotrophic hormone (ACTH) from the anterior pituitary, and, in turn, causes the release of adrenal glucocorticoids. Further activation of the HPA axis is controlled by a negative feedback mechanism wherein the increased glucocorticoid (cortisol (CORT) in humans and corticosterone (CORT) in rodents) that is released from the adrenals inhibits further activation of the axis. Importantly, the development of HPA axis and other brain circuits is incomplete during puberty [43], making these immature neural networks at great risk for limited maturation and/or flawed development due to environmental insults that affect the brain, such as binge drinking. Indeed, the peripubertal rat is responsive acute EtOH, such that acute exposure significantly increases plasma CORT levels. However, following repeated exposure to high doses of EtOH (a.k.a. binge EtOH exposure) during puberty, HPA axis activation becomes desensitized, demonstrated by significantly lower plasma CORT levels compared to rats treated with a single high dose of EtOH [44]. Furthermore, EtOH abuse during puberty alters the expression of CRH in both an immediate and long-term fashion [12,44,45]. Moreover,



adult male rats exposed to peripubertal binge EtOH exhibit different responses to acute EtOH exposure compared to adult rats never previously exposed to binge EtOH, as indicated by different plasma CORT levels following a single dose of EtOH [12]. These data provide physiologic evidence that peripubertal binge EtOH induces long-lasting dysregulation of the neuroendocrine stress response, and suggest that peripubertal binge EtOH can have long-lasting consequences on an individual's ability to respond to stress.

Interestingly, the effects of peripubertal alcohol abuse on the HPA axis are sexually dimorphic [42,44,46] and dependent on gonadal hormones [47]. Moreover, a dysregulated HPA axis is associated with mood disorders that present differentially in males and females and are often comorbid with alcohol abuse [48,49,50,51,52,53,54]. In females, fluctuating levels of circulating steroid hormones are associated with HPA axis activation [55] and the dysregulation of the HPA axis observed in patients with major depressive disorder (MDD) is associated with abnormal steroid hormone levels in women compared to healthy individuals [56]. Overall, EtOH consumption alters the adolescent stress response in a long-term manner which likely contributes to the development of anxiety and related mood disorders, however the molecular mechanisms underlying these events remain unclear. It is, therefore, possible that EtOH's effects on other brain regions such as the hippocampus, contribute to EtOH-induced HPA axis dysregulation, as the hippocampus has major neural connections to the hypothalamus.

#### *Hippocampus Malfunctioning: Ventral and Dorsal Perspectives*

The hippocampus is a functionally and structurally complex brain region. It traverses both the dorsal and ventral planes and has various connections to cortical and

limbic areas, allowing the hippocampus to contribute to numerous neural networks regulating a range of behaviors. The information flow into the hippocampus enters through a restricted area, yet the monosynaptic outputs connect to a large variety of cortical and subcortical regions and contribute to the complex network of hippocampus circuit. Anatomical and genomic evidence supports the hippocampus playing a role in regulating both memory and cognitive processing, as well as stress, emotion, sensory–motor integration and goal-directed activity via its dorsal and ventral regions, respectively [57,58,59]. Indeed, afferent and efferent connections to parahippocampal regions gradually shift along the dorsal and ventral axis and gene expression profiles differ between the two regions [60,61].

Importantly, alcohol abuse imparts particularly detrimental effects on both memory and mood functions supported by the hippocampus. For example, hippocampus-dependent memory capacity has long been evident in adults with a history of chronic heavy drinking [62] and long-term impairments in executive functioning, motor control and emotional memory capabilities following EtOH abuse are attributed to long-term alterations in neurocircuitry [63,64] and brain morphology [65,66]. Moreover, alcohol abusers often suffer from anxiety and other mood disorders [67,68,69,70,71,72], and both alcohol abuse and mood disorders are associated with a dysregulated stress response [73,74,75,76,77,78]. Therefore the effects of peripubertal binge drinking on the neuroendocrine stress response may be controlled by upstream mechanisms initiated within the hippocampus. Indeed, it is becoming more evident that heavy drinking during adolescence also damages hippocampus morphology, gene expression and behavior. For instance, adolescent rats previously exposed to EtOH perform worse on Morris water

maze, indicating that EtOH impairs spatial learning in adolescents to a greater extent than it does in adults [79]. Moreover, alcohol abuse by adolescents is also associated with a reduction in hippocampus volume [3] and EtOH abuse alters gene expression differentially in the ventral and dorsal hippocampus [8], but the mechanisms remain unclear.

Many studies have identified genes essential for hippocampus development including brain-derived neurotrophic factor (BDNF) which regulates dendritic branching, spine density, learning and memory, and neuronal survival and sirtuin 1 (SIRT1), a class III nicotinamide-adenine dinucleotide (NAD<sup>+</sup>)-dependent histone deacetylase recently implicated in the regulation of hippocampal synaptic plasticity, neurogenesis and anxiety. Importantly, both BDNF and SIRT1 expression levels are sensitive to EtOH, but what regulates this sensitivity is less understood.

#### *Hippocampal Brain-Derived Neurotrophic Factor Role in Development*

BDNF is a member of the neurotrophin family and is essential for synaptic plasticity—the alteration of the strength between two neurons—which involves functional and structural changes at the synapse [80,81,82,83,84,85]. Synaptic plasticity is required for the consolidation of hippocampal-dependent learning [24,26,27]. Indeed, heterozygous *BDNF* knockout mice (+/-) have impaired learning and memory processing which can be rescued with recombinant BDNF in the hippocampus [24,26,27]. The hippocampus expresses an abundance of BDNF, which is particularly essential to neuronal survival processes. BDNF is expressed in the astrocytes, fibroblast and various types of neurons found within the hippocampus, hypothalamus, amygdala and neocortex.

The highest expression of BDNF mRNA in the brain is found in the granule neurons of the dentate gyrus and the pyramidal neurons of areas CA3 and CA2 in the hippocampus, indicating that BDNF plays a critical role in hippocampus functions [86]. Indeed, hippocampus-dependent memory can be disrupted by EtOH [87,88,89]. For example, early postnatal pups exposed to EtOH vapor inhalation for 2 hours and 40 mins each day on PND 4-10 had significantly increased BDNF protein expression [90]. However, other studies suggest that the adolescent brain is particularly sensitive to EtOH-mediated changes in the molecular mechanism underlying learning and memory [4,5,91]. Importantly, alterations in hippocampal BDNF expression lead to reduced hippocampal volume and poor performance on declarative memory tasks and impaired fear extinction [92]. Moreover, because BDNF expression is critical for memory formation and EtOH increases BDNF levels while disrupting memory function, it is important to understand how BDNF may play a role in the mechanisms underlying EtOH-mediated effects on memory function. Importantly, EtOH's effects on various brain functions are likely mediated by multiple proteins with gene regulatory capacity, and interesting research indicates the chromatin modifying enzyme sirtuin1 (SIRT1) in the regulation of gene expression following EtOH abuse [93].

#### *Hippocampal Sirtuin 1 Role in Development*

SIRT1 is a NAD<sup>+</sup> - dependent protein deacetylase and therefore, has the potential to regulate a large variety of biological functions by silencing the expression of genes coding for a wide range of cellular processes. Indeed, sirtuins are involved in neuronal development, neuroprotection, dendritic branching and neurogenesis [94,95,96,97,98,99,100] as well as hippocampus-specific functions including memory

performance and mood regulation. For example, SIRT1 expression in the hippocampus is positively correlated with adult hippocampal-dependent learning and memory [23,25,101], however, it is also mediates anxiety in mice [102]. These data suggest that SIRT1 may play an important role in both dorsal and ventral hippocampus functions. Moreover, SIRT1 gene variants are associated with a risk for anxiety [102], which is often comorbid with alcohol abuse [67,68,69,70,71,72]. Indeed, adolescent binge drinking in rats differentially alters SIRT1 expression levels in the male rat dorsal and ventral hippocampus [8] and SIRT1 expression is altered in the hypothalamus of male mice exposed to alcohol in utero [93]. Interestingly, many of the SIRT1 functions are associated with changes in expression levels of microRNA (microRNA) [8,23], small RNA molecules with gene regulatory potential. Indeed, the importance of SIRT1 in mediating hippocampal development and functions, as well as its potential to do so via microRNAs, requires future research to elucidate these mechanisms.

## **microRNA**

### *What are microRNAs?*

Whole-genome sequencing data from a variety of species have underscored the importance of post-transcriptional and post-translational modifications needed to achieve extensive phenotypic diversity. microRNAs (microRNAs) are small, non-protein coding RNAs (~22-nt long) and contribute to the complexity of the regulation of gene expression by mediating downstream target gene expression via translational repression and/or degradation. In this fashion, they can regulate virtually all biological processes [18,19]. 50% of microRNA genes are located in intergenic regions of the genome, 40%

within gene introns, and 10% are situated in exons, and thus, some microRNA expression mimics that of their host gene(s), however, much remains to be discovered regarding how microRNA expression is regulated. Evolutionarily, the emergence and conservation of microRNA genes localized within host protein-coding genes suggests that this localization emerged in response to environmental pressures that required tight control of gene expression and coordination of specific cell functions [103]. Importantly, microRNAs expression profiles are region-specific in the brain and [8,104,105,106,107,108,109,110] play critical roles in neuronal growth and synaptic plasticity [111,112,113,114,115,116] and their dysregulation has been implicated as a causative factor in a variety of neuropathologies [117,118,119,120,121,122]. Therefore, it is important to determine the mechanisms of microRNA biogenesis, regulation and dysregulation in both healthy and pathological tissues in order to better understand the mechanisms of microRNA-based regulation of gene expression and the consequences on biological phenotypes [123,124].

#### *microRNA Biogenesis*

The biogenesis of most mature 22-24 nucleotide (nt) single stranded microRNAs involves 4 processes: 1) transcription, 2) cropping, 3) dicing and 4) loading. microRNAs are transcribed in an RNA Polymerase II-dependent fashion which produces an intergenic pri-microRNA, about 100-1000 nt in length which forms a stem-loop structure [125,126]. Cropping of microRNA is performed by the microprocessor complex, composed of DiGeorge Syndrome critical component 8 (Dgcr8), which recognizes the stem-loop structure, and the RNase III enzyme Drosha, which contains the catalytic component important for cleaving the double stranded stem, removing the loop, and generating the

pre-microRNA product [127,128]. The pre-microRNA is ~ 70-100 nt in average length and are exported from the nucleus. In the cytosol, the dicing step occurs via the RNase III enzyme Dicer, which catalytically cleaves the pre-microRNA into a double-stranded ~22 nt-long product, one strand of which is the mature microRNA. In the loading step, the single-stranded mature microRNA is transferred to the RNA-induced silencing complex (RISC), which comprises the microRNA-RISC complex (miRISC) which allows microRNAs to locate specific messenger RNA transcripts target genes and this leads to either translation repression or mRNA degradation. Interestingly, microRNAs imperfectly bind small (~6 nt) seed sequences on the 3' UTR of their gene targets, permitting promiscuity to microRNAs, as a single microRNA can target multiple genes [129]. Likewise, one gene can be targeted by multiple microRNAs. Such features enable microRNAs to mediate quick, post-transcriptional gene silencing which would be particularly useful in the regulation of synaptic plasticity.

#### *microRNAs Regulate Neurodevelopment*

The central nervous system (CNS) development requires a precise temporal orchestration of events that is uniquely suited for the fine-tuning attributes of microRNAs. The importance of microRNAs in embryonic nervous system development was originally demonstrated using transgenic animal models that manipulated microRNA biosynthetic processing enzymes, including Dicer. There is a single gene that encodes Dicer in *C. elegans*, mice, and humans, and depletion of Dicer results in severe developmental consequences. Studies in Dicer-mutant zebrafish showed that they had disrupted embryonic morphogenesis and neural differentiation [130]. More specifically, the brains lacked ventricles, neuronal positioning was disrupted suggesting migration

defects, and many neurons had defasciculated axons [130]. Strikingly, injections of microRNAs from the miR-430 family (miR-430a/b/c) reversed many of the brain morphogenic defects that resulted from Dicer deletion in the zebrafish, revealing a direct connection between mature microRNAs and Dicer during development. The partial rescue of neuronal defects in this study also provided some of the first evidence for tissue-specific effects of microRNAs [130,131]. Global Dicer deletion in mice is embryonic lethal [132] prompting the generation of tissue-specific conditional Dicer-null mouse models. In the developing neocortex, the absence of Dicer resulted in a smaller cortex, improper cortical layering, increased apoptosis, as well as an overall reduction in neural progenitor cells and oligodendrocytes [133,134,135].

microRNAs and their biogenesis enzymes are found enriched in the synpto-dendritic compartments of neurons [113,136,137,138] indicating their potential role in regulating synaptic plasticity. Indeed, rapid changes in local protein synthesis triggered by synaptic activation [139,140,141] may rely on the function of microRNAs [142]. It has been demonstrated that expression of the important synaptic plasticity molecule BDNF is regulated by miR-26a which targets the conserved sequences of the BDNF 3'UTR [143]. Moreover, miR-26a and BDNF expression have been implicated in the vulnerability and onset of schizophrenia, alcohol abuse and mood disorders in both human patients and rodent models [143,144,145,146,147,148]. In the brain, it has been demonstrated that microRNAs play a role in synaptic plasticity [113,136,137,138,149] are expressed in a sexually dimorphic manner [150] and that their expression is dependent on steroid hormones [28]. Lamina-specific expression of miR-495 was observed to be complementary to the BDNF expression in human prefrontal cortex [151].



Furthermore, miR-495 has been associated with the mechanism whereby the mood stabilizers valproate and lithium exert their neuroprotective effects [152]. Overall, microRNAs have the capacity to regulate neuronal development and maintenance—essential processes for a functioning nervous system and they also demonstrate sensitivity to mood altering substances, including EtOH.

#### *Alcohol Effects on microRNA*

The effects of alcohol on brain microRNA remains an ongoing investigation, however many recent developments have pointed towards the sensitivity of microRNA expression in response to EtOH. In early life development, fetal brain microRNA expression levels are altered by high levels of alcohol consumption by the pregnant mother, and importantly, these changes in microRNA expression following maternal EtOH abuse have been implicated in the teratology of EtOH-induced malformations in fetal brain development [153,154]. Furthermore, microRNA expression levels are known to exhibit differential expression profiles in fetal vs. postnatal brain, suggesting that microRNAs are fundamentally involved in regulating developmental processes in the brain [80,105,155,156] and suggest that alterations in microRNA expression in response to EtOH can impart damage to brain development. microRNA expression levels in the adolescent rat brain are also altered following peripubertal binge drinking [8] and in the rat prefrontal cortex in alcohol dependent rats [157]. Importantly, peripubertal or adult alcohol abuse in rats does not impart global changes in microRNA expression levels, but rather alters the expression of specific microRNAs in distinct regions of the brain. This specificity suggests that EtOH alters specific microRNA expression levels in order to damage specific brain functions. microRNA expression profiles are also distinct in

human alcoholic prefrontal cortex, providing further evidence that EtOH mediates brain damage by altering microRNA expression [158]. Identifying the effector functions of the target genes of EtOH-sensitive microRNAs is indeed a more challenging task, however, in both rat prefrontal cortex and hippocampus, BDNF has been identified as a target gene of EtOH-sensitive microRNA [157], suggesting the importance of the microRNA-mediated regulation of BDNF in the brain, and the potential for alcohol abuse to disrupt this regulation. How microRNA expression levels throughout the lifespan are expressed and what regulates their expression still remains unclear, however much evidence suggests the importance of gonadal hormones in this process. Indeed, gonadal hormones levels themselves are dynamic throughout different developmental stages of life as well as responsive to changes in the environment.

#### *Gonadal Hormone Effects on microRNA*

Steroid hormone regulation of microRNAs has been documented in a variety of sex-specific cancer models such as breast, prostate and endometrial cancer [159,160,161,162,163], however, less is known about steroid hormone regulation of microRNA expression in brain tissue. Nonetheless, studies examining the effects of sex steroid-induced microRNA regulation using in vivo models have indicated that sex-steroid regulation of microRNA expression in the brain differs across the lifespan and support the theory that sex steroid functions in the brain evolve with age. For example, blocking synthesis of E<sub>2</sub> in the neonate brain induces sexually dimorphic patterns of microRNA expression, such that males and female microRNA expression profiles become indistinguishable [29]. In the aging female brain multiple microRNAs were identified as E<sub>2</sub>-responsive in the hippocampus, central amygdala and the paraventricular

nucleus, some of which were differentially regulated by E<sub>2</sub> dependent on age [28].

Despite these informative studies demonstrating steroid-mediated microRNA expression regulation in the brain, the mechanism remains unresolved.

microRNA expression changes in response to differential hormone environments indicate that hormones may mediate changes in microRNA expression via their nuclear receptors. Steroid hormone receptors (i.e. nuclear receptors (NRs) classically function as transcription factors, by binding to promoter elements and recruiting coactivators and corepressors, together, induce gene activation or suppression. Because microRNA genes often lie within the protein-coding regions of genes, NRs can thereby regulate microRNAs in this way. Alternatively, downstream target genes regulated by NRs may, themselves, influence subsequent microRNA gene expression in an indirect fashion. Indeed, many NR target genes are transcriptionally regulated via secondary effects [164]. Another indirect manner in which microRNA expression can be regulated is via the proteins required for microRNA biogenesis and RNA interference (RNAi) activity, including the RNase III enzymes Drosha and Dicer which are responsible for the sequence-specific cleavage of complementary RNA targets. Indeed, E<sub>2</sub> has been demonstrated to mediate expression of these proteins [165], however, no such regulation has been documented in the brain.

### **Summary**

Peripubertal neurodevelopment is an important period of brain development wherein increased synaptic plasticity lends environmental exposures to play a large role in shaping how the neural circuitry develops. A common environmental exposure in the

United States is engagement in binge EtOH drinking behavior, which has been demonstrated to have sexually dimorphic, long-lasting neurological consequences in hippocampus-mediated functions. Neurological consequences of EtOH abuse have been recently demonstrated to alter microRNA expression, which have also recently been discovered to play vital regulatory roles in neuronal maintenance. Therefore, we conclude the introduction with the following hypothesis and aims to investigate microRNA expression in male and female pubertal hippocampus development as well as investigate the effects of peripubertal binge EtOH.

## CHAPTER THREE

### LONG-TERM EFFECTS OF PERIPUBERTAL BINGE ETOH EXPOSURE ON HIPPOCAMPAL MICRORNA EXPRESSION IN THE RAT

#### **Introduction**

Heavy episodic alcohol consumption (i.e. binge drinking) has been steadily increasing among adolescents in recent decades [40,166]. Indeed, data from the Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (SAMHSA 2005) showed that an alarming 90% of the alcohol consumed by youth occurs in a binge-like pattern, defined as raising the blood alcohol concentration (BAC) above the legal driving limit (0.08%) within a 2 hour time period (NIAAA, 2012). Extensive remodeling of the brain occurs during adolescence, which includes changes in cortical gray and white matter, synaptic connectivity, and increased neurogenesis [167,168,169,170,171] and alcohol exposure during this critical time can have severe detrimental effects on brain function [44,172,173,174,175,176]. Studies from our laboratory and others have demonstrated that adolescent binge-pattern alcohol exposure results in long-term dysregulation of the neuroendocrine stress response, memory impairments and behavioral deficits [12,67,177,178,179]. An altered stress response often underlies depression- and anxiety-related disorders and importantly, these conditions are commonly experienced by over 50% of alcohol-dependent patients [67,180,181]. Indeed, mood and memory impairments are often present in tandem with alcohol abuse and

neuropsychiatric illnesses that emerge post-puberty [182,183]. The hippocampus is an important brain region that mediates learning, memory, and mood and it has been well established that hippocampus structure and function is impaired by EtOH abuse [10,11,184,185,186,187,188]. Notably, pubertal EtOH abuse inhibits adult neurogenesis, impairs learning and memory in adulthood, and impairs information retention. The precise molecular mechanisms mediating the long-term effects of adolescent binge EtOH exposure are poorly understood, however short non-coding regulatory RNAs are sensitive to EtOH and have recently been recognized as critical mediators of nearly every basic cellular process [154,189]. In particular, microRNAs (microRNAs, ~22 nucleotide single-stranded non-coding RNA) regulate the translation of proteins important for neuronal development during embryogenesis, postnatal neuronal maintenance and survival, and hippocampal neurogenesis throughout life [112,113,190,191,192,193]. Moreover, disruption of mature microRNA expression and/or function has been linked to alcohol-induced neurological afflictions including addiction and fetal alcohol spectrum disorder (FASD) [194,195]. In this study we used a Wistar rat model to identify EtOH-sensitive microRNAs that target genes involved in regulating hippocampal processes, such as memory and mood. Using multiple target prediction algorithms (Targetscan: [www.targetscan.org](http://www.targetscan.org); microRNA database: [www.miRDB.org](http://www.miRDB.org)) [129,196,197], we identified brain-derived neurotrophic factor (BDNF) as a target gene of miR-10a-5p, miR-26a, miR-103, and miR-495, synapsin-2 (SYN2) as a target gene of miR-32, and sirtuin 1 (SIRT1) as a target gene of miR-26a, miR-103 and miR-495. BDNF and SYN2 are both critical regulators of synaptic plasticity. For example, BDNF is a neurotrophic factor instrumental in neurodevelopment and alterations in its expression are evidenced in

numerous psychiatric disorders, SYN2 is a synaptic vesicle phosphoprotein that modulates neurotransmitter release onto the post-synaptic membrane and its dysregulation disrupts the delicate balance of excitatory and inhibitory neurotransmission throughout the brain whereas SIRT1 is a class III protein deacetylase that has been recently associated with anxiety behavior [102,180,181,198,199].

We tested the hypothesis that mid/peripubertal binge EtOH exposure alters hippocampal microRNA expression and that this leads to changes in the expression of their target genes. Importantly, the normal expression profile of these particular microRNAs during pubertal development has not been reported in any species studied to date. Therefore, we quantified the normal developmental expression profile of miR-10a-5p, miR-26a, miR-103 and miR-495 in the rat hippocampus at three time points in pubertal development (early, peri, and late). Next, we determined how peri-pubertal binge EtOH exposure altered those normal expression levels immediately following EtOH exposure, as well as 30 days after the last EtOH dose using a binge pattern that is reliable for testing the effects of typical adolescent alcohol abuse [44,200]. Gene expression levels of microRNA processing enzymes, Droscha and Dicer, were also quantified at each time point in order to determine a possible molecular mechanism for binge EtOH effects. Lastly, to determine the downstream physiological effects of alterations in microRNA target gene transcription, we quantified the gene expression levels of putative microRNA target genes, BDNF and SIRT1 as well as their protein expression levels. Overall, our data provide evidence that peripubertal binge EtOH exposure induces long-term alterations in mature microRNA expression levels in the rat

hippocampus, which has the potential to modulate the expression of their downstream target genes.

### **Approach**

Figures 2 and 3: Male Wistar were handled daily beginning at PND 30 and given one of the following treatment paradigms starting on PND 37 (n=6/group): 1) saline control (1x/day/8days), 2) acute EtOH (saline 1x/day/7days + 1x/day/1day EtOH) or 3) binge EtOH (1x/day/3days EtOH + 1x/day/2days saline + 1x/day/3days EtOH) via intraperitoneal injection. One hour after the last dose, brains were collected, rapidly frozen, sectioned at 200  $\mu$ m on a freezing microtome and the whole hippocampus was microdissected using a 0.75 mm Palkovit's brainpunch tool. The array was performed using SABiosciences RT<sup>2</sup> miRNA PCR Array. Quantitative real-time reverse transcription PCR (qRT-PCR) was used to quantify the expression levels of six microRNAs: (miR-10a-5p, miR-26a, miR-32, miR-103, mmiR-423 and miR-495) as well as BDNF and SIRT1 mRNA.

Figures 4-13: The dorsal and ventral hippocampus were microdissected from male Wistar rats that were either untreated or treated with an EtOH paradigm administered via oral gavage beginning at PND 37 (3g/kg; 1x/day/3days EtOH, +1x/day/2days water + 1x/day/3days EtOH) . Brains collected at three different ages during pubertal development (early puberty, PND 30; mid/peripuberty, PND 44; late puberty, PND 71; N=10/group). Importantly, peripuberty in rats is considered to be ~ PND 30-45 [201,202]. Quantitative real-time reverse transcription PCR (qRT-PCR) was used to quantify the expression levels of a total of five microRNAs: (miR-10a-5p, miR-



26a, miR-32, miR-103 and miR-495) and four genes (Drosha, Dicer, BDNF and SIRT1). Western Blot was used to detect protein expression of pro-BDNF, BDNF and SIRT1.


## **Results**

### **Blood alcohol concentrations following peripubertal binge EtOH exposure**

In order to model typical pubertal binge alcohol drinking behavior, we utilized an 8-day binge EtOH exposure paradigm that has been used previously to mimic the pattern of alcohol consumption commonly observed in human adolescents (Fig. 1A) [8,44]. Experimentally, this paradigm does not affect body weight/growth curves during pubertal development in male rats [8,44,200]. Blood alcohol concentrations (BAC) were measured on the final day of treatment (day 8) 60 minutes following the last dose. The BAC in EtOH treated animals was  $190 \pm 21$  mg/dl which was consistent with our previous studies using this peripubertal binge EtOH paradigm (Fig. 1B) [8,44,200]. Overall, our observed BAC levels fall within the defined BAC threshold of binge drinking.


### **miR-10a-5p, miR-26a, miR-103, miR-32 and miR-495 expression levels are sensitive to peripubertal binge-pattern EtOH exposure in the rodent hippocampus.**

In the brain, microRNAs play an important role in neurodevelopment and neurogenesis, two processes that continue to occur throughout adolescent development [203] and microRNAs are implicated in the teratology of EtOH-induced malformations in fetal brain development [153,154]. Importantly, our understanding of microRNAs in the brain during adolescent development is severely limited and there is little to no data




**Figure 1:** Peripubertal repeated binge EtOH paradigm alters blood alcohol concentration. On PND 37, male Wistar rats are given 3g/kg EtOH (20% v/v in water), or water alone, via oral gavage at 10AM and this process is repeated according to the following schedule: 3d EtOH, 2d water, 3d EtOH (A). Blood alcohol concentrations (BAC) 1.0 h following the last dose of water (control) or peripubertal binge EtOH (B). Data expressed as mean BAC (mg/dl). \*Statistically significant difference via Student's T-test N=6 ( $p < 0.05$ ).

describing how EtOH might affect microRNAs during this critical developmental period. microRNA expression was measured in the hippocampus of male rats treated with an 8-day binge EtOH paradigm (see Fig. 1A) using a microRNA RT-PCR profiling array containing a genome-wide panel of the 88 most well-researched rat microRNAs (as annotated by the Sanger miRBase Release 14). We identified 2 novel EtOH-sensitive microRNAs (miR-10a-5p and miR-423) which demonstrated altered expression levels following an 8-day peripubertal binge-pattern EtOH exposure. Specifically, miR-10a-5p and miR-423 expression levels were increased >10-fold in the hippocampus of binge-EtOH treated rats (Fig. 2). We next designed our own primers to validate the array-generated results with qRT-PCR and when using these primers, the expression level of miR-10a-5p increased in the binge-treated animals as it did in the array (Fig. 3A). However, miR-423 expression levels did not change when using the new primers, and therefore, miR-423 was not further studied. To add depth to the study, we also quantified the expression levels of additional microRNAs, miR-26a, miR-32, miR-103 and miR-495 that were predicted to target genes which regulate hippocampus functions. These data revealed that peripubertal binge-pattern EtOH exposure increased miR-32, miR-103 and miR-495, and decreased miR-26a expression levels in the male rat whole hippocampus (Fig. 3A). Using the target prediction algorithms available via Targetscan: [www.TargetScan.org](http://www.TargetScan.org) and microRNA database: [www.miRDB.org](http://www.miRDB.org) [129,196,197], we identified BDNF and SIRT1 as a putative target genes of miR-10a-5p, miR-26a, miR-103 and miR-495. These computer algorithms identify predicted microRNA target genes



**Figure 2.** Peripubertal binge EtOH increases miR-10a-5p and miR-423 expression in the male rat whole hippocampus. Scatter plot data represents microRNA expression levels in binge EtOH treated rats relative to controls (open circles). miR-423 and miR-10a-5p (red) expression levels increased > 10-fold in binge EtOH-treated rats compared to control (one-way ANOVA, \* $p < 0.05$ ).




**Figure 3:** Peripubertal binge-pattern EtOH exposure alters microRNA and target gene expression in male rat hippocampus. microRNA (A) and target gene (B) expression levels 1.0 h following the last dose of water or EtOH. Data represent mean fold change  $\pm$  SEM as compared control. \* indicates significance between groups (one-way ANOVA,  $p < 0.05$ ). target gene. Analysis of the potential target genes for these EtOH-sensitive microRNAs,

we identified one reported binding site for miR-10a-5p, miR-26a and miR-103 within the 3'UTR of BDNF and four for miR-495. One binding site was reported within the 3'UTR of SIRT1 for miR-26a, mir-103 and miR-495. Importantly, the number of predicted binding sites is indicative of the relative specificity of the microRNA for that gene.

We next quantified the expression levels of the EtOH-sensitive microRNA gene targets, BDNF and SIRT1, which displayed strong decreasing trends in the binge EtOH-treated animals, but did not reach not statistical significance (Fig. 3B). Importantly, the male rat brain displays both age-, cell type- and region-specific microRNA expression profiles and the hippocampus consists of functionally distinct ventral and dorsal regions [57,58,105,204,205]. Therefore, we next aimed to determine the potential for peripubertal binge EtOH to induce region-specific alterations in the ventral and dorsal hippocampus.

**Mature miR-10a-5p, miR-26a, and miR-495 expression levels in the dorsal hippocampus of untreated male rats are age dependent.**

Expression levels of these specific mature microRNAs in the brain during pubertal development have not been previously described. Therefore, we first determined the normal developmental profile of mature miR-10a-5p, miR-26a, miR-103 and miR-495 expression in the dorsal hippocampus using untreated male Wistar rats. miR-32 expression levels were not significantly altered in the dorsal or ventral hippocampus by age or EtOH, and therefore, it was not further studied. Mature microRNA expression




**Figure 4.** Diagram of experimental paradigm. Diagram depicting the age of sacrifice and specific treatment paradigms for each group of male Wistar rats. 3 groups received no treatments (A). 4 groups received water or binge EtOH treatments at peripuberty, for a total of 2 control and 2 EtOH groups (B). N=10/group.

**Table 1.** Statistical Analysis of Gene Expression Levels in the Dorsal Hippocampus

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF TREATMENT</b>	<b>INTERACTION: AGE X TREATMENT</b>
miR-10a-5p	Yes: $F(2,33) = 12.293$ $p < 0.001$	No	No
miR-26a	Yes: $F(2,44) = 3.222$ $p < 0.049$	Yes: $F(1,44) = 5.212$ $p < 0.027$	Yes: $p < 0.001$
miR-103	No	No	No
miR-495	Yes: $F(2,37) = 4.923$ $p < 0.013$	Yes: $F(1,37) = 59.23$ $p < 0.001$	Yes: $p < 0.001$
<b>GENE</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF TREATMENT</b>	<b>INTERACTION: AGE X TREATMENT</b>
Drosha	Yes: $F(2,54) = 24.906$ $p < 0.001$	No	Yes: $p = 0.008$
Dicer	Yes: $F(2,54) = 18.725$ $p < 0.001$	No	Yes: $p < 0.001$
BDNF	Yes: $F(2,52) = 12.845$ $p < 0.001$	Yes: $F(1,52) = 6.574$ $p < 0.013$	No
SIRT1	No	Yes: $F(1,54) = 110.941$ $p < 0.001$	Yes: $p < 0.001$





**Figure 5.** Peripubertal binge EtOH exposure alters microRNA expression during pubertal development in the dorsal hippocampus. miR-10a-5p (A), miR-26a (B), miR-103 (C), and miR-495 (D) expression levels in untreated (solid line) and EtOH-treated (dashed line) pubertal male rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 animals. Dissimilar letters indicate a statistically significant difference between groups (2-way ANOVA,  $p < 0.05$ ).

levels were measured using qRT-PCR at three time points throughout pubertal development (early = 30 d, peri = 44 d, late = 73 d) (Fig. 4A). In the dorsal hippocampus, a two-way ANOVA revealed a main effect of age on the expression levels of all microRNAs tested, except miR-103 (Table 1). Each of the three microRNAs that showed a significant effect of age in the dorsal hippocampus had a distinct developmental pattern. For instance, miR-10a-5p expression decreased significantly between early and peripuberty, and remained lower than early puberty levels until late puberty (Fig. 5A, solid line). By contrast, miR-26a expression did not change between early and peripuberty, but significantly decreased at late puberty (Fig. 5B, solid line). Finally, a significant increase was observed in miR-495 expression between early and peripuberty and these levels remained high until late puberty (Fig. 5D, solid line).

**Binge EtOH exposure during peripuberty significantly alters normal developmental profile of microRNAs in the dorsal hippocampus.**

Next, we determined the effects of repeated binge EtOH exposure during peripuberty on these microRNA levels in the brain. Rats were administered our repeated binge-pattern EtOH exposure paradigm (Fig. 1A) and dorsal hippocampal microRNA expression of miR-10a-5p, miR-26a, miR-103 and miR-495 was compared with untreated rats/water-treated rats immediately following binge EtOH exposure and one-month post-EtOH exposure. Our results demonstrated a significant main effect of EtOH treatment on the expression of miR-26a and miR-495 and there was also a significant interaction between age and treatment, demonstrating that the effects of EtOH were age dependent (Table 1). Although there was no main effect of EtOH treatment on the expression of miR-10a-5p, and there was no immediate change following EtOH exposure at


peripuberty, by late puberty its expression was significantly increased compared to untreated rats/water-treated rats (Fig. 5A). miR-26a expression levels significantly decreased immediately following binge EtOH exposure at peripuberty, however this difference did not persist and was equivalent to those of untreated animals by late puberty (Fig. 5B). miR-103 was not significantly altered by peripubertal EtOH treatment in the dorsal hippocampus, similar to the results observed with age alone (Fig. 5C). Most striking were the results of EtOH exposure on miR-495. Similar to miR-26a, miR-495 was significantly decreased as a result of binge EtOH exposure at peripuberty. Notably however, expression levels remained significantly below normal even one-month post EtOH exposure (Fig. 5D), suggesting a potential long-term effect of pubertal binge EtOH exposure on miR-495 in the dorsal hippocampus.

#### **Mature miR-10a-5p, miR-26a, miR-103, and miR-495 expression levels in the ventral hippocampus of untreated male rats are age-dependent**

Distinct region and age-dependent expression of microRNAs has been demonstrated in the brain of a variety of species [105,206,207]. Therefore, we quantified the expression of miR-10a-5p, miR-26a, miR-103, and miR-495 in the ventral hippocampus across pubertal development in untreated rats to determine if there were region specific microRNA expression patterns in the hippocampus. In the ventral hippocampus, there was a significant main effect of age in the untreated animals on all four microRNAs tested, including miR-103, which previously did not show a significant change across pubertal development in the dorsal hippocampus (Table 2). Specifically,

**Table 2.** Statistical Analysis of microRNA Expression Levels in the Ventral Hippocampus

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF TREATMENT</b>	<b>INTERACTION: AGE X TREATMENT</b>
miR-10a-5p	Yes: $F(2,42) = 17.492$ $p < 0.001$	Yes: $F(1,42) = 8.847$ $p = 0.005$	No
miR-26a	Yes: $F(2,37) = 5.064$ $p = 0.011$	No	No
miR-103	Yes: $F(2,54) = 4.582$ $p = 0.015$	Yes: $F(1,54) = 5.739$ $p = 0.02$	Yes: $p < 0.001$
miR-495	Yes: $F(2,42) = 8.359$ $p < 0.001$	Yes: $F(1,42) = 5.998$ $p = 0.019$	Yes: $p < 0.001$
<b>GENE</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF TREATMENT</b>	<b>INTERACTION: AGE X TREATMENT</b>
Drosha	Yes: $F(2,54) = 4.650$ $p = 0.014$	No	Yes: $p < 0.001$
Dicer	Yes: $F(2,54) = 10.746$ $p < 0.001$	No	Yes: $p < 0.001$
BDNF	Yes: $F(2,41) = 13.622$ $p < 0.001$	Yes: $F(1,41) = 9.109$ $p = 0.004$	Yes: $p < 0.001$
SIRT1	Yes: $F(2,43) = 16.484$ $p < 0.001$	Yes: $F(1,43) = 14.759$ $p < 0.001$	Yes: $p < 0.001$



**Figure 6.** Peripubertal binge EtOH alters microRNA expression during pubertal development in the ventral hippocampus. miR-10a-5p (A), miR-26a (B), miR-103 (C), and miR-495 (D) expression levels in untreated (solid line) and EtOH-treated (dashed line) pubertal male rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 animals. Dissimilar letters indicate a statistically significant difference between groups (2-way ANOVA,  $p < 0.05$ ).

miR-10a-5p showed no change between early and peripuberty, but significantly increased by late puberty in the ventral hippocampus (Fig. 6A, solid line). Also, in contrast to the dorsal hippocampus miR-26a significantly decreased at peripuberty, but this change did not persist and was equivalent to early pubertal levels by late puberty (Fig. 6B, solid line). The ventral hippocampus levels of miR-103 and miR-495 had a similar profile. Both had a statistically significant decrease, or a strong trend towards decreasing, at peripuberty compared to early pubertal levels, but then the levels increased significantly above that of early pubertal levels by late puberty (Fig. 6C, D, solid line). Notably, miR-495 expression in the ventral hippocampus demonstrated the most dynamic expression profile throughout pubertal development, as it had distinct expression levels at each time point measured.

### **Repeated adolescent binge EtOH exposure differentially alters expression of miR-10a-5p, miR-26a, miR-103 and miR-495 in the ventral hippocampus**

We predicted that peripubertal binge EtOH exposure would alter the normal developmental profile of microRNA expression in the ventral hippocampus, based on the evidence obtained from the dorsal hippocampus. Indeed, there was a significant main effect of treatment and a significant interaction between age/treatment for miR-10a-5p, miR-103, and miR-495 in the ventral hippocampus (Table 2). Interestingly, the magnitude of changes in microRNA expression was overall much higher in the ventral hippocampus compared to the dorsal, with some microRNAs changing by as much as 5-fold (Fig. 6). One example of a large fold change was observed with miR-10a-5p. The mature expression levels of miR-10a-5p were significantly increased by an average of 3-5

**Table 3.** Summary of peripubertal EtOH exposure on microRNA and mRNA gene expression. Arrows indicate a statistically significant effect (increase/decrease) of EtOH compared to age-matched water-treated controls.

<b>microRNA</b>	<b>Dorsal Hipp immediate ETOH effect</b>	<b>Dorsal Hipp long-term ETOH effect</b>	<b>Ventral Hipp immediate ETOH effect</b>	<b>Ventral Hipp long-term ETOH effect</b>
10a-5p	–	↑	↑	–
26a	↓	–	–	↓
103	–	–	↑	↓
495	↓	↓	↑	↓
<b>Gene</b>	<b>Dorsal Hipp immediate ETOH effect</b>	<b>Dorsal Hipp long-term ETOH effect</b>	<b>Ventral Hipp immediate ETOH effect</b>	<b>Ventral Hipp long-term ETOH effect</b>
Drosha	↑	↑	↑	↓
Dicer	↑	↓	↑	↓
BDNF	↑	↑	↓	–
SIRT1	↑	↑	↓	–


fold following peripubertal binge EtOH exposure (Fig. 6A, dashed line). Further, the expression levels continued to show an increase at late puberty, paralleling the untreated group at that same age (Fig. 6A). There was no observed statistical effect of peripubertal EtOH treatment on miR-26a. Nevertheless, the expression levels in the EtOH-treated group did not appear to follow the normal age-dependent increase observed by late puberty (Fig. 6B). The most striking effects of peripubertal binge EtOH exposure in the ventral hippocampus were observed in miR-103 and miR-495 expression, as their normal developmental expression levels at both peripuberty and late puberty were opposite following peripubertal binge EtOH exposure. Overall, our results demonstrate both immediate and long-term effects of peripubertal binge EtOH exposure in the rat hippocampus and these effects were distinct between the dorsal and ventral regions (Table 3).

### **Mature microRNA biosynthetic processing enzymes are altered by peripubertal binge EtOH exposure in the dorsal and ventral hippocampus**

Primary microRNA transcripts are transcribed from the genome in a RNA polymerase II dependent manner and sequentially cleaved by the nuclear enzyme Drosha and the cytoplasmic enzyme Dicer to form the functionally mature single-stranded form of the microRNA [127,208,209]. We next measured mRNA levels of both Drosha and Dicer in our dorsal and ventral hippocampal tissue samples to better understand the molecular basis of altered mature microRNA expression levels. Drosha and Dicer mRNA expression levels were measured using qRT-PCR in the untreated animals at early, mid and late puberty and those levels were compared to animals that were administered our



### Enzymes in D. Hipp and V. Hipp




**Figure 7.** Peripubertal binge EtOH exposure differentially alters microRNA biosynthetic processing enzymes in the dorsal and ventral hippocampus. Drosha (A, B) and Dicer (C, D) mRNA levels in the dorsal and ventral hippocampus in untreated (solid line) and EtOH-treated (dashed line) pubertal male rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 animals. Dissimilar letters indicate a statistically significant difference between groups (2-way ANOVA,  $p < 0.05$ ).

binge EtOH treatment paradigm at peripuberty. Our results showed a main effect of age on Droscha in both the dorsal and ventral hippocampus (Tables 1, 2; Fig. 7A, B, solid line). Specifically, Droscha mRNA levels were significantly decreased between early and peripuberty in both regions of the hippocampus. The gene expression levels remained low until late puberty in the dorsal hippocampus but returned to early pubertal levels in the ventral hippocampus (Fig. 7A, B, solid line), thereby demonstrating region-specific regulation. Further, there was a significant interaction between age and EtOH on Droscha mRNA expression in both hippocampal regions and an overall main effect of EtOH treatment in the ventral hippocampus (Table 1, 2; Fig. 7A, B, dashed line). In both regions, peripubertal binge EtOH exposure significantly elevated Droscha mRNA levels immediately following EtOH exposure, suggesting a potential for increased microRNA biosynthetic processing. These increased levels persisted for up to one month (late puberty, Fig. 7A) following EtOH exposure in the dorsal hippocampus, but were significantly decreased at that same age in the ventral hippocampus (Fig. 7B). Dicer mRNA expression followed the same pattern as Droscha in untreated animals for both regions of the hippocampus and there was a statistically significant overall main effect of age (Table 1, 2; Fig. 7C, D). Dicer mRNA levels decreased between early and peripuberty in both regions and the levels remained low until late puberty in the dorsal hippocampus (Fig. 7C). By contrast, at late puberty Dicer mRNA levels were no longer significantly different from those in early puberty in the ventral hippocampus (Fig. 7D). Binge EtOH exposure in peripuberty had the same effect on Dicer mRNA expression levels in both regions of the hippocampus, with a statistically significant overall main effect of treatment and a significant interaction between age and treatment (Table 1, 2;

Fig. 7C, D). Dicer mRNA levels were immediately increased compared to untreated animals following EtOH exposure at peripuberty, but then by late puberty had decreased significantly to levels even lower than that of untreated animals at early puberty (Fig. 7C, D). In sum, these results demonstrate that both Dicer and Drosha change dynamically throughout pubertal development in the hippocampus and that these levels can be dramatically altered by peripubertal binge EtOH exposure (Table 3).


**Peripubertal Binge EtOH exposure alters putative target gene mRNA of EtOH-sensitive microRNA in the dorsal and ventral hippocampus.**

The most well understood mechanism of microRNA action is through microRNA complementary binding to the 3' untranslated region (UTR) of a primary gene transcript and its subsequent facilitation of mRNA degradation and/or inhibition of mRNA translation [210,211]. This miR-mediated degradation of mRNA target genes is attributable to observed downstream phenotypic changes. We identified two genes that were putative targets of all 4 EtOH-sensitive microRNAs in the dorsal and ventral hippocampus using publically available software target prediction programs (Targetscan: [www.targetscan.org](http://www.targetscan.org); miR database: [www.miRDB.org](http://www.miRDB.org)) [129,196,197]. Our analysis of potential targets for each EtOH-sensitive miR identified a single putative binding site in the 3'UTR of BDNF for each miR-10a-5p, miR-26a, and miR-103 (Fig. 8). Also, there were four possible binding sites on BDNF for miR-495, suggesting miR-495 might have a stronger regulatory effect on BDNF than the other microRNAs. There were no potential binding sites in the 3'UTR of SIRT1 for miR-10a-5p, but there was a single potential



**Figure 8.** Diagram depicting predicted microRNA binding sites for BDNF and SIRT1. Schematic diagram of the 3'UTR of (A) BDNF – 2,842 bp and (B) SIRT1- 1,607 bp. The putative binding sites for each microRNA were predicted using Targetscan ([www.targetscan.org](http://www.targetscan.org)) and miRDB ([www.miRDB.org](http://www.miRDB.org)) computer algorithm programs. The binding sites were predicted based on the presence of an 8-mer or 7-mer conserved microRNA seed sequence. Precise seed sequence positions are shown in parentheses.

### miR Gene Targets in D. Hipp and V. Hipp



**Figure 9.** Peripubertal binge EtOH exposure differentially alters miR target genes, BDNF and SIRT1, in the dorsal and ventral hippocampus. BDNF (A, B) and SIRT1 (C, D) mRNA levels in the dorsal and ventral hippocampus in untreated (solid line) and EtOH-treated (dashed line) pubertal male rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 animals. Dissimilar letters indicate a statistically significant difference between groups (2-way ANOVA,  $p < 0.05$ ).

binding site for each of the other microRNAs tested (miR-26a, miR-103 and miR-495; Fig. 8).


To determine whether peripubertal binge EtOH exposure altered the normal expression levels of BDNF and SIRT1 mRNA, we compared mRNA levels in untreated animals at each age (early, mid, and late puberty) to animals that had been treated with binge EtOH at peripuberty in both the dorsal and ventral hippocampus. Overall, there was a significant main effect of age on BDNF mRNA expression in both regions of the hippocampus (Tables 1, 2; Fig. 9A, B, solid line). By contrast, there was a significant main effect of age on SIRT1 mRNA expression in the ventral, but not dorsal, hippocampus (Tables 1, 2; Fig. 9C, D, solid line). There was also a significant main effect of EtOH treatment on BDNF and SIRT1 mRNA expression in both hippocampal regions (Tables 1, 2). A statistically significant interaction between age and EtOH treatment was observed for SIRT1 in the dorsal hippocampus (Table 1) and BDNF and SIRT1 in the ventral hippocampus (Table 2), demonstrating that the effects of binge EtOH exposure on BDNF and SIRT1 mRNA expression was dependent on age. Interestingly, EtOH exposure significantly increased BDNF and SIRT1 mRNA levels compared to untreated animals at peripuberty in the dorsal hippocampus (Fig. 9A, C, dashed line), while the opposite was observed in the ventral hippocampus (Fig. 9B, D, dashed line). The effects of EtOH persisted for up to one-month post-EtOH exposure for BDNF and SIRT1 in the dorsal hippocampus (Fig. 9A, C) and for BDNF in the ventral hippocampus (Fig. 9A). Although the ventral hippocampus mRNA levels of SIRT1 were not statistically different from untreated controls one-month following binge EtOH exposure, the data suggest that the normal developmental profile of SIRT1 gene

expression was retarded at a pre-pubertal phenotype as a result of peripubertal EtOH exposure (Table 3).

We next investigated the effects of peripubertal binge EtOH exposure on BDNF and SIRT1 protein expression. We quantified the dorsal and ventral hippocampus pro-BDNF (the higher molecular weight precursor of mature BDNF), mature BDNF and SIRT1 protein levels in peripubertal animals treated with water or binge EtOH one hour following the last treatment.. In the ventral hippocampus, binge EtOH-treated animals demonstrated a strong decreasing trend in expression of pro-BDNF, with a p-value = 0.05 (Fig. 10B).


#### **Binge EtOH exposure did not alter circulating testosterone levels.**

Increased gonadal steroid hormones during pubertal development can potentially modulate microRNA and/or Drosha, Dicer, BDNF and SIRT1 mRNA levels. The animals in this study were kept gonad-intact throughout puberty, however previous studies have demonstrated that EtOH can alter gonadal steroid hormones [44]. To determine the effects of binge EtOH exposure during puberty on circulating gonadal steroid hormone levels in our system, plasma testosterone (T) was measured in each age group on the day of sacrifice. As expected, circulating T levels continued to increase with age in all animals (Fig. 11), demonstrating a normal progression through pubertal development. Peripubertal EtOH exposure tended to decrease circulating T levels, but the differences between EtOH-treated animals and controls of the same age group were not statistically



**Figure 10.** Assessment BDNF and SIRT1 protein expression levels following peripubertal binge EtOH. Representative immunoblot for BDNF, pro-BDNF, SIRT1 and  $\beta$ -actin expression in the dorsal hippocampus and the % change in BDNF, pro-BDNF and SIRT1 expression normalized to  $\beta$ -actin (A). Representative immunoblot for BDNF, pro-BDNF, SIRT1 and  $\beta$ -actin expression in the ventral hippocampus and the % change in BDNF, pro-BDNF and SIRT1 expression normalized to  $\beta$ -actin (B). Quantification of densitometric analysis of protein expression calculated from at least 3 independent experiments (N=6). No significant difference between groups was observed (2-way ANOVA,  $p < 0.05$ ).





**Figure 11.** Peri-pubertal binge EtOH exposure did not affect circulating testosterone levels. Plasma concentrations of testosterone (T) 60 min. after the last treatment. Data expressed as mean  $\pm$  SEM T pg/ml. No statistically significant differences were observed.

significant (Fig. 11). Moreover, there were no apparent long-lasting effects of peripubertal EtOH treatment on circulating T levels measured at one month post EtOH treatment (Fig. 11).

## **Discussion**

Adolescent alcohol abuse has been shown to exert long-lasting detrimental effects on brain function, neuronal gene expression and behaviors, yet the precise molecular targets of EtOH remain poorly understood. Indeed, previous studies by our laboratory and others have demonstrated both immediate and long-term effects of repeated peripubertal binge EtOH exposure on genes that regulate the physiological stress response [12,44,67,212,213]. Therefore, the goals of this study were to provide a potential mechanistic explanation for EtOH-induced effects on gene expression by quantifying the expression of EtOH sensitive microRNAs (miR-10a-5p, miR-26a, miR-103 and miR-495) during normal pubertal development in the male rat hippocampus, and then elucidate how peripubertal binge EtOH exposure alters the expression of those microRNAs. Importantly, microRNAs have emerged as highly conserved critical regulators of downstream gene expression in nearly all physiological systems. Genome-wide microRNA expression profiles revealed that the microRNAs investigated in this study, miR-103 and miR-26a, are among the top 15 most abundantly expressed microRNAs in the rodent hippocampus [104]. Taken together our data revealed that the expression of miR-10a-5p, miR-26a, miR-103 and miR-495 are dynamic across pubertal development and that the developmental profiles for each microRNA are distinct between the dorsal and ventral hippocampus. Moreover, peripubertal binge EtOH exposure altered normal pubertal development expression patterns of miR-10a-5p, miR-26a, miR-103,

miR-495, Dicer, Drosha, BDNF and SIRT1 in an age and brain region-dependent manner. Most striking, our results showed that peripubertal binge EtOH exposure had significant long lasting effects on several microRNAs studied, as well as their processing enzymes and target genes. These effects were evident for as long as one-month following the last EtOH exposure, suggesting that EtOH could have lasting consequences on gene expression profiles in the male rat hippocampus through long-term regulation of microRNA expression patterns.

Quantifying EtOH-induced changes in Drosha and Dicer mRNA levels can yield insight into the mechanistic actions of EtOH by revealing specific points of EtOH-mediated perturbations along the microRNA biosynthetic pathway. The biogenesis of mature 22-24 nucleotide (nt), single-stranded microRNAs involves the following sequential processes: 1) transcription of a 100-1000 nt primary-microRNA (pri-microRNA) product, 2) cropping of the transcript by the nuclear RNase III enzyme Drosha into the preliminary-microRNA (pre-microRNA), 3) cleavage of the pre-microRNA by the cytoplasmic RNase III enzyme Dicer and 4) loading of the mature microRNA onto the RNA-induced silencing complex (RISC), which guides it to its mRNA target for degradation or translational inhibition [210,211]. Notably, the expression of microRNA biogenesis genes (i.e. Drosha, Dicer) was shown to be significantly correlated with addiction-related phenotypes [214]. Our studies revealed the interesting observation that both Drosha and Dicer mRNA significantly decreased between early and peripuberty, although these results do not necessarily reflect changes in enzyme catalytic activity. Regardless, decreased mRNA levels of microRNA biosynthetic enzymes would theoretically result in reduced mature microRNA levels

leading to increased translation of gene targets, consistent with the idea that there are global changes in overall gene expression during adolescent development. The effects of peripubertal binge EtOH exposure on Drosha and Dicer mRNA levels continued to persist for as long as one-month after the last EtOH exposure, which raises the possibility of a potential long-term EtOH-induced dysregulation of microRNA biosynthetic processing.

One of the biggest challenges since the discovery of microRNAs has been the identification of their target genes. In mammals, their imperfect base pair hybridization with mRNA targets results in promiscuous binding, such that a single microRNA can have multiple mRNA gene targets. For instance, miR-495 is predicted to target as many as 754 genes in the rat genome (miRDB). Similarly, a single mRNA transcript can be regulated by several different microRNAs and whether multiple microRNAs must act in concert to regulate a specific target gene remains unresolved. Indeed, BDNF is predicted to be targeted by 51 microRNAs (miRDB), therefore the results shown herein are not exhaustive of all potential regulators of BDNF. Nevertheless, our data demonstrating differential expression of the same microRNA in functionally distinct hippocampal regions strongly implied that the targets of these microRNAs may also be differentially expressed. Therefore, we identified putative target genes of miR-10a-5p, miR-26a, miR-103 and miR-495 that were relevant to known dorsal and ventral hippocampus functions using target prediction software programs, Targetscan and MirDB. For instance, BDNF was identified as a putative target gene for miR-10a-5p, miR-26a, miR-103 and miR-495. BDNF plays a fundamental role in guiding neurodevelopment as well as in the fine-tuning of synaptic plasticity, a critical event during adolescent brain development.

Previous studies have demonstrated that regulation of BDNF expression is mediated by miR-26a targeting the conserved BDNF 3'UTR sequence [143] and our data demonstrate inverse relationships between miR-26a and BDNF expression in the dorsal hippocampus (Fig. 5B, Fig. 9A). Together, these data suggest a role for miR-26a in the regulation of BDNF expression levels in the pubertal dorsal hippocampus. Importantly, both BDNF gene variants and miR-26a expression have been implicated in the vulnerability and onset of schizophrenia, alcohol abuse and mood disorders in both human patients and rodent models [143,144,145,146,147,148]. Moreover, these data serve as an important reminder that the computer algorithm-predicted microRNA target genes do not always translate into actual microRNA targets and that correlated changes in microRNA and target gene mRNA expression levels do not always indicate causality. Taken together, peripubertal disruption of miR-26a, miR-10a-5p and miR-495 expression following binge EtOH exposure could result in altered BDNF expression in the pubertal hippocampus.

The histone deacetylase sirtuin 1 (SIRT1) was also predicted by computer algorithms to be a putative gene target of miR-26a, mir-103 and miR-495 (Fig. 7B). Immediate responses to peripubertal binge EtOH resulted in increased SIRT1 expression in the dorsal hippocampus, and decreased SIRT1 expression in the ventral hippocampus. These data suggest that SIRT1 mRNA expression may be regulated by distinct microRNAs in the dorsal and ventral hippocampus, a hypothesis consistent with studies demonstrating that these two hippocampus regions are functionally distinct, as they have dissimilar neuronal projection patterns [57,58,205]. Moreover, the ventral and dorsal hippocampus may utilize different cellular mechanisms to mediate their respective responses to EtOH. Our data further demonstrated that peripubertal binge EtOH exposure

had long-lasting effects on SIRT1 gene expression, showing significantly increased mRNA levels lasting until late puberty in the dorsal hippocampus. Similar to BDNF, SIRT1 has recently been implicated as critical for mediating synaptic plasticity, one mechanism underlying memory formation in rodent and human cell models [23,25]. Moreover, many studies have demonstrated that EtOH exposure has long-lasting consequences on gene expression, which may be regulated, in part, by microRNAs targeting chromatin-modifying enzymes, such as SIRT1.

Regulation of gene expression mediated by microRNAs is theoretically carried out via the alterations in the microRNA target gene effector functions, i.e. the function of the protein encoded by the microRNA target gene. Effector function is best understood to occur following sequestration or degradation of the microRNA target gene mRNA. However, a single gene can be targeted and, therefore, regulated by multiple microRNAs suggesting that binding of multiple microRNAs to a single target gene's 3'UTR could be necessary for alterations in protein expression to mimic changes in microRNA target gene mRNA expression. Indeed, our data make evident that miR-10a-5p, miR-495, miR-103 and miR-26a expression are sensitive to the presence or absence of EtOH depending on age and brain region and that consequential alterations in BDNF and SIRT1 target gene expression are significantly correlated with observed EtOH-induced alterations in the expression of some of the microRNAs that target them in an immediate manner. The near significance values of BDNF and SIRT1 protein data suggest many different possibilities. One interpretation is that the protein levels of BDNF and SIRT1 following peripubertal binge EtOH may be altered to differing degrees amongst individuals. Indeed, unlike other drugs such as cocaine, alcohol alters cells membranes, ion channels,

enzymes, and receptors of neurons globally throughout the brain [215], and likely contribute to evidenced individual, sex and racial variations in alcohol metabolism and alcohol-induced neuroplastic changes [216,217] [218,219,220,221]. Importantly, the six rats whose tissue was collected for protein analysis were not from the same cohort of animals collected for RNA analysis. Furthermore, the groups of animals collected for RNA analysis had an N=10, whereas the groups collected for protein analysis had N=6. It is therefore, possible that there was not enough power to detect changes in protein expression levels, especially considering that protein levels change at a kinetically slower rate than mRNA. Alternatively, because any given miR targets multiple microRNAs, it is possible that target genes other than BDNF and SIRT1 may be altered by peripubertal binge EtOH exposure. For instance, miR-10a plays a role in regulating the homeobox developmental genes [21], and has similar expression patterns to *HOXB4* in the mouse embryo suggesting that it may have similar important functional roles in development [20,222], and therefore may have pubertal brain developmental functions. Furthermore, a member of the Nkx family of homeobox genes, thyroid transcription factor 1 (TTF1), is expressed in the postnatal hypothalamus and this expression is required for rodent pubertal onset [222]. Therefore, it is possible that miR-10a-5p targets Hox gene transcription factors expressed during pubertal development, and alterations of target genes important for morphogenesis may contribute to disruption of brain development following peripubertal binge EtOH abuse. Lastly, Overall, the molecular mechanisms underlying differences in microRNA responses to pubertal binge EtOH warrant further studies and as it is possible that changes in microRNA expression may lead to complications in downstream target gene translation in some but not all individuals

engaging in binge drinking behavior during adolescence. Therefore, given the strong supporting evidence that changes in the expression of EtOH sensitive microRNAs significantly correlate with changes in target gene mRNA, and that corresponding changes in the protein expression reach near significance, it is important that further experimentation explore the effects of peripubertal binge EtOH exposure on protein expression changes in BDNF and SIRT1.

Overall, our data reveal novel findings about the age and brain-region specific expression of miR-10a-5p, miR-26a, miR-103, and miR-495 during pubertal development in male rats. Further, we showed that mid/peripubertal binge EtOH exposure significantly alters the normal expression profile of these microRNAs, their biosynthetic processing enzymes, and two of their putative target genes, BDNF and SIRT1. However, it is important to note that these results are not necessarily predictive of microRNAs in females or other species, as several studies have demonstrated both species and sex-specific expression profiles for microRNAs [150,223,224]. Moreover, the precise molecular targets of EtOH in the biogenesis of microRNAs remain unclear and require further investigation. An important next step is the identification of specific cell types (i.e. neurons vs. glia) in which microRNAs are affected by binge EtOH exposure, as our study was limited to whole hippocampal tissue homogenates. Taken together, our data raise the possibility that EtOH modulation of these four microRNAs is a potential cellular mechanism underlying long-term changes in gene expression induced by adolescent EtOH abuse.



## CHAPTER FOUR

### SEXUALLY DIMORPHIC GENE EXPRESSION OF HIPPOCAMPUS ETOH-SENSITIVE MICRORNA, MICRORNA TARGET GENES AND MICRORNA PROCESSING ENZYMES DURING PUBERTAL DEVELOPMENT

#### **Introduction**

Adolescent binge drinking is a major public health issue [40,166] that incurs sexually dimorphic consequences on brain function and development [44,225,226,227,228,229,230]. Both male and female adolescents engage in underage drinking, and the Substance Abuse and Mental Health Services Administration (SAMHSA) has reported that males begin drinking earlier, drink more frequently and have a higher prevalence of binge drinking than their female counterparts (SAMHSA 2006) [231,232]. However, more recently the Center for Disease Control and Prevention (CDC) reported that binge drinking among females--including young girls--is also a serious, yet unrecognized, problem (CDC 2012). Moreover, studies have demonstrated that females who abuse alcohol are more prone to developing alcohol addiction, alcoholism and anxiety than men [68,72]. Sex hormones have been demonstrated to mediate, in part, sexually dimorphic responses to EtOH [233,234,235,236,237], however, a clear mechanism remains unknown.

Sex hormones can also influence microRNA expression in the brain [28,150]. microRNAs are small (~20 nt) non-coding RNA molecules encoded in the genome. Their synthesis begins in the nucleus where RNA Polymerase II or III transcribes a primary-

microRNA (pri-microRNA). The pri-microRNA is then processed by the nuclear enzyme Drosha into the precursor-microRNA (pre-microRNA) which can exit the nucleus. Once in the cytoplasm, the pre-microRNA is cleaved by the endoribonuclease Dicer into a small duplex. One strand of this duplex is the mature microRNA that then associates with the RNA-induced silencing complex (RISC) which guides the microRNA to the 3' untranslated region (UTR) of a target gene to ultimately induce gene silencing by either translational repression or mRNA cleavage. In the brain, microRNAs have been shown to regulate neuronal differentiation [238], neuronal survival [112] and synaptic plasticity [113,136]. Interestingly, both sex hormones and microRNAs have been demonstrated to govern synaptic plasticity and the development of sexually dimorphic psychiatric disorders during puberty [113,151,239,240]. These data further suggest that microRNAs, in addition to sex hormones, may work in concert to contribute to pubertal brain development.

microRNA expression is sensitive to EtOH in a variety of species [157,158,241,242], neural cultures [195,243,244], and at developmental stages [193,245,246]. We previously identified 5 microRNAs as having age and brain region-dependent expression profiles in the male rodent hippocampus that was also altered by peripubertal binge EtOH [247]. Other studies have also demonstrated that hypothalamic microRNAs are important regulators of pubertal onset in rats, and that this is true for both males and females [248]. In order to better understand how microRNA-based regulation of gene expression may play a role in the sexual dimorphic effects of EtOH abuse on brain development and behavior, it is essential to first establish the normative expression profile of microRNAs during healthy pubertal development in both sexes. In this pursuit,

the current study tested the hypothesis that during pubertal development, microRNA expression profiles of EtOH-sensitive microRNAs are sexually dimorphic. We further hypothesized that these pubertal microRNA expression profiles will be different in the ventral and dorsal hippocampus. In order to test our hypotheses, we quantified the expression levels of five EtOH-sensitive microRNAs, miR-10a-5p, miR-26a, miR-103 and miR-495 [247], two target genes of the investigated microRNAs, neurotrophin brain-derived neurotrophic factor (BDNF) and the histone deacetylase sirtuin1 (SIRT1), as well as the major microRNA biosynthetic processing enzymes, RNase III-type endonucleases Droscha and Dicer. Overall, we identified sex differences at multiple stages of normal pubertal development and in multiple brain regions in the gene expression of EtOH-sensitive microRNAs, their target genes and microRNA processing enzymes. We conclude that throughout puberty, the hippocampal miRNome of EtOH-sensitive microRNAs is dependent on sex and age, and suggest that both short and long-term developments within the hippocampus during pubertal development may rely on changes of microRNA expression over this period. Additionally, we determined that the microRNA expression profiles in both the male and female throughout pubertal development are distinct between the ventral and dorsal hippocampus.

### **Approach**

The dorsal and ventral hippocampus were microdissected from untreated male and female Wistar rats and collected at three different ages during pubertal development (early puberty, PND 30; mid/peripuberty, PND 44; late puberty, PND 71; N=10/sex/group). Importantly, peripuberty in rats is considered to be ~ PND 30-45 [201,202]. Quantitative real-time reverse transcription PCR (qRT-PCR) was used to

quantify the expression levels of a total of five microRNAs (miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495), and four genes (Drosha, Dicer, BDNF and SIRT1).

microRNAs were previously identified to be EtOH-sensitive in pubertal male Wistar rats [247], and in Chapter 2 preliminary data.


## Results

### **Dorsal hippocampus expression levels of EtOH-sensitive miR-10a-5p, miR-26a, and miR-495 in untreated rats are age and sex dependent.**

In the dorsal hippocampus, a 2-way analysis of variance (ANOVA) revealed significant main effects of age or sex on the expression of miR-10a-5p, miR-26a and miR-495 (Table 4). There was also a statistically significant interaction between age and sex for these 3 microRNAs, indicating that the sex differences are dependent on age (Table 4). On the other hand, there was a main effect of age alone on miR-32 expression (Table 4). miR-10a-5p and miR-32 expression levels demonstrated differences across puberty in females but not males (Fig. 12A, C). Specifically, early puberty expression levels of miR-10a-5p were significantly higher in females compared to males (Fig. 12A). By mid-puberty, the female levels decreased to match the expression levels of miR-10a-5p observed in males, which stayed constant from early to late puberty (Fig. 12A). Expression of miR-32 in females remained constant between early and mid-puberty, after which it increased significantly at late puberty (Fig. 12C). Interestingly, the expression of

**Table 4.** Statistical Analysis of Dorsal Hippocampus Gene Expression

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
miR-10a-5p	Yes: $F(2,39) = 12.892$ $p < 0.001$	Yes: $F(2,39) = 21.589$ $p < 0.001$	Yes: $p = 0.005$
miR-26a	Yes: $F(2,48) = 16.703$ $p < 0.001$	Yes: $F(1,48) = 138.623$ $p < 0.001$	Yes: $p < 0.001$
miR-103	No	No	No
miR-32	Yes: $F(2,53) = 3.379$ $p = 0.042$	No	No
miR-495	Yes: $F(2,44) = 11.972$ $p < 0.001$	Yes: $F(1,44) = 174.639$ $p < 0.001$	Yes: $p < 0.001$
<b>GENE</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
Drosha	Yes: $F(2,54) = 14.973$ $p < 0.001$	Yes: $F(2,54) = 45.914$ $p < 0.001$	Yes: $p < 0.001$
Dicer	Yes: $F(2,54) = 10.769$ $p < 0.001$	Yes: $F(2,54) = 145.010$ $p < 0.001$	Yes: $p < 0.001$
BDNF	Yes: $F(2,54) = 10.779$ $p < 0.001$	Yes: $F(1,54) = 90.143$ $p < 0.001$	Yes: $p < 0.001$
SIRT1	Yes: $F(2,54) = 92.432$ $p < 0.001$	Yes: $F(1,54) = 94.246$ $p < 0.001$	Yes: $p < 0.001$



**Figure 12.** Sexually dimorphic microRNA expression is evident during pubertal development in the dorsal hippocampus. miR-10a-5p (A), miR-26a (B), miR-32 (C), miR-103 (D) and miR-495 (E) expression levels in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 male animals. \* Dissimilar letters indicate a statistically significant difference between groups, two-way ANOVA ( $p < 0.05$ ).

miR-26a and miR-495 in males was significantly higher than in females throughout the entire duration of pubertal development (Fig. 12B, E). Moreover, relative to male at much lower levels (Fig. 12B, E). In males, both miR-26a and miR-495 expression levels increased between early and mid puberty, and by late puberty, miR-26a decreased significantly to reach an expression level lower than the early puberty level (Fig. 12B). However, by late puberty, miR-495 continued to increase with age in the males (Fig. 12E). miR-103 expression levels were not significantly different between the sexes or between stages of pubertal development in the dorsal hippocampus (Fig. 12D).

**Ventral hippocampus expression levels of EtOH-sensitive miR-10a-5p, miR-26a, and miR-495 in untreated rats are age and sex dependent.**

We next tested whether the expression levels of the five EtOH-sensitive microRNAs were sexually dimorphic in the ventral hippocampus. RT-PCR analysis followed by a 2-way ANOVA revealed that EtOH-sensitive microRNA expression patterns were sexually dimorphic throughout pubertal development in the ventral hippocampus. Furthermore, the sex difference patterns observed in the ventral hippocampus of these microRNAs throughout puberty differed from those observed in the dorsal hippocampus. For example, there was a significant main effect of age and sex on expression levels of miR-26a, miR-103 and miR-495, and a significant interaction between age and sex in miR-10a-5p, miR-26a, miR-103 and miR-495 (Table 5). Specifically, miR-10a-5p expression levels did not change statistically over the course of pubertal development in females, but miR-10a-5p expression did significantly increase between mid and late puberty in males (Fig. 13A). Furthermore, no other EtOH-sensitive microRNA demonstrated any change in the female ventral hippocampus over time and,

similar to the dorsal hippocampus, female expression levels of miR-26a and miR-495 were very lowly expressed compared to the levels in males (Fig. 13B, E). Male expression levels of miR-26a and miR-495 both decreased significantly between early to mid pubertal levels, followed by a stabilization of miR-26a expression at late puberty (Fig. 13B) and a significant increase in miR-495 expression between mid and late puberty, with levels exceeding those expressed at early puberty (Fig. 13E). In contrast, miR-103 expression levels in males remained stable between early to mid-puberty, followed by a significant increase at late puberty (Fig. 13D).


**Mature microRNA biosynthetic processing enzymes are altered by pubertal development in a sexually dimorphic manner in the dorsal and ventral hippocampus.**

We next assessed the expression of Drosha and Dicer in male and female dorsal and ventral hippocampus at early, mid and late puberty. We hypothesized that mature microRNA biogenesis enzyme expression levels are sexually dimorphic during puberty. Importantly, previous reports indicate that the expression of microRNA processing enzymes may be regulated, in part, by sex hormones or their nuclear receptors. For example, breast cancers can be distinguished by their estrogen receptor (ER) status (i.e. can either be ER positive (ER+) or negative (ER-) and indeed, ER $\alpha$  breast cancer cell lines and clinical mechanisms remains unknown [249]. Therefore, it is possible that Drosha and Dicer levels correlate with pubertal changes in gonadal hormone levels. We identified a significant main effect of both sex and age on Drosha and Dicer expression



**Table 5.** Statistical Analysis of Gene Expression Levels in the ventral hippocampus

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
miR-10a-5p	No	No	Yes: $p = 0.004$
miR-26a	Yes: $F(2,41) = 9.600$ $p < 0.001$	Yes: $F(2,41) = 59.915$ $p < 0.001$	Yes: $p = 0.002$
miR-32	No	No	No
miR-103	Yes: $F(2,53) = 13.038$ $p < 0.001$	Yes: $F(1,53) = 8.244$ $p = 0.006$	Yes: $p = 0.002$
miR-495	Yes: $F(2,41) = 22.776$ $p < 0.001$	Yes: $F(1,41) = 136.974$ $p < 0.001$	Yes: $p < 0.001$
<b>GENE</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
Drosha	Yes: $F(2,54) = 4.401$ $p = 0.017$	Yes: $F(2,54) = 21.607$ $p < 0.001$	Yes: $p = 0.006$
Dicer	Yes: $F(2,54) = 5.547$ $p = 0.006$	Yes: $F(2,54) = 53.759$ $p < 0.001$	Yes: $p = 0.003$
BDNF	Yes: $F(2,45) = 4.310$ $p = 0.019$	Yes: $F(1,45) = 14.607$ $p < 0.001$	Yes: $p = 0.009$
SIRT1	Yes: $F(2,47) = 28.346$ $p < 0.001$	Yes: $F(1,47) = 53.143$ $p < 0.001$	Yes: $p < 0.001$




**Figure 13.** Sexually dimorphic microRNA expression is evident during pubertal development in the ventral hippocampus. miR-10a-5p (A), miR-26a (B), miR-32 (C), miR-103 (D) and miR-495 (E) expression levels in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 male animals. Dissimilar letters indicate a statistically significant difference between groups ( $p < 0.05$ ).

levels, as well as a significant interaction between age and sex in both the dorsal and ventral hippocampus (Table 4, Table 5). Interestingly, female expression levels of Drosha and Dicer in the dorsal and ventral hippocampus were lower than the expression levels in males (Fig. 14), except for one stage of pubertal development in the ventral hippocampus wherein male and females had similar Drosha expression levels (Fig. 14B). Furthermore, significant differences across age were apparent in male Drosha and Dicer expression levels in both hippocampus regions. Between early and mid-puberty, the male Drosha and Dicer expression levels decreased and remained decreased throughout late puberty in the dorsal hippocampus (Fig. 164, C) and significantly increased between mid puberty to late puberty in the ventral hippocampus (Fig. 14B, D). In stark contrast, the female microRNA biogenesis enzymes varied very little throughout pubertal development, such that only in the ventral hippocampus, did Drosha expression levels significantly decrease between early and peripuberty, followed by a further decrease below early pubertal levels at late puberty (Fig. 14B).


**Expression levels of EtOH-sensitive microRNA target genes are differentially dependent on age and sex in the dorsal and ventral hippocampus.**

The neurotrophic factor BDNF and the histone deacetylase SIRT1 are both neuroprotective. BDNF mediates synaptic plasticity and hippocampus-dependent memory by supporting neurotransmitter release and inducing long-term potentiation [84,85,253]. SIRT1 also contributes to hippocampus-dependent learning and memory as well as regulates anxiety [23,25,102,254]. Importantly, memory and mood are partially controlled by the dorsal and ventral hippocampus, respectively. We have previously



**Figure 14.** Sexually dimorphic microRNA processing enzyme expression is evident during pubertal development in the dorsal and ventral hippocampus. Drosha (A, C) and Dicer (B, D) expression levels in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 male animals. Dissimilar letters indicate a statistically significant difference between groups ( $p < 0.05$ ).

demonstrated that alterations in the expression of microRNAs following peripubertal binge EtOH exposure in the male hippocampus significantly correlated with alterations in the expression levels of the microRNA target genes BDNF and SIRT1 [247]. The regulation of memory and mood are impaired by adolescent EtOH abuse [255,256] and such effects can also present in a sexually dimorphic manner [228,257] yet the mechanisms governing such processes remain unclear. Therefore, we hypothesized that the EtOH-sensitive microRNA expression previously reported in males would be differentially expressed in females across pubertal development and that the ventral and dorsal hippocampus would display region-specific sex differences. Our results indicate a statistically significant main effect of both age and sex on the expression levels of SIRT1 and BDNF in both the dorsal and ventral hippocampus, as well as a statistically significant interaction between age and sex (Table 4 and Table 5). In the female dorsal hippocampus, SIRT1 expression was dynamic across puberty such that it decreased between early and mid puberty and remained lowly expressed into late puberty (Fig. 15C). The same pattern throughout puberty was also observed for male ventral hippocampus SIRT1 expression, however, the level of SIRT1 was greater in males than in females at each time point (Fig. 15C). Overall, females expressed these EtOH-sensitive microRNA target genes at much lower levels relative to males and furthermore, expression levels in males demonstrated a great deal of variation over the course of pubertal development (Fig. 15). Interestingly, the expression patterns of BDNF in the ventral and dorsal hippocampus and of SIRT1 in the ventral hippocampus over puberty were similar, such that between early and mid puberty there was an increased in expression, which then lowered back to early pubertal levels at late puberty (Fig. 15A, B,



**Figure 15.** Sexually dimorphic microRNA target gene expression is evident during pubertal development in the dorsal and ventral hippocampus. BDNF (A, B) and SIRT1 (C, D) mRNA levels in the dorsal and ventral hippocampus in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to untreated PND 30 male animals. Dissimilar letters indicate a statistically significant difference between groups ( $p < 0.05$ ).

D). This similar pattern suggests that mid puberty is a possible critical window of time during which high SIRT1 and BDNF expression is critical for developmental processes in the male hippocampus.

## **Discussion**

In the current study we tested the hypothesis that EtOH-sensitive microRNAs exhibit sexually dimorphic expression profiles in the ventral and dorsal hippocampus during pubertal development. The dorsal hippocampus primarily operates learning and memory while the ventral portion regulates emotional memory and mood [57,58,258]. The continuous development of neuronal circuitry throughout pubertal development [43,259] is partially regulated by gonadal hormones [47,233,236,260,261]. Gonadal hormone signaling via nuclear receptors has also been demonstrated to influence the expression of microRNAs [165,262,263,264,265,266]. Therefore, gonadal hormone exposure may mediate hippocampal maturation during puberty via downstream microRNA effector molecules. Furthermore, the rise in sex steroids during pubertal development may contribute to a microRNA-mediated regulation of the sexually dimorphic development of memory capacities and emotional responses during this period. Bioinformatic analysis predicted that miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495 bind to the 3' UTRs of genes regulating synaptic plasticity, including SIRT1, BDNF and synapsin II (SYN2). Four of these five microRNAs have recently been identified to demonstrate dynamic expression patterns throughout healthy pubertal development in the male rat hippocampus and furthermore, their normal developmental expression profile is disrupted following peripubertal binge EtOH, a typical adolescent behavior [247].

The main goal of this study was to quantify sex differences in the expression levels of miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495, the microRNA target genes SIRT1 and BDNF and the major microRNA biosynthetic processing enzymes Drosha and Dicer in the pubertal dorsal and ventral hippocampus. Overall, this study demonstrates that miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495 are expressed in an age and sex-specific manner during intact pubertal development and that these sex differences are distinct between the dorsal and ventral hippocampus. Furthermore, we quantified, to our knowledge for the first time, sex differences in the expression levels of BDNF, SIRT1, Drosha and Dicer mRNA in the dorsal and ventral hippocampus at three stages of pubertal development, and identified significant sex differences in expression levels throughout puberty and between the ventral and dorsal hippocampus. Taken together, sexually dimorphic expression levels of EtOH-sensitive microRNAs, their gene targets and microRNA processing enzymes during pubertal development suggests that the expression of these microRNAs may be influenced by sex hormones. Furthermore, the sex, age and hippocampus region-dependent fluctuations in EtOH-sensitive microRNAs and their target genes suggest that sex differences in pubertal microRNA expression may underlie the sexual dimorphisms observed in response to stress and alcohol abuse, suggesting that they may influence the development of anxiety and depression-related disorders during this period of life.

The hippocampus plays a major role in regulating sexually dimorphic behaviors, such as learning, memory and the stress response. For instance, males perform better than females on hippocampus-dependent spatial memory acquisition tests despite the well-known positive effects that E<sub>2</sub> imparts on hippocampus-dependent function in females.



Indeed, it has been proposed that E<sub>2</sub> and T differentially influence memory functions in males and females [267,268]. It is likely that the hormonal processes of sexual differentiation during development creates organizational differences in neural substrates and that these organizational differences underlie the sexually dimorphic effects of E<sub>2</sub> and T on cognitive tasks in males and females as well as contribute to the different strategies males and females use to solve similar tasks [269]. However, it is difficult to ascertain clear sexually dimorphic effects that gonadal hormones have on hippocampus-mediated cognition in males and females because hormonal influences can differ with task type, task aspect and the degree to which the task relies on one or more brain region. In contrast, clear responses to gonadal hormone treatment can be observed when investigating morphological, anatomical and cellular aspects of hippocampus-mediated functions. For instance, synaptic plasticity in males and females relies on gonadal hormones. Gonadectomy (GDX) in males reduces spine density (structures that receive excitatory inputs and therefore serve as a measurement of synaptic plasticity and memory) and while E<sub>2</sub> treatment fails to restore this loss [270], androgen receptor activation increases spine density via up-regulation of N-methyl-D-aspartate receptors (NMDARs) [270,271,272]. In contrast, female exposure to androgens and endogenous and exogenous E<sub>2</sub> leads to increased spine density [270,272,273,274] and the electrophysiological measurement of learning and memory, long-term potentiation (LTP) [275,276]. These data demonstrate that differential mechanisms are initiated by AR and ER activation to mediate synaptic plasticity in males and females. Therefore, it is not unusual that we observed sex differences in the expression of hippocampal microRNAs (Fig. 12 and Fig. 13), as they may play a role in mediating signaling pathways

responsible for normal sexually dimorphic brain development during puberty as well as in the sexually dimorphic responses to peripubertal binge EtOH [44].

Our current understanding of how peripubertal binge EtOH affects the male and female developing brain differently is incomplete, and therefore requires further investigation into the potential molecular targets of binge-pattern of EtOH abuse. In our study we observed relatively low expression levels in the female hippocampus of the microRNAs sensitive to pubertal binge EtOH compared to males (Fig. 12 and Fig. 13). Indeed, it is possible that different microRNAs are sensitive to EtOH in males than in females. Alternatively, microRNAs may not be involved in mediating the effects of binge EtOH on the stress response or memory capacity in females, a theory in line with previous findings that binge EtOH exposure-mediated alterations in genes regulating the stress response were observed in pubertal males but not in females [44,47]. It is possible that sex differences in microRNA expression results from differences in the rates by which males and females reach sexual maturity, regarding reproductive function. Indeed, females typically develop reproductive capacity earlier than males [17,277]. While the rates of maturation for adult-like memory processing and stress responsiveness in males and females are not clearly delineated, some studies support the notion that observed sex differences in the pubertal stress response [278] are indeed established in the brain prior to pubertal onset in females [279]. These data suggest that microRNAs, microRNA biogenesis enzymes, BDNF and SIRT1 expression levels are sexually dimorphic during pubertal development and distinct between the dorsal and ventral hippocampus. Because the ventral hippocampus regulates the stress response, it is possible that the sexually dimorphic expression levels of microRNAs may play a role in sex differences in mood

and emotional memory regulation. A more completely developed female stress response and/or synaptic plasticity at pubertal onset may also underlie observed sex differences in the mechanisms employed by AR and ER activation-induced synaptic plasticity in the male and female adolescent brain, respectively. Taken together, these data underscore the need to better understand whether the observed sex differences in hippocampal microRNA expression may contribute to the sexually dimorphic adolescent responses to stressors such as binge alcohol exposure.

The biogenesis of mature microRNAs relies on the RNase III enzymes, Droscha and Dicer which generate the pre-microRNA and microRNA, respectively. In cancer cell models, E<sub>2</sub> interferes with microRNA biogenesis by acting via estrogen receptors  $\alpha$  and  $\beta$  (ER $\alpha$ , ER $\beta$ ), transcription factors of the nuclear receptor family of homeostatic regulators [161,264]. For instance, E<sub>2</sub>-induced ER $\beta$  signaling in an E<sub>2</sub>-responsive cancer cell line lead to an accumulation of pre-microRNA transcripts, possibly due to an ER $\beta$ -mediated release of Droscha from sequestration in an inhibitory chromatin complex [161]. Castellano et al. also observed alterations in pri-microRNA precursors more often than in the mature microRNAs derived from it following E<sub>2</sub>-induced nuclear receptor signaling in breast cancer cell lines, which suggests that the regulation of microRNA expression occurs at the level of microRNA biogenesis [264]. The details of how particular microRNAs are altered by nuclear receptor-mediated regulation of Droscha and/or Dicer activity remains unclear, yet these data set precedence for further investigation into the mechanisms by which mature microRNA expression can be regulated. Importantly, estrogen receptor-mediated regulation of microRNA expression has not been confirmed in the brain. However, it is possible that sex differences in hippocampal microRNA

expression throughout rat pubertal development are influenced by the regulation of the microRNA biogenesis. We demonstrate here that expression levels of Drosha and Dicer mRNA in males was overall much greater when compared to females (Fig. 14). Furthermore, the male expression levels of these enzymes fluctuated over pubertal development while the female expression levels of Drosha and Dicer had little to no changes over time (Fig. 14). The fluctuations in expression levels of Drosha and Dicer, or the lack thereof, in male and females respectively, do not correlate with the patterns of microRNA expression level changes during pubertal development in either sex. However, recent studies have repeatedly demonstrated in [280,281] multiple tissues, that models of Drosha and Dicer knockouts, knockdowns or mutants alters the expression of only subset of microRNAs [130,282,283]. For instance, homozygous Dicer mutant zebrafish demonstrate arrested growth and early death, yet ablation of Dicer does not induce complete abolishment of mature microRNA expression and the authors conclude that a maternal contribution of Dicer may maintain the biogenesis mature microRNAs [284]. An additional study examined a Dicer exon 5-deficient colorectal cell line which demonstrated once again, that only 57% of known microRNAs were downregulated in their expression in the mutant cells compared to wild type [280]. In addition, Kuehbacher et al. demonstrate that Drosha and Dicer siRNA downregulates expression of only two microRNAs (let-7f and miR-27b) out of many (members of the let-7 family, miR-21, miR-126, miR-221 and miR-222) that are highly expressed in endothelial cells [282]. Therefore, although it is surprising that Drosha and Dicer expression levels don't correlate with the observed changes in microRNA expression throughout pubertal development in our study, it is becoming clear that a lack these biogenesis enzymes may

not necessarily dictate a reduction in mature microRNA expression. Furthermore, because the *in vivo* function of Drosha remains unknown and because Drosha belongs to the same class of endonucleases as Dicer, it is not completely surprising that decreases in Drosha expression also do not correlate with decreases in microRNA expression.

microRNAs are considered master regulators of the cellular transcriptome and have been repeatedly demonstrated to play critical regulatory roles in synaptic plasticity. Indeed, the highly plastic nature of the adolescent brain underlies the substantial brain maturation that takes place at this time and plays a particular role in the formation and consolidation of spatial and emotional memory. The evolutionary advantage to a microRNA-mediated regulation of synaptic plasticity is that it enables a neuron to quickly fine-tune its gene expression and protein composition in order to adapt to different environmental stimuli. We not only focused our study on identifying the expression pattern of microRNAs that are sensitive to EtOH exposure in the pubertal hippocampus, but also on the microRNA target genes important for hippocampus functions. Indeed, the two microRNA target genes of interest, BDNF and SIRT1, are also both important mediators of synaptic plasticity. In the male ventral hippocampus we demonstrated that the significantly dynamic expression patterns of BDNF and SIRT1 correlated with the significant changes in expression levels of miR-26a, miR103 and miR-495 throughout pubertal development (Fig. 13B, D, E and Fig. 15B, D). Importantly, using target prediction software programs, Targetscan and MirDB, we identified BDNF and SIRT1 are targets of miR26a, miR-103 and miR-495. Moreover, in the male dorsal hippocampus, decreased SIRT1 expression levels during pubertal development correlated with increased miR-495 expression levels (Fig. 13E and Fig.

15C). These data suggest that miR-26a, miR-103 and miR-495 mediate important developmental roles in the pubertal male ventral and dorsal hippocampus, possibly by regulating the expression levels of BDNF and SIRT1. Consistent with this interpretation, previous studies have experimentally confirmed that miR-26a binds the 3'UTR of BDNF [143]. Also a separate study, by using massive parallel sequencing demonstrated that miR-26a is abundantly expressed in the mouse hippocampus and by using integrated genomics demonstrated that miR-26a shares a seed sequence with other microRNAs targeting genes important for hippocampus function [285]. Overall, female expression levels of EtOH-sensitive microRNAs, BDNF and SIRT1 did not correlate with one another in this study (Fig. 12, 13 and 15). Moreover, the expression levels of SIRT1 and BDNF were extremely low and nearly unchanging in the female hippocampus compared to males throughout pubertal development (Fig. 15). These “negative” data most likely do not reflect a lack of dynamic synaptic plasticity, but rather suggest that SIRT1 and BDNF are not the primary mediators of synaptic plasticity in the female pubertal hippocampus. Indeed, the female hippocampus may utilize different mechanisms responsible for synaptic plasticity. This interpretation is supported by previous work identifying that AR and ER in males and females, respectively, initiate different mechanisms to regulate synaptic plasticity [270,272,273,274,275,276]. Indeed, BDNF is well known to be regulated by gonadal hormones and it is possible that the differences we detect in pubertal male and female hippocampus BDNF expression is a reflection of respective male and female gonadal hormones partially directing different mechanisms of synaptic plasticity between the sexes. Importantly, gonadal hormone levels were not directly tested because pubertal gonadal hormone levels have been well characterized

[286,287,288]. Overall, this chapter determined that EtOH-sensitive microRNAs are expressed in different developmental and regional patterns within the male and female pubertal hippocampus, and suggests that these microRNAs may differentially regulate their gene targets in the pubertal male and female brain.

## CHAPTER FIVE

### SEX DIFFERENCES IN HIPPOCAMPUS 17 $\beta$ -ESTRADIOL-RESPONSIVE MICRORNA EXPRESSION DURING PUBERTAL DEVELOPMENT OF THE WISTAR RAT

#### **Introduction**

Adolescence is a dynamic period of life during which the brain undergoes countless physiological and psychological developments. Pubertal maturation is the most clear physiological process that transpires during adolescence, one hallmark of which is the rapid rise in the gonadal hormones 17 $\beta$ -estradiol (E<sub>2</sub>) and testosterone (T) in females and males, respectively. Puberty is initiated, in part, by activity of the neuroendocrine hypothalamo-pituitary-gonadal (HPG) axis. In turn, gonadal hormones induce physiological changes in brain structure and gene expression, which are ultimately reflected in the functional development of adultlike behaviors. Indeed, differences in circulating gonadal hormones between males and females has long been supported to underlie sexually dimorphic brain morphology [289,290,291,292], neurochemistry [293,294,295,296,297,298], and function [294,299] in most species.

Classical gonadal hormone action throughout the body and brain involves the binding of gonadal hormones to nuclear receptors which, upon activation, can regulate gene transcription. More recently, E<sub>2</sub> has been of particular interest regarding its influence on the transcription of microRNAs [19,223]. Recently, Rao et al. demonstrated that microRNA expression is altered by E<sub>2</sub> in an age-dependent manner in the



hippocampus of young and aged female rats [28], which suggests that E<sub>2</sub> can influence microRNA expression levels differentially throughout the lifespan. Indeed, experimental manipulation of puberty at different ages also alters microRNA expression in the male rodent [248]. Furthermore, alterations in brain microRNA expression are associated with sexually dimorphic psychiatric disorders that develop during adolescence [182]. For example, miR-30b expression is lower in female schizophrenic brains compared to males and this expression is associated with mutations in the human estrogen receptor gene, *Esr1* [151]. Moreover, miR-30b sensitivity to E<sub>2</sub> in the brain suggests that E<sub>2</sub>-sensitive microRNA may play a role in the sexual dimorphic etiology of psychiatric disorders. The adolescent alterations in neurocircuitry and neurochemistry which contribute to the development of adultlike behaviors undoubtedly require a high degree of finely orchestrated synaptic plasticity and regulation of gene expression during this important developmental period. It is possible that the remarkable neuronal plasticity observed in the adolescent brain is mediated, in part, due to gonadal hormone-signaling via microRNAs.

Gonadal hormones and microRNAs have both been observed to influence synaptic plasticity, however, whether they do so during adolescence is not known. In the hippocampus, E<sub>2</sub> alters synapse formation by increasing expression of pre- and post-synaptic proteins and thereby increasing dendritic spine formation *in vivo* [300,301,302,303,304]. In addition, miR-132 promotes dendrite growth in response to neuronal stimulation, regulating synaptic plasticity in a p250GAP-dependent mechanism [305]. That microRNAs regulate synaptic plasticity and are differentially expressed in psychiatric disorders suggests that they are active players in adolescent brain

development. Moreover, differences in gonadal hormone levels between males and females may reflect differences in microRNA expression.

Overall, the roles that microRNAs play in postnatal brain development and maturation still requires a great deal of investigation. Given that the hippocampus plays a critical role in synaptic plasticity, a process required for pubertal neuronal rewiring and maturation of adultlike behaviors, it is possible that the expression of microRNAs in the hippocampus are appropriately poised to play a crucial role in regulating the biological processes involved in pubertal neuronal development. To explore this theory, we hypothesized that hippocampal expression of E<sub>2</sub>-sensitive microRNAs is sexually dimorphic throughout pubertal development and that this expression is distinct between the dorsal and ventral hippocampus. To test this hypothesis, we quantified the ventral and dorsal hippocampus expression levels of 5 microRNAs discovered by Rao et al. to be altered by E<sub>2</sub> dependent on age in the female brain: let-7i, miR-7a, miR-9, miR-125a and miR-181a. The ventral and dorsal hippocampus were examined separately because the dorsal hippocampus' primary function is that of spatial memory acquisition, distinguishing it from the ventral hippocampus which regulates emotional memory and affective processing via its many projections to amygdalar nuclei [57,58,205]. Importantly, the normal developmental expression patterns these specific microRNAs have not been documented in either sex at any stage of puberty, nor between distinct functional regions of the hippocampus.

### **Approach**


The dorsal and ventral hippocampus were microdissected from untreated male and female Wistar rats and collected at three different ages during pubertal development

(early puberty, PND 30; mid/peripuberty, PND 44; late puberty, PND 71; N=10/sex/group). Importantly, peripuberty in rats is considered to be ~ PND 30-45 [201,202]. Quantitative real-time reverse transcription PCR (qRT-PCR) was used to quantify the expression levels of a total of five microRNAs: (let-7i, miR-7a, miR-9, miR-125a, and miR-181a). microRNAs were previously identified to be 17 $\beta$ -estradiol-sensitive in female rats by Rao et al. [247].

## Results

### **let-7i, miR-7a, miR-9, miR-125a, and miR-181a expression levels throughout pubertal development in the dorsal hippocampus are dependent on age and sex.**

microRNA expression can be regulated by E<sub>2</sub> in whole body homogenates [306] as well as a variety of other tissues [165,307,308]. In the brain, expression levels of mature let-7i, miR-7a, miR-9, miR-125a, and miR-181a are altered by E<sub>2</sub> dependent on age in the adult female rat [28], however, the normal physiological expression profile of these microRNAs throughout puberty, when circulating E<sub>2</sub> levels increase dramatically in females, is unknown. Therefore, we measured the expression levels of 5 E<sub>2</sub>-responsive microRNAs in the dorsal hippocampus using qRT-PCR at three time points throughout pubertal development (early = 30 d, peri = 44 d, late = 73 d) in untreated male and female rats (Fig. 16). In the dorsal hippocampus, a two-way ANOVA revealed a main effect of sex in 4 out of the 5 microRNAs tested including miR-7a, miR-9, miR-125a, and miR-




**Figure 16. Diagram of experimental paradigm.** Diagram depicting the age of sacrifice for each group of male and female Wistar rats. N=10/group.

**Table 6.** Statistical Analysis of microRNA Expression Levels in the Dorsal Hippocampus

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
let-7i	No	No	No
miR-7a	No	Yes: $F(1,46) = 10.138$ $p = 0.003$	Yes $p = 0.015$
miR-9	Yes: $F(2,48) = 20.038$ $p < 0.001$	Yes: $F(1,48) = 13.933$ $p < 0.001$	Yes $p = 0.003$
miR-125a	Yes: $F(2,48) = 4.984$ $p = 0.011$	Yes: $F(1,48) = 10.476$ $p = 0.002$	Yes $p = 0.004$
miR-181a	Yes: $F(2,45) = 10.076$ $p < 0.001$	Yes: $F(1,54) = 8.206$ $p p = 0.006$	Yes $p < 0.001$

181a, and 3 of these microRNAs, (miR-9, miR-125a and miR-181a) demonstrated a main effect of age (Table 6). A statistically significant interaction between sex and age was observed for miR-7a, miR-9, miR-125a, and miR-181a expression in the dorsal hippocampus (Table 6), demonstrating that the sex differences in microRNA expression is dependent on age. Interestingly, the only E<sub>2</sub>-responsive microRNA that was expressed in a manner singly dependent on sex, miR-7a, demonstrated an expression pattern throughout puberty that was different from the other 3 sex-dependent E<sub>2</sub>-responsive microRNAs. Specifically, the expression levels of miR-9, miR-125a and miR-181a significantly decreased in females between early and peripuberty, and remain decreased until late puberty (Fig. 17C, D, E). In contrast, male expression levels of miR-9, miR-181a and miR-125a remained constant. On the other hand, miR-7a expression did not change between early and peripuberty, but significantly increased and decreased between peripuberty and late puberty in females and males, respectively (Fig. 17B). These data suggest that miR-9, miR-125a, miR-181a may play important roles in dorsal hippocampus sexual dimorphisms during early puberty and that miR-7a may be important for mediating sex differences in brain function during late puberty. Overall, dorsal hippocampus microRNA expression levels that are dependent on sex and age decrease over the course of puberty whereas the microRNAs dependent on sex alone demonstrate divergent expression patterns between males and females.

**Sexually dimorphic expression patterns of E<sub>2</sub>-responsive microRNAs throughout pubertal development in the ventral hippocampus are distinct from the dorsal hippocampus.**



**Figure 17.** Sexually dimorphic microRNA expression is evident during pubertal development in the dorsal hippocampus. let-7i (A), miR-7a (B), miR-9 (C), miR-125a (D) and miR-181a (E) expression levels in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to PND 30 male animals. Dissimilar letters indicate a statistically significant difference between groups based on a 2-way ANOVA ( $p < 0.05$ ).


microRNA expression levels in the brain can be dependent on age, sex and hormone levels [309,310,311]. However, sex differences in microRNA expression during puberty have not been compared between the anatomically and functionally distinct dorsal and ventral hippocampus regions [57,58,205]. Therefore, we next quantified the expression levels of the E<sub>2</sub>-responsive microRNAs let-7i, miR-7a, miR-9, miR-125a, and miR-181 [28] in the ventral hippocampus to determine if they are differentially expressed from the dorsal hippocampus during pubertal development. In the ventral hippocampus there was a significant main effect of sex, independent of age, in 4 out of the 5 microRNAs, miR-7a, miR-9, miR-125a and miR-181a (Table 7). This is in stark contrast to the alterations in E<sub>2</sub>-responsive microRNAs in the dorsal hippocampus, in which these microRNAs were altered by sex dependent on age (Table 6). Interestingly, miR-7a, miR-125a and miR-181a expression levels remained constant throughout pubertal development in both males and females, yet female microRNA expression levels were significantly greater than males throughout this time period (Fig. 18B, D, E). The expression level of miR-9 however, was dynamic across pubertal development in males and not in females. Specifically, during early puberty, miR-9 expression levels were significantly greater in males compared to females, after which miR-9 significantly decreased below female levels at peripuberty and remained lowered until late puberty (Fig. 18C). Finally, no interaction was detected between age and sex for the E<sub>2</sub>-responsive microRNAs in the ventral hippocampus which demonstrated sexually dimorphic expression levels, suggesting that these observed sex differences are not dependent on E<sub>2</sub>.

## **Discussion**



**Table 7.** Statistical Analysis of Gene Expression Levels in the Ventral Hippocampus

<b>microRNA</b>	<b>MAIN EFFECT OF AGE</b>	<b>MAIN EFFECT OF SEX</b>	<b>INTERACTION: AGE X SEX</b>
let-7i	No	No	No
miR-7a	No	Yes: $F(2,53) = 5.319$ $p = 0.008$	No
miR-9	No	Yes: $F(1,53) = 25.571$ $p < 0.001$	No
miR-125a	No	Yes: $F(1,47) = 71.900$ $p < 0.001$	No
miR-181a	No	Yes: $F(1,45) = 20.266$ $p < 0.001$	No



**Figure 18.** Sexually dimorphic microRNA expression is evident during pubertal development in the ventral hippocampus. let-7i (A), miR-7a (B), miR-9 (C), miR-125a (D) and miR-181a (E) expression levels in untreated male (grey line) and untreated female (pink line) pubertal rats. N=10/group. Data represent mean fold change  $\pm$  SEM as compared to PND 30 male animals. Dissimilar letters indicate a statistically significant difference between groups based on a 2-way ANOVA ( $p < 0.05$ ).

Two novel findings emerged from these studies. First, 4 E<sub>2</sub>-responsive microRNAs, miR-7a, miR-9, miR-125a and miR-181a, are differentially expressed between males and females throughout pubertal development. Second, the degree to which microRNA expression levels vary by sex differs between brain regions. To our knowledge, these are the first studies to identify sexually dimorphic microRNA expression within the ventral and dorsal hippocampus. Moreover, the regional expression patterns of microRNAs in the hippocampus suggest that these microRNAs have distinct roles in the ventral and dorsal hippocampus. Our study extends these findings by determining that miR-7a, miR-9, miR-125a and miR-181a expression levels are dependent on sex and age during pubertal development. Overall, we determined that E<sub>2</sub>-sensitive microRNAs expression levels are distinct between three stages of pubertal development and that these expression profiles are sexually dimorphic and brain region-dependent.

Our data support the concept that microRNA expression levels are dynamic during periods of significant hormone fluctuations (such as puberty), although more experiments would be required to ascertain whether gonadal hormones directly regulate these microRNA expression profiles. Importantly, E<sub>2</sub> and T levels are well characterized, such that they increase during pubertal development [286,287,288] and recent studies also indicate that periods of shifting gonadal hormone levels are associated with shifts in microRNA profiles in the brain [28,29,312]. For example, perinatal exposure to T associated with brain masculinization dramatically changes microRNA expression patterns [29] and E<sub>2</sub> treatment in the aged female brain leads to different patterns of brain microRNA expression in the aged compared to young female rats [28]. A recent miR RT-

PCR array also revealed distinct microRNA expression profiles in the prefrontal cortex of pre-pubertal (PND 28) male and female mice, providing further evidence that sexually dimorphic gonadal hormones and/or sex can influence miR expression patterns in the brain [312]. Therefore, our results which revealed that miR-7a, miR-9, miR-125a and miR-181a are differentially expressed between males and females throughout pubertal development suggests that the expression of these microRNAs may be regulated by gonadal hormones.

The primary purpose of this study was to determine the effect of pubertal development (i.e. age) on the expression levels of E<sub>2</sub>-responsive microRNAs [28] in male and female rats, yet our data reveal that microRNA expression levels in the ventral hippocampus are dependent on sex alone throughout pubertal development. This is in contrast to the dorsal hippocampus, where none of the E<sub>2</sub>-responsive microRNAs were dependent on sex alone. Therefore, these data reveal that sex is a critical factor in determining the microRNA expression profile in the pubertal rat ventral hippocampus, suggesting that these microRNAs may regulate the development of the ventral hippocampus and the sexually dimorphic mood and memory behaviors that it mediates.

Our data also demonstrate regional differences in the expression levels of sexually dimorphic E<sub>2</sub>-responsive microRNAs during puberty. For instance, in the dorsal hippocampus, the expression levels of 3 microRNAs (miR-9, miR-125a and miR-181a) are altered by sex dependent on age, whereas only one microRNA (miR-7a) is dependent on age alone. These data indicate that the expression levels of miR-9, miR-125a and miR-181a are likely to regulate key sexually dimorphic developmental processes in the dorsal hippocampus occurring at distinct stages of puberty. This further suggests pubertal

microRNA expression levels can be sensitive to alterations in hormonal signaling and synaptic activation, two constantly evolving processes that shape the developing adolescent brain [313]. Interestingly, the expression level of let-7i is not dependent on sex or age in either the ventral and dorsal hippocampus, despite previous findings indicating its E<sub>2</sub>-responsiveness in the hippocampus of older rats [28]. This discrepancy underscores the concept that microRNAs can have very specific functions at different stages of development and/or different stages of hormonal environments in the brain and suggest that regional differences in ventral and dorsal hippocampus neurodevelopment during puberty may be mediated, in part, by differential E<sub>2</sub>-responsive microRNA expression

Currently, there is not much information regarding the functions of the microRNAs measured in this study, particularly in the brain. Indeed, many efforts in the microRNA field are focused on identifying the potential target genes of microRNAs in different biological systems in order to elucidate their functions. However, most of these studies rely largely on imperfect computer algorithms to predict gene targets, making it difficult to draw compelling conclusions from such experimentally-derived data. For example, current microRNA target prediction programs cannot determine cell-type specificity nor indicate whether multiple microRNAs—as opposed to a singly identifiable microRNA—may be required to act in concert to regulate the expression of a given target gene, and indeed, complete mechanisms regarding microRNAs' regulatory potential remain to be elucidated. Of the 5 microRNAs Rao et al. found to be E<sub>2</sub>-responsive, miR-9 is by far the most widely researched regarding its neuronal functions including neurogenesis and neuronal outgrowth. For instance, undifferentiated human [314] and

mouse-derived [315] stem cells rely on miR-9 expression during neural progenitor cell differentiation and miR-9 can also convert human fibroblasts into neurons [316,317]. miR-9 plays an instructive role in mediating neuronal cell fate by targeting transcription factors [318,319], suggesting that miR-9 may influence hippocampal neurogenesis during puberty. Moreover, miR-9 contributes to neuronal communication by regulating dendritic branching [320]. Together, these studies reveal that miR-9 regulates the generation and architecture of neuronal networks, linking miR-9 function to synaptic plasticity.

However, our limited understanding of microRNAs lends functional studies to appear far-reaching. Therefore, documenting the developmental profiles of microRNAs in all biological systems is a logical starting point towards the goal of identifying functional capacity. Towards this aim, the current study identified both temporal and sex-specific expression patterns of E<sub>2</sub>-responsive microRNA in the adolescent brain, and provided insight into their potential functions as being mediators of sexually dimorphic pubertal brain development. Taken together, the data presented here establish fundamental rationale for future studies aiming to classify the effector (i.e target gene) functions of these microRNAs.

## CHAPTER SIX

### GENERAL DISCUSSION

#### **Summary**

The previous chapters have described the following four main findings: 1) peripubertal binge EtOH exposure induces long-term alterations in mature microRNA expression levels in the male rat hippocampus, and has the potential to modulate the expression of their downstream target genes, 2) expression profiles of EtOH-sensitive microRNAs, miR-10a-5p, miR-26a, miR-32, miR-103 and miR-495, and their target genes, are dependent on sex and age in the pubertal rat hippocampus and therefore may contribute to sexually dimorphic hippocampus neurodevelopment, 3) expression profiles of E<sub>2</sub>-responsive microRNAs, miR-7a, miR-9, miR-125a and miR-181a, are differentially dependent on sex and age throughout pubertal development, suggesting that they play distinct developmental roles during puberty and 4) EtOH-sensitive and E<sub>2</sub>-responsive microRNAs have distinct expression profiles in the dorsal and ventral hippocampus throughout pubertal development, suggesting that their respective functions are region-specific. Taken together, these data chapters significantly contribute to the scientific literature that the expression levels of microRNAs can vary throughout pubertal development in brain region and sex-specific manner, and that distinct microRNAs are sensitive to peripubertal binge EtOH. Global analysis of our findings suggest the following two overarching possibilities that 1) peripubertal binge EtOH-mediated alteration of microRNA expression and/or processing underlies long-term alcohol abuse-

induced central nervous system impairments, and 2) that microRNAs may play a regulatory role in sexually dimorphism of adolescents. Chapters 3-5 have delivered data generating a framework from which one can further study the above two theories/hypotheses.

### **Key Findings**

- **Chapter III:**

- A) Characterization of EtOH-sensitive microRNAs:**

- dorsal hippocampus miR-10a-5p expression decreases between early and mid puberty, remaining decreased until late puberty
- dorsal hippocampus miR-26a expression levels decrease between mid and late puberty
- dorsal hippocampus miR-103 expression levels do not change throughout pubertal development
- dorsal hippocampus miR-495 expression levels increase between early and mid puberty, remaining increased until late puberty
- peripubertal binge EtOH increases miR-10a-5p expression levels between mid and late puberty in the dorsal hippocampus
- peripubertal binge EtOH decreases miR-26a expression levels immediately following the last treatment at mid puberty in the dorsal hippocampus
- peripubertal binge EtOH does not alter miR-103 expression levels during puberty in the dorsal hippocampus



- peripubertal binge EtOH decreases miR-495 expression levels immediately following binge, remaining decreased until late puberty in the dorsal hippocampus
- ventral hippocampus miR-10a-5p expression levels increase between mid and late puberty
- ventral hippocampus miR-26a expression levels decrease between early and mid puberty, and increase again by late puberty
- ventral hippocampus miR-103 expression levels increase between mid and late puberty
- ventral hippocampus miR-495 expression levels decrease between early and mid puberty and increase beyond early puberty levels by late puberty
- peripubertal binge EtOH increases miR-10a-5p expression levels immediately following the last treatment in the ventral hippocampus
- peripubertal binge EtOH does not alter miR-26a expression levels in the ventral hippocampus
- peripubertal binge EtOH increases miR-103 and miR-495 expression levels immediately following the last treatment, and the levels are decreased by late puberty in the ventral hippocampus

**B) Characterization of microRNA biogenesis enzymes:**

- dorsal hippocampus Drosha and Dicer expression levels decrease between early and mid puberty, remaining decreased until late puberty
- ventral hippocampus Drosha and Dicer expression levels decrease between early and mid puberty, and increase by late puberty
- peripubertal binge EtOH increases Drosha expression levels in an immediate and long-term manner in the dorsal hippocampus
- peripubertal binge EtOH increases dorsal hippocampus Dicer and ventral hippocampus Drosha and Dicer expression levels immediately following treatment and these levels are decreased beyond untreated levels by late puberty

**C) Characterization of microRNA target genes BDNF and SIRT1:**

- dorsal hippocampus BDNF expression levels increase between early and mid puberty, then decrease by late puberty
- dorsal hippocampus SIRT1 expression levels decrease between early and mid puberty, and increase by late puberty
- ventral hippocampus BDNF and SIRT1 expression levels increase between early and mid puberty then decrease by late puberty
- peripubertal binge EtOH increases BDNF and SIRT1 expression levels in immediate and long-term manners in the dorsal hippocampus

- peripubertal binge EtOH decreases BDNF expression levels between early and mid puberty, and levels increase beyond untreated levels by late puberty in the dorsal hippocampus
- peripubertal binge EtOH decreases SIRT1 expression levels in an immediate and long-term manner in the ventral hippocampus

▪ **Chapter IV:**

**A) Characterization of sex differences in EtOH-sensitive microRNAs:**

- dorsal hippocampus miR-10a-5p expression levels are increased at early puberty in the female relative to male
- dorsal hippocampus miR-26a and miR-495 expression levels are decreased throughout pubertal development in the female relative to male
- dorsal hippocampus miR-32 and miR-103 expression levels are not sexually dimorphic
- ventral hippocampus miR-10a-5p expression levels are increased at early puberty, equivalent at mid puberty and decreased at late puberty in the female relative to male
- ventral hippocampus miR-26a and miR-495 expression levels are decreased throughout pubertal development in the female relative to male
- ventral hippocampus miR-103 expression levels are increased at mid puberty and decreased at late puberty in the female relative to male

- ventral hippocampus miR-32 expression levels are not sexually dimorphic

**B) Characterization of sex differences in miR biogenesis enzymes and EtOH-sensitive miR target gene expression:**

- Dorsal hippocampus Drosha and Dicer and ventral hippocampus Dicer expression levels are decreased in the females relative to males throughout pubertal development
- Ventral hippocampus Drosha expression levels are decreased at early and late puberty, but equivalent at mid puberty in females relative to males
- Dorsal hippocampus BDNF and SIRT1 expression levels are decreased throughout pubertal development in females relative to males
- Ventral hippocampus BDNF and SIRT1 expression levels are equivalent at early and late puberty but decreased at mid puberty in females relative to males

▪ **Chapter V:**

**A) Characterization of sex differences in E<sub>2</sub>-sensitive microRNAs:**

- Dorsal and ventral hippocampus let-7i expression levels are not sexually dimorphic, nor do they change, throughout pubertal development

- Dorsal hippocampus miR-7a expression levels increase and decrease in females and males, respectively, between mid and late puberty
- Dorsal hippocampus miR-9, miR-125a and miR-181a expression levels are increased at early puberty in females relative to males; male expression levels do not change throughout puberty
- Ventral hippocampus miR-7a, miR-125a and miR-181a expression levels are not dynamic throughout puberty in males or females, however, females expression higher levels relative to males throughout pubertal development
- Ventral hippocampus miR-9 expression levels in females is increased at early puberty and decreased at mid and late puberty relative to males

## **Final Remarks**

### *Selection of Model Organism*

Rats demonstrate markers unique to an adolescent growth spurt such as hyperphagia (developmental overeating) and accelerated growth [321], making them a valuable model organism of choice for this study. Memory processes are more adversely affected in adolescent rodents abusing alcohol than in the adults abusing the same dose [5,79,322], highlighting the important need to study how alcohol alters the adolescent brain, and the hippocampus in particular. Importantly, our lab has demonstrated that no differences in body weight results from administering our binge EtOH paradigm in adolescent Wistar rats and that the blood alcohol concentration does not significantly

differ when administering EtOH via gavage or intraperitoneal injection [8,12,44].

Moreover, our binge EtOH exposure paradigm has been shown to be a reliable model for administering amounts of EtOH typically consumed amongst adolescents [200].

#### *microRNA Functions in the Brain: Relevance to Neuronal Development*

The discrete biological functions of individual microRNAs during neuronal development are not well understood. Lsy-6 was the first specific microRNA recognized to have a role in nervous system development *in vivo*, where it was shown to regulate left/right asymmetrical patterning of the taste receptor neurons in *C. elegans* [323]. Two other microRNAs, miR-9 and miR-10, are highly expressed in the brain and have been shown to play important roles in the brain development of many species including humans, rodents, zebrafish, and drosophila, demonstrating a high degree of evolutionary conservation among these microRNAs. Specifically, miR-9 promotes migration and proliferation in human neural progenitor cells by targeting stathmin, a gene required for microtubule assembly [314] and peripheral nervous system sensory organ development in drosophila [324]. Further, miR-9 is significantly reduced in the presenilin-1 null mouse model, which exhibits severe CNS developmental defects, during specific stages of development compared with wild-type mice [325]. Although our microRNA RT-PCR array experiment did not identify miR-9 as EtOH sensitive following peripubertal binge EtOH, it is possible that by examining the whole hippocampus, we were unable to delineate ventral vs. dorsal hippocampus EtOH-mediated alteration in miR-9 expression. This is further supported by our experiments identifying ventral vs. dorsal differences in miR-9 expression levels in untreated peripubertal rats. Another important microRNA during development is miR-10, which targets members of the HOX gene family, a highly

conserved group of transcription factors that coordinate anterior-posterior body axis alignment in zebrafish and other species during development [326]. These studies have revealed that specific microRNAs are critical for proper gene expression and brain function throughout development, however, the role of microRNAs beyond early developmental periods is just beginning to be investigated. Mature microRNA expression is age-dependent, and accordingly, microRNAs regulate both early developmental gene expression changes as well as those that occur throughout the lifespan in various species [43, 46, 47, 111]. In our studies, miR-10a-5p was sensitive to EtOH in both a short and long-term manner in the ventral and dorsal hippocampus, respectively. Moreover, as microRNAs can regulate the expression of numerous target genes, it is extremely likely that targets change as a function of age. Indeed, microRNAs have been demonstrated as important regulators neuronal stem cell fate determination such as the SOX family of transcription factors which control NSC differentiation and gliogenesis [327,328].

The origin of oligodendroglial fate determination has been historically controversial in the literature, and interesting recent studies implicate microRNAs in this process. For example, in a mouse model of Dicer1 KO specific to oligodendrocytes, miR-219 and miR-338 expression is absent, whereas they are abundantly expressed in wildtype oligodendrocytes [329,330]. Moreover, manipulation of these microRNAs in oligodendrocyte precursor cells demonstrates that they are essential for the differentiation of oligodendrocyte precursors into myelinating oligodendrocytes. Myelination of neurons by oligodendrocytes is essential for the process of salutatory nerve conduction and many studies have implicated reduced myelination as a mediator of neurological insults during

adolescence. Therefore, the control of oligodendrocyte maturation via microRNAs highlights yet another important role of microRNAs in brain development.

How microRNA expression changes over time is another active area of research in the microRNA field. A deep sequencing study recently highlighted 75 microRNAs that were differentially expressed in the brain with age [206]. Interestingly, let-7 microRNAs are inhibited by the RNA binding protein Lin28 in the reproductive areas of the neonate rat brain, yet, upon progression towards puberty, this expression pattern is reversed with increasing and decreasing expression of let-7 and Lin28, respectively, in the rat hypothalamus [248]. These data suggest a role for microRNA in the regulation of pubertal onset and highlight the potential for microRNAs to regulate pubertal brain development. Moreover, our data demonstrating that in the healthy adolescent brain, microRNA expression is dynamic at 3 distinct time points during pubertal development suggests that microRNAs are important for brain development at this time. Much still remains to be discovered regarding the role of microRNAs in regulating brain development, however, compelling data suggests that negative effects of alcohol on the developing brain may be mediated by alterations in brain microRNA expression and/or the mechanisms of their transcriptional regulation.

#### *The Role of microRNAs in Ethanol Neurotoxicity During Development*

Neurological consequences of alcohol abuse (repetitive and heavy drinking) occur throughout the lifespan, and an emerging role for microRNAs in mediated alcohol's effects on addiction, toxicity and teratology is becoming evident. The long-term consequences of alcohol exposure during important neurodevelopmental time periods has



recently become widely discussed, particularly with regards to prenatal EtOH exposure leading to fetal alcohol spectrum disorder (FASD). FASD describes a wide range of effects that a mother drinking alcohol during pregnancy can inflict upon the developing fetus and symptoms of FASD range from physical abnormalities, identified as fetal alcohol syndrome (FAS), to neurobehavioral alterations including depression, hyperactivity, learning disabilities and psychosis [331,332,333]. microRNAs have been implicated in the development of FASD. For example, a murine miR-9 (KO) model results in FASD phenotypes and growth retardation presumably by inhibiting the miR-9-mediated inhibition of *Foxg1*, a gene that promotes proliferation and prevents differentiation and whose mutation also leads to fetal growth retardation, microcephaly and mental retardation [318,334]. This miR-9 KO leads to increased *Foxg1* expression and inhibits maturation of neuronal populations. Interestingly, a subset of EtOH-sensitive microRNAs that play a role in cell and tissue maturation, miR-10a/b, miR-21 and miR-335, demonstrate developmental stage-specificity in their sensitivity to EtOH in neuronal stem cells (NSCs) [193]. This finding provides support to our observation that expression of a subset of microRNAs is dependent on age, sex and EtOH. As mentioned previously, miR-9 was not altered by peripubertal binge EtOH in males however it is possible that in females, miR-9 may be sensitive to EtOH. For instance, we observe sex differences in steady-state miR-9 expression in our untreated peripubertal rats and it is possible that miR-9 operates the peripubertal neuronal differentiation (i.e. peripubertal neuronal growth) in female rats. If so, the sex-differences observed in binge EtOH-exposed rats may be explained by a sensitivity of miR-9 to EtOH in females and not males. Future studies can investigate this possibility further with binge-EtOH-exposed female rodents.

Interestingly, Sathyan et al. demonstrated that miR-10a demonstrated sensitivity to EtOH in NSC's as we did in peripubertal male rats. However, different target genes are implicated in these two EtOH studies. This difference may be due to EtOH-sensitive microRNAs identified in these two studies being responsible for regulating unique sets of developmental genes, supported by the fact that 5p- and 3p-microRNAs are expected to be complementary to different target genes.

The neurochemical mechanisms responsible for such EtOH-induced anatomical and behavioral alterations remain unclear. It is compelling to suggest that the mechanism of binge EtOH effects on the pubertal brain—such as an increased risk of adult alcohol abuse [335], spatial learning impairments [79], reduced hippocampus volume [3]—are carried out by microRNA-mediated regulation of gene expression. Our studies demonstrate that peripubertal binge EtOH leads to long-term alteration in microRNA and BDNF expression, suggesting a potential long-term effect of peripubertal binge drinking on microRNA-mediated regulation of synaptic plasticity (of which BDNF is a primary mediator). Moreover, differential effect of EtOH on microRNAs and target gene BDNF expression between the ventral and dorsal hippocampus in the peripubertal male rat suggest that EtOH may disrupt both ventral and dorsal hippocampus functions. In support of this possibility, others have demonstrated that EtOH alters mood and memory in response to peripubertal binge EtOH and have also described conflicting data regarding expression levels of BDNF mRNA and protein expression following EtOH. For instance, one study investigated early postnatal pups exposed to EtOH vapor inhalation for 2 hours and 40 mins each day on PND 4-10 and reported a significant increased BDNF protein expression [90]. Other studies also describe that BDNF increases in its expression in the

hippocampus following EtOH exposure and suggest that these effects lead to changes in synaptic plasticity leading to behavioral responses to EtOH [147]. Indeed, ethanol has been shown to increase neuronal synaptic adaptation [336] but also decrease hippocampal LTP [4,91,337] and CA3 BDNF mRNA expression [338]. Based on our data, such inconsistencies in the literature may be due to different responses of the ventral and dorsal hippocampus to EtOH, however, factors including experimental design, model organisms, EtOH paradigm, age and sex of test subjects cannot be ruled out. Controversy also exists regarding whether an ethanol-induced increase or decrease in hippocampal BDNF expression during developmental periods is neuroprotective or neurotoxic. Our data also indicate that this may also depend on whether the ventral or dorsal region is under consideration, as our data indicate that EtOH increases and decreases BDNF expression in a long-term manner differently in the dorsal and ventral hippocampus, respectively. Although we do not know from our studies whether these results directly result in reduced hippocampal functional capacity, it is compelling to suggest that our observed increases in the dorsal hippocampus BDNF and decreases in the ventral may correlate with the known impairment in mood and memory following EtOH abuse. It is important to continue to tease out the mechanisms behind EtOH-mediated effects and the potential role of BDNF expression in the hippocampus.

It is also possible that microRNA processing is broadly influenced by EtOH. Our data demonstrate that that Drosha and Dicer expression are altered in both short and long-term fashions following binge EtOH. It has also been previously demonstrated in zebrafish embryos that pre-miR-9-3 transcripts accumulate in response to EtOH [194]. In our study, it is possible that the pri- and pre-microRNA transcripts are differentially

altered by binge EtOH and leads to altered mature microRNA expression. However, because binge EtOH doesn't alter global microRNA expression, it remains unclear whether microRNA processing is a mechanism by which EtOH alters expression of mature microRNAs.

Chapter 3 data introduces the novel finding that pubertal miR expression profiles for miR 10a-5p, miR-26a, miR-103 and miR-495 are altered by peripubertal binge EtOH exposure differentially in the dorsal and ventral hippocampus. The hippocampus is involved in learning and in consolidation of explicit memories from short-term to long-term memory storage in the cortex. Importantly, stress impairs hippocampal-dependent memory in both human and rats [339,340,341]. Its sensitivity to stress is thought to be due to its high expression of glucocorticoid receptors through which it participates in the termination of stress responses via the glucocorticoid-mediated negative feedback of the hypothalamo-pituitary-adrenal (HPA) axis [342,343]. Interestingly, studies have demonstrated a dissociation of stress-induced increases in corticosterone levels and the ability to induce hippocampal long-term potentiation (LTP) (the cellular process underlying memory), suggesting stress-induced effects on the hippocampus may extend beyond cellular LTP. Indeed, other studies have implicated that peripubertal binge EtOH imparts dysregulation of the neuroendocrine stress axis in a long-term manner [12]. The adolescent stress response is still maturing, evidenced by enhanced stress reactivity [344](Dahl RE and Gunnar, 2009, [345,346,347]), suggesting that an altered development of the stress response following adolescent alcohol abuse may increase the risk of developing mood disorders commonly associated with alcohol abuse. Our studies suggest that the physiological stress of adolescent binge EtOH may incur long-lasting

stress-induced molecular alterations via changes in the expression levels of microRNAs that target genes important for synaptic plasticity. However, it has long been noted that changes in long-term potentiation LTP and/or dendritic spine density in the hippocampus are believed to demonstrate stress-induced hippocampal effects at the anatomical and cellular level [348,349,350,351]. Taken together, these studies demonstrate that high levels of alcohol consumption during adolescence lead to permanent adolescent-like brain function, such that the brain does not reach full maturity or that its development is altered.

*EtOH-Induced Changes in Hippocampal microRNA Correlate with Altered BDNF and SIRT1 mRNA: Proposed Role of Ethanol Metabolism*

The hippocampus is particularly sensitive to the negative effects of EtOH [62,352,353] and oxidative stress has been implicated in mediating EtOH's neurotoxicity [354], however, the complete neurobiological mechanism remains unclear. In this study we observe binge EtOH-mediated changes in hippocampal microRNA expression, and therefore, it is possible that this may be due to increased oxidative stress of hippocampal mitochondria. For example, ethanol metabolism begins with its conversion into acetaldehyde, which in the brain of binge drinkers, is accomplished largely by microsomal enzyme cytochrome (CYP2E1) [355]. This is because alcohol dehydrogenase (ALDH) becomes saturated under conditions of heavy alcohol consumption. CYP2E1, in turn, generates superoxide and hydrogen peroxide reactive oxygen species (ROS) leading to EtOH-induced oxidative DNA damage [356,357]. Mitochondrial oxidative stress disrupts the permeability of the inner and outer mitochondrial membrane, which leads to an increase in cytochrome c release and overproduction of ROS, which together, increase

DNA damage and programmed cell death or apoptosis [358]. It is unlikely that our paradigm of binge drinking increases programmed cell death because we do not see global decreases in microRNA expression and moreover, one would also expect that the control gene, U6, would also be decreased under conditions of global apoptosis. Based on evidence that EtOH induces mitochondrial oxidative stress response in the brain [359], it is possible that EtOH-mediated oxidative stress may be a mechanism through which microRNAs regulate expression of SIRT1 and BDNF in our study. For example, it has been demonstrated that microRNAs can be primarily expressed in the mitochondrial genome [360], that mature microRNAs can even localize to the mitochondria [361] and that these mitochondrial microRNAs may play a role in apoptosis. Under such circumstances, EtOH-mediated oxidative stress in neuronal mitochondria could release mature microRNAs into the cytoplasm where they can regulate the expression of target genes. Our observation that microRNAs increase following binge EtOH in the ventral hippocampus may do so via mitochondrial pore formation following an oxidative stress response, allowing microRNAs to leak into the cytosol where they can associate with RISC and regulation expression of BDNF and SIRT1 mRNA. In support of this theory, mitochondrial disruption has previously been reported to mediate translational repression, one of the mechanisms through which microRNAs postranscriptionally regulate gene expression [362]. Overall, our data identifying EtOH-sensitive microRNAs in the hippocampus and correlative alterations in BDNF and SIRT1 expression may be specific to repeated heavy (binge) EtOH consumption leading to an oxidative metabolic response. To further investigate this theory, one could fractionate cell lysate and run expression analysis on mitochondrial RNA. In summary, EtOH metabolism disrupting mitochondrial

membrane permeability demonstrates a targeted approach in which heavy doses of EtOH can disrupt gene regulation of a variety of cellular processes, as it may unleash the mitochondrial genome-containing epigenetic regulatory agents such as microRNAs.

However, it remains necessary to identify whether our observed EtOH-mediated microRNA changes directly alter the expression of BDNF and SIRT1. One way to test this is to assay for direct binding of these microRNAs to the target gene 3'UTR following EtOH exposure, as successfully demonstrated previously by Caputo et al. using luciferase reporter assays for the binding of miR-26a to the BDNF 3' UTR [143]. If direct microRNA-3'UTR binding is indicated, then to take this research one step further and test the functional effect of altered BDNF and SIRT1 mRNA in response to peripubertal binge EtOH. For example, performing behavioral assays which measure learning, anxiety and memory behavior may indicate whether CNS impairments occur as an extension of with EtOH-induced alterations in microRNA and BDNF and SIRT1 expression in our peripubertal binge EtOH paradigm.

Finally, given the developmental nature of the peripubertal period and the promiscuous nature of microRNAs, it is plausible that EtOH-sensitive microRNAs regulate the expression of a host of additional developmental genes. On that order, a microarray can efficiently identify potential additional target genes. However, an alternate hypothesis to EtOH-induced alterations in microRNA expression regulating the correlative changes in BDNF and SIRT1 mRNA is that these microRNAs are performing unconventional regulatory functions. For instance, it is becoming clear in the literature that microRNA functions are not restricted to mRNA degradation and translational repression but also orchestrate alternate gene regulatory mechanisms. For example,

epigenetic gene regulation by microRNAs include binding to the 5'UTR of mRNA targets to enhance translation [363], tethering Argonaute proteins to promoter regions to enhance or inactivate gene transcription [364,365], secretion into plasma to alter target cell behavior [366,367] which suggest microRNAs as potential endocrine factors. These additional functions of microRNAs demonstrate that microRNAs are not limited to regulating gene expression in the typical method of 3'UTR binding and therefore future experiments testing for BDNF and SIRT1 3'UTR binding will not exhaust all of the possible ways in which EtOH-sensitive microRNAs may regulate these gene targets.

*Possible Role for Gonadal Hormones in Mediating Sexually Dimorphic Pubertal miR Expression*

Puberty is a period of life associated with an increased risk of developing psychiatric disorders, most likely due to the incomplete maturation state of the brain in addition to increased risky behaviors common amongst adolescents. We have identified that the expression levels of 10 microRNAs (let-7i, miR-7a, miR-9, miR-10a-5p, miR-26a, miR-32, miR-103, miR-125a, miR-181a and miR-495) are specific to sex, stage of puberty and brain region, indicating that they play specific roles in the brain development of males and females during puberty. Moreover, normal fluctuations of brain microRNAs during puberty and their apparent sensitivity to EtOH, targets them as mediums through which noxious environments or genome may disrupt gene expression. To that end, it is possible that the regulation of miR expression levels may not only play a large role in normative brain maturation, but may also underlie the molecular mechanisms leading to psychiatric disorders. In agreement with this theory, we have recently shown that peripubertal exposure to repeated episodes of binge EtOH alters the normal



developmental profiles of microRNAs in the hippocampus [247], a brain region susceptible to alcohol-abuse and suggested to be damaged in a host of neuropathologies [11,338,358,368,369,370]. Furthermore, many studies in rodents and humans demonstrate that substance abuse during puberty leads to altered adult gene expression, memory dysfunction and continued substance abuse and dependence [4,5,10,11,12,79,247,371]. Therefore, microRNAs altered by EtOH may have a variety of negative (or positive) neurological consequences including impaired memory and stress response regulation, however the mechanisms remain unclear.

Sexually dimorphic expression levels of microRNAs may be due to differences in circulating gonadal hormones between males and females. It has been recently reported that gonadal hormone-mediated generation of sexually dimorphic physiology is, in part, due to their ability to influence miR expression. microRNAs are important regulators of the genome, but what controls their regulation is not well understood. Studies using both *in vivo* and *in vitro* systems have demonstrated that miR expression profiles (miRNomes) can be altered by T, E<sub>2</sub> and their respective nuclear receptors (androgen receptor and estrogen receptor ) in healthy and cancerous tissues, as well as during different stages of development [28,150,151,161,165,264,372,373]. These studies suggest that microRNAs may contribute to the development of typical adolescent sexual dimorphisms.

Gonadal steroids can also influence the miR target genes we tested in our studies. For instance, female levels of circulating E<sub>2</sub> is much higher in females than in males and E<sub>2</sub> replacement following GDX restores levels of BDNF mRNA and protein in the postnatal developing rat hippocampus, indicating that E<sub>2</sub> regulates the expression of BDNF factor mRNA and protein in the rat hippocampus [374]. How E<sub>2</sub> regulates the

expression of BDNF is not known, but it is possible that E<sub>2</sub> regulates peripubertal BDNF expression in a sexually dimorphic manner. For instance, in the second postnatal week male rodents express higher levels of aromatase, the enzyme that converts T to E<sub>2</sub>, as well as ER $\alpha/\beta$  in the hippocampus, suggesting a greater effect of E<sub>2</sub> signaling in males than in females at this developmental period [375]. Indeed, BDNF mRNA in the hippocampus is increased by E<sub>2</sub> [376], and peripubertal males may have greater local E<sub>2</sub> synthesis in hippocampal neurons and this may increase the levels of BDNF during the mid-phase of pubertal development. Furthermore, BDNF mRNA is known to fluctuate across the estrous cycle, in correspondence with the changing levels of E<sub>2</sub> [377] and it is possible that the lack of BDNF mRNA changes observed in our females compared to males may have occurred due to the females being in diestrus stage of their menstrual cycle when E<sub>2</sub> levels are lower. The consequences of differences in circulating E<sub>2</sub> levels, is that both E<sub>2</sub> and BDNF positively affect synaptic plasticity, and therefore, the dynamic expression of microRNAs throughout pubertal development in males may correspond with alterations in the BDNF and/or SIRT1 target genes and ultimately play a role in mediating the formation of long-term alterations in brain development during puberty.

The role of gonadal hormones and their influence on growth-promoting signals may underlie the necessity of peripubertal exposure to E<sub>2</sub> and T for development of adultlike mating, social and cognitive behaviors. For instance, ovariectomized (OVX) reduces mRNA and protein expression of (BDNF) and this is reversed with E<sub>2</sub> replacement [378,379]. The hippocampus is also a rare site wherein *de novo* synthesis of E<sub>2</sub> occurs outside of the gonads, and its expression is implicated in regulating local synaptic plasticity in dendritic spines [380]. Indeed, our data demonstrate clear sex

differences in the miR expression throughout pubertal development in multiple brain regions. In stark contrast to E<sub>2</sub> effects on BDNF and synaptic plasticity, gonadectomy (GDX) in adult male rats increases BDNF protein expression and synaptic plasticity in hippocampus mossy fibers [381] and the same pattern was observed in PND 90 male rats that gonadectomized at early puberty (PND 30) [86], suggesting that T represses hippocampal BDNF protein during puberty. Indeed, the author concludes the effects of gonadectomy are independent of pubertal status, however, because the immediate effects of PND 30 GDX on BDNF expression was not measured, it is impossible to know whether levels of BDNF were increased during peripuberty. Given that GDX incurs a reduction in BDNF protein measured at 4, 7 and 10 days post-GDX [374], it is clear that GDX at different developmental ages can have differential effects on BDNF protein expression. Furthermore, our data reveal that BDNF mRNA in male dorsal and ventral hippocampus is significantly increased at peripuberty compared to early puberty, suggesting that the rising levels of T correlate with rising BDNF mRNA. Our observed correlation in levels of T and BDNF are in contrast to experiments demonstrating that GDX increases BDNF protein levels. However, Solum and Handa have previously observed that neonatal GDX decreases BDNF mRNA yet increases BDNF protein levels in the same animals [374], demonstrating that BDNF mRNA levels do not always correspond with protein levels. Variable measurements of BDNF expression may also reflect differences in expression within the whole hippocampus vs. the isolated hippocampal subregions, the ventral and dorsal hippocampus. Indeed, differences in brain region expression levels of BDNF are to be expected, evidenced by T levels in the pelvic ganglia being correlated with BDNF protein expression following adult male GDX, such

that GDX decreased the number of BDNF-immunoreactive neurons [382]. In addition, T has been shown to contribute to the anxiogenic effect of exposure to a novel environment observed in male rodents as well as contribute to their social behavior [383,384], whereas in male Syrian hamsters, pubertal exposure to T contributes to aggressive, sexual and communication behaviors [385]. Lastly, T exposure during puberty in humans is necessary for the sex differences observed in spatial memory, performance of which is primarily controlled by the dorsal hippocampus [386,387]. Overall, the hippocampus plays a clear role in contributing to the gonadal hormone-mediated alterations in synaptic plasticity throughout pubertal development and as microRNAs can regulate gonadal hormone receptors, it is possible that microRNAs help to orchestrate the role that gonadal hormones play in the development of the adolescent brain.

Overall, female expression levels of EtOH-sensitive microRNAs and their target genes, BDNF and SIRT1, did not correlate with one another in this study. Moreover, the expression levels of SIRT1 and BDNF were extremely low and nearly unchanging in the female hippocampus compared males throughout pubertal development. These data most likely do not reflect a lack of dynamic synaptic plasticity. Indeed, the female expression levels of EtOH-sensitive microRNAs, BDNF and SIRT1 demonstrating no correlation whereas they did in the males (Chapter 4, Fig. 12, 153 and 14), it is important to point out that these EtOH-sensitive microRNAs were identified in male specimens in (see Chapter 3, Fig. 3). Therefore, it is possible that future studies using the same approach which was used in Chapter 3 males (aka the microRNA array) in females, would indeed discover a set of different EtOH-sensitive microRNAs that is specific to females. Moreover, due to

the promiscuous nature by which microRNAs bind their mRNA target genes, it is very likely that potential female-specific EtOH-sensitive would also target BDNF and SIRT1.

Gonadal hormones also influence behavioral systems required for learning and memory. For instance, it has been demonstrated in multiple species that T acts via NMDAR circuits to alter synaptic plasticity and learning [388,389,390]. Specifically in a rodent model, males that are not exposed to androgen receptor (AR) activation during puberty exhibit altered hippocampal CA1 synaptic plasticity, as measured by social recognition memory [388]. Together these data suggest that organizational events mediated by T activation of ARs during puberty are required for the development of adultlike social behaviors and effective hippocampal-dependent learning and memory processes.

Adolescent EtOH abuse-induced disruption of neurodevelopment occurs in both males and females [391,392,393], although often times to different magnitudes [46], as well as in one sex and not the other. These studies implicate a role for the involvement of circulating gonadal steroids, yet the causality of these sexually dimorphic effects remain unclear. Evidence from both rodent and human studies supports the theory that gonadal hormones play a role in mediating the sex differences observed in EtOH consumption, a typical adolescent behavior. Indeed, T moderates EtOH intake in male rodents, such that pre- or post-pubertal castration increases adult EtOH intake and dihydrotestosterone (DHT) or T replacement following castration sufficiently increases EtOH intake back to intact, GDX, and sham-GDX male and female controls [236,394]. Furthermore, this increased EtOH consumption in adult male rats that were GDX either pre- or post-puberty, is not due a weakened aversion to the taste of EtOH over time as conditioned

taste aversion (CTA) experiments demonstrated an increased CTA in males GDX in pre-puberty compared to post-puberty [395], whereas GDX at either time point resulted in a similar increase in EtOH intake [236,260]. Interestingly, in adult male rodents bred selectively for high EtOH preference compared to those bred to not prefer EtOH, high T levels are strongly correlated with a preference for EtOH throughout the day when not intoxicated [396]. In human male twin studies, following individual analysis and adjustment for pubertal stage, a higher T levels correlated with a greater number of alcohol symptoms as well as diagnosis for alcohol dependence [397]. Indeed, T is positively correlated with EtOH consumption not only in males but also in females. In self-report studies, higher levels of T and E<sub>2</sub> in male and female adolescents, was positively associated with alcohol use [398]. Furthermore, female rodent consumption also varies across the estrous cycle, such that during proestrus (slowly rising E<sub>2</sub>) and estrus (high E<sub>2</sub>) there is a decrease in total alcohol consumption [399,400]. Potential neural mechanisms underlying sex differences in alcohol drinking behavior involves dopamine release, which is greater in the female rat nucleus accumbens [401]. These data underlie the importance of distinguishing sex differences regarding substance abuse. Although steroid hormones are likely not an exclusive factor, an important next step is to determine whether E<sub>2</sub> and/or T are responsible for mediating the sex-specific differences in EtOH-sensitive miR expression.

It is clear that gonadal hormones contribute to the exceedingly plastic nature of the adolescent brain as they contribute to neurite outgrowth, increases in neurogenesis and alterations in steroid receptor expression [240]. Moreover, the effects of sex hormones on synaptic plasticity are sexually dimorphic [402], yet the mechanism remains

unclear. It is possible that microRNA-mediated regulation of gene expression has evolved to adapt to changing hormonal environments during puberty (and menstrual cycle), however, contemporary lifestyles of adolescents including experimentation with binge drinking and the use of oral contraceptives may have detrimental effects on microRNA expression that is critical for adolescent brain development. Chapter 5 data demonstrate that there are clear sex differences in the expression of 5 E<sub>2</sub>-responsive microRNAs, let-7i, miR-7a, miR-9, miR-125a and miR-181a, throughout pubertal development. We also demonstrated that their expression is distinct between the functionally dimorphic hippocampal regions, the ventral and dorsal hippocampus. In conclusion, we have identified that males and females express E<sub>2</sub>-responsive microRNAs in an age and sex dependent manner throughout pubertal development and that the expression patterns are unique between the dorsal and ventral hippocampus. However, it remains unclear whether the pubertal expression levels of E<sub>2</sub>-responsive microRNAs are correlated with gonadal hormone levels in a sexually dimorphic manner. Therefore, it will be important to determine how pre-pubertal removal of endogenous hormones and subsequent exogenous treatment with E<sub>2</sub> alters the expression of these microRNAs in the male and female dorsal and ventral hippocampus in future studies. It will be important to monitor the stage of the estrus cycle in female rats and the analysis of expression levels of microRNAs should be separate for females in different estrous cycle stages. It is important to note that symptoms of depression, schizophrenia and anxiety fluctuate across the menstrual cycle [403,404,405,406,407,408,409], and puberty is period of life wherein there is an increased risk for developing anxiety and other related mood disorders. Moreover, the course and prevalence of mood disorders are sexually

dimorphic, and it is possible that microRNAs expressed differentially between males and females in the mood-regulating ventral hippocampus target genes important for regulating mood. In addition, gonadectomy (GDX) followed by T treatment would be important to determine T effects on microRNA expression levels as gonadal androgen exposure during puberty is required for male-typical behaviors associated with the stress response such as anxiety-related decrease in social behavior in an unfamiliar environment [384]. Lastly, target genes of these microRNAs, such as GABA, SIRT1, BDNF and GR, could be measured in male and female hippocampi to identify potential mood-regulating target genes that these microRNAs may control the expression of during pubertal development. Indeed, puberty is a period of continuous sexual differentiation in the brain [410] and pubertal development is controlled by functionally connected networks coordinated by epigenetic mechanisms [411], indicating that miR expression levels can influence the establishment of neural networks during puberty which can have long-lasting consequences on adult brain function.

The dorsal hippocampus reduction of microRNA expression in females contrasts rising levels of circulating  $E_2$  in females at this time. Furthermore, the dorsal hippocampus microRNA expression in males remains stable and indeed, males exhibit an increase in T throughout puberty but not  $E_2$ . These data would suggest that the decrease in these female microRNAs during pubertal development is  $E_2$ -dependent, however, one would also expect the early puberty levels of male microRNAs to be similar to the female early puberty levels. Overall, the lack of change in microRNA expression levels in males mimics their lack of exposure to changes in circulating  $E_2$  throughout puberty. Of note, the early pubertal expression level of these 3 microRNAs is significantly increased in



females, but this may be due to the fact that it is not typical for males and females to exhibit similar levels of circulating E<sub>2</sub> at this time age [286,287,288]. Indeed, activation of the HPG axis in females can occur before males, and could explain the differences observed in E<sub>2</sub>-responsive miR expression at early puberty between males and females. However, reduced miR expression over time as E<sub>2</sub> levels rise in females does not agree with the sex differences observed at early puberty such that female miR expression is high as their E<sub>2</sub> levels begin to rise. It is possible that ligand-independent effects can be regulating miR expression levels before the rise in gonadal hormones at pubertal onset, such that the estrogen receptor (ER) acts on the miR promoters to increase their transcription. Indeed, ER $\beta$  has been shown to have transcriptional activity in rodent neuronal cell lines [412,413] and protein-protein interactions in the female rodent ventral hippocampus [414] that are altered in the presence of E<sub>2</sub>, suggesting that ER $\beta$  has ligand-independent functions in the female brain. Overall, it is possible that E<sub>2</sub> regulation of microRNAs can be dynamic throughout pubertal development, however, more experiments are required to adequately address this theory.

#### *Take Home Message*

In summary, adolescent alcohol abuse is strongly associated with multiple other health risk behaviors including alcohol-related motor-vehicle accidents, alcohol-related sexual assault, suicide and weakened learning and memory capacities [415,416,417,418] supporting the role that adolescence is a sensitive period with regards to brain organization. The mechanisms regulating EtOH abuse-induced brain changes remains unclear. These studies provide important data that peripubertal binge EtOH alters microRNA expression in both an immediate and long-term fashion. Our data also suggest

that BDNF and SIRT1 are potential effector molecules of microRNAs altered by EtOH. However, the mechanisms regarding miR biogenesis as well as their regulation of target genes expression remains controversial [223,419], and thus, the mechanisms underlying EtOH-mediated regulation of miR expression as well as whether changes in miR target gene expression plays a role in altering mood or memory function requires more studies. Future studies are important in order to increase our understanding of the molecular events leading to the maturation of an organized and functional brain and therefore, increase the capacity for human livelihood.

#### *Future Directions*

One of the most interesting findings regarding microRNAs gene regulatory capacity is that their mechanisms of gene regulation and even their biogenesis can be mediated by different mechanisms. In particular, the potential for microRNAs to function as endocrine molecules is an exciting new prospect. In this respect, circulating microRNAs are suggested to mediate cell to cell communication in a long-distance fashion to various cells and organs throughout the body. Moreover, the potential for microRNAs to be able to function as biomarkers and as tools for disease prevention. Just in this past year, studies have highlighted microRNAs as biomarkers for various neurological impairments including but not limited to autism [420], glioblastoma [421], neurodegeneration [422], stroke [423,424] and neurotoxicity [425]. Therefore, microRNAs provide promising opportunities for better understanding and potentially treating or preventing ETOH abuse-mediated neurological impairments.

## CHAPTER SEVEN

### GENERAL METHODS

#### **Animals**

All Wistar rats were purchased from Charles River Laboratories (Wilmington, MA) at weaning (postnatal day (PND) 23)) and allowed to acclimate for 7 days, after which they were randomly distributed into 3 groups: early pubertal (PND 30), peripubertal (PND 37), and late pubertal (PND 73). For Chapter 2 and 3 experiments there were *male and female* early, peripubertal and late pubertal grouped animals which were left untreated (N=10/age group) until euthanized at PND 30, PND 44 and PND 73, respectively. For Chapter 3 *whole hippocampus* experiments, there were 2 male groups, one administered treatments of peripubertal water (control) at peripubertal age and the other administered treatments of peripubertal binge EtOH (see methods below) (N= 6/treatment group; total of 12 animals). These 2 groups of animals were euthenized 1 hr. following the last treatment. For Chapter 3 *dorsal and ventral hippocampus experiments*, there were 3 male groups left untreated and euthenized at early (PND 30), mid (PND 44) and late (PND 73) puberty. In addition, there were 2 male groups administered treatments of peripubertal water (control) at peripubertal age (N= 10/treatment group; total of 20 water-treated animals) and 2 male groups administered treatments of peripubertal binge EtOH (N= 10/treatment group; total of 20 EtOH-treated animals) (Fig. 19B). The water and EtOH groups were handled for 5 minutes once/day beginning at PND 30 to eliminate non-specific effects of stress associated with handling. Peripubertal water or EtOH

treatments (see methods below) began on PND 37, an age which has been previously defined as peripuberty based on circulating gonadotropin levels and stages of spermatogenesis [286,287,288]. One group each from of the mid/peri-pubertal water and EtOH treated animals were sacrificed 60 min. following the last water and EtOH treatment, respectively, at PND 44 (N= 10 water + N= 10 EtOH). The remaining 2 groups (N= 10 water + N= 10 EtOH) were left undisturbed following the last EtOH treatment in their home cage until sacrificed at late puberty (PND 73). All animals were pair-housed on a 12:12 light/dark cycle with lights on at 07.00 h. Food and water were available ad libitum.

### **Ethics Statement**

All animal protocols were approved by the Institutional Animal Care and Use Committee at Loyola University Chicago permit #2011002. All measures were taken to minimize animal numbers and suffering.

### **Binge Exposure Paradigm and Treatment Design**

For Chapter 3 *whole hippocampus* experiments, peripubertal (PND 37) male animals were randomly assigned to either 1) peripubertal water (N = 6) or 2) peripubertal binge EtOH (N = 6) treatment groups, such that a total of 12 animals were used in these experiments. For Chapter 3 *dorsal and ventral hippocampus* experiments, peripubertal (PND 37) male animals were randomly assigned to either 1) peripubertal water (N = 20) or 2) peripubertal binge EtOH (N = 20) treatment groups. These animals were compared to the untreated groups (N=10) at each age, such that a total of 70 animals were used in these studies.

All animals in the binge EtOH group received the following 8-day peripubertal repeated binge-pattern EtOH paradigm: on PND 37, animals are given 3g/kg EtOH (20% v/v in water), or water alone, via oral gavage at 10.00 hr. This process is repeated according to the following schedule: 3d EtOH, 2d water, 3d EtOH (i.e. a total of 6 EtOH treatments over the course of 8 days). This binge exposure paradigm has been used previously to mimic the pattern of binge alcohol consumption typically observed in adolescents [44,200]. Moreover, our previous studies have shown that this repeated binge-pattern EtOH paradigm did not affect body weight/growth curves during pubertal development and consistently resulted in similar blood alcohol concentrations (BAC) [8,44,200]. The water group was administered room temperature tap water via oral gavage once/day for 8 consecutive days. The animals were anesthetized with inhalation of isoflurane then euthanized by rapid decapitation 60 minutes following EtOH treatment (PND 44, “immediate EtOH effects”) or [in the Chapter 3 *ventral and dorsal* hippocampus experiments] 30 days following the last day of treatment at late puberty (PND 73, “long-term EtOH effects”). The blood alcohol concentration in EtOH treated animals was  $190 \pm 21$  mg/dl, which is consistent with our previous reports using this peripubertal binge EtOH paradigm [12]. It is important to note that the untreated and the water-treated mid/peri- and late-pubertal groups were not statistically different for any parameter measured and were therefore, combined into one group for further statistical analyses.

### **Tissue Collection**

Trunk blood and brains were collected immediately following decapitation. Trunk blood was collected on ice into heparinized glass tubes, centrifuged at 4000 rpm for 10

minutes, plasma separated and stored at -20°C until processed for testosterone levels using enzyme immunoassay (EIA, see below). Brains were rapidly dissected, flash-frozen in isopentane (-35°C) on dry ice, and stored at -80°C until further processing. Frozen brains were sectioned at 200 µm on a freezing microtome, mounted onto glass slides, and the ventral and dorsal hippocampi were microdissected using a 0.75 mm Palkovit's brain punch tool (Stoelting Co., Wood Dale, IL). The coordinates of the microdissected regions was determined using *The Rat Brain in Stereotaxic Coordinates, Fourth Edition Atlas* (G. Paxinos and C. Watson). The ventral hippocampus was defined as being located between 1.8 mm to 3.8 mm posterior to Bregma, 3 mm below the top of the brain and 6.6 mm from the bottom of the brain. The dorsal hippocampus was defined as being located between 4.16 mm and 6.05 mm posterior to Bregma, 3 mm below the top of the brain and 2 mm above the bottom of the brain. Brain tissue punch samples were collected on ice into microcentrifuge tubes containing 1 ml of TriZol reagent (Invitrogen, Inc., Carlsbad, CA). Tissue samples were sonicated on ice prior to total RNA isolation.

### **RNA Isolation**

Total RNA was isolated from micropunched tissue samples using Trizol reagent (Invitrogen Inc., Carlsbad, CA) according to the manufacturer's directions. All RNA samples were analyzed for quality by Nanodrop spectrophotometry and by visualization of the RNA on a 1.5% agarose gel.

### **Rat RT<sup>2</sup> microRNA PCR Array**

The Rat RT<sup>2</sup> microRNA PCR array (SABiosciences) was performed according to manufacturer's instructions. Briefly, the cDNA was combined with RT<sup>2</sup> SYBR Green

qPCR Master Mix, RT<sup>2</sup> microRNA universal primer and water. Next, the mixture was aliquoted across the 96-well RT<sup>2</sup> microRNA PCR Array platform containing a panel of primers for 88 well-researched microRNAs in the rat genome followed by quantitative reverse transcription PCR quantification of gene expression using the  $\Delta\Delta C_t$  method [426]. Data analysis was performed using the free Web and Excel based microRNA PCR Array Data Analysis Software. (Importantly, the RNA population used for this experiment was a microRNA population isolated from total RNA using Qiagen's MiRNeasy Mini Kit and cDNA synthesis was performed using SABiosciences RT<sup>2</sup> microRNA First Strand Kit (MA-03) according to the manufacturer's instructions).

### **Quantitative Reverse Transcription PCR (qRT-PCR)**

Following RNA isolation, 1.0  $\mu\text{g}$  total RNA was reverse transcribed using the First Strand Synthesis SuperMix for qRT-PCR (Invitrogen, Inc., Carlsbad, CA) for mRNA quantification, and 1.0  $\mu\text{g}$  total RNA was used for NCode microRNA First-Strand cDNA Synthesis Kit for microRNA quantification. microRNA and mRNA qRT-PCR was performed with Fast Start Universal SYBR Green Master Mix (Roche) on an Eppendorf Realplex4 with a silver block. Forward primers for specific microRNAs were designed as described in the Ncode™ microRNA First-Strand cDNA synthesis kit handbook (Invitrogen, Inc., Carlsbad, CA) and using miRBase 18 as a sequence reference. The small RNA, U6 and housekeeping gene hypoxanthine guanine phosphoribosyl transferase 1 (HPRT) were used as a loading control and to normalize the data for microRNA and mRNA analysis, respectively, as neither were altered by EtOH treatment [44]. The following thermocycler program was used for mRNA target genes: 1) 95°C for 10 minutes, 2) 95°C for 30 seconds, 3) 59°C for 30 seconds, 4) 72°C for 30 seconds, and

melting curve analysis. The following thermocycler program was used for microRNA: 1) 95°C for 10 minutes, 2) 95°C for 30 seconds, 3) 65.3°C for 20 seconds, 4) 72°C for 12 seconds. Quantification of the gene expression was achieved using the  $\Delta\Delta C_t$  method [426]. Importantly, all miR and mRNA fold change values were generated using the normalized expression levels of untreated male PND 30 rats as a baseline value. The following intron-spanning primers were used for analysis of selected microRNA target genes and for microRNA processing enzymes:

SIRT1: 5'GCGGCCGCGGATAGGTCCATA, 3'TCCCACAGGAGACAGAAACCCCA,

BDNF: 5'AGCCTCCTCTGCTCTTTCTGCTGGA,

3'CTTTTGTCTATGCCCTGCAGCCTT,

Drosha: 5'GAAGTCACCGTGGAGCTGAGTA,

3'ATCATTGCATGCTGACAGACATC,

Dicer: 5'GGGAAAGTCTGCAGAACAAAC AND

3'GGCTGTCTGAGCTCTTAGTTC.

The following forward primers were used for analysis of selected mature microRNA along with a universal reverse primer provided in the NCode microRNA First-Strand cDNA Synthesis Kit:

miR-10a-5p: 5'CGCTACCCTGTAGATCCGAATTTGTG,

miR-26a: 5'CCGGTTCAAGTAATCCAGGATAGGC,

miR-103: 5'GGAGCAGCATTGTACAGGGCTATGA,

miR-495: 5'CGCGAAACAAACATGGTGCACCTTCTT.



let-7i: 5'CGCGTGAGGTAGTAGTTTGTGCTGTT

miR-7a: 5'GCGCTGGAAGACTAGTGATTTTGTCT

miR-9: 5' CGCGTCTTTGGTTATCTAGCTGTATG

miR-125a: 5'CGTCCCTGAGACCCTTTAACCTGTGA

miR-181a: 5'CGAACATTCAACGCTGTCGGTGAGT

### **Hormone Measurements**

Plasma levels of testosterone were measured using a commercially available EIA kit (Cayman Chemical, Ann Arbor, MI) according to manufacturer's instructions. The range of detection was between 3.9 and 500pg/ml and the intra-assay CVs was 2.2. Briefly, blood samples were collected in heparinized tubes and centrifuged at 3,000 rpm for 10 min. at 4°C, and plasma was stored at -20°C. Plasma samples were combined with a testosterone-acetylcholinesterase (AChE) conjugate (testosterone tracer) as well as testosterone EIA antiserum and incubated in a 96-well IgG-coated plate for 2.0 hr. at room temperature. Samples were washed 5 times with provided wash buffer then combined with Ellman's reagent containing the substrate for AChE, and the plate was developed for 60 minutes, shaking and covered, at room temperature. Absorbance was read at 412 nm on a multimode Synergy HT plate reader (BioTek Instruments, Inc., Winooski, VT).

### **Blood Alcohol Concentration Assay**

Trunk blood samples were collected into heparinized tubes, centrifuged at 3000 rpm for 10 min. at 4°C; and plasma stored at -20°C. Blood alcohol levels were

determined by measuring the change in absorbance at 340 nm following enzymatic oxidation of EtOH to acetaldehyde (Point Scientific Alcohol Reagent Kit). Assay range is 0 to 400 mg/dl and intra and interassay CV = 6.4% and 7.9%, respectively.

### **Western Blot**

Total protein is isolated using Tissue Protein Extraction Reagent (Thermo Scientific, Waltham, MA) supplemented with protease inhibitor cocktail (Roche catalogue #04693159001; 7x stock solution), according to manufacturer's instructions and boiled at 95<sup>o</sup> C for 5 mins. 10 µg of protein was loaded onto 10% SDS-PAGE gel and then transferred onto a PVDF membrane (Millipore, Billerica, MA). The membrane was blocked with 5% bovine serum albumin (BSA) for 0.5 hr. and incubated with the following primary antibodies: SIRT1 (Santa Cruz Biotechnology, H-300) at 1:200 or BDNF (Santa Cruz Biotechnology, N-20) at 1:200 in TBST for 2 hr. at 4°C. Following primary antibody incubation, the membrane was washed three times for 10 min. each in 10 ml TBST (Tris Base Solution containing 0.1% Tween 20), incubated in secondary antibody (HRP conjugated goat anti rabbit IgG, Santa Cruz Biotechnology) at 1:5000 concentration in TBST for 1.5 hr., and washed three times for 10 min. each in TBST. Stripping blots was performed with 6 M GnHCl, 0.2% Triton 100, 20mM Tris-HCl pH8, two times at RT for 5 min. each followed by two 10 min. washes with TBSTr. In order to control loading efficiency, blots are stripped as described above, re-blocked with 5% BSA and incubated in primary rabbit β-actin antibody (Cell signaling, 4970S) at 1:3000 dilution in TBST for 1.0 hr., washed three times for 10 min. each in TBST, incubated in HRP conjugated goat α rabbit IgG (Santa Cruz Biotechnology, sc-2004) at 1:5000 dilution in TBST for 1.5 hr., and washed three times for 10 min. each in TBST.

Following antibody applications and washes, each blot was imaged on the Biorad Chemidoc XRS+ imager using ECL Chemiluminescent substrate (Pierce).

### **Statistical Analysis**

Statistical analyses were performed by the Biostatistics Core Facility at Loyola University Stritch School of Medicine in consultation with Dr. James Sinacore. Data obtained from qRT-PCR or blood alcohol and hormone concentration assays were analyzed by a one-way Analysis of Variance (ANOVA) when EtOH was a single variable factor or by a two-way ANOVA when age and EtOH or age and sex were the two variable factors followed by Tukey's posthoc test for all pair-wise comparisons when there was a significant main effect and interaction. The above tests were performed using SigmaStat Statistical Analysis Software. A p-value of less than 0.05 was designated as significant. Exclusion criteria comprised outliers greater than or equal to 2 times the standard deviation of the mean. Please note that all qRT-PCR age-dependent miR and mRNA fold change values were generated using the normalized expression levels of untreated male PND 30 rats as a baseline value. Importantly, the untreated and the water-treated peri and late pubertal groups were not statistically different for any parameter measured and were therefore, combined into one group for statistical analyses.

Data obtained from western blotting were subject to densitometry analysis using ImageLab software. Lanes were detected manually and bands were detected using the 'high sensitivity' detection limit and lane-based background subtraction was applied. Statistical significance was analyzed by a two-way ANOVA followed by Tukey's post-hoc test using an average of 3 independent blots containing samples from 6 different

animals per treatment group ( $N=3$ ,  $p<0.05$ ). Western blotting data analysis tests were performed using SigmaStat Statistical Analysis Software. A p-value of less than 0.05 was designated as significant. Exclusion criteria comprised outliers greater than or equal to 3 times the standard deviation of the mean.

## REFERENCE LIST

1. Brown SA, Tapert SF, Granholm E, Delis DC (2000) Neurocognitive functioning of adolescents: effects of protracted alcohol use. *Alcohol Clin Exp Res* 24: 164-171.
2. Crews FT, Braun CJ, Hoplight B, Switzer RC, 3rd, Knapp DJ (2000) Binge ethanol consumption causes differential brain damage in young adolescent rats compared with adult rats. *Alcohol Clin Exp Res* 24: 1712-1723.
3. De Bellis MD, Clark DB, Beers SR, Soloff PH, Boring AM, et al. (2000) Hippocampal volume in adolescent-onset alcohol use disorders. *Am J Psychiatry* 157: 737-744.
4. Swartzwelder HS, Wilson WA, Tayyeb MI (1995) Age-dependent inhibition of long-term potentiation by ethanol in immature versus mature hippocampus. *Alcohol Clin Exp Res* 19: 1480-1485.
5. Swartzwelder HS, Wilson WA, Tayyeb MI (1995) Differential sensitivity of NMDA receptor-mediated synaptic potentials to ethanol in immature versus mature hippocampus. *Alcohol Clin Exp Res* 19: 320-323.
6. Tapert SF, Brown SA (1999) Neuropsychological correlates of adolescent substance abuse: four-year outcomes. *J Int Neuropsychol Soc* 5: 481-493.
7. White AM, Swartzwelder HS (2005) Age-related effects of alcohol on memory and memory-related brain function in adolescents and adults. *Recent Dev Alcohol* 17: 161-176.
8. Walker BM, Ehlers CL (2009) Age-related differences in the blood alcohol levels of Wistar rats. *Pharmacol Biochem Behav* 91: 560-565.
9. Morris SA, Kelso ML, Liput DJ, Marshall SA, Nixon K (2010) Similar withdrawal severity in adolescents and adults in a rat model of alcohol dependence. *Alcohol* 44: 89-98.
10. Morris SA, Eaves DW, Smith AR, Nixon K (2010) Alcohol inhibition of neurogenesis: a mechanism of hippocampal neurodegeneration in an adolescent alcohol abuse model. *Hippocampus* 20: 596-607.
11. Richardson HN, Chan SH, Crawford EF, Lee YK, Funk CK, et al. (2009) Permanent impairment of birth and survival of cortical and hippocampal proliferating cells following excessive drinking during alcohol dependence. *Neurobiol Dis* 36: 1-10.

12. Przybycien-Szymanska MM, Mott NN, Paul CR, Gillespie RA, Pak TR (2011) Binge-pattern alcohol exposure during puberty induces long-term changes in HPA axis reactivity. *PLoS ONE* 6: e18350.
13. Giedd JN (2004) Structural magnetic resonance imaging of the adolescent brain. *Ann N Y Acad Sci* 1021: 77-85.
14. Arnold AP, Breedlove SM (1985) Organizational and activational effects of sex steroids on brain and behavior: a reanalysis. *Horm Behav* 19: 469-498.
15. Schulz KM, Molenda-Figueira HA, Sisk CL (2009) Back to the future: The organizational-activational hypothesis adapted to puberty and adolescence. *Horm Behav* 55: 597-604.
16. Romeo RD (2003) Puberty: a period of both organizational and activational effects of steroid hormones on neurobehavioural development. *J Neuroendocrinol* 15: 1185-1192.
17. Spear LP (2000) The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 24: 417-463.
18. Kim VN (2005) Small RNAs: classification, biogenesis, and function. *Mol Cells* 19: 1-15.
19. Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116: 281-297.
20. Mansfield JH, Harfe BD, Nissen R, Obenaus J, Srineel J, et al. (2004) MicroRNA-responsive 'sensor' transgenes uncover Hox-like and other developmentally regulated patterns of vertebrate microRNA expression. *Nat Genet* 36: 1079-1083.
21. Ohuchida K, Mizumoto K, Lin C, Yamaguchi H, Ohtsuka T, et al. (2012) MicroRNA-10a is overexpressed in human pancreatic cancer and involved in its invasiveness partially via suppression of the HOXA1 gene. *Ann Surg Oncol* 19: 2394-2402.
22. Hunsberger JG, Fessler EB, Chibane FL, Leng Y, Maric D, et al. (2013) Mood stabilizer-regulated miRNAs in neuropsychiatric and neurodegenerative diseases: identifying associations and functions. *Am J Transl Res* 5: 450-464.
23. Gao J, Wang WY, Mao YW, Graff J, Guan JS, et al. (2010) A novel pathway regulates memory and plasticity via SIRT1 and miR-134. *Nature* 466: 1105-1109.
24. Liu IY, Lyons WE, Mamounas LA, Thompson RF (2004) Brain-derived neurotrophic factor plays a critical role in contextual fear conditioning. *J Neurosci* 24: 7958-7963.
25. Michan S, Li Y, Chou MM, Parrella E, Ge H, et al. (2010) SIRT1 is essential for normal cognitive function and synaptic plasticity. *J Neurosci* 30: 9695-9707.

26. Patterson SL, Abel T, Deuel TA, Martin KC, Rose JC, et al. (1996) Recombinant BDNF rescues deficits in basal synaptic transmission and hippocampal LTP in BDNF knockout mice. *Neuron* 16: 1137-1145.
27. Tyler WJ, Alonso M, Bramham CR, Pozzo-Miller LD (2002) From acquisition to consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-dependent learning. *Learn Mem* 9: 224-237.
28. Rao YS, Mott NN, Wang Y, Chung WC, Pak TR (2013) MicroRNAs in the aging female brain: a putative mechanism for age-specific estrogen effects. *Endocrinology* 154: 2795-2806.
29. Morgan CP, Bale TL (2011) Early prenatal stress epigenetically programs dysmasculinization in second-generation offspring via the paternal lineage. *J Neurosci* 31: 11748-11755.
30. Baumrind D (1987) A developmental perspective on adolescent risk taking in contemporary America. *New Dir Child Dev*: 93-125.
31. Arain M, Haque M, Johal L, Mathur P, Nel W, et al. (2013) Maturation of the adolescent brain. *Neuropsychiatr Dis Treat* 9: 449-461.
32. Giedd JN, Clasen LS, Lenroot R, Greenstein D, Wallace GL, et al. (2006) Puberty-related influences on brain development. *Mol Cell Endocrinol* 254-255: 154-162.
33. <pubertal onset in primates.pdf>.
34. Bourgeois JP, Rakic P (1993) Changes of synaptic density in the primary visual cortex of the macaque monkey from fetal to adult stage. *J Neurosci* 13: 2801-2820.
35. Benes FM, Taylor JB, Cunningham MC (2000) Convergence and plasticity of monoaminergic systems in the medial prefrontal cortex during the postnatal period: implications for the development of psychopathology. *Cereb Cortex* 10: 1014-1027.
36. Nunez JL, Lauschke DM, Juraska JM (2001) Cell death in the development of the posterior cortex in male and female rats. *J Comp Neurol* 436: 32-41.
37. Andersen SL, Thompson AT, Rutstein M, Hostetter JC, Teicher MH (2000) Dopamine receptor pruning in prefrontal cortex during the periadolescent period in rats. *Synapse* 37: 167-169.
38. Gerocs K, Rethelyi M, Halasz B (1986) Quantitative analysis of dendritic protrusions in the medial preoptic area during postnatal development. *Brain Res* 391: 49-57.
39. Lewis DA, Cruz D, Eggen S, Erickson S (2004) Postnatal development of prefrontal inhibitory circuits and the pathophysiology of cognitive dysfunction in schizophrenia. *Ann N Y Acad Sci* 1021: 64-76.

40. White AM, Kraus CL, Swartzwelder H (2006) Many college freshmen drink at levels far beyond the binge threshold. *Alcohol Clin Exp Res* 30: 1006-1010.
41. White AM, Bae JG, Truesdale MC, Ahmad S, Wilson WA, et al. (2002) Chronic-intermittent ethanol exposure during adolescence prevents normal developmental changes in sensitivity to ethanol-induced motor impairments. *Alcohol Clin Exp Res* 26: 960-968.
42. Ogilvie KM, Rivier C (1997) Gender difference in hypothalamic-pituitary-adrenal axis response to alcohol in the rat: activational role of gonadal steroids. *Brain Res* 766: 19-28.
43. Romeo RD, Bellani R, Karatsoreos IN, Chhua N, Vernov M, et al. (2006) Stress history and pubertal development interact to shape hypothalamic-pituitary-adrenal axis plasticity. *Endocrinology* 147: 1664-1674.
44. Przybycien-Szymanska MM, Rao YS, Pak TR (2010) Binge-pattern alcohol exposure during puberty induces sexually dimorphic changes in genes regulating the HPA axis. *Am J Physiol Endocrinol Metab* 298: E320-328.
45. Rivier C, Lee S (1996) Acute alcohol administration stimulates the activity of hypothalamic neurons that express corticotropin-releasing factor and vasopressin. *Brain Res* 726: 1-10.
46. Rivier C (1993) Female rats release more corticosterone than males in response to alcohol: influence of circulating sex steroids and possible consequences for blood alcohol levels. *Alcohol Clin Exp Res* 17: 854-859.
47. Przybycien-Szymanska MM, Gillespie RA, Pak TR (2012) 17beta-Estradiol is required for the sexually dimorphic effects of repeated binge-pattern alcohol exposure on the HPA axis during adolescence. *PLoS One* 7: e32263.
48. Herman JP, Figueiredo H, Mueller NK, Ulrich-Lai Y, Ostrander MM, et al. (2003) Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. *Front Neuroendocrinol* 24: 151-180.
49. Barden N (2004) Implication of the hypothalamic-pituitary-adrenal axis in the physiopathology of depression. *J Psychiatry Neurosci* 29: 185-193.
50. O'Toole SM, Sekula LK, Rubin RT (1997) Pituitary-adrenal cortical axis measures as predictors of sustained remission in major depression. *Biol Psychiatry* 42: 85-89.
51. Muller MB, Holsboer F (2006) Mice with mutations in the HPA-system as models for symptoms of depression. *Biol Psychiatry* 59: 1104-1115.



52. Bradley AJ, Dinan TG (2010) A systematic review of hypothalamic-pituitary-adrenal axis function in schizophrenia: implications for mortality. *J Psychopharmacol* 24: 91-118.
53. Tsigos C, Chrousos GP (2002) Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *J Psychosom Res* 53: 865-871.
54. Landgraf R, Wigger A, Holsboer F, Neumann ID (1999) Hyper-reactive hypothalamo-pituitary-adrenocortical axis in rats bred for high anxiety-related behaviour. *J Neuroendocrinol* 11: 405-407.
55. Goldstein JM, Jerram M, Abbs B, Whitfield-Gabrieli S, Makris N (2010) Sex differences in stress response circuitry activation dependent on female hormonal cycle. *J Neurosci* 30: 431-438.
56. Holsen LM, Lee JH, Spaeth SB, Ogden LA, Klibanski A, et al. (2012) Brain hypoactivation, autonomic nervous system dysregulation, and gonadal hormones in depression: a preliminary study. *Neurosci Lett* 514: 57-61.
57. Moser MB, Moser EI (1998) Functional differentiation in the hippocampus. *Hippocampus* 8: 608-619.
58. Fanselow MS, Dong HW (2010) Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* 65: 7-19.
59. Lau JY, Goldman D, Buzas B, Hodgkinson C, Leibenluft E, et al. (2010) BDNF gene polymorphism (Val66Met) predicts amygdala and anterior hippocampus responses to emotional faces in anxious and depressed adolescents. *Neuroimage* 53: 952-961.
60. Dong HW, Swanson LW, Chen L, Fanselow MS, Toga AW (2009) Genomic-anatomic evidence for distinct functional domains in hippocampal field CA1. *Proc Natl Acad Sci U S A* 106: 11794-11799.
61. Thompson CL, Pathak SD, Jeromin A, Ng LL, MacPherson CR, et al. (2008) Genomic anatomy of the hippocampus. *Neuron* 60: 1010-1021.
62. Grant I (1987) Alcohol and the brain: neuropsychological correlates. *J Consult Clin Psychol* 55: 310-324.
63. Pfefferbaum A, Rosenbloom M, Deshmukh A, Sullivan E (2001) Sex differences in the effects of alcohol on brain structure. *Am J Psychiatry* 158: 188-197.
64. Sullivan EV, Harding AJ, Pentney R, Dlugos C, Martin PR, et al. (2003) Disruption of frontocerebellar circuitry and function in alcoholism. *Alcohol Clin Exp Res* 27: 301-309.

65. Pfefferbaum A, Sullivan EV, Hedehus M, Adalsteinsson E, Lim KO, et al. (2000) In vivo detection and functional correlates of white matter microstructural disruption in chronic alcoholism. *Alcohol Clin Exp Res* 24: 1214-1221.
66. Pfefferbaum A, Lim KO, Desmond JE, Sullivan EV (1996) Thinning of the corpus callosum in older alcoholic men: a magnetic resonance imaging study. *Alcohol Clin Exp Res* 20: 752-757.
67. Conway KP, Compton W, Stinson FS, Grant BF (2006) Lifetime comorbidity of DSM-IV mood and anxiety disorders and specific drug use disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions. *J Clin Psychiatry* 67: 247-257.
68. Epstein EE, Fischer-Elber K, Al-Otaiba Z (2007) Women, aging, and alcohol use disorders. *J Women Aging* 19: 31-48.
69. Rohde P, Lewinsohn PM, Seeley JR (1996) Psychiatric comorbidity with problematic alcohol use in high school students. *J Am Acad Child Adolesc Psychiatry* 35: 101-109.
70. Ross HE, Glaser FB, Germanson T (1988) The prevalence of psychiatric disorders in patients with alcohol and other drug problems. *Arch Gen Psychiatry* 45: 1023-1031.
71. Silberman Y, Bajo M, Chappell AM, Christian DT, Cruz M, et al. (2009) Neurobiological mechanisms contributing to alcohol-stress-anxiety interactions. *Alcohol* 43: 509-519.
72. Zilberman ML, Tavares H, Blume SB, el-Guebaly N (2003) Substance use disorders: sex differences and psychiatric comorbidities. *Can J Psychiatry* 48: 5-13.
73. Blume A, Bosch OJ, Miklos S, Torner L, Wales L, et al. (2008) Oxytocin reduces anxiety via ERK1/2 activation: local effect within the rat hypothalamic paraventricular nucleus. *Eur J Neurosci* 27: 1947-1956.
74. Madeira MD, Andrade JP, Lieberman AR, Sousa N, Almeida OF, et al. (1997) Chronic alcohol consumption and withdrawal do not induce cell death in the suprachiasmatic nucleus, but lead to irreversible depression of peptide immunoreactivity and mRNA levels. *J Neurosci* 17: 1302-1319.
75. Merlo Pich E, Lorang M, Yeganeh M, Rodriguez de Fonseca F, Raber J, et al. (1995) Increase of extracellular corticotropin-releasing factor-like immunoreactivity levels in the amygdala of awake rats during restraint stress and ethanol withdrawal as measured by microdialysis. *J Neurosci* 15: 5439-5447.
76. Ogilvie KM, Lee S, Rivier C (1997) Role of arginine vasopressin and corticotropin-releasing factor in mediating alcohol-induced adrenocorticotropin and vasopressin secretion in male rats bearing lesions of the paraventricular nuclei. *Brain Res* 744: 83-95.

77. Rivier C, Bruhn T, Vale W (1984) Effect of ethanol on the hypothalamic-pituitary-adrenal axis in the rat: role of corticotropin-releasing factor (CRF). *J Pharmacol Exp Ther* 229: 127-131.
78. Tabakoff B, Jafee RC, Ritzmann RF (1978) Corticosterone concentrations in mice during ethanol drinking and withdrawal. *J Pharm Pharmacol* 30: 371-374.
79. Markwiese BJ, Acheson SK, Levin ED, Wilson WA, Swartzwelder HS (1998) Differential effects of ethanol on memory in adolescent and adult rats. *Alcohol Clin Exp Res* 22: 416-421.
80. Kang HJ, Schuman EM (1995) Neurotrophin-induced modulation of synaptic transmission in the adult hippocampus. *J Physiol Paris* 89: 11-22.
81. Kang H, Schuman EM (1995) Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. *Science* 267: 1658-1662.
82. Kang H, Jia LZ, Suh KY, Tang L, Schuman EM (1996) Determinants of BDNF-induced hippocampal synaptic plasticity: role of the Trk B receptor and the kinetics of neurotrophin delivery. *Learn Mem* 3: 188-196.
83. Kang H, Schuman EM (1996) A requirement for local protein synthesis in neurotrophin-induced hippocampal synaptic plasticity. *Science* 273: 1402-1406.
84. Schuman EM (1999) Neurotrophin regulation of synaptic transmission. *Curr Opin Neurobiol* 9: 105-109.
85. Schinder AF, Poo M (2000) The neurotrophin hypothesis for synaptic plasticity. *Trends Neurosci* 23: 639-645.
86. Scharfman HE, MacLusky NJ (2014) Differential regulation of BDNF, synaptic plasticity and sprouting in the hippocampal mossy fiber pathway of male and female rats. *Neuropharmacology* 76 Pt C: 696-708.
87. Sutherland RJ, McDonald RJ, Savage DD (1997) Prenatal exposure to moderate levels of ethanol can have long-lasting effects on hippocampal synaptic plasticity in adult offspring. *Hippocampus* 7: 232-238.
88. Titterness AK, Christie BR (2012) Prenatal ethanol exposure enhances NMDAR-dependent long-term potentiation in the adolescent female dentate gyrus. *Hippocampus* 22: 69-81.
89. Sickmann HM, Patten AR, Morch K, Sawchuk S, Zhang C, et al. (2014) Prenatal ethanol exposure has sex-specific effects on hippocampal long-term potentiation. *Hippocampus* 24: 54-64.

90. Heaton MB, Mitchell JJ, Paiva M, Walker DW (2000) Ethanol-induced alterations in the expression of neurotrophic factors in the developing rat central nervous system. *Brain Res Dev Brain Res* 121: 97-107.
91. Pyapali GK, Turner DA, Wilson WA, Swartzwelder HS (1999) Age and dose-dependent effects of ethanol on the induction of hippocampal long-term potentiation. *Alcohol* 19: 107-111.
92. Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, et al. (2003) The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 112: 257-269.
93. Agapito MA, Zhang C, Murugan S, Sarkar DK (2014) Fetal Alcohol Exposure Disrupts Metabolic Signaling in Hypothalamic Proopiomelanocortin Neurons via a Circadian Mechanism in Male Mice. *Endocrinology* 155: 2578-2588.
94. Codocedo JF, Allard C, Godoy JA, Varela-Nallar L, Inestrosa NC (2012) SIRT1 regulates dendritic development in hippocampal neurons. *PLoS One* 7: e47073.
95. Chen J, Zhou Y, Mueller-Steiner S, Chen LF, Kwon H, et al. (2005) SIRT1 protects against microglia-dependent amyloid-beta toxicity through inhibiting NF-kappaB signaling. *J Biol Chem* 280: 40364-40374.
96. Kim D, Nguyen MD, Dobbin MM, Fischer A, Sananbenesi F, et al. (2007) SIRT1 deacetylase protects against neurodegeneration in models for Alzheimer's disease and amyotrophic lateral sclerosis. *EMBO J* 26: 3169-3179.
97. Sakamoto J, Miura T, Shimamoto K, Horio Y (2004) Predominant expression of Sir2alpha, an NAD-dependent histone deacetylase, in the embryonic mouse heart and brain. *FEBS Lett* 556: 281-286.
98. Parker JA, Arango M, Abderrahmane S, Lambert E, Tourette C, et al. (2005) Resveratrol rescues mutant polyglutamine cytotoxicity in nematode and mammalian neurons. *Nat Genet* 37: 349-350.
99. Saharan S, Jhaveri DJ, Bartlett PF (2013) SIRT1 regulates the neurogenic potential of neural precursors in the adult subventricular zone and hippocampus. *J Neurosci Res* 91: 642-659.
100. Libert S, Cohen D, Guarente L (2008) Neurogenesis directed by Sirt1. *Nat Cell Biol* 10: 373-374.
101. Heyward FD, Walton RG, Carle MS, Coleman MA, Garvey WT, et al. (2012) Adult mice maintained on a high-fat diet exhibit object location memory deficits and reduced hippocampal SIRT1 gene expression. *Neurobiol Learn Mem* 98: 25-32.
102. Libert S, Pointer K, Bell EL, Das A, Cohen DE, et al. (2011) SIRT1 activates MAO-A in the brain to mediate anxiety and exploratory drive. *Cell* 147: 1459-1472.

103. O'Carroll D, Schaefer A (2013) General principals of miRNA biogenesis and regulation in the brain. *Neuropsychopharmacology* 38: 39-54.
104. Juhila J, Sipila T, Icaý K, Nicorici D, Ellonen P, et al. (2011) MicroRNA expression profiling reveals miRNA families regulating specific biological pathways in mouse frontal cortex and hippocampus. *PLoS One* 6: e21495.
105. Olsen L, Klausen M, Helboe L, Nielsen FC, Werge T (2009) MicroRNAs Show Mutually Exclusive Expression Patterns in the Brain of Adult Male Rats. *PLoS ONE* 4: e7225.
106. Amar L, Benoit C, Beaumont G, Vacher CM, Crepin D, et al. (2012) MicroRNA expression profiling of hypothalamic arcuate and paraventricular nuclei from single rats using Illumina sequencing technology. *J Neurosci Methods* 209: 134-143.
107. Bak M, Silaharoglu A, Moller M, Christensen M, Rath MF, et al. (2008) MicroRNA expression in the adult mouse central nervous system. *RNA* 14: 432-444.
108. He M, Liu Y, Wang X, Zhang MQ, Hannon GJ, et al. (2012) Cell-type-based analysis of microRNA profiles in the mouse brain. *Neuron* 73: 35-48.
109. Hua YJ, Tang ZY, Tu K, Zhu L, Li YX, et al. (2009) Identification and target prediction of miRNAs specifically expressed in rat neural tissue. *BMC Genomics* 10: 214.
110. Parsons MJ, Grimm CH, Paya-Cano JL, Sugden K, Nietfeld W, et al. (2008) Using hippocampal microRNA expression differences between mouse inbred strains to characterise miRNA function. *Mamm Genome* 19: 552-560.
111. Konopka W, Kiryk A, Novak M, Herwerth M, Parkitna JR, et al. (2010) MicroRNA loss enhances learning and memory in mice. *J Neurosci* 30: 14835-14842.
112. Schaefer A, O'Carroll D, Tan CL, Hillman D, Sugimori M, et al. (2007) Cerebellar neurodegeneration in the absence of microRNAs. *Journal of Experimental Medicine* 204: 1553-1558.
113. Schratt GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, et al. (2006) A brain-specific microRNA regulates dendritic spine development. *Nature* 439: 283-289.
114. Lambert TJ, Storm DR, Sullivan JM (2010) MicroRNA132 modulates short-term synaptic plasticity but not basal release probability in hippocampal neurons. *PLoS One* 5: e15182.
115. Magill ST, Cambronne XA, Luikart BW, Lioy DT, Leighton BH, et al. (2010) microRNA-132 regulates dendritic growth and arborization of newborn neurons in the adult hippocampus. *Proc Natl Acad Sci U S A* 107: 20382-20387.

116. Wu D, Raafat A, Pak E, Clemens S, Murashov AK (2012) Dicer-microRNA pathway is critical for peripheral nerve regeneration and functional recovery in vivo and regenerative axonogenesis in vitro. *Exp Neurol* 233: 555-565.
117. Baudry A, Mouillet-Richard S, Schneider B, Launay JM, Kellermann O (2010) miR-16 targets the serotonin transporter: a new facet for adaptive responses to antidepressants. *Science* 329: 1537-1541.
118. Haramati S, Navon I, Issler O, Ezra-Nevo G, Gil S, et al. (2011) MicroRNA as repressors of stress-induced anxiety: the case of amygdalar miR-34. *J Neurosci* 31: 14191-14203.
119. Miller BH, Zeier Z, Xi L, Lanz TA, Deng S, et al. (2012) MicroRNA-132 dysregulation in schizophrenia has implications for both neurodevelopment and adult brain function. *Proc Natl Acad Sci U S A* 109: 3125-3130.
120. Moreau MP, Bruse SE, David-Rus R, Buyske S, Brzustowicz LM (2011) Altered microRNA expression profiles in postmortem brain samples from individuals with schizophrenia and bipolar disorder. *Biol Psychiatry* 69: 188-193.
121. Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL (2008) The bifunctional microRNA miR-9/miR-9\* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* 28: 14341-14346.
122. Sethi P, Lukiw WJ (2009) Micro-RNA abundance and stability in human brain: specific alterations in Alzheimer's disease temporal lobe neocortex. *Neurosci Lett* 459: 100-104.
123. Kawahara H, Imai T, Okano H (2012) MicroRNAs in Neural Stem Cells and Neurogenesis. *Front Neurosci* 6: 30.
124. Treiber T, Treiber N, Meister G (2012) Regulation of microRNA biogenesis and function. *Thromb Haemost* 107: 605-610.
125. Lee Y, Kim M, Han J, Yeom KH, Lee S, et al. (2004) MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* 23: 4051-4060.
126. Lee Y, Jeon K, Lee JT, Kim S, Kim VN (2002) MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J* 21: 4663-4670.
127. Lee Y, Ahn C, Han J, Choi H, Kim J, et al. (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425: 415-419.
128. Han J, Lee Y, Yeom KH, Kim YK, Jin H, et al. (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* 18: 3016-3027.

129. Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120: 15-20.
130. Giraldez AJ (2005) MicroRNAs Regulate Brain Morphogenesis in Zebrafish. *Science* 308: 833-838.
131. Plasterk RHA (2006) Micro RNAs in Animal Development. *Cell* 124: 877-881.
132. Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, et al. (2003) Dicer is essential for mouse development. *Nature Genetics* 35: 215-217.
133. De Pietri Tonelli D, Pulvers JN, Haffner C, Murchison EP, Hannon GJ, et al. (2008) miRNAs are essential for survival and differentiation of newborn neurons but not for expansion of neural progenitors during early neurogenesis in the mouse embryonic neocortex. *Development* 135: 3911-3921.
134. Kawase-Koga Y, Otaegi G, Sun T (2009) Different timings of dicer deletion affect neurogenesis and gliogenesis in the developing mouse central nervous system. *Developmental Dynamics* 238: 2800-2812.
135. McLoughlin HS, Fineberg SK, Ghosh LL, Tecedor L, Davidson BL (2012) Dicer is required for proliferation, viability, migration and differentiation in corticoneurogenesis. *Neuroscience* 223: 285-295.
136. Siegel G, Obernosterer G, Fiore R, Oehmen M, Bicker S, et al. (2009) A functional screen implicates microRNA-138-dependent regulation of the depalmitoylation enzyme APT1 in dendritic spine morphogenesis. *Nat Cell Biol* 11: 705-716.
137. Lugli G, Torvik VI, Larson J, Smalheiser NR (2008) Expression of microRNAs and their precursors in synaptic fractions of adult mouse forebrain. *J Neurochem* 106: 650-661.
138. Kye MJ, Liu T, Levy SF, Xu NL, Groves BB, et al. (2007) Somatodendritic microRNAs identified by laser capture and multiplex RT-PCR. *RNA* 13: 1224-1234.
139. Huber KM, Kayser MS, Bear MF (2000) Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* 288: 1254-1257.
140. Martin KC, Casadio A, Zhu H, Yaping E, Rose JC, et al. (1997) Synapse-specific, long-term facilitation of aplysia sensory to motor synapses: a function for local protein synthesis in memory storage. *Cell* 91: 927-938.
141. Sutton MA, Schuman EM (2006) Dendritic protein synthesis, synaptic plasticity, and memory. *Cell* 127: 49-58.

142. Kim J, Krichevsky A, Grad Y, Hayes GD, Kosik KS, et al. (2004) Identification of many microRNAs that copurify with polyribosomes in mammalian neurons. *Proc Natl Acad Sci U S A* 101: 360-365.
143. Caputo V, Sinibaldi L, Fiorentino A, Parisi C, Catalanotto C, et al. (2011) Brain derived neurotrophic factor (BDNF) expression is regulated by microRNAs miR-26a and miR-26b allele-specific binding. *PLoS One* 6: e28656.
144. Hansen T, Olsen L, Lindow M, Jakobsen KD, Ullum H, et al. (2007) Brain expressed microRNAs implicated in schizophrenia etiology. *PLoS One* 2: e873.
145. Karege F, Perret G, Bondolfi G, Schwald M, Bertschy G, et al. (2002) Decreased serum brain-derived neurotrophic factor levels in major depressed patients. *Psychiatry Res* 109: 143-148.
146. Krebs MO, Guillin O, Bourdell MC, Schwartz JC, Olie JP, et al. (2000) Brain derived neurotrophic factor (BDNF) gene variants association with age at onset and therapeutic response in schizophrenia. *Mol Psychiatry* 5: 558-562.
147. McGough NN, He DY, Logrip ML, Jeanblanc J, Phamluong K, et al. (2004) RACK1 and brain-derived neurotrophic factor: a homeostatic pathway that regulates alcohol addiction. *J Neurosci* 24: 10542-10552.
148. Montag C, Basten U, Stelzel C, Fiebach CJ, Reuter M (2010) The BDNF Val66Met polymorphism and anxiety: Support for animal knock-in studies from a genetic association study in humans. *Psychiatry Research* 179: 86-90.
149. Park CS, Tang SJ (2009) Regulation of microRNA expression by induction of bidirectional synaptic plasticity. *J Mol Neurosci* 38: 50-56.
150. Morgan CP, Bale TL (2012) Sex differences in microRNA regulation of gene expression: no smoke, just miRs. *Biol Sex Differ* 3: 22.
151. Mellios N, Huang HS, Grigorenko A, Rogaev E, Akbarian S (2008) A set of differentially expressed miRNAs, including miR-30a-5p, act as post-transcriptional inhibitors of BDNF in prefrontal cortex. *Hum Mol Genet* 17: 3030-3042.
152. Hunsberger JG, Fessler EB, Wang Z, Elkahloun AG, Chuang DM (2012) Post-insult valproic acid-regulated microRNAs: potential targets for cerebral ischemia. *Am J Transl Res* 4: 316-332.
153. Miranda RC (2012) MicroRNAs and Fetal Brain Development: Implications for Ethanol Teratology during the Second Trimester Period of Neurogenesis. *Frontiers in Genetics* 3.
154. Miranda RC, Pietrzykowski AZ, Tang Y, Sathyan P, Mayfield D, et al. (2010) MicroRNAs: master regulators of ethanol abuse and toxicity? *Alcohol Clin Exp Res* 34: 575-587.



155. Chan AW, Kocerha J (2012) The Path to microRNA Therapeutics in Psychiatric and Neurodegenerative Disorders. *Front Genet* 3: 82.
156. Kapsimali M, Kloosterman WP, de Bruijn E, Rosa F, Plasterk RHA, et al. (2007) MicroRNAs show a wide diversity of expression profiles in the developing and mature central nervous system. *Genome Biology* 8: R173.
157. Tapocik JD, Barbier E, Flanigan M, Solomon M, Pincus A, et al. (2014) microRNA-206 in rat medial prefrontal cortex regulates BDNF expression and alcohol drinking. *J Neurosci* 34: 4581-4588.
158. Lewohl JM, Nunez YO, Dodd PR, Tiwari GR, Harris RA, et al. (2011) Up-regulation of microRNAs in brain of human alcoholics. *Alcohol Clin Exp Res* 35: 1928-1937.
159. Lam EW, Shah K, Brosens JJ (2012) The diversity of sex steroid action: the role of micro-RNAs and FOXO transcription factors in cycling endometrium and cancer. *J Endocrinol* 212: 13-25.
160. Kuokkanen S, Chen B, Ojalvo L, Benard L, Santoro N, et al. (2010) Genomic profiling of microRNAs and messenger RNAs reveals hormonal regulation in microRNA expression in human endometrium. *Biol Reprod* 82: 791-801.
161. Paris O, Ferraro L, Grober OM, Ravo M, De Filippo MR, et al. (2012) Direct regulation of microRNA biogenesis and expression by estrogen receptor beta in hormone-responsive breast cancer. *Oncogene* 31: 4196-4206.
162. Waltering KK, Porkka KP, Jalava SE, Urbanucci A, Kohonen PJ, et al. (2011) Androgen regulation of micro-RNAs in prostate cancer. *Prostate* 71: 604-614.
163. Bhat-Nakshatri P, Wang G, Collins NR, Thomson MJ, Geistlinger TR, et al. (2009) Estradiol-regulated microRNAs control estradiol response in breast cancer cells. *Nucleic Acids Res* 37: 4850-4861.
164. Kininis M, Kraus WL (2008) A global view of transcriptional regulation by nuclear receptors: gene expression, factor localization, and DNA sequence analysis. *Nucl Recept Signal* 6: e005.
165. Yamagata K, Fujiyama S, Ito S, Ueda T, Murata T, et al. (2009) Maturation of microRNA is hormonally regulated by a nuclear receptor. *Mol Cell* 36: 340-347.
166. Read JP, Beattie M, Chamberlain R, Merrill JE (2008) Beyond the "Binge" threshold: heavy drinking patterns and their association with alcohol involvement indices in college students. *Addict Behav* 33: 225-234.
167. Benes FM, Turtle M, Khan Y, Farol P (1994) MYelination of a key relay zone in the hippocampal formation occurs in the human brain during childhood, adolescence, and adulthood. *Archives of General Psychiatry* 51: 477-484.

168. Giedd JN, Blumenthal J, Jeffries NO, Castellanos FX, Liu H, et al. (1999) Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci* 2: 861-863.
169. Gogtay N, Nugent TF, Herman DH, Ordonez A, Greenstein D, et al. (2006) Dynamic mapping of normal human hippocampal development. *Hippocampus* 16: 664-672.
170. Paus T, Zijdenbos A, Worsley K, Collins DL, Blumenthal J, et al. (1999) Structural maturation of neural pathways in children and adolescents: in vivo study. *Science* 283: 1908-1911.
171. Toga AW, Thompson PM, Sowell ER (2006) Mapping brain maturation. *Trends in Neurosciences* 29: 148-159.
172. Bava S, Tapert S (2010) Adolescent Brain Development and the Risk for Alcohol and Other Drug Problems. *Neuropsychology Review* 20: 398-413.
173. Coleman Jr LG, He J, Lee J, Styner M, Crews FT (2011) Adolescent Binge Drinking Alters Adult Brain Neurotransmitter Gene Expression, Behavior, Brain Regional Volumes, and Neurochemistry in Mice. *Alcoholism: Clinical and Experimental Research* 35: 671-688.
174. Criado JR, Wills DN, Walker BM, Ehlers CL (2008) Electrophysiological Effects of Dizocilpine (MK-801) in Adult Rats Exposed to Ethanol During Adolescence. *Alcoholism: Clinical and Experimental Research* 32: 1752-1762.
175. Parada M, Corral M, Caamano-Isorna F, Mota N, Crego A, et al. (2011) Binge drinking and declarative memory in university students. *Alcohol Clin Exp Res* 35: 1475-1484.
176. Silvers JM, Tokunaga S, Mittleman G, Matthews DB (2003) Chronic Intermittent Injections of High-Dose Ethanol During Adolescence Produce Metabolic, Hypnotic, and Cognitive Tolerance in Rats. *Alcoholism: Clinical and Experimental Research* 27: 1606-1612.
177. Allen CD, Lee S, Koob GF, Rivier C (2011) Immediate and prolonged effects of alcohol exposure on the activity of the hypothalamic–pituitary–adrenal axis in adult and adolescent rats. *Brain, Behavior, and Immunity* 25, Supplement 1: S50-S60.
178. Ehlers CL, Criado JR, Wills DN, Liu W, Crews FT (2011) Periadolescent ethanol exposure reduces adult forebrain ChAT+IR neurons: correlation with behavioral pathology. *Neuroscience* 199: 333-345.
179. Schulteis G, Archer C, Tapert SF, Frank LR (2008) Intermittent binge alcohol exposure during the periadolescent period induces spatial working memory deficits in young adult rats. *Alcohol* 42: 459-467.

180. Autry AE, Monteggia LM (2012) Brain-derived neurotrophic factor and neuropsychiatric disorders. *Pharmacol Rev* 64: 238-258.
181. Martinowich K, Manji H, Lu B (2007) New insights into BDNF function in depression and anxiety. *Nat Neurosci* 10: 1089-1093.
182. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE (2005) Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62: 617-627.
183. Rosellini AJ, Rutter LA, Bourgeois ML, Emmert-Aronson BO, Brown TA (2013) The Relevance of Age of Onset to the Psychopathology of Social Phobia. *J Psychopathol Behav Assess* 35: 356-365.
184. Crews FT, Mdzinarishvili A, Kim D, He J, Nixon K (2006) Neurogenesis in adolescent brain is potently inhibited by ethanol. *Neuroscience* 137: 437-445.
185. Matthews DB, Morrow AL (2000) Effects of acute and chronic ethanol exposure on spatial cognitive processing and hippocampal function in the rat. *Hippocampus* 10: 122-130.
186. Oscar-Berman M, Bowirrat A (2005) Genetic influences in emotional dysfunction and alcoholism-related brain damage. *Neuropsychiatr Dis Treat* 1: 211-229.
187. Pfefferbaum A, Sullivan EV, Mathalon DH, Shear PK, Rosenbloom MJ, et al. (1995) Longitudinal Changes in Magnetic Resonance Imaging Brain Volumes in Abstinent and Relapsed Alcoholics. *Alcoholism: Clinical and Experimental Research* 19: 1177-1191.
188. Ros-Simó C, Moscoso-Castro M, Ruiz-Medina J, Ros J, Valverde O (2013) Memory impairment and hippocampus specific protein oxidation induced by ethanol intake and 3, 4-Methylenedioxymethamphetamine (MDMA) in mice. *Journal of Neurochemistry*: n/a-n/a.
189. Bartel DP (2009) MicroRNAs: Target Recognition and Regulatory Functions. *Cell* 136: 215-233.
190. Chang S, Johnston RJ, Jr., Hobert O (2003) A transcriptional regulatory cascade that controls left/right asymmetry in chemosensory neurons of *C. elegans*. *Genes Dev* 17: 2123-2137.
191. Strickland ER, Hook MA, Balaraman S, Huie JR, Grau JW, et al. (2011) MicroRNA dysregulation following spinal cord contusion: implications for neural plasticity and repair. *Neuroscience* 186: 146-160.
192. Uchida N (2010) MicroRNA-9 Controls a Migratory Mechanism in Human Neural Progenitor Cells. *Cell Stem Cell* 6: 294-296.

193. Sathyan P, Golden HB, Miranda RC (2007) Competing interactions between microRNAs determine neural progenitor survival and proliferation after ethanol exposure: evidence from an ex vivo model of the fetal cerebral cortical neuroepithelium. *J Neurosci* 27: 8546-8557.
194. Pappalardo-Carter DL, Balaraman S, Sathyan P, Carter ES, Chen WJ, et al. (2013) Suppression and Epigenetic Regulation of MiR-9 Contributes to Ethanol Teratology: Evidence from Zebrafish and Murine Fetal Neural Stem Cell Models. *Alcohol Clin Exp Res*.
195. Pietrzykowski AZ, Friesen RM, Martin GE, Puig SI, Nowak CL, et al. (2008) Posttranscriptional regulation of BK channel splice variant stability by miR-9 underlies neuroadaptation to alcohol. *Neuron* 59: 274-287.
196. Wang G, Wang X, Wang Y, Yang JY, Li L, et al. (2008) Identification of transcription factor and microRNA binding sites in responsible to fetal alcohol syndrome. *BMC Genomics* 9 Suppl 1: S19.
197. Wang X (2008) miRDB: a microRNA target prediction and functional annotation database with a wiki interface. *RNA* 14: 1012-1017.
198. Feliciano P, Andrade R, Bykhovskaia M (2013) Synapsin II and Rab3a cooperate in the regulation of epileptic and synaptic activity in the CA1 region of the hippocampus. *J Neurosci* 33: 18319-18330.
199. Medrihan L, Cesca F, Raimondi A, Lignani G, Baldelli P, et al. (2013) Synapsin II desynchronizes neurotransmitter release at inhibitory synapses by interacting with presynaptic calcium channels. *Nat Commun* 4: 1512.
200. Lauing K, Himes R, Rachwalski M, Strotman P, Callaci JJ (2008) Binge alcohol treatment of adolescent rats followed by alcohol abstinence is associated with site-specific differences in bone loss and incomplete recovery of bone mass and strength. *Alcohol* 42: 649-656.
201. Laviola G, Macri S, Adriani W, Morley Fletcher S (2002) [Psychobiological determinants of risk behavior in adolescence]. *Ann Ist Super Sanita* 38: 279-287.
202. Spear LP, Brake SC (1983) Periadolescence: age-dependent behavior and psychopharmacological responsivity in rats. *Dev Psychobiol* 16: 83-109.
203. Presutti C, Rosati J, Vincenti S, Nasi S (2006) Non coding RNA and brain. *BMC Neurosci* 7 Suppl 1: S5.
204. Jovicic A, Roshan R, Moiso N, Pradervand S, Moser R, et al. (2013) Comprehensive expression analyses of neural cell-type-specific miRNAs identify new determinants of the specification and maintenance of neuronal phenotypes. *J Neurosci* 33: 5127-5137.

205. Czerniawski J, Yoon T, Otto T (2009) Dissociating space and trace in dorsal and ventral hippocampus. *Hippocampus* 19: 20-32.
206. Inukai S, de Lencastre A, Turner M, Slack F (2012) Novel microRNAs differentially expressed during aging in the mouse brain. *PLoS ONE* 7: e40028.
207. Podolska A, Kaczkowski B, Kamp Busk P, Sokilde R, Litman T, et al. (2011) MicroRNA expression profiling of the porcine developing brain. *PLoS One* 6: e14494.
208. Hutvagner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, et al. (2001) A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* 293: 834-838.
209. Ketting RF, Fischer SE, Bernstein E, Sijen T, Hannon GJ, et al. (2001) Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes Dev* 15: 2654-2659.
210. Guo H, Ingolia NT, Weissman JS, Bartel DP (2010) Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature* 466: 835-840.
211. Selbach M, Schwanhauser B, Thierfelder N, Fang Z, Khanin R, et al. (2008) Widespread changes in protein synthesis induced by microRNAs. *Nature* 455: 58-63.
212. Allen CD, Rivier CL, Lee SY (2011) Adolescent alcohol exposure alters the central brain circuits known to regulate the stress response. *Neuroscience* 182: 162-168.
213. Przybycien-Szymanska MM, Mott NN, Pak TR (2011) Alcohol dysregulates corticotropin-releasing-hormone (CRH) promoter activity by interfering with the negative glucocorticoid response element (nGRE). *PLoS One* 6: e26647.
214. Mulligan MK, Dubose C, Yue J, Miles MF, Lu L, et al. (2013) Expression, covariation, and genetic regulation of miRNA Biogenesis genes in brain supports their role in addiction, psychiatric disorders, and disease. *Front Genet* 4: 126.
215. Valenzuela CF (1997) Alcohol and neurotransmitter interactions. *Alcohol Health Res World* 21: 144-148.
216. Lorenzo A, Auguet T, Vidal F, Broch M, Olona M, et al. (2006) Polymorphisms of alcohol-metabolizing enzymes and the risk for alcoholism and alcoholic liver disease in Caucasian Spanish women. *Drug Alcohol Depend* 84: 195-200.
217. Toth R, Pocsai Z, Fiatal S, Szeles G, Kardos L, et al. (2010) ADH1B\*2 allele is protective against alcoholism but not chronic liver disease in the Hungarian population. *Addiction* 105: 891-896.
218. Palmer RH, McGeary JE, Francazio S, Raphael BJ, Lander AD, et al. (2012) The genetics of alcohol dependence: advancing towards systems-based approaches. *Drug Alcohol Depend* 125: 179-191.

219. Lin P, Hartz SM, Wang JC, Agrawal A, Zhang TX, et al. (2012) Copy number variations in 6q14.1 and 5q13.2 are associated with alcohol dependence. *Alcohol Clin Exp Res* 36: 1512-1518.
220. Schuckit MA, Tapert S, Matthews SC, Paulus MP, Tolentino NJ, et al. (2012) fMRI differences between subjects with low and high responses to alcohol during a stop signal task. *Alcohol Clin Exp Res* 36: 130-140.
221. Crabbe JC, Phillips TJ, Belknap JK (2010) The complexity of alcohol drinking: studies in rodent genetic models. *Behav Genet* 40: 737-750.
222. Mastronardi C, Smiley GG, Raber J, Kusakabe T, Kawaguchi A, et al. (2006) Deletion of the *Ttf1* gene in differentiated neurons disrupts female reproduction without impairing basal ganglia function. *J Neurosci* 26: 13167-13179.
223. Carthew RW, Sontheimer EJ (2009) Origins and Mechanisms of miRNAs and siRNAs. *Cell* 136: 642-655.
224. Pak T, Rao Y, Prins S, Mott N (2013) An emerging role for microRNAs in sexually dimorphic neurobiological systems. *Pflügers Archiv - European Journal of Physiology* 465: 655-667.
225. Medina KL, McQueeney T, Nagel BJ, Hanson KL, Schweinsburg AD, et al. (2008) Prefrontal cortex volumes in adolescents with alcohol use disorders: unique gender effects. *Alcohol Clin Exp Res* 32: 386-394.
226. Squeglia LM, Spadoni AD, Infante MA, Myers MG, Tapert SF (2009) Initiating moderate to heavy alcohol use predicts changes in neuropsychological functioning for adolescent girls and boys. *Psychol Addict Behav* 23: 715-722.
227. Squeglia LM, Sorg SF, Schweinsburg AD, Wetherill RR, Pulido C, et al. (2012) Binge drinking differentially affects adolescent male and female brain morphometry. *Psychopharmacology (Berl)* 220: 529-539.
228. Squeglia LM, Schweinsburg AD, Pulido C, Tapert SF (2011) Adolescent binge drinking linked to abnormal spatial working memory brain activation: differential gender effects. *Alcohol Clin Exp Res* 35: 1831-1841.
229. Mann K, Batra A, Gunthner A, Schroth G (1992) Do women develop alcoholic brain damage more readily than men? *Alcohol Clin Exp Res* 16: 1052-1056.
230. Schweinsburg BC, Alhassoon OM, Taylor MJ, Gonzalez R, Videen JS, et al. (2003) Effects of alcoholism and gender on brain metabolism. *Am J Psychiatry* 160: 1180-1183.
231. O'Malley PM, Johnston LD (2002) Epidemiology of alcohol and other drug use among American college students. *J Stud Alcohol Suppl*: 23-39.

232. Faden VB (2006) Trends in initiation of alcohol use in the United States 1975 to 2003. *Alcohol Clin Exp Res* 30: 1011-1022.
233. Varlinskaya EI, Vetter-O'Hagen CS, Spear LP (2013) Puberty and gonadal hormones: role in adolescent-typical behavioral alterations. *Horm Behav* 64: 343-349.
234. Vetter CS, Doremus-Fitzwater TL, Spear LP (2007) Time course of elevated ethanol intake in adolescent relative to adult rats under continuous, voluntary-access conditions. *Alcohol Clin Exp Res* 31: 1159-1168.
235. Vetter-O'Hagen CS, Spear LP (2012) The effects of gonadectomy on sex- and age-typical responses to novelty and ethanol-induced social inhibition in adult male and female Sprague-Dawley rats. *Behav Brain Res* 227: 224-232.
236. Vetter-O'Hagen CS, Spear LP (2011) The effects of gonadectomy on age- and sex-typical patterns of ethanol consumption in Sprague-Dawley rats. *Alcohol Clin Exp Res* 35: 2039-2049.
237. Sherrill LK, Berthold C, Koss WA, Juraska JM, Gulley JM (2011) Sex differences in the effects of ethanol pre-exposure during adolescence on ethanol-induced conditioned taste aversion in adult rats. *Behav Brain Res* 225: 104-109.
238. Cheng LC, Pastrana E, Tavazoie M, Doetsch F (2009) miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. *Nat Neurosci* 12: 399-408.
239. Feng J, Sun G, Yan J, Noltner K, Li W, et al. (2009) Evidence for X-chromosomal schizophrenia associated with microRNA alterations. *PLoS One* 4: e6121.
240. Romeo RD, Waters EM, McEwen BS (2004) Steroid-induced hippocampal synaptic plasticity: sex differences and similarities. *Neuron Glia Biol* 1: 219-229.
241. Tal TL, Franzosa JA, Tilton SC, Philbrick KA, Iwaniec UT, et al. (2012) MicroRNAs control neurobehavioral development and function in zebrafish. *FASEB J* 26: 1452-1461.
242. Gorini G, Nunez YO, Mayfield RD (2013) Integration of miRNA and protein profiling reveals coordinated neuroadaptations in the alcohol-dependent mouse brain. *PLoS One* 8: e82565.
243. Guo Y, Chen Y, Carreon S, Qiang M (2012) Chronic Intermittent Ethanol Exposure and Its Removal Induce a Different miRNA Expression Pattern in Primary Cortical Neuronal Cultures. *Alcoholism: Clinical and Experimental Research* 36: 1058-1066.
244. Yadav S, Pandey A, Shukla A, Talwelkar SS, Kumar A, et al. (2011) miR-497 and miR-302b regulate ethanol-induced neuronal cell death through BCL2 protein and cyclin D2. *J Biol Chem* 286: 37347-37357.

245. Wang LL, Zhang Z, Li Q, Yang R, Pei X, et al. (2008) Ethanol exposure induces differential microRNA and target gene expression and teratogenic effects which can be suppressed by folic acid supplementation. *Human Reproduction* 24: 562-579.
246. Mantha K, Laufer BI, Singh SM (2014) Molecular changes during neurodevelopment following second-trimester binge ethanol exposure in a mouse model of fetal alcohol spectrum disorder: from immediate effects to long-term adaptation. *Dev Neurosci* 36: 29-43.
247. Prins SA, Przybycien-Szymanska MM, Rao YS, Pak TR (2014) Long-term effects of peripubertal binge EtOH exposure on hippocampal microRNA expression in the rat. *PLoS One* 9: e83166.
248. Sangiao-Alvarellos S, Manfredi-Lozano M, Ruiz-Pino F, Navarro VM, Sanchez-Garrido MA, et al. (2013) Changes in hypothalamic expression of the Lin28/let-7 system and related microRNAs during postnatal maturation and after experimental manipulations of puberty. *Endocrinology* 154: 942-955.
249. Bettgowda A, Lee KB, Smith GW (2008) Cytoplasmic and nuclear determinants of the maternal-to-embryonic transition. *Reprod Fertil Dev* 20: 45-53.
250. Faggad A, Budczies J, Tchernitsa O, Darb-Esfahani S, Sehouli J, et al. (2010) Prognostic significance of Dicer expression in ovarian cancer-link to global microRNA changes and oestrogen receptor expression. *J Pathol* 220: 382-391.
251. Cochrane DR, Cittelly DM, Howe EN, Spoelstra NS, McKinsey EL, et al. (2010) MicroRNAs link estrogen receptor alpha status and Dicer levels in breast cancer. *Horm Cancer* 1: 306-319.
252. Cheng C, Fu X, Alves P, Gerstein M (2009) mRNA expression profiles show differential regulatory effects of microRNAs between estrogen receptor-positive and estrogen receptor-negative breast cancer. *Genome Biol* 10: R90.
253. Bolton MM, Lo DC, Sherwood NT (2000) Long-term regulation of excitatory and inhibitory synaptic transmission in hippocampal cultures by brain-derived neurotrophic factor. *Prog Brain Res* 128: 203-218.
254. Leal G, Comprido D, Duarte CB (2014) BDNF-induced local protein synthesis and synaptic plasticity. *Neuropharmacology* 76 Pt C: 639-656.
255. Courtney KE, Polich J (2009) Binge drinking in young adults: Data, definitions, and determinants. *Psychol Bull* 135: 142-156.
256. Jacobus J, Tapert SF (2013) Neurotoxic effects of alcohol in adolescence. *Annu Rev Clin Psychol* 9: 703-721.



257. Caldwell LC, Schweinsburg AD, Nagel BJ, Barlett VC, Brown SA, et al. (2005) Gender and adolescent alcohol use disorders on BOLD (blood oxygen level dependent) response to spatial working memory. *Alcohol Alcohol* 40: 194-200.
258. Bast T, Feldon J (2003) Hippocampal modulation of sensorimotor processes. *Prog Neurobiol* 70: 319-345.
259. Romeo RD (2005) Neuroendocrine and behavioral development during puberty: a tale of two axes. *Vitam Horm* 71: 1-25.
260. Vetter-O'Hagen CS, Sanders KW, Spear LP (2011) Evidence for suppressant effects of testosterone on sex-typical ethanol intake in male Sprague-Dawley rats. *Behav Brain Res* 224: 403-407.
261. Vetter-O'Hagen CS, Spear LP (2012) Hormonal and physical markers of puberty and their relationship to adolescent-typical novelty-directed behavior. *Dev Psychobiol* 54: 523-535.
262. Pandey DP, Picard D (2009) miR-22 inhibits estrogen signaling by directly targeting the estrogen receptor alpha mRNA. *Mol Cell Biol* 29: 3783-3790.
263. Pandey DP, Picard D (2010) Multidirectional interplay between nuclear receptors and microRNAs. *Curr Opin Pharmacol* 10: 637-642.
264. Castellano L, Giamas G, Jacob J, Coombes RC, Lucchesi W, et al. (2009) The estrogen receptor-alpha-induced microRNA signature regulates itself and its transcriptional response. *Proc Natl Acad Sci U S A* 106: 15732-15737.
265. Liu WH, Yeh SH, Lu CC, Yu SL, Chen HY, et al. (2009) MicroRNA-18a prevents estrogen receptor-alpha expression, promoting proliferation of hepatocellular carcinoma cells. *Gastroenterology* 136: 683-693.
266. Di Leva G, Gasparini P, Piovan C, Ngankeu A, Garofalo M, et al. (2010) MicroRNA cluster 221-222 and estrogen receptor alpha interactions in breast cancer. *J Natl Cancer Inst* 102: 706-721.
267. Fugger HN, Cunningham SG, Rissman EF, Foster TC (1998) Sex differences in the activational effect of ERalpha on spatial learning. *Horm Behav* 34: 163-170.
268. Gibbs RB (2005) Testosterone and estradiol produce different effects on cognitive performance in male rats. *Horm Behav* 48: 268-277.
269. Raber J (2008) AR, apoE, and cognitive function. *Horm Behav* 53: 706-715.
270. Leranath C, Petnehazy O, MacLusky NJ (2003) Gonadal hormones affect spine synaptic density in the CA1 hippocampal subfield of male rats. *J Neurosci* 23: 1588-1592.

271. Romeo RD, Staub D, Jasnow AM, Karatsoreos IN, Thornton JE, et al. (2005) Dihydrotestosterone increases hippocampal N-methyl-D-aspartate binding but does not affect choline acetyltransferase cell number in the forebrain or choline transporter levels in the CA1 region of adult male rats. *Endocrinology* 146: 2091-2097.
272. Hajszan T, MacLusky NJ, Leranth C (2008) Role of androgens and the androgen receptor in remodeling of spine synapses in limbic brain areas. *Horm Behav* 53: 638-646.
273. Spencer JL, Waters EM, Milner TA, McEwen BS (2008) Estrous cycle regulates activation of hippocampal Akt, LIM kinase, and neurotrophin receptors in C57BL/6 mice. *Neuroscience* 155: 1106-1119.
274. McEwen BS, Milner TA (2007) Hippocampal formation: shedding light on the influence of sex and stress on the brain. *Brain Res Rev* 55: 343-355.
275. Daniel JM (2006) Effects of oestrogen on cognition: what have we learned from basic research? *J Neuroendocrinol* 18: 787-795.
276. Luine VN (2008) Sex steroids and cognitive function. *J Neuroendocrinol* 20: 866-872.
277. Euling SY, Selevan SG, Pescovitz OH, Skakkebaek NE (2008) Role of environmental factors in the timing of puberty. *Pediatrics* 121 Suppl 3: S167-171.
278. Stroud LR, Papandonatos GD, Williamson DE, Dahl RE (2004) Sex differences in the effects of pubertal development on responses to a corticotropin-releasing hormone challenge: the Pittsburgh psychobiologic studies. *Ann N Y Acad Sci* 1021: 348-351.
279. Evuarherhe O, Leggett J, Waite E, Kershaw Y, Lightman S (2009) Reversal of the hypothalamo-pituitary-adrenal response to oestrogens around puberty. *J Endocrinol* 202: 279-285.
280. Cummins JM, He Y, Leary RJ, Pagliarini R, Diaz LA, Jr., et al. (2006) The colorectal microRNAome. *Proc Natl Acad Sci U S A* 103: 3687-3692.
281. Yang JS, Maurin T, Robine N, Rasmussen KD, Jeffrey KL, et al. (2010) Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc Natl Acad Sci U S A* 107: 15163-15168.
282. Kuehbacher A, Urbich C, Zeiher AM, Dimmeler S (2007) Role of Dicer and Drosha for endothelial microRNA expression and angiogenesis. *Circ Res* 101: 59-68.
283. Fukagawa T, Nogami M, Yoshikawa M, Ikeno M, Okazaki T, et al. (2004) Dicer is essential for formation of the heterochromatin structure in vertebrate cells. *Nat Cell Biol* 6: 784-791.

284. Wienholds E, Koudijs MJ, van Eeden FJ, Cuppen E, Plasterk RH (2003) The microRNA-producing enzyme Dicer1 is essential for zebrafish development. *Nat Genet* 35: 217-218.
285. Zovoilis A, Agbemenyah HY, Agis-Balboa RC, Stilling RM, Edbauer D, et al. (2011) microRNA-34c is a novel target to treat dementias. *EMBO J* 30: 4299-4308.
286. Ketelslegers JM, Hetzel WD, Sherins RJ, Catt KJ (1978) Developmental changes in testicular gonadotropin receptors: plasma gonadotropins and plasma testosterone in the rat. *Endocrinology* 103: 212-222.
287. Sodersten P, Damassa DA, Smith ER (1977) Sexual behavior in developing male rats. *Horm Behav* 8: 320-341.
288. Sodersten P (1975) Receptive behavior in developing female rats. *Horm Behav* 6: 307-317.
289. Gorski RA (1978) Sexual differentiation of the brain. *Hosp Pract* 13: 55-62.
290. Calaresu FR, Henry JL (1971) Sex difference in the number of the sympathetic neurons in the spinal cord of the cat. *Science* 173: 343-344.
291. Swaab DF, Hofman MA (1984) Sexual differentiation of the human brain. A historical perspective. *Prog Brain Res* 61: 361-374.
292. Pakkenberg B, Gundersen HJ (1997) Neocortical neuron number in humans: effect of sex and age. *J Comp Neurol* 384: 312-320.
293. Balthazart J, Ball GF (1998) New insights into the regulation and function of brain estrogen synthase (aromatase). *Trends Neurosci* 21: 243-249.
294. Bao AM, Swaab DF (2011) Sexual differentiation of the human brain: relation to gender identity, sexual orientation and neuropsychiatric disorders. *Front Neuroendocrinol* 32: 214-226.
295. Bloch GJ, Eckersell C, Mills R (1993) Distribution of galanin-immunoreactive cells within sexually dimorphic components of the medial preoptic area of the male and female rat. *Brain Res* 620: 259-268.
296. Simerly RB, Swanson LW, Gorski RA (1985) The distribution of monoaminergic cells and fibers in a periventricular preoptic nucleus involved in the control of gonadotropin release: immunohistochemical evidence for a dopaminergic sexual dimorphism. *Brain Res* 330: 55-64.
297. Gu GB, Simerly RB (1997) Projections of the sexually dimorphic anteroventral periventricular nucleus in the female rat. *J Comp Neurol* 384: 142-164.

298. Hutton LA, Gu G, Simerly RB (1998) Development of a sexually dimorphic projection from the bed nuclei of the stria terminalis to the anteroventral periventricular nucleus in the rat. *J Neurosci* 18: 3003-3013.
299. Swaab DF, Chung WC, Kruijver FP, Hofman MA, Ishunina TA (2001) Structural and functional sex differences in the human hypothalamus. *Horm Behav* 40: 93-98.
300. Lee SJ, Duong JK, Stern DF (2004) A Ddc2-Rad53 fusion protein can bypass the requirements for RAD9 and MRC1 in Rad53 activation. *Mol Biol Cell* 15: 5443-5455.
301. Murphy DD, Segal M (1996) Regulation of dendritic spine density in cultured rat hippocampal neurons by steroid hormones. *J Neurosci* 16: 4059-4068.
302. Akama KT, McEwen BS (2003) Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway. *J Neurosci* 23: 2333-2339.
303. Choi JM, Romeo RD, Brake WG, Bethea CL, Rosenwaks Z, et al. (2003) Estradiol increases pre- and post-synaptic proteins in the CA1 region of the hippocampus in female rhesus macaques (*Macaca mulatta*). *Endocrinology* 144: 4734-4738.
304. Zhao L, Chen S, Ming Wang J, Brinton RD (2005) 17beta-estradiol induces Ca<sup>2+</sup> influx, dendritic and nuclear Ca<sup>2+</sup> rise and subsequent cyclic AMP response element-binding protein activation in hippocampal neurons: a potential initiation mechanism for estrogen neurotrophism. *Neuroscience* 132: 299-311.
305. Wayman GA, Davare M, Ando H, Fortin D, Varlamova O, et al. (2008) An activity-regulated microRNA controls dendritic plasticity by down-regulating p250GAP. *Proc Natl Acad Sci U S A* 105: 9093-9098.
306. Cohen A, Shmoish M, Levi L, Cheruti U, Levavi-Sivan B, et al. (2008) Alterations in micro-ribonucleic acid expression profiles reveal a novel pathway for estrogen regulation. *Endocrinology* 149: 1687-1696.
307. Kovalchuk O, Tryndyak VP, Montgomery B, Boyko A, Kutanzi K, et al. (2007) Estrogen-induced rat breast carcinogenesis is characterized by alterations in DNA methylation, histone modifications and aberrant microRNA expression. *Cell Cycle* 6: 2010-2018.
308. Dai R, Phillips RA, Zhang Y, Khan D, Crasta O, et al. (2008) Suppression of LPS-induced Interferon-gamma and nitric oxide in splenic lymphocytes by select estrogen-regulated microRNAs: a novel mechanism of immune modulation. *Blood* 112: 4591-4597.
309. Kim J, Inoue K, Ishii J, Vanti WB, Voronov SV, et al. (2007) A MicroRNA Feedback Circuit in Midbrain Dopamine Neurons. *Science* 317: 1220-1224.
310. Klinge CM (2012) miRNAs and estrogen action. *Trends Endocrinol Metab* 23: 223-233.

311. Koturbash I, Zemp F, Kolb B, Kovalchuk O (2011) Sex-specific radiation-induced microRNAome responses in the hippocampus, cerebellum and frontal cortex in a mouse model. *Mutat Res* 722: 114-118.
312. Morrison KE, Rodgers AB, Morgan CP, Bale TL (2014) Epigenetic mechanisms in pubertal brain maturation. *Neuroscience* 264: 17-24.
313. Paus T (2013) How environment and genes shape the adolescent brain. *Horm Behav* 64: 195-202.
314. Delaloy C, Liu L, Lee JA, Su H, Shen F, et al. (2010) MicroRNA-9 coordinates proliferation and migration of human embryonic stem cell-derived neural progenitors. *Cell Stem Cell* 6: 323-335.
315. Krichevsky AM, Sonntag KC, Isacson O, Kosik KS (2006) Specific microRNAs modulate embryonic stem cell-derived neurogenesis. *Stem Cells* 24: 857-864.
316. Yoo AS, Sun AX, Li L, Shcheglovitov A, Portmann T, et al. (2011) MicroRNA-mediated conversion of human fibroblasts to neurons. *Nature* 476: 228-231.
317. Ambasudhan R, Talantova M, Coleman R, Yuan X, Zhu S, et al. (2011) Direct reprogramming of adult human fibroblasts to functional neurons under defined conditions. *Cell Stem Cell* 9: 113-118.
318. Shibata M, Nakao H, Kiyonari H, Abe T, Aizawa S (2011) MicroRNA-9 regulates neurogenesis in mouse telencephalon by targeting multiple transcription factors. *J Neurosci* 31: 3407-3422.
319. Zhao C, Sun G, Li S, Shi Y (2009) A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol* 16: 365-371.
320. Dajas-Bailador F, Bonev B, Garcez P, Stanley P, Guillemot F, et al. (2012) microRNA-9 regulates axon extension and branching by targeting Map1b in mouse cortical neurons. *Nat Neurosci*.
321. Kennedy GC (1969) The relation between the central control of appetite, growth and sexual maturation. *Guys Hosp Rep* 118: 315-327.
322. Spear LP (2004) Adolescent brain development and animal models. *Ann N Y Acad Sci* 1021: 23-26.
323. Johnston RJ, Hobert O (2003) A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. *Nature* 426: 845-849.
324. Li Y, Wang F, Lee JA, Gao FB (2006) MicroRNA-9a ensures the precise specification of sensory organ precursors in *Drosophila*. *Genes Dev* 20: 2793-2805.

325. Krichevsky AM, King KS, Donahue CP, Khrapko K, Kosik KS (2003) A microRNA array reveals extensive regulation of microRNAs during brain development. *RNA* 9: 1274-1281.
326. Woltering JM, Durston AJ (2008) MiR-10 represses HoxB1a and HoxB3a in zebrafish. *PLoS ONE* 3: e1396.
327. Bian S, Xu TL, Sun T (2013) Tuning the cell fate of neurons and glia by microRNAs. *Curr Opin Neurobiol* 23: 928-934.
328. Wen S, Li H, Liu J (2009) Dynamic signaling for neural stem cell fate determination. *Cell Adh Migr* 3: 107-117.
329. Dugas JC, Cuellar TL, Scholze A, Ason B, Ibrahim A, et al. (2010) Dicer1 and miR-219 Are required for normal oligodendrocyte differentiation and myelination. *Neuron* 65: 597-611.
330. Zhao X, He X, Han X, Yu Y, Ye F, et al. (2010) MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* 65: 612-626.
331. McGee CL, Riley EP (2006) Brain imaging and fetal alcohol spectrum disorders. *Ann Ist Super Sanita* 42: 46-52.
332. Olney JW (2004) Fetal alcohol syndrome at the cellular level. *Addict Biol* 9: 137-149; discussion 151.
333. Riley EP, McGee CL (2005) Fetal alcohol spectrum disorders: an overview with emphasis on changes in brain and behavior. *Exp Biol Med (Maywood)* 230: 357-365.
334. Kortum F, Das S, Flindt M, Morris-Rosendahl DJ, Stefanova I, et al. (2011) The core FOXG1 syndrome phenotype consists of postnatal microcephaly, severe mental retardation, absent language, dyskinesia, and corpus callosum hypogenesis. *J Med Genet* 48: 396-406.
335. Spear NE, Molina JC (2005) Fetal or infantile exposure to ethanol promotes ethanol ingestion in adolescence and adulthood: a theoretical review. *Alcohol Clin Exp Res* 29: 909-929.
336. Saal D, Dong Y, Bonci A, Malenka RC (2003) Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron* 37: 577-582.
337. White AM, Best PJ (2000) Effects of ethanol on hippocampal place-cell and interneuron activity. *Brain Res* 876: 154-165.
338. Nona CN, Guirguis S, Nobrega JN (2013) Susceptibility to ethanol sensitization is differentially associated with changes in pCREB, trkB and BDNF mRNA expression in the mouse brain. *Behav Brain Res* 242: 25-33.

339. Sapolsky RM (1992) Do glucocorticoid concentrations rise with age in the rat? *Neurobiol Aging* 13: 171-174.
340. McEwen BS (2000) The neurobiology of stress: from serendipity to clinical relevance. *Brain Res* 886: 172-189.
341. Kim JJ, Diamond DM (2002) The stressed hippocampus, synaptic plasticity and lost memories. *Nat Rev Neurosci* 3: 453-462.
342. McEwen BS, Sapolsky RM (1995) Stress and cognitive function. *Curr Opin Neurobiol* 5: 205-216.
343. Reul JM, de Kloet ER (1985) Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* 117: 2505-2511.
344. Dahl RE, Gunnar MR (2009) Heightened stress responsiveness and emotional reactivity during pubertal maturation: implications for psychopathology. *Dev Psychopathol* 21: 1-6.
345. McCormick CM, Mathews IZ (2007) HPA function in adolescence: role of sex hormones in its regulation and the enduring consequences of exposure to stressors. *Pharmacol Biochem Behav* 86: 220-233.
346. McCormick CM, Mathews IZ (2010) Adolescent development, hypothalamic-pituitary-adrenal function, and programming of adult learning and memory. *Prog Neuropsychopharmacol Biol Psychiatry* 34: 756-765.
347. Romeo RD (2010) Pubertal maturation and programming of hypothalamic-pituitary-adrenal reactivity. *Front Neuroendocrinol* 31: 232-240.
348. Foy MR, Stanton ME, Levine S, Thompson RF (1987) Behavioral stress impairs long-term potentiation in rodent hippocampus. *Behav Neural Biol* 48: 138-149.
349. McEwen BS, Magarinos AM (1997) Stress effects on morphology and function of the hippocampus. *Ann N Y Acad Sci* 821: 271-284.
350. Holderbach R, Clark K, Moreau JL, Bischofberger J, Normann C (2007) Enhanced long-term synaptic depression in an animal model of depression. *Biol Psychiatry* 62: 92-100.
351. Joels M, Karst H, Alfarez D, Heine VM, Qin Y, et al. (2004) Effects of chronic stress on structure and cell function in rat hippocampus and hypothalamus. *Stress* 7: 221-231.
352. Bleich S, Sperling W, Wiltfang J, Maler JM, Kornhuber J (2003) [Excitatory neurotransmission in alcoholism]. *Fortschr Neurol Psychiatr* 71 Suppl 1: S36-44.

353. Agartz I, Momenan R, Rawlings RR, Kerich MJ, Hommer DW (1999) Hippocampal volume in patients with alcohol dependence. *Arch Gen Psychiatry* 56: 356-363.
354. Wang J, Du H, Jiang L, Ma X, de Graaf RA, et al. (2013) Oxidation of ethanol in the rat brain and effects associated with chronic ethanol exposure. *Proc Natl Acad Sci U S A* 110: 14444-14449.
355. Obernier JA, White AM, Swartzwelder HS, Crews FT (2002) Cognitive deficits and CNS damage after a 4-day binge ethanol exposure in rats. *Pharmacol Biochem Behav* 72: 521-532.
356. Adachi M, Ishii H (2002) Role of mitochondria in alcoholic liver injury. *Free Radic Biol Med* 32: 487-491.
357. Higuchi H, Adachi M, Miura S, Gores GJ, Ishii H (2001) The mitochondrial permeability transition contributes to acute ethanol-induced apoptosis in rat hepatocytes. *Hepatology* 34: 320-328.
358. Manzo-Avalos S, Saavedra-Molina A (2010) Cellular and mitochondrial effects of alcohol consumption. *Int J Environ Res Public Health* 7: 4281-4304.
359. Reddy VD, Padmavathi P, Kavitha G, Saradamma B, Varadacharyulu N (2013) Alcohol-induced oxidative/nitrosative stress alters brain mitochondrial membrane properties. *Mol Cell Biochem* 375: 39-47.
360. Kren BT, Wong PY, Sarver A, Zhang X, Zeng Y, et al. (2009) MicroRNAs identified in highly purified liver-derived mitochondria may play a role in apoptosis. *RNA Biol* 6: 65-72.
361. Sripada L, Tomar D, Prajapati P, Singh R, Singh AK (2012) Systematic analysis of small RNAs associated with human mitochondria by deep sequencing: detailed analysis of mitochondrial associated miRNA. *PLoS One* 7: e44873.
362. Gagliardi D, Stepien PP, Temperley RJ, Lightowlers RN, Chrzanowska-Lightowlers ZM (2004) Messenger RNA stability in mitochondria: different means to an end. *Trends Genet* 20: 260-267.
363. Orom UA, Nielsen FC, Lund AH (2008) MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation. *Mol Cell* 30: 460-471.
364. Place RF, Li LC, Pookot D, Noonan EJ, Dahiya R (2008) MicroRNA-373 induces expression of genes with complementary promoter sequences. *Proc Natl Acad Sci U S A* 105: 1608-1613.
365. Gonzalez S, Pisano DG, Serrano M (2008) Mechanistic principles of chromatin remodeling guided by siRNAs and miRNAs. *Cell Cycle* 7: 2601-2608.



366. Zhang Y, Jia Y, Zheng R, Guo Y, Wang Y, et al. (2010) Plasma microRNA-122 as a biomarker for viral-, alcohol-, and chemical-related hepatic diseases. *Clin Chem* 56: 1830-1838.
367. Hunter MP, Ismail N, Zhang X, Aguda BD, Lee EJ, et al. (2008) Detection of microRNA expression in human peripheral blood microvesicles. *PLoS One* 3: e3694.
368. Kruman, II, Henderson GI, Bergeson SE (2012) DNA damage and neurotoxicity of chronic alcohol abuse. *Exp Biol Med (Maywood)* 237: 740-747.
369. Sullivan EV, Marsh L, Mathalon DH, Lim KO, Pfefferbaum A (1995) Anterior hippocampal volume deficits in nonamnesic, aging chronic alcoholics. *Alcohol Clin Exp Res* 19: 110-122.
370. Beresford TP, Arciniegas DB, Alfors J, Clapp L, Martin B, et al. (2006) Hippocampus volume loss due to chronic heavy drinking. *Alcohol Clin Exp Res* 30: 1866-1870.
371. DeWit DJ, Adlaf EM, Offord DR, Ogborne AC (2000) Age at first alcohol use: a risk factor for the development of alcohol disorders. *Am J Psychiatry* 157: 745-750.
372. Torley KJ, da Silveira JC, Smith P, Anthony RV, Veeramachaneni DN, et al. (2011) Expression of miRNAs in ovine fetal gonads: potential role in gonadal differentiation. *Reprod Biol Endocrinol* 9: 2.
373. Nassa G, Tarallo R, Giurato G, De Filippo MR, Ravo M, et al. (2014) Post-transcriptional regulation of human breast cancer cell proteome by unliganded estrogen receptor beta via microRNAs. *Mol Cell Proteomics* 13: 1076-1090.
374. Solum DT, Handa RJ (2002) Estrogen regulates the development of brain-derived neurotrophic factor mRNA and protein in the rat hippocampus. *J Neurosci* 22: 2650-2659.
375. Ivanova T, Beyer C (2000) Ontogenetic expression and sex differences of aromatase and estrogen receptor-alpha/beta mRNA in the mouse hippocampus. *Cell Tissue Res* 300: 231-237.
376. Singh M, Meyer EM, Simpkins JW (1995) The effect of ovariectomy and estradiol replacement on brain-derived neurotrophic factor messenger ribonucleic acid expression in cortical and hippocampal brain regions of female Sprague-Dawley rats. *Endocrinology* 136: 2320-2324.
377. Gibbs RB (1998) Levels of trkA and BDNF mRNA, but not NGF mRNA, fluctuate across the estrous cycle and increase in response to acute hormone replacement. *Brain Res* 810: 294.

378. Scharfman HE, MacLusky NJ (2006) Estrogen and brain-derived neurotrophic factor (BDNF) in hippocampus: complexity of steroid hormone-growth factor interactions in the adult CNS. *Front Neuroendocrinol* 27: 415-435.
379. Gibbs RB, Burke AM, Johnson DA (1998) Estrogen replacement attenuates effects of scopolamine and lorazepam on memory acquisition and retention. *Horm Behav* 34: 112-125.
380. Fester L, Prange-Kiel J, Jarry H, Rune GM (2011) Estrogen synthesis in the hippocampus. *Cell Tissue Res* 345: 285-294.
381. Skucas VA, Duffy AM, Harte-Hargrove LC, Magagna-Poveda A, Radman T, et al. (2013) Testosterone depletion in adult male rats increases mossy fiber transmission, LTP, and sprouting in area CA3 of hippocampus. *J Neurosci* 33: 2338-2355.
382. Squillacioti C, De Luca A, Paino G, Mirabella N (2008) Effects of castration on the immunoreactivity to NGF, BDNF and their receptors in the pelvic ganglia of the male rat. *Eur J Histochem* 52: 101-106.
383. Primus RJ, Kellogg CK (1989) Pubertal-related changes influence the development of environment-related social interaction in the male rat. *Dev Psychobiol* 22: 633-643.
384. Primus RJ, Kellogg CK (1990) Gonadal hormones during puberty organize environment-related social interaction in the male rat. *Horm Behav* 24: 311-323.
385. Schulz KM, Zehr JL, Salas-Ramirez KY, Sisk CL (2009) Testosterone programs adult social behavior before and during, but not after, adolescence. *Endocrinology* 150: 3690-3698.
386. Hier DB, Crowley WF, Jr. (1982) Spatial ability in androgen-deficient men. *N Engl J Med* 306: 1202-1205.
387. Mueller SC, Temple V, Oh E, VanRyzin C, Williams A, et al. (2008) Early androgen exposure modulates spatial cognition in congenital adrenal hyperplasia (CAH). *Psychoneuroendocrinology* 33: 973-980.
388. Hebbard PC, King RR, Malsbury CW, Harley CW (2003) Two organizational effects of pubertal testosterone in male rats: transient social memory and a shift away from long-term potentiation following a tetanus in hippocampal CA1. *Exp Neurol* 182: 470-475.
389. Harley CW, Malsbury CW, Squires A, Brown RA (2000) Testosterone decreases CA1 plasticity in vivo in gonadectomized male rats. *Hippocampus* 10: 693-697.
390. White SA, Livingston FS, Mooney R (1999) Androgens modulate NMDA receptor-mediated EPSCs in the zebra finch song system. *J Neurophysiol* 82: 2221-2234.

391. Cicero TJ, Adams ML, O'Connor L, Nock B, Meyer ER, et al. (1990) Influence of chronic alcohol administration on representative indices of puberty and sexual maturation in male rats and the development of their progeny. *J Pharmacol Exp Ther* 255: 707-715.
392. Dees WL, Skelley CW (1990) Effects of ethanol during the onset of female puberty. *Neuroendocrinology* 51: 64-69.
393. Dees WL, Skelley CW, Hiney JK, Johnston CA (1990) Actions of ethanol on hypothalamic and pituitary hormones in prepubertal female rats. *Alcohol* 7: 21-25.
394. Almeida OF, Shoaib M, Deicke J, Fischer D, Darwish MH, et al. (1998) Gender differences in ethanol preference and ingestion in rats. The role of the gonadal steroid environment. *J Clin Invest* 101: 2677-2685.
395. Morales M, Spear LP (2013) Differences in sensitivity to ethanol-induced conditioned taste aversions emerge after pre- or post-pubertal gonadectomy in male and female rats. *Behav Brain Res* 240: 69-75.
396. Apter SJ, Eriksson CJ (2003) The effect of alcohol on testosterone concentrations in alcohol-preferring and non-preferring rat lines. *Alcohol Clin Exp Res* 27: 1190-1193.
397. Eriksson CJ, Kaprio J, Pulkkinen L, Rose RJ (2005) Testosterone and alcohol use among adolescent male twins: testing between-family associations in within-family comparisons. *Behav Genet* 35: 359-368.
398. Martin CA, Mainous AG, 3rd, Curry T, Martin D (1999) Alcohol use in adolescent females: correlates with estradiol and testosterone. *Am J Addict* 8: 9-14.
399. Forger NG, Morin LP (1982) Reproductive state modulates ethanol intake in rats: effects of ovariectomy, ethanol concentration, estrous cycle and pregnancy. *Pharmacol Biochem Behav* 17: 323-331.
400. Roberts AJ, Smith AD, Weiss F, Rivier C, Koob GF (1998) Estrous cycle effects on operant responding for ethanol in female rats. *Alcohol Clin Exp Res* 22: 1564-1569.
401. Blanchard BA, Steindorf S, Wang S, Glick SD (1993) Sex differences in ethanol-induced dopamine release in nucleus accumbens and in ethanol consumption in rats. *Alcohol Clin Exp Res* 17: 968-973.
402. Woolley CS, McEwen BS (1993) Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *J Comp Neurol* 336: 293-306.
403. Bergemann N, Parzer P, Runnebaum B, Resch F, Mundt C (2007) Estrogen, menstrual cycle phases, and psychopathology in women suffering from schizophrenia. *Psychol Med* 37: 1427-1436.

404. Hendrick V, Altshuler LL, Burt VK (1996) Course of psychiatric disorders across the menstrual cycle. *Harv Rev Psychiatry* 4: 200-207.
405. Grigoriadis S, Kennedy SH (2002) Role of estrogen in the treatment of depression. *Am J Ther* 9: 503-509.
406. Grigoriadis S, Seeman MV (2002) The role of estrogen in schizophrenia: implications for schizophrenia practice guidelines for women. *Can J Psychiatry* 47: 437-442.
407. Fink G, Sumner BE, Rosie R, Grace O, Quinn JP (1996) Estrogen control of central neurotransmission: effect on mood, mental state, and memory. *Cell Mol Neurobiol* 16: 325-344.
408. Fink G, Sumner BE (1996) Oestrogen and mental state. *Nature* 383: 306.
409. Mora S, Dussaubat N, Diaz-Veliz G (1996) Effects of the estrous cycle and ovarian hormones on behavioral indices of anxiety in female rats. *Psychoneuroendocrinology* 21: 609-620.
410. Juraska JM, Sisk CL, DonCarlos LL (2013) Sexual differentiation of the adolescent rodent brain: hormonal influences and developmental mechanisms. *Horm Behav* 64: 203-210.
411. Lomniczi A, Loche A, Castellano JM, Ronnekleiv OK, Bosch M, et al. (2013) Epigenetic control of female puberty. *Nat Neurosci* 16: 281-289.
412. Pak TR, Chung WC, Hinds LR, Handa RJ (2007) Estrogen receptor-beta mediates dihydrotestosterone-induced stimulation of the arginine vasopressin promoter in neuronal cells. *Endocrinology* 148: 3371-3382.
413. Pak TR, Chung WC, Roberts JL, Handa RJ (2006) Ligand-independent effects of estrogen receptor beta on mouse gonadotropin-releasing hormone promoter activity. *Endocrinology* 147: 1924-1931.
414. Mott NN, Pak TR (2013) Estrogen signaling and the aging brain: context-dependent considerations for postmenopausal hormone therapy. *ISRN Endocrinol* 2013: 814690.
415. Brown SA, Tapert SF (2004) Adolescence and the trajectory of alcohol use: basic to clinical studies. *Ann N Y Acad Sci* 1021: 234-244.
416. Miller JW, Naimi TS, Brewer RD, Jones SE (2007) Binge drinking and associated health risk behaviors among high school students. *Pediatrics* 119: 76-85.
417. Hingson R, Heeren T, Winter M, Wechsler H (2005) Magnitude of alcohol-related mortality and morbidity among U.S. college students ages 18-24: changes from 1998 to 2001. *Annu Rev Public Health* 26: 259-279.

418. Chatterji P, Dave D, Kaestner R, Markowitz S (2004) Alcohol abuse and suicide attempts among youth. *Econ Hum Biol* 2: 159-180.
419. Stroynowska-Czerwinska A, Fiszler A, Krzyzosiak WJ (2014) The panorama of miRNA-mediated mechanisms in mammalian cells. *Cell Mol Life Sci* 71: 2253-2270.
420. Mundalil Vasu M, Anitha A, Thanseem I, Suzuki K, Yamada K, et al. (2014) Serum microRNA profiles in children with autism. *Mol Autism* 5: 40.
421. Godlewski J, Krichevsky AM, Johnson MD, Chiocca EA, Bronisz A (2014) Belonging to a network-microRNAs, extracellular vesicles, and the glioblastoma microenvironment. *Neuro Oncol*.
422. Grasso M, Piscopo P, Confaloni A, Denti MA (2014) Circulating miRNAs as biomarkers for neurodegenerative disorders. *Molecules* 19: 6891-6910.
423. Selvamani A, Williams MH, Miranda RC, Sohrabji F (2014) Circulating miRNA profiles provide a biomarker for severity of stroke outcomes associated with age and sex in a rat model. *Clin Sci (Lond)* 127: 77-89.
424. Sepramaniam S, Tan JR, Tan KS, DeSilva DA, Tavintharan S, et al. (2014) Circulating microRNAs as biomarkers of acute stroke. *Int J Mol Sci* 15: 1418-1432.
425. Ogata K, Sumida K, Miyata K, Kushida M, Kuwamura M, et al. (2014) Circulating miR-9\* and miR-384-5p as Potential Indicators for Trimethyltin-induced Neurotoxicity. *Toxicol Pathol*.
426. Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the  $2^{-\Delta\Delta CT}$  Method. *Methods* 25: 402-408.

## VITA

Sarah A. Prins was born in Riverside, CA on July 23rd, 1986 to Peter and Enid Prins as the youngest of six children. In high school she developed a curiosity for neuroscience and earned a governor's scholarship for high academic achievement. Upon graduating in 2004, she moved to Santa Cruz, CA to major in Neuroscience and Behavior at University of California, Santa Cruz (UCSC). While at UCSC, she engaged in academic research in the laboratory of Dr. Yi Zuo, studying the morphological dynamics of dendritic spines during learning and memory formation. In 2009, Sarah earned her Bachelor of Science degree and moved to Chicago, IL to join the graduate program in Biomedical Sciences at Loyola University Chicago, Stritch School of Medicine.

In 2010, Sarah joined Dr. Toni Pak's laboratory in the Department of Cell and Molecular Physiology to study the effects of adolescent binge drinking on the developing brain. During her training, Sarah studied how hippocampal microRNAs are altered throughout pubertal development and following repeated adolescent binge drinking. She was awarded three years of support for her research with a National Institute on Alcohol Abuse and Alcoholism fellowship from Loyola's Alcohol Research Program and two years of support to attend the annual Society for Neuroscience meeting with Charles Robert Schuster travel awards from Loyola's Neuroscience Research Institute. At Loyola she also served with the Graduate Student Council and conducted assistant teaching for the Medical Neuroscience course.

Sarah actively offered neuroscientific services by teaching at Pritzker Elementary School's annual science night, reviewing manuscript submissions for the journal of Central Nervous System Agents in Medicinal Chemistry and volunteering with the International Drug Abuse Research Society. On April 15th, 2014, she and Jonathan Kulpit had their first child, Raymond. She plans to continue studying and communicating neuroscience research in her career.