

Comparison of musculoskeletal disorders (MSDs) in military versus civilian workers of the ministry of defense of spain

SanzBustillo-Aguirre, Beatriz; Higuera-Gomez, Loreto; Miralles-Muñoz, Fernando; Angulo-Diaz-Parreño, Santiago; Orr, Rob Marc; Lopez-Mojares, Luis

Licence:
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

Recommended citation(APA):

SanzBustillo-Aguirre, B., Higuera-Gomez, L., Miralles-Muñoz, F., Angulo-Diaz-Parreño, S., Orr, R. M., & Lopez-Mojares, L. (2021). *Comparison of musculoskeletal disorders (MSDs) in military versus civilian workers of the ministry of defense of spain*. Poster session presented at XXII World Congress on Safety and Health , Canada.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Comparison of Musculoskeletal Disorders (MSDs) in Military *versus* Civilian Workers of the Ministry of Defense of Spain

Beatriz SanzBustillo-Aguirre^{1,2}, Loreto A. Higuera-Gomez¹, Fernando Miralles-Muñoz², Santiago Angulo-Diaz-Parreño², Robin Orr³, Luis M. Lopez-Mojares^{1,4}

¹Ministerio de Defensa, Madrid, Spain; ²Universidad San Pablo-CEU, CEU Universities, Madrid, Spain; ³Tactical Research Unit, Bond University, Gold Coast, Queensland, Australia; ⁴Universidad Europea Madrid, Madrid, Spain.

Background

Work-related Musculoskeletal Disorders (MSDs) impact on workers:

- Safety & health
- Functional ability
- Sick-leave


Injury incidence: military > general population

Purpose

To compare proportions of self-reported MSDs between military and civilian workers of the central headquarters of the Ministry of Defense of Spain.

Materials and Methods

Study sample: 100 participants (n= 50 military; n=50 civilian, (50% each gender) completed validated Spanish version of standardised Kuorinka Nordic Questionnaire.



Adapted from: Kuorinka (1987)

Ethics Approval Institution:

Hospital Central de la Defensa “Gómez Ulla” (Madrid, Spain).

Results

1 Symptoms (last 12 months)

Cervical Region

	♂ (n = 50)	♀ (n = 50)
Military	10 (40%)	21 (84%)
Civilian	15 (60%)	20 (80%)

Dorsolumbar Region

	♂ (n = 50)	♀ (n = 50)
Military	15 (60%)	18 (72%)
Civilian	11 (44%)	21 (84%)

Shoulder Region

	♂ (n=50)	♀ (n = 50)
Military	8 (32%)	13 (52%)
Civilian	10 (40%)	16 (64%)

2 Predominant duration of symptoms/body region (last 12 m)

	♂ (n=50)	♀ (n = 50)
Military	50% Cervical: 1-7 days Shoulder: >30 non-consecutive days	44.4% Dorsolumbar: > 30 non-consecutive days
Civilian	45.5% Dorsolumbar: 8-30 days	40% Cervical: > 30 non-consecutive days

3 Main symptomatic body area preventing from work (last 12 m)

	♂ (n=50)	♀ (n = 50)
Military	6.7% Dorsolumbar	16.8% Dorsolumbar
Civilian	25% Dorsolumbar	10% Cervical

Contact details

Presenting Author: Beatriz SanzBustillo-Aguirre
beabustillo@telefonica.net

Reference Number: 690
 Type: Poster



Conclusions

- The three most common symptom sites reported for military and civilian personnel were the same, with no statistical differences across both populations ($p > 0.05$).
- Female personnel represented a greater proportion of results in all three main body regions ($p < 0.05$) when compared to male personnel.
 - Female personnel are known to more likely report an injury.
- While reported symptoms for the cervical and shoulder regions were higher in the civilian population, military personnel reported higher rates of longer lasting MSDs episodes than their civilian counterparts.
 - Underreporting of minor injuries is common in the military.
- The spinal region, notably the dorsolumbar, was the main symptomatic site preventing from work.

Recommendations

- Identification of factors leading to the higher representation of female personnel reporting symptoms.
- Improve reporting of minor injuries for military personnel.
- Risk mitigation strategies targeting the spinal and shoulder regions (e.g., workplace ergonomics) be investigated and trialled.

References

- Kuorinka I, Jonsson B, Kilbom A, et al. *Standardised Nordic Questionnaire for the analysis of musculoskeletal symptoms*. Appl Ergon 1987;18:233-237.
- Pope R, Orr R. *Incidence rates for work health and safety incidents and injuries in Australian Army Reserve vs full time soldiers, and a comparison of reporting systems*. Journal of Military and Veteran’s Health 2017; 25:16-25.
- Protocol for systematic reviews of exposure to occupational ergonomic risk factors and of the effect of exposure to occupational ergonomic risk factors on osteoarthritis of hip or knee and selected other musculoskeletal diseases*. Hulshof CTJ, Colosio C, Daams JG, et al. Environment International. 2019;125:554-566.