

Bond University
Research Repository



Alexithymia, rash impulsiveness, and reward sensitivity in relation to symptoms of exercise dependence in physically active young adults

Lyvers, Michael F.; Sweetnam, Taylor; Thorberg, Fred

Published in:
Australian Journal of Psychology

DOI:
[10.1080/00049530.2021.1981747](https://doi.org/10.1080/00049530.2021.1981747)

Licence:
Other

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Lyvers, M. F., Sweetnam, T., & Thorberg, F. (2021). Alexithymia, rash impulsiveness, and reward sensitivity in relation to symptoms of exercise dependence in physically active young adults. *Australian Journal of Psychology*. <https://doi.org/10.1080/00049530.2021.1981747>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

Alexithymia, Rash Impulsiveness, and Reward Sensitivity in Relation to Symptoms
of Exercise Dependence in Physically Active Young Adults

¹Michael Lyvers, Ph.D.

¹Taylor Sweetnam, Graduate Diploma of Psychological Science

²Fred Arne Thorberg, Ph.D.

¹School of Psychology, Bond University, Gold Coast Qld Australia

²School of Psychology, University of Queensland, Brisbane Qld Australia

Corresponding author: Michael Lyvers, School of Psychology, Bond University, Gold Coast
Qld Australia. Phone: +61755952565 Email: mlyvers@bond.edu.au

Abstract

Objective. Exercise dependence refers to excessive exercise accompanied by addiction-like symptoms such as craving, tolerance, withdrawal, impaired control, and disruption of life domains. The present study investigated whether personality traits linked to substance and behavioral addictions show similar associations with symptoms of exercise dependence.

Method. Alexithymia and two forms of impulsivity, rash impulsiveness and reward sensitivity, were assessed in relation to exercise dependence symptoms in a sample of 99 physically active young adult men and women. **Results.** All three traits showed significant positive correlations with exercise dependence symptoms and were significant predictors of such symptoms in a hierarchical regression model. Alexithymia was the strongest predictor and fully mediated the contribution of rash impulsiveness according to bootstrapped mediation analysis. **Conclusions.** Findings suggest similar associations of addiction-linked traits with symptoms of exercise dependence and are discussed in terms of potential mechanisms.

Keywords: exercise psychology; sport psychology; personality; health; addiction

The physical and psychological benefits of regular exercise are well-established (WHO, 2010), yet there can be negative effects such as acute or chronic injuries, especially when exercise becomes excessive. A minority of those who exercise regularly may become psychologically dependent on exercise and exhibit symptoms similar to addictions such as craving, impaired control, tolerance, withdrawal, engaging in the behavior despite negative consequences such as injuries, and interference with other life domains (Hausenblas & Symons Downs, 2002a). A review by Sussman et al. (2011a) concluded that the prevalence rate of exercise dependence in the general population is approximately 3%, though a prevalence of nearly 9% was reported in a sample of gym attendees (Manfredi & Gambarini, 2015). As a relatively new concept, exercise dependence has also been referred to as exercise addiction, obsessive exercise, compulsive exercise, exercise abuse, obligatory exercise and problematic practice of physical exercise (Egorov & Szabo, 2013).

Exercise dependence has been reported to show comorbidity with other compulsive behaviors, most commonly eating disorders; this has justified a distinction between “primary” and “secondary” exercise dependence with the latter defined by a reportedly reciprocal relationship with eating disorders and weight loss motivation (Cook & Hausenblas, 2008; Cunningham et al., 2016; Di Lodovico et al., 2018; Meyer et al., 2011). Irrespective of typology, negative psychological symptoms are common correlates of exercise dependence (Back et al., 2019; Costa et al., 2013; Grandi et al., 2011; Meyer et al., 2011; Weinstein et al., 2015). For example, Li et al. (2015) reported that compared to non-dependent students, those with exercise dependence reported higher levels of negative moods such as depression and anxiety, similar to those with substance or eating disorders (e.g., Harrison et al., 2010; Lyvers, Hinton et al., 2014; Thorberg et al., 2017). Also similar to other disorders characterized by compulsive behaviors are the disruptions to social life and personal relationships reported by those with exercise dependence, in this case due to an overriding

commitment to exercise regimens. A unique aspect of exercise dependence is the increased risk of musculoskeletal injuries, as individuals who exercise compulsively may exercise to the point of injuring themselves and continue to exercise despite the injury (Freimuth et al., 2011; Lichtenstein et al., 2017).

The neurophysiological basis of exercise dependence may be similar to that of substance use disorders. According to Dietrich's (2006) transient hypofrontality hypothesis, exercise improves mood by temporarily inhibiting various brain regions including prefrontal cortex; such hypofrontality can also be induced by addictive drugs such as alcohol, cocaine and heroin (Lyvers, 2000). The mood-enhancing effects of exercise suggest a link to reward drive in exercise dependence, similar to substance addictions (e.g., Beck et al., 2009; Lyvers, Hinton et al., 2014). Freimuth et al. (2011) outlined how an individual may progress from healthy involvement in exercise to dependence. In Freimuth et al.'s interpretation, adopting an exercise regimen typically begins as a healthy activity without negative consequences. However, when the individual recognizes the psychological benefits of exercise such as mood elevation and stress reduction, they may start to organize their life around exercise, tend to exercise alone, and suffer negative consequences such as injuries or persistent muscle soreness. Eventually, addiction symptoms such as craving, tolerance and withdrawal may develop. Exercise dependence symptoms thus present at the extreme end of a continuum with normal exercise regimens. Though such hypotheses provide plausible explanations for the progression and maintenance of exercise dependence, they do not explain why some people who exercise regularly become dependent on exercise whereas most do not, similar to the question of why only a minority of substance users or gamblers become addicted (Egorov & Szabo, 2013).

Personality Traits as Risk Factors

Some personality traits have been invoked as potential risk factors for exercise

dependence, including extraversion, narcissism, and neuroticism (Bircher et al., 2017; Cook et al., 2020; Costa & Oliva, 2012; Di Lodovico et al., 2018; Lichtenstein et al., 2014).

Alexithymia and impulsivity are additional traits that have been widely linked to substance and behavioral addictions as risk factors (Cruise & Becerra, 2018; Kandri et al., 2014; Lyvers, Hinton et al., 2014; Mahapatra & Sharma, 2018; Marchetti et al., 2019; Rose & Segrist, 2012; Toneatto et al., 2009), hence the present study investigated their potential associations with symptoms of exercise dependence in a young adult sample.

Alexithymia is a multifaceted personality construct characterized by utilitarian and concrete cognition and difficulties with experiencing and describing feelings (Bagby et al., 2020). It has been regarded as a unique trait dimension that does not show significant overlap with other well-known personality constructs such as the Big Five (Wise et al., 1992). In Australia, high alexithymia as defined by established cut-off score on the Toronto Alexithymia Scale (TAS-20; Bagby et al., 1994b) showed a prevalence of 46% in psychiatric samples versus 12% in community samples (McGillivray et al., 2016). Longitudinal research has indicated that alexithymia tends to be stable over time, showing little change from baseline testing to up to 11-years follow up (Hiirola et al., 2017; Salminen et al., 2006; Tolmunen et al., 2011). A large-scale twin study by Jorgensen et al. (2007) suggested relatively low heritability of alexithymia, thus subsequent research has suggested that high alexithymia may originate in adverse childhood experiences such as neglect, trauma, inadequate maternal bonding and insecure attachment (Lyvers et al., 2019; Thorberg et al., 2011a).

The reported prevalence of high alexithymia in substance abusing samples ranges from 30-50% (Cruise & Becerra, 2018; Lyvers, Hinton et al., 2014; Thorberg et al., 2009), commonly interpreted as reflecting use of substances for emotional self-regulation, which is deficient in alexithymia. For example, in young adults, alexithymia was positively associated

with risky or problematic alcohol or cannabis use and with the use of such substances to cope with negative moods (Lyvers et al., 2018). Alexithymia has also been linked to behavioral addictions and other compulsive behaviors including compulsive buying (Rose & Segrist, 2012), internet addiction (Kandri et al., 2014; Lyvers et al., 2016; Mahapatra & Sharma, 2018), pathological gambling (Marchetti et al., 2019; Toneatto et al., 2009) and eating disorders (Marsero et al., 2011). However, relatively few studies have examined alexithymia in the context of exercise. A qualitative study of male swimmers by Allegre et al. (2007) suggested that expert swimmers were more alexithymic than amateur swimmers. Manfredi and Gambarini (2015) reported a positive relationship between scores on the TAS-20 and the Exercise Dependence Scale (EDS-21; Hausenblas & Symons Downs, 2002b) in an Italian sample of gym attendees; EDS-21 criteria identified that nearly 9% of the sample was exercise dependent. More recently, Van Landeghem et al. (2019) found that exercise dependence symptoms, as measured by the EDS-21, were positively associated with alexithymia as measured by the Bermond-Vorst Alexithymia Questionnaire (BVAQ; Vorst & Bermond, 2001) in an undergraduate student sample. The present study used the EDS-21 to measure exercise dependence symptoms in a sample of physically active young adults.

Like alexithymia, trait impulsivity has also been linked to alcohol abuse and dependence (Littlefield & Sher, 2010), eating disorders (Beck et al., 2009; Claes et al., 2002; Kane et al., 2004) and internet addiction (Dalbudak et al., 2013; Lyvers et al., 2016; Ryu et al., 2018). Impulsivity is a multifaceted construct that Dawe et al. (2004) suggested is best conceptualized in the context of substance abuse by a two-factor model, the 2-Component Approach to Reinforcing Substances (2-CARS) model (Gullo & Dawe, 2008; Gullo et al., 2010; Stautz et al., 2017), incorporating rash impulsiveness and reward drive. An individual with a high level of rash impulsiveness would tend to act without thinking, disregard risk and fail to consider the future consequences of their actions. Rash impulsiveness has a stronger

genetic component than alexithymia (e.g., Anokhin et al., 2015), however both impulsivity (Littlefield & Sher, 2010; Spinella, 2004) and alexithymia (Chester et al., 2015; Kano & Fukudo, 2013; Lyvers, Makin et al., 2014) have been linked to generalized executive dysfunction and deficient prefrontal cortical activation during tasks that normally activate that brain region. Given the higher heritability of rash impulsiveness, one possibility is that high inherent levels of this trait may predispose to development of alexithymia in response to childhood neglect, trauma or inadequate parental bonding (Gullo & Dawe, 2008; Lyvers et al., 2019), perhaps accounting for the reported positive association between the two traits (Shishido et al., 2013). Dawe et al. suggested that rash impulsiveness can be measured by total scores on the Barratt Impulsiveness Scale (BIS-11; Patton et al., 1995), which includes items assessing motor, attentional, and non-planning impulsiveness as facets. As BIS-11 scores have shown positive relationships with symptoms of a variety of substance and behavioral addictions (Dalbudak et al., 2013; Lyvers, Hinton et al., 2014; Ryu et al., 2018), the BIS-11 was used as an index of rash impulsiveness in the present study of exercise dependence symptoms.

The other form of impulsivity relevant to addictions in the 2-CARS model, as proposed by Dawe et al. (2004), is sensitivity to reward (SR), which the authors suggested can be measured by the SR subscale of the Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ; Torrubia et al., 2001). Like rash impulsiveness, SR or reward-driven impulsivity has been linked to a variety of compulsive behaviors as a likely risk factor (e.g., Beck et al., 2009; Lyvers, Hinton et al., 2014). The SPSRQ was thus used in the present study to measure SR in relation to exercise dependence symptoms, and was expected to be positively related to such symptoms given the well-known mood enhancing effects of exercise.

Considering the support for the 2-CARS model of impulsivity in relation to other excessive behaviors, the present study sought to extend this paradigm to symptoms of exercise dependence. For example, those at risk of exercise dependence may be particularly sensitive to the positive reward-based incentives of exercise, whether this be winning in competition, achievement of mastery, improvement in appearance, or direct positive mood effects of exercise. Rash impulsiveness may be manifested by continuing to exercise despite signs of over-exertion or impending injury (Freimuth et al., 2011; Lichtenstein et al., 2017). The present study therefore investigated the 2-CARS model of impulsivity proposed by Dawe et al. (2004) in relation to the EDS-21 measure of exercise dependence symptoms to assess whether either or both forms of impulsivity may be risk factors for exercise dependence as has been reported for other forms of excessive behavior. The prevalence of full exercise dependence by EDS-21 criteria was anticipated to be low in the current non-clinical sample of physically active young adults, who were not recruited from gyms or other venues likely to be frequented by exercise dependent individuals (Manfredi & Gamberini, 2015). However, addictive behaviors are recognized as existing at the extreme end of a continuum (APA, 2013; SAMHSA, 2016), such that lower levels of symptoms can progress to higher levels in susceptible individuals; thus relationships of the traits of interest with exercise dependence symptoms were anticipated in the present sample.

Specifically, scores on the measures of alexithymia, rash impulsivity and SR were expected to be significantly positively correlated with EDS-21 scores, and were anticipated to be significant predictors in a hierarchical regression model. Sex was controlled as a covariate given that some studies have reported sex differences on the traits of interest, particularly alexithymia (e.g., Hiirola et al., 2017; Levant et al., 2009; Salminen et al., 2006), and that so-called primary exercise dependence, which is not associated with a comorbid eating disorder, has been reported to be more common in males (Costa et al., 2013; Cunningham et al., 2016).

The contributions of alexithymia and rash impulsiveness were expected to show some degree of overlap in accounting for variance in exercise dependence symptoms, based on previous evidence of a positive association of rash impulsiveness with alexithymia in the context of substance use (Lyvers et al., 2013; Shishido et al., 2013) as well as the evidence cited earlier suggesting that high levels of either trait reflect generalized executive/prefrontal dysfunction. The degree of independence of alexithymia and rash impulsiveness in relation to symptoms of exercise dependence was assessed by regression followed by a planned bootstrapped mediation analysis.

Method

Participants

The initial sample consisted of 119 young adults recruited from two university campuses in southeast Queensland, Australia, and from other states via Qualtrics Panels. Due to the covid-19 pandemic and resulting low student presence at the two universities at the time of the study, most participants were recruited online via Qualtrics Panels ($n = 84$); they were contacted via email by the recruitment team at Qualtrics and were provided points-based incentives. Participants from one university ($n = 23$) were recruited through an online undergraduate psychology student participation noticeboard and were offered course credit as an incentive. At the other university, participants ($n = 12$) were recruited via campus-wide email and were not offered any incentive. The advertisement for the study asked for participants who were 18-30 years old and who regularly (i.e., at least weekly) engage in exercise or sport, with no history of traumatic brain injury and no current use of medication for a psychiatric or neurological disorder. After excluding cases that did not meet one or more inclusion criteria, the final sample of 99 participants aged 18 to 30 years ($M = 24.53$, $SD = 3.56$) was obtained, including 56 females and 43 males. G*power 3.1.9.7 indicated that for a small effect size (.20), power of .90, alpha of .05, and five predictors in multiple regression, a

minimum sample size of 89 was required; the final sample of 99 was thus adequate for present purposes.

Scoring protocols for the EDS-21 index of exercise dependence symptoms (described below) indicated a high rate of exercise dependence symptoms in the sample, such that 71% were classified as “symptomatic,” though only a further 2% could be classified as fully exercise dependent (which was close to the estimated prevalence in the general population; Sussman et al., 2011a). Nearly all participants (92%) reported exercising at least 2-3 times per week; 84% reported that their primary exercise consisted of fitness-oriented activities, with the remainder reporting engaging in either team sports or recreational non-team sports as their primary exercise.

Materials

Demographics Questionnaire. This questionnaire requested information from the participants regarding their age, sex, education, study status, exercise habits and any history of brain injury or psychiatric/neurological conditions (for screening purposes).

Toronto Alexithymia Scale 20 (TAS-20; Bagby et al., 1994a). The TAS-20 is a self-report questionnaire designed to measure alexithymia via items assessing difficulty identifying feelings (DIF), difficulty describing feelings (DDF), and externally oriented thinking (EOT). It is the most widely used measure of alexithymia (Bagby et al., 2020). The scale consists of 20 items (e.g., “I am often puzzled by sensations in my body”), five of which are reverse scored (e.g., “Being in touch with emotions is essential”); response options are anchored on a five-point Likert scale from 1 (strongly disagree) to 5 (strongly agree). Item responses are summed to yield a total score, where higher scores indicate higher alexithymia. There are cutoff scores for high alexithymia (61+) and low to no alexithymia (< 52), with scores in between considered borderline. The present study used the total score as

recommended by the authors of the TAS-20 (see Sekely et al., 2018). Internal consistency reliability was acceptable in the current sample ($\alpha = .83$).

Barratt Impulsiveness Scale (BIS-11; Patton et al., 1995). The BIS-11 is a self-report measure of rash impulsiveness (Dawe et al., 2004) via items assessing attentional impulsiveness (inability to focus attention or concentrate), motor impulsiveness (acting without thinking), and non-planning impulsiveness (lack of future time orientation). The scale consists of 30 items (e.g., “I say things without thinking”), 11 of which are reverse scored (e.g., “I plan tasks carefully”); response options are scored on a five-point Likert scale from 1 (rarely/never) to 5 (almost always/always). Total scores are obtained by summing item responses, with a higher total score indicating a higher level of rash impulsiveness (Dawe et al., 2004). Internal consistency reliability was acceptable for the current sample ($\alpha = .83$) and was the same as previously reported by Stanford et al. (2009).

Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ; Torrubia et al., 2001). The SPSRQ is a self-report measure designed to assess everyday influences of the Behavioral Approach and Behavioral Inhibition systems on an individual’s behavior, based on Gray’s (1987) Reinforcement Sensitivity Theory. There are two corresponding subscales, Sensitivity to Reward (SR) and Sensitivity to Punishment (SP). These subscales consist of 24 items each, with response options of “yes” or “no.” Even numbered items tap SR; e.g., “Do you like displaying your physical abilities even though this may involve danger?” Odd numbered items tap SP; e.g., “Whenever possible, do you avoid demonstrating your skills for fear of being embarrassed?” Subscale scoring is completed by adding all “yes” responses, with a higher score indicating higher SR or SP. In the current sample, internal consistency reliability was acceptable for both SP ($\alpha = .82$) and SR ($\alpha = .77$).

Exercise Dependence Scale Revised (EDS-21; Hausenblas & Symons Downs, 2002b). The EDS-21 is a 21 item self-report questionnaire assessing symptoms of exercise

dependence based on the DSM-IV (APA, 2000) criteria for substance dependence. Each of the seven criteria indicative of exercise dependence (tolerance, withdrawal, intention effects, lack of control, time, reduction in other activities, and continuance) is assessed by three items. Response options are anchored on a six-point Likert scale, with options ranging from 1 (never) to 6 (always). Examples of items include “I exercise to avoid feeling irritable” and “I exercise when injured.” Item responses can be summed to provide an overall score where higher scores indicate higher levels of exercise dependence symptoms. Criteria are also provided for classification of respondents as non-dependent, symptomatic, or dependent. In the current sample internal consistency reliability of the total EDS-21 was high ($\alpha = .96$).

Procedure

Ethics approval was obtained from the Bond University Human Research Ethics Committee (approval number TS02912) prior to data collection. The internet survey hosting service Qualtrics was used to post the measures online. On clicking the link provided, the first page presented an explanatory statement which provided contact details of the researchers, described the general purpose of the study in lay terms, and explained that all information collected was anonymous. Participants were asked to tick a box to provide informed consent. Participants then completed the demographics questionnaire, after which the other questionnaires were presented in uniquely randomized orders for each participant. Each question required a response, such that participants could not continue to the next question until they had responded to the previous one. Completion of all measures took approximately 30 minutes.

Results

Data are available on request and were collected via Qualtrics and imported to an IBM SPSS Statistics (Version 26.0) datafile for analysis via JASP 0.14.1. Data screening indicated that for the variables of interest there were no missing values, no univariate or multivariate

outliers, and no evidence of multicollinearity; skewness and kurtosis were within the acceptable range (Tabachnik & Fidell, 2014). Comparison of males and females on all variables of interest using multivariate analysis of variance indicated no significant sex differences; however, TAS-20 alexithymia scores approached significance. Males scored nonsignificantly higher on the TAS-20 ($M = 58.35$, $SD = 10.83$) than females ($M = 54.21$, $SD = 11.81$), $p = .08$, hence sex was included as a covariate in regression and mediation analyses. No other measures approached significance.

Correlational Analysis

Descriptive statistics and Pearson correlations for continuous variables are shown in Table 1. As predicted, significant positive correlations were obtained between participants' EDS-21 and TAS-20, BIS-11, and SPSRQ-SR scores. SPSRQ-SP scores showed small to moderate positive correlations with BIS-11 and SR, respectively, but were not correlated with EDS-21; nevertheless SP was included in subsequent analyses given its correlations with the other predictors. Age was not correlated with any of the variables of interest and was thus not entered as a covariate in regression.

Regression on Exercise Dependence Symptoms

Hierarchical regression was conducted on EDS-21 scores. Sex was entered at step 1 as a covariate for reasons described earlier; the impulsivity-linked trait measures BIS-11 and SR, as well as SP, were entered at step 2, followed by the TAS-20 alexithymia measure at step 3. At step 1, sex accounted for only 2% of variance and was not significant, $F(1, 97) = 1.77$, $p = .19$. At step 2, SR, SP, and BIS-11 explained 15% of additional variance, which was significant, $\Delta F(3, 94) = 5.74$, $p = .001$; as expected, SR and BIS-11 were significant predictors (see Table 2). At step 3, TAS-20 scores explained 16% of additional variance, which was significant, $\Delta F(1, 93) = 21.78$, $p < .001$; TAS-20 and SR were significant predictors, whereas

BIS-11 was no longer significant. The final model was significant and accounted for 33% of variance in EDS-21 scores, $F(5, 93) = 9.06, p < .001$.

Mediation Analysis

As anticipated based on previously cited evidence of high rash impulsiveness associated with alexithymia in the context of substance use, the contribution of BIS-11 scores became nonsignificant when TAS-20 scores were entered into the regression model, suggesting that the relationship between BIS-11 and EDS-21 scores was mediated by alexithymia. The mediation hypothesis was tested using bootstrapping with 1000 bias-corrected replications, controlling for sex, SR and SP as covariates. The 95% confidence interval for the indirect effect of rash impulsiveness through alexithymia did not include zero (.07, .42). The model as depicted in Figure 1 indicated full mediation by alexithymia, $p < .001$. By contrast, reversing the mediation model, with rash impulsiveness as the presumed mediator between alexithymia and exercise dependence symptoms while controlling for sex, SR and SP, did not show a significant indirect effect ($p = .41$).

Discussion

All predictions were supported in the present sample of physically active male and female young adults. The two forms of impulsivity specified by the 2-CARS model of Dawe et al. (2004) as inherent risk factors for addictions, i.e., reward sensitivity and rash impulsiveness, were both positively correlated with scores on the EDS-21 index of exercise dependence symptoms and predicted EDS-21 scores in a regression model. However, entry of alexithymia into the model explained more variance in exercise dependence symptoms and fully mediated the contribution of rash impulsiveness. Interestingly, there was no reverse mediation; rash impulsiveness did not mediate the association of alexithymia with exercise dependence symptoms. Alexithymia was thus the trait most strongly associated with exercise dependence symptoms in the present sample. The findings add to the traits implicated as

potential risk factors for exercise dependence, and reinforce previous reports of a positive relationship between alexithymia and exercise dependence symptoms in gym attendees (Manfredi & Gambarini, 2015) and undergraduate students (Van Landeghem et al., 2019).

As discussed earlier, both alexithymia and impulsivity have been identified as traits linked to a variety of addictions and other excessive behaviors. However, there has been comparatively little research assessing alexithymia or impulsivity in relation to exercise dependence symptoms, a gap the present study sought to address. Using Dawe et al.'s (2004) 2-CARS paradigm of two forms of impulsivity relevant to addictions, present results suggested largely independent contributions of alexithymia and reward sensitivity to exercise dependence symptoms, whereas the contribution of rash impulsiveness was indirect through its association with alexithymia. The role of reward sensitivity would presumably reflect the numerous benefits of exercise, including positive mood changes and stress relief. Indeed, engaging in exercise is often recommended as a means to aid recovery from substance use disorders, with some empirical support (e.g., Smith & Lynch, 2012). However, alexithymia was by far the strongest predictor of exercise dependence symptoms in the present sample, which raises the question as to why.

The links between alexithymia and various excessive behaviors have often been attributed to self-medication of the negative moods commonly associated with alexithymia such as depression and anxiety (Ghalehban & Besharat, 2011; Lyvers, Hinton et al., 2014; Lyvers et al., 2018, 2020; Thorberg et al., 2009, 2011b, 2017). A different interpretation was offered by Brewer et al. (2016), who suggested that deficient interoceptive awareness accounts for excessive substance use or eating disorders in alexithymia. Present results are consistent with both interpretations. The ability of exercise to relieve stress and regulate mood is well known, hence exercise can be used as a means of coping with negative affect (Dietrich, 2006; Egorov & Szabo, 2013; Freimuth et al., 2011; Sussman et al., 2011b), which as noted

earlier is prevalent in high alexithymia (Thorberg et al., 2009). In addition, deficient interoception may impair recognition of internal cues of over-exertion or impending bodily injury during exercise, with the result that exercise continues despite negative health consequences. The insecure attachment style and dissatisfaction with personal relationships reported by those with high alexithymia (Besharat et al., 2014; Humphreys et al., 2009; Thorberg et al., 2011b) may also encourage solo activities such as exercise at the expense of social interactions and responsibilities. Future work could evaluate the potential mechanisms of the relationship between alexithymia and exercise dependence symptoms, e.g., by including measures of negative mood, interoceptive awareness and attachment style as potential mediators.

The present study had several limitations. The cross-sectional design did not allow interpretation of relationships among variables in terms of causation, hence longitudinal studies are needed. On the other hand, as noted earlier the traits assessed in the present study appear to be relatively stable over time and likely have an early onset, although impulsivity appears to have a stronger genetic basis than alexithymia. Both impulsivity and alexithymia have been interpreted as different manifestations of generalized executive dysfunction that may predispose to addictions and other excessive behaviors (Lyvers, Hinton et al., 2014; Thorberg et al., 2017). The use of a non-clinical sample of young adults with a low rate of exercise dependence (2%), based on EDS-21 criteria, was also a potential limitation though there was a high rate of “symptomatic” participants (71%) in the sample. The present approach was premised on the currently accepted notion that addictive behaviors present on a continuum of symptoms with behavior that is not excessive (APA, 2013; SAMHSA, 2016), and that risk factors may lead susceptible individuals to progress to addiction over time; nevertheless, the present findings cannot be generalized to clinical samples.

A further issue concerns the reported comorbidity of exercise dependence with eating disorders, which has led to a distinction between primary and secondary exercise dependence; the latter type is defined by this comorbidity and is more common in females as discussed earlier. In the present sample there were no sex differences on any of the measures of interest, though there was a trend for higher alexithymia in males, and in any case the potential influence of sex differences was controlled by covariate analysis. However, there was no measure of disordered eating. Future research may consider comparing primary and secondary exercise dependence in relation to alexithymia, though given the known association of alexithymia with eating disorders (Marsero et al., 2011), the results might be anticipated to be similar for both types of exercise dependence.

The prevalence of high alexithymia in the current sample (37%), as defined by established TAS-20 cutoff score (Bagby et al., 1994b), fell between the estimated Australian prevalence of 12% in non-clinical samples and 46% in clinical samples (McGillivray et al., 2016). The prevalence in the present sample was thus surprisingly high, although a similar prevalence was reported in a recent study of frequent Facebook users in Australia (38% prevalence; Lyvers et al., 2020). In the latter study this was attributed to online recruitment of participants, based on evidence that individuals with high alexithymia spend more time on the internet and are thus more likely to be represented in samples recruited online. In the present study, most participants were similarly recruited online due to the covid-19 pandemic, which may limit generalizability of the findings. Another unusual feature of the current sample was the moderate positive correlation between SR and SP. This was surprising given that SR and SP are theorised to reflect the influences of distinct brain systems in Gray's (1987) BIS/BAS theory (Behavioral Inhibition or amygdala system and Behavioral Approach or dopaminergic reward system, respectively), which the SPSRQ scales are designed to assess. In any case, this unexpected relationship was controlled for in the regression and mediation analyses.

Despite the aforementioned limitations, the present findings reinforce the notion that excessive exercise shares features in common with other excessive behaviors, in this case traits presumed to be risk factors for substance and behavioral addictions. Future research should focus on elucidating the complex relationships between trait factors as potential contributors to risk of exercise dependence, so that the roles of inherent risk factors are better understood and can be targeted effectively when needed.

References

- Allegre, B., Souville, M., Noel-Jorand, M., Pellegrin, L., & Therme, P. (2007). Intensive physical activity and alexithymia: Results from swimmers' discourse analysis. *Psychological Reports, 100*(3), 1129-1139. <https://doi.org/10.2466/pr0.100.4.1129-1139>
- American Psychiatric Association (APA) (2000). *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR (4th ed., text revision)*. Washington, DC: APA.
- American Psychiatric Association (APA) (2013). *Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.)*. Washington DC: APA.
- Anokhin, A.P., Grant, J.D., Mulligan, R.C., & Heath, R.C. (2015). The genetics of impulsivity: Evidence for the heritability of delay discounting. *Biological Psychiatry, 77*(10), 887-94. <https://doi.org/10.1016/j.biopsych.2014.10.022>
- Back, J., Josefsson, T., Ivarsson, A., & Gustafsson, H. (2019). Psychological risk factors for exercise dependence. *International Journal of Sport and Exercise Psychology, 19*(4), 1-12. <https://doi.org/10.1080/1612197x.2019.1674902>
- Bagby, R.,M., Parker, J., & Taylor, G. (1994a). The twenty-item Toronto Alexithymia Scale—I. Item selection and cross-validation of the factor structure. *Journal of Psychosomatic Research, 38*(1), 23-32. [https://doi.org/10.1016/0022-3999\(94\)90005-1](https://doi.org/10.1016/0022-3999(94)90005-1)
- Bagby, R. M., Parker, J. D. A., & Taylor, G. J. (2020). Twenty-five years with the 20-item Toronto Alexithymia Scale. *Journal of Psychosomatic Research, 131*, 109940 <https://doi.org/10.1016/j.jpsychores.2020.109940>
- Bagby, R.M., Taylor, G., & Parker, J. (1994b). The twenty-item Toronto Alexithymia Scale—II. Convergent, discriminant, and concurrent validity. *Journal of Psychosomatic Research, 38*(1), 33-40. [https://doi.org/10.1016/0022-3999\(94\)90006-x](https://doi.org/10.1016/0022-3999(94)90006-x)

- Beck, I., Smits, D., Claes, L., Vandereycken, W., & Bijttebier, P. (2009). Psychometric evaluation of the Behavioral Inhibition/Behavioral Activation System Scales and the Sensitivity to Punishment and Sensitivity to Reward Questionnaire in a sample of eating disordered patients. *Personality and Individual Differences, 47*(5), 407-412. <https://doi.org/10.1016/j.paid.2009.04.007>
- Besharat, M. A., Naghshineh, N., Ganji, P., & Tavalaeayan, F. (2014). The moderating role of attachment styles on the relationship of alexithymia and fear of intimacy with marital satisfaction. *International Journal of Psychological Studies, 6*(3), 106-117. doi:10.5539/ijps.v6n3p106
- Bircher, J., Griffiths, M., Kasos, K., Demetrovics, Z., & Szabo, A. (2017). Exercise addiction and personality: A two-decade systematic review of the empirical literature (1995–2016). *Baltic Journal of Sport and Health Sciences, 3*(106), 19-33. <https://doi.org/10.33607/bjshs.v3i106.30>
- Brewer, R., Cook, R., & Bird, G. (2016). Alexithymia: A general deficit of interoception. *Royal Society Open Science, 3*(10), 150664. doi: 10.1098/rsos.150664
- Chester, D. S., Pond, R. S., Jr., & DeWall, N. C. (2015). Alexithymia is associated with blunted anterior cingulate response to social rejection: Implications for daily rejection. *Social Cognitive & Affective Neuroscience, 10*(4), 517-522. doi:10.1093/scan/nsu082
- Claes, L., Vandereycken, W., & Vertommen, H. (2002). Impulsive and compulsive traits in eating disordered patients compared with controls. *Personality and Individual Differences, 32*(4), 707-714. [https://doi.org/10.1016/s0191-8869\(01\)00071-x](https://doi.org/10.1016/s0191-8869(01)00071-x)
- Cook, R., Griffiths, M., & Pontes, H. (2020). Personality factors in exercise addiction: A pilot study exploring the role of narcissism, extraversion, and agreeableness. *International Journal of Mental Health and Addiction, 18*(1), 89-102. <https://doi.org/10.1007/s11469-018-9939-z>

- Cook, B., & Hausenblas, H. A. (2008). The role of exercise dependence for the relationship between exercise behavior and eating pathology. *Journal of Health Psychology, 13*(4), 495-502. <https://doi.org/10.1177/1359105308088520>
- Costa, S., Hausenblas, H. A., Oliva, P., Cuzzocrea, F., & Larcán, R. (2013). The role of age, gender, mood states and exercise frequency on exercise dependence. *Journal of Behavioral Addictions, 2*(4), 216-223. <https://doi.org/10.1556/jba.2.2013.014>
- Costa, S., & Oliva, P. (2012). Examining relationship between personality characteristics and exercise dependence. *Review of Psychology, 19*(1), 5-12.
http://mjesecc.ffzg.hr/revija.psi/vol%2019%20no%201%202012/Costa_Oliva_2012_v19-1.pdf
- Cruise, K., & Becerra, R. (2018). Alexithymia and problematic alcohol use: A critical update. *Addictive Behaviors, 77*, 232-246. <https://doi.org/10.1016/j.addbeh.2017.09.025>
- Cunningham, H., Pearman, S., & Brewerton, T. (2016). Conceptualizing primary and secondary pathological exercise using available measures of excessive exercise. *International Journal of Eating Disorders, 49*(8), 778-792.
<https://doi.org/10.1002/eat.22551>
- Dalbudak, E., Evren, C., Topcu, M., Aldemir, S., Coskun, K., Bozkurt, M., Evren, B., & Canbal, M. (2013). Relationship of internet addiction with impulsivity and severity of psychopathology among Turkish university students. *Psychiatry Research, 210*(3), 1086-1091. <https://doi.org/10.1016/j.psychres.2013.08.014>
- Dawe, S., Gullo, M., & Loxton, N. (2004). Reward drive and rash impulsiveness as dimensions of impulsivity: Implications for substance misuse. *Addictive Behaviors, 29*(7), 1389-1405. <https://doi.org/10.1016/j.addbeh.2004.06.004>
- Dietrich, A. (2006). Transient hypofrontality as a mechanism for the psychological effects of

exercise. *Psychiatry Research*, 145(1), 79-83.

<https://doi.org/10.1016/j.psychres.2005.07.033>

Di Lodovico, L., Dubertret, C., & Ameller, A. (2018). Vulnerability to exercise addiction, socio-demographic, behavioral and psychological characteristics of runners at risk for eating disorders. *Comprehensive Psychiatry*, 81, 48-52.

<https://doi.org/10.1016/j.comppsy.2017.11.006>

Egorov, A., & Szabo, A. (2013). The exercise paradox: An interactional model for a clearer conceptualization of exercise addiction. *Journal of Behavioral Addictions*, 2(4), 199-208. <https://doi.org/10.1556/jba.2.2013.4.2>

Freimuth, M., Moniz, S., & Kim, S. (2011). Clarifying exercise addiction: Differential diagnosis, co-occurring disorders, and phases of addiction. *International Journal of Environmental Research and Public Health*, 8(10), 4069-4081.

<https://doi.org/10.3390/ijerph8104069>

Ghalehban, M., & Besharat, M. (2011). Examination and comparison of alexithymia and self regulation in patients with substance abuse disorder and normal individuals. *Procedia - Social and Behavioral Sciences*, 30, 38-42.

<https://doi.org/10.1016/j.sbspro.2011.10.008>

Grandi, S., Clementi, C., Guidi, J., Benassi, M., & Tossani, E. (2011). Personality characteristics and psychological distress associated with primary exercise dependence: An exploratory study. *Psychiatry Research*, 189(2), 270-275.

<https://doi.org/10.1016/j.psychres.2011.02.025>

Gray, J. A. (1987). *The psychology of fear and stress*. Cambridge University Press.

Gullo, M. J., & Dawe, S. (2008). Impulsivity and adolescent substance use: Rashly dismissed as “all-bad”? *Neuroscience and Biobehavioral Reviews*, 32(8), 1507–1518.

<https://doi.org/10.1016/j.neubiorev.2008.06.003>

- Gullo, M. J., Dawe, S., Kambouropoulos, N., Staiger, P. K., & Jackson, C. J. (2010). Alcohol expectancies and drinking refusal self-efficacy mediate the association of impulsivity with alcohol misuse. *Alcoholism: Clinical and Experimental Research, 34*(8), 1386–1399. <https://doi.org/10.1111/j.1530-0277.2010.01222.x>
- Harrison, A., Sullivan, S., Tchanturia, K., & Treasure, J. (2010). Emotional functioning in eating disorders: Attentional bias, emotion recognition, and emotion regulation. *Psychological Medicine, 40*(11), 1887-1897. doi:10.1017/S0033291710000036
- Hausenblas, H. A., & Symons Downs, D. (2002a). Exercise dependence: A systematic review. *Psychology of Sport and Exercise, 3*(2), 89-123. [https://doi.org/10.1016/S1469-0292\(00\)00015-7](https://doi.org/10.1016/S1469-0292(00)00015-7)
- Hausenblas, H.A., & Symons Downs, D. (2002b). How much is too much? The development and validation of the Exercise Dependence Scale. *Psychology & Health, 17*(4), 387-404. <https://doi.org/10.1080/0887044022000004894>
- Hiirola, A., Pirkola, S., Karukivi, M., Markkula, N., Bagby, R.M., Joukamaa, M., Jula, A., Kronholm, E., Saarijarvi, S., Salminen, J.R., Suvisari, J., Taylor, G., & Mattila, A.K. (2017). An evaluation of the absolute and relative stability of alexithymia over 11 years in a Finnish general population. *Journal of Psychosomatic Research, 95*, 81-87. <https://doi.org/10.1016/j.jpsychores.2017.02.007>
- Humphreys, T. P., Wood, L. M., & Parker, J. D. A. (2009). Alexithymia and satisfaction in intimate relationships. *Personality and Individual Differences, 46*, 43-47. doi:10.1016/j.paid.2008.09.002
- Jorgensen, M., Zachariae, R., Skytthe, A., & Kyvik, K. (2007). Genetic and environmental factors in alexithymia: A population-based study of 8,785 Danish twin pairs. *Psychotherapy and Psychosomatics, 76*(6), 369-375. <https://doi.org/10.2307/48511040>
- Kandri, T., Bonotis, K., Floros, G., & Zafiropoulou, M. (2014). Alexithymia components in

- excessive internet users: A multi-factorial analysis. *Psychiatry Research*, 220(1-2), 348-355. <https://doi.org/10.1016/j.psychres.2014.07.066>
- Kane, T., Loxton, N., Staiger, P., & Dawe, S. (2004). Does the tendency to act impulsively underlie binge eating and alcohol use problems? An empirical investigation. *Personality and Individual Differences*, 36(1), 83-94. [https://doi.org/10.1016/s0191-8869\(03\)00070-9](https://doi.org/10.1016/s0191-8869(03)00070-9)
- Kano, M., & Fukudo, S. (2013). The alexithymic brain: The neural pathways linking alexithymia to physical disorders. *Biopsychosocial Medicine*, 7(1), 1. <https://doi.org/10.1186/1751-0759-7-1>
- Levant, R., Hall, R., Williams, C., & Hasan, N. (2009). Gender differences in alexithymia. *Psychology of Men and Masculinity*, 10(3), 190-203. <https://doi.org/10.1037/a0015652>
- Li, M., Nie, J., & Ren, Y. (2015). Effects of exercise dependence on psychological health of Chinese college students. *Psychiatria Danubina*, 27(4), 413-419. http://www.psychiatriadanubina.com/UserDocsImages/pdf/dnb_vol27_no4/dnb_vol27_no4_413.pdf
- Lichtenstein, M., Christiansen, E., Elklit, A., Bilenberg, N., & Støving, R. (2014). Exercise addiction: A study of eating disorder symptoms, quality of life, personality traits and attachment styles. *Psychiatry Research*, 215(2), 410-416. <https://doi.org/10.1016/j.psychres.2013.11.010>
- Lichtenstein, M., Hinze, C., Emborg, B., Thomsen, F., & Hemmingsen, S. (2017). Compulsive exercise: Links, risks and challenges faced. *Psychology Research and Behavior Management*, 10, 85-95. <https://doi.org/10.2147/prbm.s113093>
- Littlefield, A., & Sher, K. (2010). The multiple, distinct ways that personality contributes to alcohol use disorders. *Social and Personality Psychology Compass*, 4(9), 767-782. <https://doi.org/10.1111/j.1751-9004.2010.00296.x>

- Lyvers, M. (2000). "Loss of control" in alcoholism and drug addiction: A neuroscientific interpretation. *Experimental and Clinical Psychopharmacology*, 8(2), 225-249.
<https://doi.org/10.1037//1064-1297.8.2.225>
- Lyvers, M., Cutinho, D., & Thorberg, F.A. (2020). Alexithymia, impulsivity, disordered social media use, mood and alcohol use in relation to Facebook self-disclosure. *Computers in Human Behavior*, 103, 174-180.
<https://doi.org/10.1016/j.chb.2019.09.004>
- Lyvers, M., Hanigan, C., & Thorberg, F.A. (2018). Social interaction anxiety, alexithymia, and drinking motives in Australian university students. *Journal of Psychoactive Drugs*, 50(5), 402-410. <https://doi.org/10.1080/02791072.2018.1517228>
- Lyvers, M., Hinton, R., Gotsis, S., Roddy, M., Edwards, M., & Thorberg, F.A. (2014). Traits linked to executive and reward systems functioning in clients undergoing residential treatment for substance dependence. *Personality and Individual Differences*, 70, 194-199. <https://doi.org/10.1016/j.paid.2014.07.004>
- Lyvers, M., Jamieson, R., & Thorberg, F.A. (2013). Risky cannabis use is associated with alexithymia, frontal lobe dysfunction, and impulsivity in young adult cannabis users. *Journal of Psychoactive Drugs*, 45(5), 394-403.
<https://doi.org/10.1080/02791072.2013.844525>
- Lyvers, M., Karantonis, J., Edwards, M. S., Thorberg, F.A. (2016). Traits associated with internet addiction in young adults: Potential risk factors. *Addictive Behaviors Reports*, 3, 56-60. <https://dx.doi.org/10.1016/j.abrep.2016.04.001>
- Lyvers, M., Makin, C., Toms, E., Thorberg, F.A., & Samios, C. (2014). Trait mindfulness in relation to alexithymia, emotion regulation, mood and everyday executive functioning. *Mindfulness*, 5, 619-625. <https://doi.org/10.1007/s12671-013-0213-y>

- Lyvers, M., Mayer, K., Needham, K., & Thorberg, F.A. (2019). Parental bonding, adult attachment, and theory of mind: A developmental model of alexithymia and alcohol-related risk. *Journal of Clinical Psychology, 75*(7), 1288-1304. DOI: 10.1002/jclp.22772
- Mahapatra, A., & Sharma, P. (2018). Association of internet addiction and alexithymia – A scoping review. *Addictive Behaviors, 81*, 175-182. <https://doi.org/10.1016/j.addbeh.2018.02.004>
- Manfredi, P., & Gambarini, A. (2015). Exercise addiction and alexithymia. *Journal of Psychology and Behavioral Science, 3*(1), 61-70 <https://doi.org/10.15640/jpbs.v3n1a7>
- Marchetti, D., Verrocchio, M., & Porcelli, P. (2019). Gambling problems and alexithymia: A systematic review. *Brain Sciences, 9*(8), 191. <https://doi.org/10.3390/brainsci9080191>
- Marsero, S., Ruggiero, G., Scarone, S., Bertelli, S., & Sassaroli, S. (2011). The relationship between alexithymia and maladaptive perfectionism in eating disorders: A mediation moderation analysis methodology. *Eating and Weight Disorders, 16*(3), 182-187. <https://doi.org/10.1007/bf03325130>
- McGillivray, L., Becerra, R., & Harms, C. (2016). Prevalence and demographic correlates of alexithymia: A comparison between Australian psychiatric and community samples. *Journal of Clinical Psychology, 73*(1), 76-87. <https://doi.org/10.1002/jclp.22314>
- Meyer, C., Taranis, L., Goodwin, H., & Haycraft, E. (2011). Compulsive exercise and eating disorders. *European Eating Disorders Review, 19*(3), 174-189. <https://doi.org/10.1002/erv.1122>
- Patton, J., Stanford, M., & Barratt, E. (1995). Factor structure of the Barratt Impulsiveness Scale. *Journal of Clinical Psychology, 51*(6), 768-774. [https://doi.org/10.1002/1097-4679\(199511\)51:6<768::aid-jclp2270510607>3.0.co;2-1](https://doi.org/10.1002/1097-4679(199511)51:6<768::aid-jclp2270510607>3.0.co;2-1)

- Rose, P., & Segrist, D. (2012). Difficulty identifying feelings, distress tolerance and compulsive buying: Analyzing the associations to inform therapeutic strategies. *International Journal of Mental Health and Addiction*, *10*(6), 927-935.
<https://doi.org/10.1007/s11469-012-9389-y>
- Ryu, H., Lee, J., Choi, A., Park, S., Kim, D., & Choi, J. (2018). The relationship between impulsivity and internet gaming disorder in young adults: Mediating effects of interpersonal relationships and depression. *International Journal of Environmental Research and Public Health*, *15*(3), 458. <https://doi.org/10.3390/ijerph15030458>
- Salminen, J., Saarijärvi, S., Toikka, T., Kauhanen, J., & Äärelä, E. (2006). Alexithymia behaves as a personality trait over a 5-year period in Finnish general population. *Journal of Psychosomatic Research*, *61*(2), 275-278.
<https://doi.org/10.1016/j.jpsychores.2006.01.014>
- Sekely, A., Bagby, R. M., & Porcelli, P. (2018). Assessment of the alexithymia construct. In O. Luminet, R. M. Bagby & G. J. Taylor (Eds.), *Alexithymia: Advances in research, theory, and clinical practice* (pp. 17-32). Cambridge, UK: Cambridge University Press.
<https://doi.org/10.1017/9781108241595.004>
- Shishido, H., Gaher, R.M., & Simons, J.S. (2013). I don't know how I feel, therefore I act: Alexithymia, urgency, and alcohol problems. *Addictive Behaviors*, *38*(4), 2014-2017.
<https://doi.org/10.1016/j.addbeh.2012.12.014>
- Smith, M.A., & Lynch, W.J. (2012). Exercise as a potential treatment for drug abuse: Evidence from preclinical studies. *Frontiers in Psychiatry*, *2*, 82.
doi: 10.3389/fpsyt.2011.00082
- Spinella, M. (2004). Neurobehavioral correlates of impulsivity: Evidence of prefrontal involvement. *International Journal of Neuroscience*, *114*(1), 95-104. doi:
10.1080/00207450490249347

- Stanford, M., Mathias, C., Dougherty, D., Lake, S., Anderson, N., & Patton, J. (2009). Fifty years of the Barratt Impulsiveness Scale: An update and review. *Personality and Individual Differences, 47*(5), 385-395. <https://doi.org/10.1016/j.paid.2009.04.008>
- Stautz, K., Dinc, L., & Cooper, A. J. (2017). Combining trait models of impulsivity to improve explanation of substance use behaviour. *European Journal of Personality, 31*(1), 118–132. <https://doi.org/10.1002/per.2091>
- Substance Abuse and Mental Health Services Administration (SAMHSA); Office of the Surgeon General (2016). *Facing addiction in America*: Washington, D.C.: US Department of Health and Human Services.
- Sussman, S., Lisha, N., & Griffiths, M. (2011a). Prevalence of the addictions: A problem of the majority or the minority? *Evaluation and the Health Professions, 34*(1), 3-56. <https://doi.org/10.1177/0163278710380124>
- Sussman, S., Leventhal, A., Bluthenthal, R., Freimuth, M., Forster, M., & Ames, S. (2011b). A framework for the specificity of addictions. *International Journal of Environmental Research and Public Health, 8*(8), 3399-3415. <https://doi.org/10.3390/ijerph8083399>
- Tabachnick, B., & Fidell, L. (2014). *Using multivariate statistics* (6th ed.). Pearson.
- Thorberg, F.A., Young, R. McD., Lyvers, M., Connor, J. P., Tyssen, R., London, E. D., & Feeney, GFX. (2017). Alexithymia in alcohol-dependent patients is mediated by stress, anxiety and loss of self-control over drinking. *Drug and Alcohol Dependence, 171*, e201-e202. <https://doi.org/10.1016/j.drugalcdep.2016.08.551>
- Thorberg, F.A., Young, R., Sullivan, K., & Lyvers, M. (2009). Alexithymia and alcohol use disorders: A critical review. *Addictive Behaviors, 34*(3), 237-245. <https://doi.org/10.1016/j.addbeh.2008.10.016>
- Thorberg, F.A., Young, R., Sullivan, K.A., & Lyvers, M. (2011a). Parental bonding and

- alexithymia: A meta-analysis. *European Psychiatry*, 26(3), 187-193.
<https://doi.org/10.1016/j.eurpsy.2010.09.010>
- Thorberg, F. A., Young, R. M., Sullivan, K. A., Lyvers, M., Connor, J. P. & Feeney, G. F. (2011b). Attachment security and alexithymia in a heavy drinking sample. *Addiction Research and Theory*, 19(6), 566-570. <https://doi.org/10.3109/16066359.2011.580065>
- Tolmunen, T., Heliste, M., Lehto, S., Hintikka, J., Honkalampi, K., & Kauhanen, J. (2011). Stability of alexithymia in the general population: An 11-year followup. *Comprehensive Psychiatry*, 52(5), 536-541.
<https://doi.org/10.1016/j.comppsy.2010.09.007>
- Toneatto, T., Lecce, J., & Bagby, M. (2009). Alexithymia and pathological gambling. *Journal of Addictive Diseases*, 28(3), 193-198. <https://doi.org/10.1080/10550880903014775>
- Torrubia, R., Ávila, C., Moltó, J., & Caseras, X. (2001). The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and impulsivity dimensions. *Personality and Individual Differences*, 31(6), 837-862.
[https://doi.org/10.1016/s0191-8869\(00\)00183-5](https://doi.org/10.1016/s0191-8869(00)00183-5)
- Van Landeghem, C., Jakobson, L.S., & Keough, M.T. (2019). Risk of exercise dependence in university students: A subtyping study utilizing latent profile analysis, *Psychology of Sport and Exercise*, 45, 101575.
<https://doi.org/10.1016/j.psychsport.2019.101575>
- Vorst, H.C.M., & Bermond, B. (2001). Validity and reliability of the Bermond–Vorst Alexithymia Questionnaire. *Personality and Individual Differences*, 30(3), 413-434.
[https://doi.org/10.1016/S0191-8869\(00\)00033-7](https://doi.org/10.1016/S0191-8869(00)00033-7)
- Weinstein, A., Maayan, G., & Weinstein, Y. (2015). A study on the relationship between compulsive exercise, depression and anxiety. *Journal of Behavioral Addictions*, 4(4), 315-318. <https://doi.org/10.1556/2006.4.2015.034>

Wise, T., Mann, L., & Shay, L. (1992). Alexithymia and the five-factor model of personality.

Comprehensive Psychiatry, 33(3), 147-151. <https://doi.org/10.1016/0010->

440x(92)90023-j

World Health Organization (2010). *Global recommendations on physical activity for health*.

World Health Organization. <https://apps.who.int/iris/handle/10665/44399>

Table 1

Descriptive Statistics and Intercorrelations Among Variables (N = 99)

| Variable | <i>M (SD)</i> | 1 | 2 | 3 | 4 | 5 | 6 |
|---------------------------|---------------|------|--------|-------|--------|-----|---|
| 1. Age | 24.53 (3.56) | - | | | | | |
| 2. Alexithymia | 56.01 (11.53) | .05 | - | | | | |
| 3. Rash Impulsiveness | 70.23 (11.24) | .09 | .47*** | - | | | |
| 4. Reward Sensitivity | 12.04 (4.98) | -.03 | .27** | .27** | - | | |
| 5. Punishment Sensitivity | 13.92 (5.16) | -.08 | .17 | .29** | .50*** | - | |
| 6. Exercise Dependence | 54.93 (19.27) | -.02 | .53*** | .31** | .28** | .03 | - |

$p < .05$. ** $p < .01$. *** $p < .001$.

Table 2

Hierarchical Multiple Regression on Exercise Dependence Symptoms with sex and scores on rash impulsiveness (BIS-11), sensitivity to reward (SR), sensitivity to punishment (SP), and alexithymia (TAS-20) as predictors.

| Predictor | ΔR^2 | β | <i>B</i> | <i>SE B</i> | 95% CI for <i>B</i> |
|-----------|--------------|---------|----------|-------------|---------------------|
| Step 1 | .02 | | | | |
| Sex | | .13 | 5.18 | 3.89 | [-2.54, 12.91] |
| Step 2 | .15*** | | | | |
| Sex | | .09 | 3.50 | 3.76 | [-3.98, 10.97] |
| BIS-11 | | .29** | .50 | .17 | [.16, .84] |
| SR | | .27* | 1.05 | .44 | [.18, 1.92] |
| SP | | -.18 | -.66 | .42 | [-1.49, .17] |
| Step 3 | .33*** | | | | |
| Sex | | .01 | .18 | 3.48 | [-6.73, 7.09] |
| BIS-11 | | .08 | .14 | .17 | [-.20, .48] |
| SR | | .22* | .84 | .40 | [.05, 1.63] |
| SP | | -.18 | -.67 | .38 | [-1.42, .08] |
| TAS-20 | | .47*** | .78 | .17 | [.45, 1.11] |

Note. SE B = standard error of unstandardized coefficient; CI = confidence interval.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Figure 1. Unmediated and mediated models of the association between rash impulsivity and exercise dependence: support for full mediation by alexithymia. Standardised beta coefficients are shown for all paths. Sex, SR, and SP were controlled as covariates.

