

PHYSICAL ACTIVITY AND BODY MASS INDEX IN COLOMBIAN SCHOOLCHILDREN. MULTICENTER STUDY

José Armando Vidarte Claros*
Consuelo Vélez Álvarez**
José Hernán Parra Sánchez***

Recibido en febrero 11 de 2018, aceptado en mayo 09 de 2018

Citar este artículo así:

Autor. Physical activity and body mass index in Colombian schoolchildren. Multicenter study. *Hacia Promoc. Salud.* 2019; 24(1): 44-55. DOI: 10.17151/hpsal.2019.24.1.5



Abstract



Objective: To establish the relationship between levels of physical activity and body mass index in Colombian students. **Materials and methods:** Correlational descriptive study, performed with 2,651 children between 10-12 years of age, educated in both public and private institutions in six Colombian cities. Prior informed consent was signed by parents, and permission was given by student participants. A survey was applied, in order to collect sociodemographic variables, as was a physical activity form and lifestyle questionnaire (PAQ-C) **Results:** There was a 1:1 male-female ratio. Of the participants in the study, 36.1% were 10-year-old children. In 87% of overweight participants, physical activity levels were evaluated as low or very low. Of those who had normal body mass indexes, one in 10 performed moderate physical activity, and in underweight children, 81.6% had low or very low physical activity levels. Statistically significant differences in physical activity levels by both sex and age $p < 0.05$ were revealed. Low levels of significant inverse correlation were found between physical activity and body mass index (-0.059 for the group total, and -0.070 and -0.046 for boys and girls, respectively). **Conclusions:** Physical activity and body mass index are significantly, differentially, and inversely associated. Physical activity should be promoted in children, as it directly impacts their body mass index, and both variables predict adult cardiovascular health.

Key words

Motor activity, sedentary lifestyle, child, body mass index, school health (*Source: MeSH, NLM*).

* Licenciado en Educación Física. PhD en Ciencias de la Actividad Física y el Deporte. Universidad Autónoma de Manizales. Manizales, Colombia. Autor para correspondencia. Correo electrónico: jovida@autónoma.edu.co.  orcid.org/0000-0002-7982-3848 

** Enfermera. Doctora en Salud Pública. Universidad Autónoma de Manizales. Manizales, Colombia. Correo electrónico: cva@autonoma.edu.co.  orcid.org/0000-0001-7274-7304 

*** Contador Público. Mg en Ciencias Económicas. Universidad Nacional de Colombia. Manizales, Colombia Correo electrónico: jhparrasa@unal.edu.co.  orcid.org/0000-0003-0070-2718 



ACTIVIDAD FÍSICA E ÍNDICE DE MASA CORPORAL EN ESCOLARES COLOMBIANOS. ESTUDIO MULTICÉNTRICO

Resumen

Objetivo: establecer la relación entre los niveles de actividad física y el índice de masa corporal en escolares colombianos. **Materiales y métodos:** estudio descriptivo correlacional, realizado en 2651 niños entre 10 y 12 años escolarizados en instituciones públicas y privadas de seis ciudades de Colombia, previa firma del consentimiento informado de los padres y asentimiento de los estudiantes. Se aplicó una encuesta para variables sociodemográficas, el formato de práctica de actividad física y cuestionario de estilo de vida (PAQ-C) **Resultados:** la relación hombre mujer fue de 1:1, el 36,1% fueron niños de 10 años, en el 87% de los participantes con sobrepeso la actividad física fue evaluada como baja y muy baja. En quienes presentaron índice de masa corporal normal, 1 de cada 10 realizan actividad física moderada, en los niños de bajo peso el 81,6% realizan actividad física baja y muy baja. El nivel de actividad física presentó diferencias estadísticamente significativas por sexo y edad $p < 0,05$. Se encontró correlación baja e inversa, significativa entre la actividad física y el índice de masa corporal (-0,059 para el grupo total y -0,070 y -0,046 para niños y niñas respectivamente). **Conclusiones:** la actividad física y el índice de masa corporal se asociaron de manera significativa y diferencial, la relación encontrada fue inversa. La actividad física debe promoverse en los niños, ya que ésta impacta directamente en su índice de masa corporal, y ambas variables son variables predictoras de problemas cardiovasculares en la edad adulta.

Palabras clave

Actividad motora, estilo de vida sedentario, niño, índice de masa corporal, salud escolar (*Fuente: DeCS, BIREME*).

ATIVIDADE FÍSICA E ÍNDICE DE MASSA CORPORAL EM ESCOLARES COLOMBIANOS ESTUDO MULTICÊNTRICO

Resumo

Objetivo: estabelecer a relação entre os níveis de atividade física e o índice de massa corporal em escolares colombianos. **Materiais e métodos:** estudo descritivo correlacional, realizado em 2651 crianças entre 10 e 12 anos escolarizados em instituições públicas e particulares de seis cidades de Colômbia, previa assinatura do consentimento informado dos pais e aprovação dos estudantes. Aplicou-se uma enquete para variáveis sócio demográficas, o formato de prática de atividade física e questionário de estilo de vida (PAQ-C) **Resultados:** a relação homem mulher foi de 1:1, o 36,1% foram crianças de 10 anos, no 87% dos participantes com sobrepeso a atividade física foi avaliada como baixa e muito baixa. Em quem apresentaram índice de massa corporal normal, 1 de cada 10 realizam atividade física moderada, nas crianças de baixo peso o 81,6% realizam atividade física baixa e muito baixa. O nível de atividade física apresentou diferenças estatisticamente significativas por sexo e idade $p < 0,05$. Encontrou-se correlação baixa e inversa, significativa entre a atividade física e o índice de massa corporal (-0,059 para o grupo total e -0,070 e -0,046 para crianças e meninas respectivamente). **Conclusões:** a atividade física e o índice de massa corporal se associaram de maneira significativa e diferencial, a relação encontrada foi inversa. A atividade física deve promover-se nas crianças, já que está impacta diretamente em seu índice de massa corporal, e ambas variáveis são variáveis de previsão de problemas cardiovasculares na idade adulta.

Palavras chave

Atividade motora, estilo de vida sedentário, criança, índice de massa corporal, saúde escolar (*Fonte: DeCS, BIREME*).

INTRODUCTION

Physical activity plays an important role in the prevention of non-communicable chronic disease and lowers obesity risk. It regulates energy balance and preserves or maintains lean mass instead of fat mass (1, 2).

Generally, childrens' activities, such as playing, and the development of their fundamental movement patterns of movement, which include jumping, running, dancing, and riding a bicycle, provide a wide variety and large number of vigorous activities from which to choose. However, certain studies have revealed that, because the level of physical activity during free time, at these ages, has been significantly reduced (4), the above activities are no longer regularly performed (3).

The prevalence of excess weight and obesity in the United States in the last decade affected 37.1% of children aged 6-11 years old, and 34% of those aged 12-19 years old (5). Childhood obesity is a condition related to an increase in HBP, hypercholesterolemia, hyperinsulinemia, and metabolic syndrome, a decrease in growth hormone, respiratory changes, and bone problems (5), in addition to psychological and social changes (6). Further, obesity frequently continues into adult life. The numbers indicate that 40% of children who are overweight at seven years old continue on to become obese adults (7), and that physical activity is among the causal factors in their development.

Today's lifestyles, which keep children inactive, are predisposing factors to excess weight gain (less time dedicated to physical activity, decrease in aerobic capabilities, and more time dedicated to sedentary activities such as watching television, playing videogames, and surfing the internet) (8, 9).

In Colombia, the National Survey of Nutritional Status (NSNS) (2010) indicates that virtually half of the adult population is heavy for their

stature, 32.3% is overweight, and 13.8% is obese, according to both American tables and WHO criteria (10). However, Colombian studies offer little information on the prevalence of excess weight and obesity in the child population, and their relationship with factors such as eating habits, physical activity levels, physical condition, and sedentary activities. Furthermore, there is minimal data regarding those modifiable factors which contribute to excess weight gain, obesity, inactive lifestyles, or low weight, which are fundamental for the implementation of intervention measures aimed at the prevention of excess weight, obesity, low weight, and the consequences thereof (10, 11).

Body Mass Index (BMI) is a fundamental indicator, in epidemiological studies, which allows for the up-to-date definition of a child or adult's nutritional status as underweight, overweight, or obese (12, 13). With its universal use, the BMI has been adopted by the WHO as an anthropometric norm which serves as a reference in the monitoring and classification of growth, development, and health status, associated with either a lack or excess of nutrients (14, 15). To this effect, the present investigation seeks to establish the relationship between physical activity and the BMI in Colombian school children by way of a multicenter study.

MATERIALS AND METHODS

Design

Descriptive study with a correlational phase, carried out with adolescents between 10-12 years of age, educated in public and private institutions in the cities of Manizales (Caldas), Riohacha (Guajira), Popayán (Cauca), Guadalajara de Buga (Valle del Cauca), Yarumal (Antioquia), and Zarzal (Valle del Cauca), Colombia. A random, stratified sample was used, with uniform allocation. A margin of error of 0.3 kg/m² was employed, as was 95% reliability, for a total sample of 2,651 children distributed

thus: (Manizales, 448, Popayán 442, Riohacha 441, Yarumal, 440, Buga 440, and Zarzal 440). For the variance estimate, a study entitled, “Exposición a pantallas, sobrepeso y desacondicionamiento físico en niños y niñas” was used as base (16). Inclusion criteria for enrollment in the educational institutions were as follows: being between 10-12 years old, participating voluntarily, having the informed consent of a parent or guardian, and possessing student permission.

Procedure

Once the final version of the sociodemographic variable form had been obtained, authorization was requested from the municipalities, ministries of education, and educational institutions, in order to administer said form. This was applied both by investigators and interviewers who had been previously trained by investigators, and with whom the process was repeated several times, in order to reduce bias in information collection. They then proceeded to present the study, explain the objective, and accompany participants in filling out the collection form.

Measurements

Observation techniques were used together with the survey. The tools employed included the sociodemographic variables survey form, with structured questions, and the physical activity form and PAQ-C lifestyle questionnaire. Said questionnaire has been validated for Colombia (17).

The PAQ-C measures physical activity levels, from moderate to vigorous, in the last seven days of the school year. It consists of ten questions with multiple-choice answers, on a scale of five points. These questions inquire about the activities performed during one's free time, the physical activities performed in physical education class, during recess, lunch, just after school, in the afternoons, and on the weekends. It also elicits information about physical activity performed on

the weekend and the frequency with which physical activity is performed on each day of the week (18, 19). To calculate the final score, the average of the first nine questions is calculated. Question 10 is not used as part of the total score. A score of one indicates low levels of physical activity, and five indicates high levels of physical activity. Internal consistency was determined using the Cronbach's alpha coefficient and the reliability test, by way of the intraclass correlation coefficient. The internal consistency of the questionnaire revealed a Crocker reliability of 80 (19) and internal consistency values between 0.79 and 0.89 (20). The PAQ-C is thus a questionnaire which contributes to the evaluation of physical activity in boys and girls during the school year.

Through use of the Tanita BC-543 scale, participant weight was obtained, and size information was gathered using a measuring rod (Seca portable stadiometer). During this process, the children were barefoot, and minimally clothed. Weight was attained with 0.1 kg precision, and height with 0.1 cm precision. The scale was calibrated after each 100 measurements.

For management of the information obtained, the BMI was classified in accordance with that established by the WHO, 2004 (21), which provides standard deviations (SD) that differentiate those who are: at risk of malnourishment (-1.9 SD, -1 SD), malnourished (<-2 SD), eutrophic (-0.9 SD, 1 SD), overweight (1 SD, 1.9 SD), obese (2 SD, 2.9 SD), morbidly obese (>3 SD), at risk of small stature (-1.9 SD, -1 SD), chronically malnourished, or small for their age (size <-2 SD).

Statistical analysis

The data were processed and analyzed in the SPSS Version 21.0 program, licensed by the Universidad Autónoma de Manizales. A univariate analysis was performed, which made the presentation of data in tables of absolute and relative frequencies possible. Central tendency and variability or

dispersion measurements were calculated next. For the bivariate analysis, associations were established between study variables. In order to determine the linear relationship between variables, the Pearson correlation coefficient was used. Statistical significance was established with <0.05 p values.

Ethical considerations

The entire study was framed within the current provisions for the protection of those human subjects who participated in the investigation. These are contained in the Colombian Ministry of Health's Resolution 8430 of 1993, and the Helsinki Declaration, classifying study "with minimal risk", in accordance with letter b of Article 11 (22). The approval of the Universidad Autónoma de Manizales' Investigation Ethics Committee was also obtained.

RESULTS

Exactly 2,651 schoolchildren between the ages of 10-12 years old participated in this study. The male-female ratio was 1:1, with 36.1% 10-year-old, 32.7% 11-year-old, and 31.2% 12-year-old boys.

In order to compare physical activity levels by both sex and age, previous verification of normality and homoscedasticity assumptions was necessary. It was shown that normality was not present, and that statistically significant differences occur between the sexes, $p<0.05$ according to the U Mann-Whitney test. Statistically significant differences were revealed, by age, in the average of the three groups, with significantly fewer differences than in the group of 10-year-olds, which had the lowest physical activity level score obtained, ($p<0.05$) according to the Kruskal-Wallis test (Table 1).

Table 1. Difference of means of physical activity by sex and age.

Variable	Half	n	D.E	Test Statistics	
Sex				U de Mann-Whitney	p Value
Man	2,5129	1308	0,51745		
Woman	2,4157	1343	0,56309	785741,5	0,000*
Total	2,4637	2651	0,54313		
Age				Kruskal Wallis	p Value
10 years	2,4042	958	0,57530		
11 years	2,5253	867	0,51132	23,775	0,000*
12 years	2,4680	826	0,53015		
Total	2,4637	2651	0,54313		

Source: Author elaboration.

(*) Statistically significant association

In both the male and female groups, the greatest percentage of participants presented normal BMIs, although statistically significant differences were found between these variables ($p < 0.05$), even though the coefficient value supposes that there is no dependence relationship there between.

When comparing BMI averages by age and sex, it was observed that, in all age groups, and with both

sexes, the greatest percentage of schoolchildren enjoyed normal BMIs. It is important to highlight that, in the group of 10 and 11-year-olds, malnourished children were present, and the risk of malnutrition in all age groups was higher in males than in females. Statistically significant differences between the sexes were not found in the BMI nutritional state $p < 0.05$ (Table 2).

Table 2. Nutritional classification by BMI, age, and sex.

AGE (years)	Classification by BMI	SEX				Total N	X ²	P value
		Males		Females				
		n	%	n	%			
10	Malnourished	1	0.2	10	2.0	11	8.551*	0.128
	At risk of malnourishment	89	19.3	90	18.1	179		
	Normal	321	69.6	355	71.4	676		
	Overweight	35	7.6	29	5.8	64		
	Obese	9	2.0	7	1.4	16		
	Morbidly obese	6	1.3	6	1.2	12		
	Total	461	100.0	497	1.0	958		
11	Malnourished	1	0.2	2	0.5	3	5.101*	0.404
	At risk of malnourishment	66	15.3	51	11.7	117		
	Normal	288	67.0	321	73.5	609		
	Overweight	55	12.8	46	10.5	101		
	Obese	16	3.7	13	3.0	29		
	Morbidly obese	4	0.9	4	0.9	8		
	Total	430	100.0	437	100.0	867		
12	Malnourished	0	0	0	0	0	5.831*	0.212
	At risk of malnourishment	46	11.0	39	9.5	85		
	Normal	301	72.2	288	70.4	589		
	Overweight	53	12.7	68	16.6	121		
	Obese	16	3.8	10	2.4	26		
	Morbidly obese	1	0.2	4	1.0	5		
	Total	417	100.0	409	100.0	826		
General total		1308		1343		2651		

Source: Author elaboration

(*) Chi Square statistic

Table 3. Association between physical activity and participant BMI.

Classification of physical activity	Nutritional classification by BMI (kg/m ²)														X ²	P value	Coefficient
	Malnourished		At risk of malnourishment		Normal		Overweight		Obese		Morbidly obese		Total				
	n	%	n	%	n	%	n	%	n	%	n	%	n	%			
Very low	9	64.3	76	19.9	381	20.3	68	23.8	14	19.7	6	24.0	554	20.9			
Low	4	28.6	233	61.2	1173	62.6	180	62.9	47	66.2	17	68.0	1654	62.4	22.779	0.012	-0.019
Moderate	1	7.1	72	18.9	320	17.1	38	13.3	10	14.1	2	8.0	443	16.7			
Total	14	100.0	381	100.0	1874	100.0	286	100.0	71	100.0	25	100.0	2651	100.0			

Source: Author elaboration.

The correlation coefficient between physical activity and BMI was quite low. In other words, the greater the physical activity, the lower the BMI. This association was statistically significant in males ($p < 0.05$) (Table 4).

Table 4. Correlation matrix.

P value statistic		Physical activity	BMI
Physical activity	Pearson correlation	1	-0.059
	P value		0,002
Sex of child evaluated		Physical activity	BMI
Male	Pearson correlation	1	-0.070
	P value		0.011
Female	Pearson correlation	1	-0.046
	P value		0.09

Source: Author elaboration.

DISCUSSION

In the population studied, 70.7% of the children were of normal weight, and 10.8% were overweight, data which contrasts with the results of Bojórquez's investigation, in which 7.3% of children who participated were underweight, and 42% were overweight and obese (23). It also contrasts with that reported by Aviña et al., whose study reveals that, in the schoolchildren studied, 4.05% were underweight, and 45.3% were either overweight or obese (24). Similarly, Fernández finds that 3.7% of the children in their study were underweight, 79.5% were of normal weight, 7.9% were at risk of excess weight, and 8.8% overweight in their study (25).

Regarding sex and age, statistically significant differences were found with the degree of physical activity. Similar results were reported in a study of 3,742 children, with a significant difference in favor of males over females (26.27%). In another study, carried out in Madrid, which calculated the number of minutes that children dedicated to physical activity during the week, in a population of 253 students between 10-13 years old, it was found that the time that boys spent being physically active was significantly higher than that of girls ($p < 0.001$) (28). Yet another study reported that just 46.1% of girls performed extracurricular physical sporting activities, in contrast to 60.3% of boys, (29). Similarly, Ruiz evaluated 120 schoolchildren from 12-14 years of age, and found a significant difference between sexes $p = 0.000$ (30). This reveals that girls perform less physical activity than boys during both pre-adolescence and adolescence, which could affect their motor development (31). More recent studies evaluated the possible differences between anthropometric variables and the physical condition of schoolchildren in primary school and showed a difference in the level of physical activity between sexes (32).

In the population studied in the present investigation, the higher the amount of physical

activity, the lower the BMI. This data differs from the results of by Chacón et al., who report that, about physical activity and the degree of obesity, the data do not correlate ($p = 0.595$), and that statistically significant differences were not found (33). In this same sense, Albán et al. (2011) did not discover any statistically significant difference between BMI and the type of physical activity (34). Additionally, one study carried out in Belgium, with 2,517 students ranging from 5-13 years old, concluded that the weight of the child negatively influenced future physical activity levels, and especially affected certain motor skills, such as gross motor coordination and vice-versa (35). Similarly, transversal studies have demonstrated negative correlations between BMI, gross motor skills, and coordination in the normal development of children and young adults. One investigation performed in Portugal, on children between 10-14 years old, revealed that all correlations between the motor coefficient (MC) and BMI were negative, and varied between 0.05 and 0.49 (36). The authors recognize that those restrictions which are inherent to descriptive study may limit the present investigation. However, the sample framework suggests the presence of important evidence regarding the problem studied. From these results, new analytical and/or experimental studies, that permit a focus on the suggested causal relationships, may be formulated.

CONCLUSION

The fact that analysis of the relationship between physical activity and BMI in schoolchildren reveals that this association, apart from being statistically significant, is also inverse, should motivate physical activity in family, school, and social contexts. This is because it directly impacts their nutritional status. Both variables are considered to be predicting variables of adult cardiovascular problems, which, in the context of world public health, occupy the first places in causes of morbidity and mortality in different population groups. Similarly, the results of this

investigation should be considered in training curriculums, in order to incentivize professionals in the importance of interdisciplinary work for the promotion of physical activity from childhood.

ACKNOWLEDGEMENTS

The authors thank the students from the Universidad Autónoma de Manizales' Master's program in

Comprehensive Athlete Treatment (VII cohort), as well as those children and educational institutions who participated in this study.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest in the investigation presented.

BIBLIOGRAPHIC REFERENCES

1. Santana JC. Impacto social del ejercicio físico en la prevención y tratamiento de enfermedades crónicas no transmisibles. *Revista Conrado* [seriada en línea]. 2013; 9 (36): 38-42.
2. Salomon JA, Vos T, Hogan DR, Gagnon M, Naghavi M, Mokdad A, et al. Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2129-2143.
3. Cordova A, Villa G, Sureda A, Rodriguez-Marroyo JA, Sánchez-Collado MP. Actividad física y factores de riesgo cardiovascular de niños españoles de 11-13 años. *Rev Esp Cardiol*. 2012; 65(7):620-626.
4. Trost S. Medición de la actividad física en los niños y adolescentes. *Lifestyle Am J Med*. 2007; 1(4):299 -314.
5. Sassi F, Devaux M. OECD Obesity Update 2012. Disponible en: <http://www.oecd.org/dataoecd/1/61/49716427.pdf>.
6. Pompa E, Montoya BI. Evaluación de la manifestación de ansiedad y depresión en niños con sobrepeso y obesidad en un campo de verano. *Psicología y Salud*. 2011; 21(1): 119-124.
7. Acosta JA, Medrano G, Duarte G, González SR. Malos hábitos alimentarios y falta de actividad física. Principales factores desencadenantes de sobrepeso y obesidad en los niños escolares. *CULCyT*. 2014; 54(1):81-90.
8. Dos Santos MF, Osuna CA, Bernal J. Las horas pantalla se asocian al consumo de alimentos de elevada densidad calórica, sobrepeso, obesidad y sedentarismo en niños venezolanos. *Rev Esp Nutr Comunitaria*. 2014; 20(3):78-84.
9. Tremblay M, Leblanc A, Janssen I, Kho M, Hicks A, Murumets K et al. Canadian sedentary behavior guidelines for children and youth. *Appl Physiol Nutr Metab*. 2011; 36 (1): 59-64.
10. República de Colombia Ministerio de la Protección Social Instituto Colombiano de Bienestar Familiar. Encuesta Nacional de la situación nutricional en Colombia ENSIN. [Internet]; [citado 3 abril 2016]. Disponible en: <http://www.minsalud.gov.co/.../Base 20de 20datos 20ENSIN 20-20.2010>.
11. Zayas V, Benítez M, Frey NG, Miranda D. (2016). Diagnóstico nutricional y su relación con la merienda escolar. Nutritional diagnosis and its relationship with school meals. *Revista Virtual de Posgrado*. 2016; 1(1): 15-26.
12. Piña CE, Fernández ML, Fonte N, Piña LK. Instrumento predictor de obesidad en adolescentes en el Policlínico Universitario “Manuel González Díaz”, 2013-2014, *Boletín Médico del Hospital Infantil de México*. 2015; 72 (1): 34-44.
13. Fernandes T, Gallo P, Advíncula A. Anthropometric assessment in preschool children in Mogi-Guacu, State of Sao Paulo: a support for public health policies. *Rev. Bras. Saude Mater. Infant*. 2006, 6 (2): 217-222.
14. González R, Llapur R, Díaz M, Illa MR, Yee E, Pérez D. Estilos de vida, hipertensión arterial y obesidad en adolescentes. *Rev Cubana Pediatr* [Internet]. 2015 Sep [citado 2016 Abr 03]; 87(3): 273-284. Disponible en: http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S0034-75312015000300003&lng=es.
15. Lopes VP, Maia JAR, Rodrigues LP, Robert Malina R. Motor coordination, physical activity and fitness as predictors of longitudinal change in adiposity during childhood. *European Journal of Sport Science*. 2012; 12 (4): 384-391.

16. Duque IL y Parra JH. Exposición a pantallas, sobrepeso y desacondicionamiento físico en niños y niñas. *Rev. latinoam. cienc. soc. niñez juv.* 2012;10 (2): 971-981.
17. Herazo Y, Domínguez R. Confiabilidad del cuestionario de actividad física en niños colombianos. *Rev. salud pública.* 2012; 14 (5): 802-809.
18. Martínez GD, Martínez DV, Pozo T, Welk GJ, Villagra A, Calle ME, et al. Fiabilidad y validez del cuestionario de actividad física PAQ-A en adolescentes españoles. *Rev. Esp Salud Pública.* 2009; 83 (3): 427-439.
19. Crocker PR, Bailey DA, Faulkner RA, Kowalski KC, McGrath R. Measuring general levels of physical activity: preliminary evidence for the Physical Activity Questionnaire for Older Children. *Med Sci Sports Exerc.* 1997; 29:1344-1349.
20. Janz KF, Lutuchy EM, Wenthe P, Levy SM. Measuring Activity in Children and Adolescents Using Self-Report: PAQ-C and PAQ-A. *Med. Sci. Sports Exerc.* 2008; 40 (4): 767-772.
21. World Health Organization. AnthroPlus for personal computers manual: Software for assessing growth of the world's children and adolescents. Geneva: World Health Organization, 2007. Disponible en: <http://www.who.int/childgrowth/software/en/>.
22. Colombia. Ministerio de Salud. Resolución No. 008430 de 1993, por la cual se establecen las normas científicas, técnicas y administrativas para la investigación en salud [Internet]; Colombia, 1993 [citado 13 de junio de 2013]. Disponible en: <http://comitedeetica.clinicalili.org/docs/RESL8430.pdf>.
23. Bojórquez CI, Angulo CM, Reynozo L. Factores de riesgo de hipertensión arterial en niños de primaria. *Psicología y Salud.* 2011; 21(2), 245-252.
24. Aviña MA, Castillo O, Vázquez F, Perales A, Alemán S. Evaluación nutricional de escolares en una ciudad fronteriza entre Estados Unidos y México. *Rev Med Chile.* 2016; 144(3):347-354.
25. Fernández J, Ruíz F. Estudio transversal de crecimiento de los escolares bogotanos: valores de estatura, peso e índice de masa corporal de los siete a los dieciocho años. *Rev Univ Salud.* 2012; 15(1): 21-41.
26. Lopes V, Maia J, Silva R, Seabra A y Morais F. Estudo do nível de desenvolvimento da coordenação motora da população escolar (6 a 10 anos de idade) da Região Autónoma dos Açores. *Rev Port Cien Desp.* 2003; 3 (1):47-60.
27. Torralba MA, Vieira MB, Lleixà T y Gorla JI. Evaluación de la Coordinación Motora en Educación Primaria de Barcelona y Provincia. *Rev int med cienc act fis deporte.* 2014; 16(62):355-371.
28. Montil M, Barriopedro MI, Oliva J. El sedentarismo en la infancia. Los niveles de actividad física en niños/as de la Comunidad Autónoma de Madrid. *Apunts, Educación Física y Deportes* 2005; 4(82): 5-11.
29. Luengo C. Actividad físico-deportiva extraescolar en alumnos de primaria. *Rev int med cienc act fis deporte.* 2007; 7(27):174-184.
30. Ruiz LM, Gómez M, Jiménez PJ, Ramón I, Peñaloza R. ¿Debemos preocuparnos por la coordinación corporal de los escolares de la Educación Secundaria Obligatoria? *Rev. Pediatría Ante Primaria.* 2015; 17(66): e109-e116.
31. Santos LB. Desarrollo de las habilidades motoras fundamentales en función del sexo y del índice de masa corporal en escolares. *CuadPsicol del Deport* 2013; 13:63-71.
32. Luque GT, Rivera EC, Sánchez AJ, Sánchez ML. Niveles de condición física de escolares de educación primaria en relación a su nivel de actividad física y al género. *Retos: nuevas tendencias en educación física, deporte y recreación.* 2014;(25 (1):17-22.

33. Chacón R, Espejo T, Cabrera Á, Castro M, López JF, Zurita F. “Exergames” para la mejora de la salud en niños y niñas en edad escolar: estudio a partir de hábitos sedentarios e índices de obesidad. *Rev. latinoam. tecnol. educ.* 2015; 14(2), 39-50.
34. Albán A, Álvarez M, Bacuilima P. Evaluación del estado nutricional en adolescentes de 15 a 18 años, que asisten al Colegio Julio María Matovelle, Cantón Cuenca. México; 2011: Universidad de Cuenca. Disponible en: <http://dspace.ucuenca.edu.ec/bitstream/123456789/3454/1/MED89.pdf>
35. D’Hondt E, Deforche B, Gentier I, Verstuyf J, Vaeyens R, de Bourdeaudhuij I, et al. A longitudinal study of gross motor coordination and weight status in children. *Obesity.* 2014; 22(6):1505–1511.
36. Lopes VP, Stodden DF, Bianchi MM, Maia J a R, Rodrigues LP. Correlation between BMI and motor coordination in children. *J Sci Med Sport.* 2012; 15(1):38-43.