



ISSN: (Print) 1942-0889 (Online) Journal homepage: <https://www.tandfonline.com/loi/kcib20>

Anesthetics disrupt brain development via actions on the mTOR pathway

Jing Xu, Eunchai Kang & C. David Mintz

To cite this article: Jing Xu, Eunchai Kang & C. David Mintz (2018) Anesthetics disrupt brain development via actions on the mTOR pathway, *Communicative & Integrative Biology*, 11:2, 1-4, DOI: [10.1080/19420889.2018.1451719](https://doi.org/10.1080/19420889.2018.1451719)

To link to this article: <https://doi.org/10.1080/19420889.2018.1451719>



© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 09 Apr 2018.



Submit your article to this journal [↗](#)



Article views: 680



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 8 View citing articles [↗](#)

Anesthetics disrupt brain development via actions on the mTOR pathway

Jing Xu ^{a,b}, Eunchai Kang^c and C. David Mintz^{b*}

^aDepartment of Anesthesiology, Second Affiliated Hospital of Xi'an Jiaotong University School of Medicine, Xi'an, Shaanxi, China; ^bDepartment of Anesthesiology, The Johns Hopkins University School of Medicine, Baltimore, MD, USA; ^cDepartment of Neuroscience and Mahoney Institute for Neurosciences, Perelman School for Medicine, University of Pennsylvania, Philadelphia, PA, USA

ABSTRACT

Experiments conducted in non-human primates have recently provided new evidence supporting a longstanding concern that exposure to general anesthesia during late intrauterine life or early childhood can cause lasting cognitive deficits through harmful effects on brain development. The mammalian target of rapamycin (mTOR) signaling system plays a key role in both normal brain development and in a wide range of developmental disorders that are characterized by cognitive deficits. Intriguingly, our recently published work shows that anesthetics can chronically alter mTOR signaling in the hippocampal dentate gyrus and that normalization of mTOR signaling can prevent anesthesia-induced perturbation of structure and function. In this addendum, we briefly discuss the putative role of mTOR in developmental anesthetic neurotoxicity.

ARTICLE HISTORY

Received 31 December 2017
Accepted 8 March 2018

KEYWORDS

mTOR; cell signaling;
anesthesia; neurogenesis;
neurotoxicity

1. Introduction

The United States Food and Drug Administration has recently required that 12 commonly used anesthetic and sedative agents with mechanisms of action on NMDA and GABA receptors carry labels warning that repeated or lengthy exposure to these drugs between the third trimester and the first three years of life may result in adverse consequences for brain development [1]. These safety concerns are based on a combination of findings from human subjects research and animal model studies including rodents and primates. While the only prospective studies of pediatric patients conducted to date indicate that a short, single exposure to general anesthesia does not result in cognitive deficits [2,3], several large retrospective studies have found an association between anesthesia/surgery exposure in young children and worsened cognitive function at later ages [4–6]. The interpretation of the medical relevance of these studies is complex at best, and it may never be possible to design an ethical clinical study that answers fundamental questions about what kind of harm may come from anesthetic exposure. Multiple independent investigations have concluded that early postnatal exposure to general anesthesia in rodents can cause lasting deficits on behavioral tests on learning and memory e.g. [7–10]. It should be noted that some

studies in rodents have found no effects of anesthetics on cognitive function [11] and others have even found neuroprotective effects [12,13]. There is considerable variability between models of early developmental anesthesia which complicates the interpretation of the rodent literature, and any rodent model is potentially complicated by the need for high doses to cause general anesthesia, the difficulty in selecting an appropriate control condition, and the possibility of physiologic perturbation such as hypoxia, hypercarbia, hypotension, and hypothermia. The most compelling evidence in support of developmental anesthetic neurotoxicity comes from a recent study in non-human primates by Baxter and co-workers that showed clear deficits in visual recognition memory resulting from clinically relevant exposures to general anesthesia. These data are free from the confounds of surgery and co-morbid disease which complicate the interpretation of clinical investigations and the model for anesthesia exposure much more closely resembles human pediatric general anesthesia than the published rodent models [14]. The mechanisms by which general anesthetics might disrupt brain development have yet to be clearly defined. Early work on this topic identified molecular evidence of upregulation of pro-apoptotic pathways by general anesthetic drugs in the developing brain [15–17], but

CONTACT C. David Mintz  cmintz2@jhmi.org  The Johns Hopkins University School of Medicine, 720 Rutland Ave, Ross Bldg 370, Baltimore, MD, 21205, USA.

Addendum to: Kang E, Jiang D, Ryu YK, et al. Early postnatal exposure to isoflurane causes cognitive deficits and disrupts development of newborn hippocampal neurons via activation of the mTOR pathway. *PLoS Biol* 2017; 15:e2001246.

© 2018 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

more recent studies have cast doubt on whether cell death is the principal mechanism of damage in anesthetic neurotoxicity [18–20]. Thus, it is of interest to consider other potential mechanisms of injury, particularly those which relate developmental anesthetic neurotoxicity to better-studied disorders of brain development. In this addendum to our recently published work, Kang et al 2017, we briefly discuss mTOR dysregulation in neurodevelopmental disease and we place our findings in the context of two other recent studies related to anesthetic neurotoxicity.

2. The mTOR pathway and neurodevelopmental disease

The mTOR protein is a highly conserved protein kinase which is the key participant in two protein complexes, mTOR complex 1 and 2 that play a critical role in growth, metabolism, and homeostasis in almost every eukaryotic cell type. mTOR complexes regulate the integrated response of multiple signaling pathways to a wide array of intracellular sensing cues and extracellular signals [21]. In the developing nervous system, mTOR plays a key role in a wide variety of normal processes, including neural stem cell regulation, cellular migration, and dendrite, axon and synapse development [22]. Thus, it is unsurprising that perturbations in mTOR signaling are potentially pathological, although given the complexity of the pathway it can be difficult to determine whether changes in mTOR signaling are the cause of pathology or a consequence of it. One of the best-studied examples of the role of aberrant mTOR activity in the pathology of neurodevelopmental disease is in tuberous sclerosis, a multisystem genetic disorder that can include intellectual disability, developmental delay, and autism spectrum disorder (ASD). Tuberous sclerosis results from mutations in TSC1 and TSC2, which are direct negative regulators of the mTORC1 complex [23,24]. Transgenic mouse models of tuberous sclerosis such as *Tsc2*^{+/-} and *Tsc1*^{+/-} exhibit increased activity in the mTOR pathway, which is accompanied by deficits in long-term potentiation and spatial learning, which can be reversed by chronic treatment with a rapamycin, a mTOR pathway inhibitor [25]. Another well-characterized example is Fragile X syndrome, the most common developmental cause of mental retardation and a common cause of ASD, which results from the transcriptional silencing of the FMR1 gene. Elevated mTOR signaling is observed in the *Fmr1*^{-/-} mouse model of Fragile X Syndrome e.g.[26], however, the connection between *Fmr1* and mTOR has not been

fully elucidated. While later treatment with rapamycin does not necessarily reverse pathology associated with the *Fmr1* knockout [26], a double knockout that also deletes S6 Kinase does not show the synaptic and behavioral deficits associated with *Fmr1*^{-/-} alone [27]. A role for mTOR has also been postulated for a variety of neuropsychiatric disorders, including other causes of ASD, schizophrenia, drug addiction and depression [28]. While changes in the mTOR pathway in these disorders are not simple or uniform, the trend is generally that inappropriately increased activity is associated with pathology.

3. mTOR as a target in developmental anesthetic neurotoxicity

Because the mTOR pathway is involved in many processes brain development, any pharmacologic action on this signaling pathway has the potential to cause a pathologic state. The first report that a widely used general anesthetic acts on the mTOR pathway came from a study conducted in the human umbilical vein endothelial cell line, which found that isoflurane exposure increased levels of phospho-Akt, phospho-mTOR, and phospho-GSK3B and that effects of isoflurane on hypoxia-inducible factor 1 can be blocked with rapamycin [29]. Two manuscripts published this past year have provided evidence that increased activity in the mTOR pathway may contribute to the phenotype of developmental anesthetic neurotoxicity. Na and co-workers found that rapamycin treatment prevented the loss of chemotaxis that results from developmental exposure of *C. elegans* to isoflurane. Based on this data and the results of a forward genetics screen that identified endoplasmic reticulum stress pathways as a target in anesthetic-induced neurotoxicity, that isoflurane acts on mitochondria to create reactive oxygen species that in turn activate the mTOR pathway [30]. In our recently published manuscript, we described a study in a mouse model of developmental anesthetic neurotoxicity which revealed that anesthetic effects on mTOR signaling profoundly disrupted development of newborn dentate gyrus granule cells in the hippocampus. Isoflurane exposure was found to cause a lasting increase in phospho-S6 (pS6) levels that was accompanied by a reduction in mushroom spines. Both the spine loss and behavioral deficits in learning induced by anesthetic exposure could be reversed by treatment with rapamycin. We concluded that isoflurane caused a chronic upregulation of mTOR that disrupted the developmental program in this vulnerable cohort of neurons, which are critical for learning functions [10].

Of note, several previous reports suggest that anesthetics may act to suppress mTOR activity in the hippocampus in rodent models of anesthetic neurotoxicity, a

finding that may or may not be at odds with ours given differences in study design and in what was measured. For example, work in both primary cultured hippocampal neurons and in the intact hippocampus, Li et al. found that isoflurane exposure during development acutely reduced phospho-S6 via inhibition of the Insulin-like growth factor 1(IGF-1)/ Phosphoinositide 3-kinase (PI3K)/ Protein Kinase B (Akt) pathway [31]. Furthermore, they found that that treatment with IGF-1 on the day of exposure and for two days afterward restored performance on behavioral tests of learning that were impaired with isoflurane exposure in their model. There are, of course, differences in the exposure paradigm between this report and ours, but perhaps the most striking difference is that we measured pS6 levels at 12 and 42 days after the exposure, rather than immediately after the exposure. Liu et al., working in an aged rodent model rather than during development, found that sevoflurane exposure resulted in a decrease in phospho-mTOR levels in the hippocampus that persisted for 7 days after exposure [32]. However, no later time points were measured, and intriguingly they found that sevoflurane exposure did not result in any change in pS6 levels between 1 and 7 days of exposure. The model in this study differs very substantially from our own in terms of the aged versus developing hippocampus, but certainly these findings do not support a universal or uniform effect of anesthetics on the mTOR pathway.

It is not year clear how to reconcile the currently published studies currently on the effects of anesthetics on the mTOR pathway. Some of the difficulty results from a broader problem in the field of developmental anesthetic neurotoxicity where there is currently no standard model of anesthetic exposure nor any agreed upon standard for measurable outcomes and time points at which these outcomes are considered relevant. Nevertheless, there is now preliminary evidence suggesting that anesthetics can have unintended and harmful effects on the mTOR pathway, and given the importance of this pathway for brain development, we feel that this new avenue of investigation deserves further consideration.

Author contributions

Jing Xu wrote for the initial draft and participated in revision; Eunchai Kang wrote for the initial draft; C. David Mintz developed the concept and wrote for the initial and revised draft.

Conflicts of interest

The authors do not have any financial, commercial, legal, or professional conflicts of interest related to this addendum to

their previously published manuscript “Isoflurane Causes Cognitive Deficits and Disrupts Brain Circuitry Development via Activation of the mTOR Pathway” on PLoS Biol 2017;7:e2001246.

Funding

This work was supported by an ACCM StAAR Award and NIH 1R01GM120519-01 to C.D.M, NARSAD to E.K., a grant from the Chinese Scholarship Council to J.X.

ORCID

Jing Xu  <http://orcid.org/0000-0002-6004-4713>

References

- [1] FDA Drug Safety Communication: FDA approves label changes for use of general anesthetic and sedation drugs in young children. Available online: <https://www.fda.gov/Drugs/DrugSafety/ucm554634.htm> (Accessed on April 28th, 2017).
- [2] Sun LS, Li G, Miller TL, et al. Association Between a Single General Anesthesia Exposure Before Age 36 Months and Neurocognitive Outcomes in Later Childhood. *Jama*. 2016;315(21):2312–20.
- [3] Davidson AJ, Disma N, de Graaff JC, et al. Neurodevelopmental outcome at 2 years of age after general anaesthesia and awake-regional anaesthesia in infancy (GAS): an international multicentre, randomised controlled trial. *Lancet (London, England)*. 2016;387(10015):239–50.
- [4] Ing C, Hegarty MK, Perkins JW, et al. Duration of general anaesthetic exposure in early childhood and long-term language and cognitive ability. *Br J Anaesth*. 2017;119(3):532–40.
- [5] DiMaggio C, Sun LS, Li G. Early childhood exposure to anesthesia and risk of developmental and behavioral disorders in a sibling birth cohort. *Anesthes Analg*. 2011;113(5):1143–51.
- [6] Hu D, Flick RP, Zaccariello MJ, et al. Association between Exposure of Young Children to Procedures Requiring General Anesthesia and Learning and Behavioral Outcomes in a Population-based Birth Cohort. *Anesthesiology*. 2017;127(2):227–40.
- [7] Jevtovic-Todorovic V, Hartman RE, Izumi Y, et al. Early exposure to common anesthetic agents causes widespread neurodegeneration in the developing rat brain and persistent learning deficits. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2003;23(3):876–82.
- [8] Lee BH, Chan JT, Hazarika O, et al. Early exposure to volatile anesthetics impairs long-term associative learning and recognition memory. *PloS One*. 2014;9(8):e105340.
- [9] Stratmann G, Sall JW, May LD, et al. Beyond anesthetic properties: the effects of isoflurane on brain cell death, neurogenesis, and long-term neurocognitive function. *Anesth Analg*. 2010;110(2):431–7.
- [10] Kang E, Jiang D, Ryu YK, et al. Early postnatal exposure to isoflurane causes cognitive deficits and disrupts

- development of newborn hippocampal neurons via activation of the mTOR pathway. *PLoS Biology*. 2017;15(7):e2001246.
- [11] Rosenholm M, Paro E, Antila H, et al. Repeated brief isoflurane anesthesia during early postnatal development produces negligible changes on adult behavior in male mice. *PloS one*. 2017;12(4):e0175258.
- [12] Leikas JV, Kohtala S, Theilmann W, et al. Brief isoflurane anesthesia regulates striatal AKT-GSK3beta signaling and ameliorates motor deficits in a rat model of early-stage Parkinson's disease. *J Neurochem*. 2017;142(3):456–63.
- [13] Jiang M, Sun L, Feng DX, et al. Neuroprotection provided by isoflurane pre-conditioning and post-conditioning. *Medical Gas Res*. 2017;7(1):48–55.
- [14] Alvarado MC, Murphy KL, Baxter MG. Visual recognition memory is impaired in rhesus monkeys repeatedly exposed to sevoflurane in infancy. *Br J Anaesth*. 2017;119(3):517–23.
- [15] Sun L. Early childhood general anaesthesia exposure and neurocognitive development. *Br J Anaesthesia*. 2010;105 Suppl 1:i61–8.
- [16] Loftis GK, Collins S, McDowell M. Anesthesia-induced neuronal apoptosis during synaptogenesis: a review of the literature. *AANA J*. 2012;80(4):291–8.
- [17] Colon E, Bittner EA, Kussman B, et al. Anesthesia, brain changes, and behavior: Insights from neural systems biology. *Progress in neurobiology*. 2017;153:121–60.
- [18] Jackson WM, Gray CD, Jiang D, et al. Molecular Mechanisms of Anesthetic Neurotoxicity: A Review of the Current Literature. *Journal of neurosurgical anesthesiology*. 2016;28(4):361–72.
- [19] Yu D, Li L, Yuan W. Neonatal anesthetic neurotoxicity: Insight into the molecular mechanisms of long-term neurocognitive deficits. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. 2017;87:196–9.
- [20] Vutskits L, Xie Z. Lasting impact of general anaesthesia on the brain: mechanisms and relevance. *Nat Rev Neurosci*. 2016;17(11):705–17.
- [21] Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell*. 2012;149(2):274–93.
- [22] Takei N, Nawa H. mTOR signaling and its roles in normal and abnormal brain development. *Frontiers in molecular neuroscience*. 2014;7:28.
- [23] Ehninger D, Li W, Fox K, et al. Reversing neurodevelopmental disorders in adults. *Neuron*. 2008;60(6):950–60.
- [24] Zhou J, Parada LF. PTEN signaling in autism spectrum disorders. *Current opinion in neurobiology*. 2012;22(5):873–9.
- [25] Ehninger D, Han S, Shilyansky C, et al. Reversal of learning deficits in a *Tsc2*^{+/-} mouse model of tuberous sclerosis. *Nat Med*. 2008;14(8):843–8.
- [26] Sharma A, Hoeffler CA, Takayasu Y, et al. Dysregulation of mTOR signaling in fragile X syndrome. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2010;30(2):694–702.
- [27] Bhattacharya A, Kaphzan H, Alvarez-Dieppa AC, et al. Genetic removal of p70 S6 kinase 1 corrects molecular, synaptic, and behavioral phenotypes in fragile X syndrome mice. *Neuron*. 2012;76(2):325–37.
- [28] Costa-Mattioli M, Monteggia LM. mTOR complexes in neurodevelopmental and neuropsychiatric disorders. *Nat Neurosci*. 2013;16(11):1537–43.
- [29] Sun Y, Li QF, Zhang Y, et al. Isoflurane preconditioning increases survival of rat skin random-pattern flaps by induction of HIF-1alpha expression. *Cellular physiology and biochemistry: international journal of experimental cellular physiology, biochemistry, and pharmacology*. 2013;31(4–5):579–91.
- [30] Na HS, Brockway NL, Gentry KR, et al. The genetics of isoflurane-induced developmental neurotoxicity. *Neurotoxicol Teratol*. 2017;60:40–49.
- [31] Li G, Xue Q, Luo Y, et al. S6 inhibition contributes to isoflurane neurotoxicity in the developing brain. *Toxicol Lett*. 2015;233(2):102–13.
- [32] Liu Y, Ma L, Jiao L, et al. Mammalian target of rapamycin/p70 ribosomal S6 protein kinase signaling is altered by sevoflurane and/or surgery in aged rats. *Mol Med Rep*. 2015;12(6):8253–60.