
MYCOTAXON

<http://dx.doi.org/10.5248/123.281>

Volume 123, pp. 281–284

January–March 2013

A new species of *Heliocephala* from Vietnam

VADIM A. MEL'NIK¹, RAFAEL F. CASTAÑEDA-RUIZ²
& MILAGRO GRANADOS³*

¹V. L. Komarov Botanical Institute of the Russian Academy of Sciences,
Prof. Popov Street 2, 197376, St. Petersburg, Russia

²Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT),
Académico Titular de la Academia de Ciencias de Cuba,
Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

³Centro de Investigaciones en Protección de Cultivos (CIPROC), Universidad de Costa Rica

*CORRESPONDENCE TO: milagranados@gmail.com

ABSTRACT — *Heliocephala vietnamensis* sp. nov., a hyphomycetous fungus collected on decaying leaves of an unidentified plant, is described and illustrated. It is distinguished by determinate apical cluster of conidiogenous cells and obclavate to sub-navicular, 3-septate, pale brown conidia.

KEY WORDS — anamorphic fungi, systematics, leaf litter

Several samples of dead plant material colonized by anamorphic fungi were collected in a dense monsoon tropical forest in Vietnam. Among these samples was a conspicuous *Heliocephala* species, which is herein described and illustrated.

Taxonomy

Heliocephala vietnamensis Melnik & R.F. Castañeda, sp. nov.

FIGS 1,2

MYCOBANKMB801539

Differs from *Heliocephala elegans* and *H. triseptata* by conidial shape, number of septa and size.

TYPE: Vietnam, Dong Nai, Cat Tien National Park, right bank of the Dong Nai River, near to *Ficus* sp. plot, a dense monsoon tropical forest, 11°26'N 107°25'E, on decaying leaves of unidentified plant, 2 January 2011, coll. Yu. Novozhilov, (**Holotype:** LE 261853).

ETYMOLOGY: Latin, *vietnamensis*, refers to the country where the fungus was found.

COLONIES on the natural substrate effuse, hairy, amphigenous, brown. Mycelium mostly superficial composed of septate, branched, dark brown, smooth-walled



FIG 1. *Helioccephala vietnamensis* (ex holotype LE 261853). A. Conidia. B. Conidiophore, conidiogenous cell and attached conidia. Scale bars = 10 µm.

hyphae, 1.8–2.8 µm diam. CONIDIOPHORES distinct, single, cylindrical, erect, straight, 210–340 × 6–8 µm, 14–16 µm wide at the base, 7–12-septate, bearing a conidiogenous apparatus at the apex consisting of 3–6 irregular-campanulate




FIG 2. *Heliocephala vietnamensis* (ex holotype LE 261853).
Conidiogenous cells and conidia. Scale bar = 10 μ m.

to irregular-doliiform or cuneiform primary and secondary branches (metula-like), pale brown, $3-4 \times 3.5-4.0 \mu\text{m}$. CONIDIOGENOUS CELLS monoblastic, discrete, determinate, delicate, lageniform, $1.7-3.0 \times 1.5-2.5 \mu\text{m}$, pale brown, smooth, forming a compact cluster at the ends of the branches. CONIDIAL SECESSION schizolytic. CONIDIA solitary, acrogenous, obclavate to navicular, 3-septate, pale brown, $14-17 \times 2.8-3.8 \mu\text{m}$, smooth-walled, dry. Teleomorph unknown.

NOTE: The genus *Heliocephala* V. Rao et al. (Rao et al. 1984) was based on *H. proliferans*, which was distinguished by the production of compact clusters of discrete, lageniform, monoblastic conidiogenous cells arising more less radially or closely fasciculate from short secondary or tertiary branches (metula-like) at the apex of the conidiophores. The conidia are obclavate, long or short rostrate, septate, pale brown, with an unusual germination of the apical cell to form a secondary cluster of conidiogenous cells.

Heredia et al. (2011), who emended *Heliocephala* after molecular and morphological studies of some *Heliocephala* species and the closely similar *Holubovaniella* R.F. Castañeda (Castañeda 1985), considered *Holubovaniella* a synonym of *Heliocephala*. They provided a key to the species and expanded the generic concept of *Heliocephala* to accommodate taxa with indeterminate conidiophores with several extensions of the main axis and solitary conidia produced by the same kind of conidiogenesis. Following Heredia et al. (2011), *Heliocephala* comprised six species: *H. elegans* (R.F. Castañeda) R.F. Castañeda & Unter., *H. gracilis* (R.F. Castañeda) R.F. Castañeda & Unter., *H. natarajanii* Kumaresan & M. Sriniv., *H. proliferans* V. Rao et al., *H. triseptata* Heredia et al., and *H. zimbabweensis* Decock et al.

Among the previously described *Heliocephala* species, only *H. elegans* and *H. triseptata* are similar to *H. vietnamensis*, but *H. elegans* (Castañeda-Ruiz 1985) has indeterminate conidiophores with several clusters of short branches and intercalary conidiogenous cells on the axis before each extension and olivaceous conidia that are obclavate to lecythiform with a shortly rostrate apical cell, mostly 2-septate, $8\text{--}25 \times 3\text{--}4 \mu\text{m}$. *H. triseptata* (Heredia et al. 2011) has conidia that are obclavate or navicular to broadly fusiform, rostrate, $15\text{--}27 \times 3.5\text{--}4.5 \mu\text{m}$ with a $10\text{--}14 \mu\text{m}$ long rostrum. Neither species can be confused with *H. vietnamensis*.

Acknowledgments

The authors express their sincere gratitude to Dr. Bryce Kendrick and Dr. De-Wei Li for their critical review of the manuscript. We are deeply indebted to Dr. Yuri Novozhilov for providing the specimen collected. In part this study has been done in frame of Research Program of the Russian-Vietnamese Tropical Research and Technological Center is also appreciated. We also thank Dr. Gabriela Heredia for facilities during microscopic observations and photographs. The authors are deeply indebted to the Russian Academy of Sciences and the Cuban Ministry of Agriculture for facilities. We also thank Mirtha Caraballo and Beatriz Ramos for technical assistance. We acknowledge the facilities provided by Dr. P.M. Kirk and Drs. V. Robert, G. Stegehuis and A. Decock through the maintenance of the Index Fungorum and Mycobank websites. Dr. Lorelei L. Norvell's editorial review and Dr. Shaun Pennycook's nomenclatural review are greatly appreciated.

Literature cited

- Castañeda-Ruiz RF. 1985. Deuteromycotina de Cuba. Hyphomycetes. 3. Instituto de Investigaciones Fundamentales en Agricultura Tropical "Alejandro de Humboldt". La Habana, Cuba. 72 p.
- Heredia Abarca G, Castañeda-Ruiz RF, Arias Mota RM, Becerra Hernández CI, Gómez S, Bogale M, Untereiner WA. 2011. A new species of *Heliocephala* from Mexico with an assessment of the systematic positions of the anamorph genera *Heliocephala* and *Holubovaniella*. *Mycologia* 103: 631–640. <http://dx.doi.org/10.3852/10-230>
- Rao V, Reddy KA, de Hoog GS. 1984. *Heliocephala*, a new genus of dematiaceous hyphomycetes. *Persoonia* 12: 239–242.