



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Functional and structural neuroplastic changes related to sensitization proxies in patients with Osteoarthritis

*a systematic review*

Pedersini, P; Gobbo, M; Bishop, M D; Arendt-Nielsen, L; Villafañe, J H

*Published in:*  
Pain Medicine

*DOI (link to publication from Publisher):*  
[10.1093/pm/pnab301](https://doi.org/10.1093/pm/pnab301)

*Publication date:*  
2022

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Pedersini, P., Gobbo, M., Bishop, M. D., Arendt-Nielsen, L., & Villafañe, J. H. (2022). Functional and structural neuroplastic changes related to sensitization proxies in patients with Osteoarthritis: a systematic review. *Pain Medicine*, 23(3), 488–498. <https://doi.org/10.1093/pm/pnab301>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Functional and structural neuroplastic changes related to sensitization proxies in patients with Osteoarthritis: a systematic review

Pedersini P<sup>1</sup>, MSc; Gobbo M<sup>2</sup>, MD; Bishop MD<sup>3</sup>, PhD; Arendt-Nielsen L<sup>4</sup>, MD; Villafañe JH<sup>1</sup>, PhD

## Affiliation:

1. IRCCS Fondazione Don Carlo Gnocchi, Milan, Italy
2. Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy
3. Department of Physical Therapy, University of Florida, USA.
4. Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, School of Medicine, Aalborg University, Aalborg, Denmark

## Correspondence:

Name	Paolo Pedersini
Institution	IRCCS Fondazione Don Carlo Gnocchi, Milan
Country	Italy
Tel	+39.338.8782284
Fax	+39-0119065495
Email	<a href="mailto:pedersini93@gmail.com">pedersini93@gmail.com</a>

**Original Paper for:** Pain Medicine

**Category:** Review

**Running title:** Neuroplastic brain changes in OA patients

**Funding sources:** None

**Conflict of interest statement:** The authors have no conflicts of interest to declare.

**Number of Text Pages:** 26

**Number of Figures:** 1

**Number of Tables:** 2

## Abstract

**Objective:** Several reports in literature have identified sensitization as a possible basis for the enhanced pain reactions associated with Osteoarthritis (OA). The aim of this current systematic review is to summarize functional and structural brain changes associated with surrogate sensitization parameters assessed in patients with OA-related pain.

**Design:** Systematic review.

**Subjects:** Patients with OA related pain.

**Methods:** A literature search was conducted systematically in MEDLINE, CINAHL, EMBASE databases for human studies up to December 2019. Articles were included if they assessed brain imaging and sensitisation parameters (quantitative sensory testing and questionnaires) in adults with OA related pain. Methodological quality was assessed using the Methodological Index for Non-Randomized Studies (MINORS) score.

**Results:** Five studies reporting on 138 patients were included in this review. The MINORS scale yielded mean scores of 8.5/16 and 12.3/24, for the cohort and case-control studies respectively. Four low-quality studies suggest a greater pain matrix activation associated with clinical measures of sensitization in patients with OA, while another study underlined the presence of structural changes (reduced gray matter volume) in the cortical areas involved in the nociceptive processing possible also related to sensitization.

**Conclusion:** This review shows conflicting evidence for structural and functional neuroplastic brain changes related to sensitization proxies in patients with OA.

**Keywords:** Osteoarthritis; sensitization; pain; brain imaging; MRI; fMRI.

## Introduction

Understanding and managing osteoarthritis (OA) pain is challenging given the several OA phenotypes, differentiated clinical presentations, and the sensory and psychological factors that modulate pain (1, 2). The socioeconomical burden of the OA is growing with about 300 million people affected around the world. The incidence of OA has risen 60% since 1990 (3) with a 31.4% increase in incidence during the period 2007 to 2017. This epidemiologic transition led to growing interest in the study of the sequelae related to OA pain and the mechanism involved in its generation and maintenance (4).

OA symptoms are often considered the results of the chronic overload and impaired biomechanics of the joint which in parts may lead to destruction of the articular cartilage and eventually inflammation. Latest evidence, about the pain-structure relationship in mice, has shown that loading induces an initial stress reaction in the joint and local inflammation, but these processes are not directly responsible for the nociceptive phenotype observed in mice (5). Human radiographic measures of pathologic joint changes have shown modest associations with clinical pain in patients with OA (6). Supporting this, OA patients may be divided in subgroups constructed by dichotomizing clinical knee pain scores and OA grade scores, revealed heightened pain sensitivity in the high pain/low OA grade group, while the low pain/high OA grade group can be less pain sensitive (7).

Recent evidence suggests that this discrepancy may be explained by the propensity of some patients to develop sensitization (8-10). In conditions, such as fibromyalgia, low back pain, tension type headache and persistent pelvic pain, aberration in pain processing mechanisms have been investigated as the basis of chronic pain maintenance (11-14). Proxies of central mechanisms have shown to be manifested across many chronic pain conditions (15) including patients with OA (16). Strong evidence for pain sensitization was reported in a recent systematic review and meta-analysis about the manifestations of pain

1  
2  
3 sensitization in people with knee OA based on meta-analyses of quantitative sensory testing  
4 (QST). The analysis indicated that the pain reported by people with knee OA in those studies  
5  
6 was associated with one aspect of QST, pressure pain thresholds (17).  
7  
8  
9

10 Combinations of QST tools can provide a battery for assessing pain threshold,  
11 responses to repeated noxious stimuli and evaluation of sensitivity through mechanical,  
12 chemical, electrical, and/or thermal testing, (9, 18) and have been suggested as proxies for  
13 individual aspects of the manifestations of sensitization. For example, a person with OA may  
14 have reduced thresholds to pressure pain (i.e. more sensitive) locally at the affected joint  
15 compared to a pain-free individual and similar thresholds at other body sites. This could be  
16 interpreted as peripheral (local) sensitization. If the person with OA has reduced thresholds  
17 at the affected joint and reduced thresholds at other body sites, this might be considered a  
18 result of additional central changes. Other more dynamic QST assessments of facilitatory  
19 and inhibitory processes provide additional insight to levels of sensitization. QST has  
20 revealed marked heterogeneity in nociceptive facilitation and inhibition in patients with OA,  
21 suggesting that different adaptations of the nociceptive system are present even within the  
22 same condition (19). Despite these findings, few studies have investigated if brain-related  
23 measures may possibly be connected to facilitated nociceptive processing and/or joint  
24 structural changes.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 Several neuroimaging studies have discussed the role of specific brain regions in the  
45 sensory, affective and cognitive aspects of pain experience (20). Structural and functional  
46 magnetic resonance imaging (MRI) may provide data on key central neural adaptations and  
47 assist in identifying subgroups OA might and possibly those susceptible to the development  
48 of severe chronic pain (21) or chronic postoperative pain after surgery (22). A systematic  
49 review on fibromyalgia investigating both functional and structural changes in the brain  
50 related to sensitization parameters found conflicting evidence for decreased gray matter  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 volume in specific brain area and showed an increased activation in the pain matrix  
4  
5 (cerebellum, insula, anterior and posterior cingulate, inferior parietal lobe and primary and  
6  
7 secondary somatosensory cortex) related to central sensitization in patients with  
8  
9 fibromyalgia (23). Such a review has not been performed to identify potential brain-related  
10  
11 changes in people with OA-related pain.  
12  
13

14 The aim of this current systematic review, therefore, is to summarize functional and  
15  
16 structural brain changes associated with surrogate sensitization parameters (e.g. QST)  
17  
18 assessed in patients with OA-related pain.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Method

This is a systematic literature review of studies investigating or reporting neuroplastic changes related to sensitization proxies in patients with OA. PRISMA guidelines were followed during the design, search and reporting stages of this systematic review. The protocol for this systematic review was registered on PROSPERO (CRD42020156007).

### *Systematic Literature Search*

Our literature search aimed at identifying all available studies that evaluated brain changes related to clinical manifestations of sensitization in the OA population. Electronic literature searches were conducted in the following databases from their inception until December 2019: MEDLINE, CINAHL and EMBASE. Additional records were searched through other sources to complement the database findings; for example, manual searches of reference lists of relevant literature reviews and indexes of peer-reviewed journals were used. Two authors (P.P. and S.M.) performed the search and evaluated the abstracts independently for potential eligibility and subsequently full-text publications for eligibility. A third author (J.H.V.) resolved discrepancies. Each researcher reviewed the title and abstract of all the articles, selecting the relevant ones according to inclusion and exclusion criteria. The references list of each article was also screened in order to find any additional original articles.

### *Population*

The participants in the selected studies had to be adults (18 years of age or older) with a diagnosis of symptomatic OA (osteoarthritis pain)

### *Exposure*

Sensitization. Clinical assessment tools such as questionnaires and/or relative sensory function differences using QST. QST aimed at assessing threshold ratios, provoked

1  
2  
3 hyperalgesia/allodynia, temporal summation (wind-up like pain), spatial summation, reflex  
4  
5 receptive fields, descending pain modulation and referred pain areas were included.  
6

### 7 *Comparator*

8  
9  
10 People with OA and sensitization compared to 1) people with OA and no sensitization,  
11  
12 2) pain-free people.  
13

### 14 *Outcome*

15  
16 Structural and functional MRI. For this aim, the search strategy included:  
17  
18 neuroimaging OR functional neuroimaging OR Brain imaging OR fMRI OR rs-fMRI OR  
19  
20 Voxel-Based Morphometry OR VBM.  
21  
22  
23  
24  
25

### 26 *Study selection*

27  
28 The search included observational studies (cohort studies, case–control studies and  
29  
30 case series) with human subjects and without restrictions regarding date of publication. The  
31  
32 decision to include case series was taken because from the beginning of the review we  
33  
34 expected few papers to be included and given the objective of summarizing the literature on  
35  
36 this topic could be useful to consider this type of studies. We excluded from the analysis all  
37  
38 repeated articles, case reports, letters to editor, pilot studies, editorials, technical notes and  
39  
40 review articles. Also *in vitro*, preclinical and animal studies were excluded. The participants  
41  
42 in the selected studies had to be adults (18 years of age or older) with a diagnosis of  
43  
44 symptomatic OA (osteoarthritis pain).  
45  
46  
47  
48

### 49 *Data extraction*

50  
51 All relevant articles from the aforementioned datasets were identified by two  
52  
53 reviewers (P.P. and S.M.) who conducted the data extraction independently. A third author  
54  
55 (J.H.V.) resolved discrepancies. Reviewers were not masked to any pieces of information  
56  
57 regarding the authors, the journal or the outcomes for each article reviewed. A standardized  
58  
59  
60



1  
2  
3 form was used to extract data concerning study design, number and mean age of  
4 participants, year and country of publication, setting, brain area involved, follow-up timing,  
5 clinical outcome measures and reported findings. The form was developed according to the  
6 directions of the *Cochrane Handbook for Systematic Reviews of Interventions* – Version  
7 5.1.0. This form was pilot-tested for reliability using a representative sample of the studies  
8 to be reviewed.  
9

### 16 *Quality assessment*

17  
18 The methodological index for non-randomized studies (MINORS) was used to assess the  
19 risk of bias and the methodological quality of the included studies (Slim et al., 2003) (24).  
20 This scoring system includes eight items for non-randomized studies and four additional  
21 items for comparative studies. Each item is scored between 0 and 2, and the maximum  
22 possible score is 16 and 24 for non-randomized studies and comparative studies,  
23 respectively. Two authors independently answered the questions with 0 (not reported), 1  
24 (reported but inadequate) or 2 (reported and adequate). Any disagreement was resolved by  
25 discussion, and if consensus was not reached an external review author was consulted and  
26 then, a decision was made.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

### 42 *Data analysis plan*

43  
44 We planned to perform a systematic review by descriptively presenting the results of  
45 the retrieved studies. A meta-analysis was not planned because the main objective of the  
46 study was to perform a descriptive analysis to present a state of the art of the topic with the  
47 aim of stimulating research in this area.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Results

### *Study selection*

Initially, 1676 studies were identified through database searching. After removing duplicates and screening titles and abstracts of all remaining unique articles, 14 full-text articles were assessed to verify their eligibility for the inclusion in the present study. Nine of these manuscripts were excluded (N= 8 articles did not clinically assess sensitization; N=1 Brain imaging not specifically for pain). Thus, five studies were finally selected for this review (**Figure 1**). A total of 146 patients with OA (113 knee OA, 20 hip OA and 13 hand OA) and 67 healthy subjects were included in these studies. The basic characteristics of the included studies are summarized in **Table 1**. The type of the included studies were 3 case-control studies (19, 25, 26) and 2 cohort studies (27, 28). The five included studies were conducted in Europe (4) and Oceania (1) and published from 2009 to 2019.

### *Risk of bias within and across the studies*

The MINORS Scale was used to score cohort studies and case-control studies, which yielded mean scores of 8.5 out of 16 and 12.3 out of 24, respectively (**table 2**). In general, the cohort (2) and case-control (3) studies were of poor quality. In five of the included studies, there was a discrepancy between outcomes listed in the method section and the result section and/or an unclear definition of outcomes in the study.

### *Data from studies*

#### *Association between cortical structure and sensitization proxies*

Lewis et al.(19) investigated brain structure in people with knee OA before and after total knee arthroplasty (TKA) and the relationships between these findings and QST (pressure pain threshold, temporal summation to repeated pressure stimulation and conditioned pain modulation). Twenty-nine patients with knee OA were compared with 18

1  
2  
3 pain-free subjects at presurgical baseline and 6 months following surgery. Brain structure  
4 was assessed through MRI Voxel based morphometry (VBM) and the QST was considered  
5 the nociceptive system outcome. Before TKA, there was reduced gray matter volume in  
6 areas associated with nociceptive processing, compared with the control group. In the  
7 longitudinal comparison of the knee OA group, the bilateral amygdala, contralateral  
8 hippocampus, and contralateral periaqueductal gray (PAG) were significantly larger at  
9 postsurgical time compared with presurgical (peak P-value <0.05). There were no areas that  
10 were significantly larger in presurgical OA patients compared with healthy controls, while  
11 healthy subjects shown significantly larger gray matter areas bilaterally in the Nucleus  
12 Accumbens and Amygdala and in the ipsilateral primary somatosensory cortex compared  
13 with presurgical OA patients (peak P-value <0.05). A significant relationship was found in a  
14 partial correlation between MRI and clinical variables in the knee OA group before surgery:  
15 a larger ipsilateral primary somatosensory cortex volume was associated with a lower knee  
16 pressure pain threshold on the knee ( $p=0.006$ ). VBM analyses indicated that the pain-free  
17 subjects had larger gray matter volume bilaterally in the nucleus accumbens, amygdala and  
18 in the ipsilateral primary somatosensory cortex compared with preoperative OA patients.  
19 The MINORS quality score of the study was 14/24.

#### 20 21 22 *Association between cortical activation and sensitization proxies*

23 Pujol et al. (27) studied pain sensitisation in patients with knee OA. Based on the evidence  
24 for spreading sensitization to a pressure stimulus modality and temporal summation to  
25 repeated pressure stimulation, 60 patients with knee OA were stratified into non-sensitised  
26 (27 participants) and sensitised (33 participants) groups and assessed in a single screening  
27 at baseline; 24 control subjects without pain or OA were evaluated to compare the results.  
28 fMRI was used to assess the evoked brain response during 3 tests (T): T1, pressure  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 stimulation on the articular line; T2, pressure stimulation on the anterior surface of the tibial  
4 region; T3, during a painful heat stimulation on the forearm. The PainDETECT score and  
5 specific pain assessment (spreading sensitization and temporal summation) was considered  
6 for nociceptive system assessment. No significant differences in brain activation during T1  
7 between sensitized and non-sensitized patients were found. Across the whole patient  
8 sample, brain activation during T2, correlated significantly with clinical measurements of  
9 pain sensitization. No significant differences in brain activation during T3 between sensitised  
10 and non-sensitised patients and between patients and control subjects were found. T3 did  
11 not discriminate the sensitization phenomenon. T1 robustly activated all of the neural  
12 elements typically involved in pain perception. T2 evoked greater activation in sensitized  
13 patients in regions typically involved in pain perception. T3 evoked a pattern of brain  
14 activation mostly involving bilateral frontoparietal opercula, insula and basal ganglia.  
15 Activations were also identified in the medial frontal cortex and premotor cortex. The quality  
16 score of the study was 9/16.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 Sofat et al.(26) assessed if sensitization mediates pain perception in hand OA.  
39 Thirteen patients with hand OA were compared with 13 pain-free subjects in a single  
40 session. fMRI was used to assess brain evoked response to pain induced by a finger flexion-  
41 extension (FFE). QST was used assess for signs of sensitization (assessed as the reduced  
42 pressure pain threshold) in the considered hand OA patients in a previous published study  
43 (Wajed et al., 2012) (29) on the same cohort. Analysis of fMRI data showed increased  
44 activation in the thalamus, cingulate cortex, frontal and somatosensory cortex ( $p<0.05$ )  
45 during FFE in the group with hand OA compared with pain-free subjects. Regions of  
46 activation were mapped to Brodmann areas 3, 4, 6, 9, 13, 22, 24 and 44. The quality score  
47 of the study was 11/24.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Gwilym et al. (25) investigated the supraspinal influences that underlie clinical  
4 manifestations (assessed as punctate stimulus detection threshold, punctate hyperalgesia,  
5 change in thermal perception thresholds and thermal pain threshold level) that could be  
6 considered indicative of possible sensitization. Twelve patients with hip OA were compared  
7 with 12 pain-free subjects in a single session. fMRI was used to assess evoked brain  
8 responses during punctate and cold stimuli. The PainDETECT score and QST were  
9 considered the nociceptive system assessments. Patients were found to have significantly  
10 lower threshold perception to punctate stimuli and were hyperalgesic to the noxious  
11 punctate stimulus in their areas of referred pain ( $p < 0.001$ ). fMRI data illustrated significantly  
12 greater activation in the brainstem of patients with OA in response to punctate stimulation  
13 of their referred pain areas compared with healthy controls, and the magnitude of BOLD  
14 response of this activation positively correlated with the extent of neuropathic-like elements  
15 to the patient's pain, as indicated by the PainDETECT score. There were no significant  
16 differences in activations of patients versus controls in response to the cold stimulus.  
17 Considering the whole brain, increased activity in the PAG in response to punctate  
18 stimulation of the referred pain area was detected in patients with OA ( $p < 0.05$ ). Considering  
19 punctate stimulation, the analysis between groups revealed greater activation in the OA  
20 group in the following regions: the anterior cingulate cortex, the right dorsolateral prefrontal  
21 cortex, the left middle frontal gyrus, and the left lateral occipital cortex. The quality score of  
22 the study was 12/24.

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49 The study of Soni et al. (28) aimed at identifying sensitization using neuroimaging  
50 and relating it to arthroplasty outcomes. Twenty-four knee patients with OA, waiting for  
51 arthroplasty, were stratified in two groups based on PainDETECT (PainDETECT score  $< 13$ :  
52 nociceptive group; PainDETECT score  $\geq 13$ : Neuropathic like pain group) and assessed  
53 before surgery and at 12 months follow-up. fMRI was used to assess evoked brain response  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 during punctate stimuli and cold stimuli. PainDETECT and QTS (mechanical pain threshold,  
4 cold detection threshold and cold pain threshold) were considered for the nociceptive  
5 system assessment. Patients with neuropathic-like pain before surgery (n = 14) reported  
6 significantly higher pain in response to punctate stimuli and cold stimuli near the affected  
7 joint. Neural activity in patients with neuropathic-like pain, compared to those with  
8 nociceptive pain, was significantly lower in the rostral anterior cingulate cortex and higher in  
9 the rostral ventromedial medulla during punctate stimulation, with significant functional  
10 connectivity between these two areas ( $r = 0.49$ ,  $P = 0.018$ ). There were no areas in which  
11 activation was significantly higher in the neuropathic-like pain group than in the nociceptive  
12 pain group ( $p < 0.05$ ). There was no significant difference in activation in the PAG. The  
13 punctate stimuli evoked increased brain activity bilaterally in the secondary somatosensory  
14 cortex (S2), anterior and posterior insula, and supplementary motor area, as well as in the  
15 mid–anterior cingulate cortex. The cold paradigm was associated with activation in the  
16 following areas bilaterally: S2 cortex, caudate, thalamus, cerebellum, and contralateral  
17 insula and putamen. The quality score of the study was 8/16.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Discussion

### *Findings*

This systematic review explored the literature about neuroplastic changes in the brain associated or co-occurring with sensitization proxies in patients with OA using specific functional and structural brain imaging. The number of included studies in this systematic review was relatively small (1 structural and 4 functional imaging) and the cohorts of patients likewise small and heterogeneous. The vast majority of the excluded studies did not consider the presence of sensitization and its implication in chronic pain maintenance or were based on clinical case reports and expert opinions, which we excluded from this review. As the quality scores of the included studies were not high and multiple heterogeneous approaches were used, the low quality evidence suggests that conflicting evidence exists regarding a relationship between neuroplastic changes and sensitization in people with OA, and depended on the different methodology employed in the studies.

Neuroplastic changes in the brain areas of patients with OA, associated with sensitization reported in the included studies, were in regions associated with the “pain matrix” (30). The signs of cortical changes in patients that clinically show sensitization suggest that these variations are probably related to central pain mechanisms. However, because of the lack of prospective studies performed in this area of study, it is not possible to establish whether the reported neuroplastic changes were a cause or a result of pain.

### *MRI and OA: People with OA versus pain-free people.*

Several cortical structural and functional differences were identified between people with OA and pain free people in this review. First, pain free people showed significantly greater gray matter bilaterally in affective brain regions (Nucleus Accumbens and Amygdala)

1  
2  
3 and in the ipsilateral primary somatosensory cortex compared with people with OA awaiting  
4 joint replacement surgery. The primary somatosensory cortex is of interest because of its  
5 major role in the localization and discrimination of pain alteration (31). Indeed, the  
6 neuroplastic changes of this brain area are thought to be one of the causes for chronic pain  
7 and not merely a passive phenomenon following tissue/nerve injury as previously assumed  
8 (32).  
9

10  
11  
12  
13  
14  
15  
16  
17 Structural alterations of gray matter is thought to possibly result from plasticity that  
18 occurs in the context of structural remodeling and reorganization of synapses, cells and  
19 circuits, potentially contributing to the long-term nature of chronic pain (33, 34). Structural  
20 brain biomarkers for chronic pain have been reported in literature for other conditions  
21 suggesting numerous patterns of structural changes that are specific to several conditions  
22 related to pain (35, 36).  
23  
24  
25  
26  
27  
28  
29

30  
31 Functionally, increased activation in the thalamus, cingulate cortex, frontal and  
32 somatosensory cortex occur during movement evoked pain in people with hand OA  
33 compared with pain-free subjects (26). These cortical regions are implicated in central  
34 sensitization. Additionally, greater activation in the brainstem, anterior cingulate cortex, the  
35 right dorsolateral prefrontal cortex, the left middle frontal gyrus, and the left lateral occipital  
36 cortex was also reported (25) in patients compared to pain free people in response to  
37 punctate stimuli. This activation was correlated with the extent of neuropathic pain  
38 symptoms reported by patients.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 *MRI and OA: People with sensitization versus people without*

52  
53  
54 In the study by Pujol (27), pressure stimulation on the anterior surface of the tibial  
55 region (i.e a site remote to the reported painful joint) evoked greater activation  
56 “somatosensory cortices, supramarginal gyrus, insula, visual and auditory” areas in patients  
57  
58  
59  
60



1  
2  
3 classified as “sensitized” compared to the non-sensitized group. Interestingly, the authors  
4 indicate that these identified areas overlap with those activated in people with fibromyalgia,  
5 a condition associated with primary sensitization. In separate work (28), patients with  
6 neuropathic-like pain before surgery reported significantly higher pain in response to  
7 punctate and cold stimuli near the affected joint than those without neuropathic pain. Neural  
8 activity bilaterally in the secondary somatosensory cortex (S2), anterior and posterior insula,  
9 and supplementary motor area, as well as in the mid–anterior cingulate cortex in these  
10 patients was significantly lower in the rostral anterior cingulate cortex and higher in the  
11 rostral ventromedial medulla during punctate stimulation, with significant functional  
12 connectivity between these two areas. Interestingly there was no significant difference in  
13 activation in the PAG possibly suggesting that differences in endogenous inhibitory capacity  
14 may not have differed between groups.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 *MRI: structural differences and QST*

34  
35 Lewis et al.(19) explored structural brain changes in patients with OA before and after  
36 TKA, showing their relationship with the QST. In particular, they identified a significant  
37 correlation between imaging data and clinical PPT such that a larger ipsilateral primary  
38 somatosensory cortex volume was associated with a lower knee pressure pain threshold on  
39 the knee.  
40  
41  
42  
43  
44  
45  
46

47 In contrast, an emerging study that analysed the individual difference in sensitivity in  
48 a large sample of people in pain strongly suggests an absence of associations between PPT  
49 and gray matter volume (37). Previous studies have reported that PPTs at the knee and  
50 distant sites are reduced in people with knee OA, and the presence of widespread  
51 hyperalgesia and enhanced spatial summation observed as indicative for sensitization (38).  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Widespread sensitization was detected also in patients with pain after TKA revision and  
4 underlined the importance of ongoing nociceptive input for the chronification process (39).  
5  
6  
7  
8  
9  
10

### 11 *MRI: function (activation) differences and QST*

12  
13  
14 The observation that several brain areas are activated by transient painful stimuli,  
15 and that the magnitude of this activity is often graded with pain intensity, has prompted  
16 researchers to extract features of brain activity that could serve as biomarkers to measure  
17 the pain matrix activity. An fMRI study by Soni et al.(28) analyzed functionally linked regions  
18 associated with pain processing, specifically related to sensitization assessed as  
19 mechanical and cold pain threshold and cold detection threshold. These neuroimaging data  
20 suggest that a subset of patients with OA possible have sensitization showing up as  
21 increased brain activity bilaterally in the secondary somatosensory cortex, anterior and  
22 posterior insula, and supplementary motor area, as well as in the mid–anterior cingulate  
23 cortex.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36

37  
38 Emerging evidence suggests that pain sensitization in people with knee OA may be  
39 associated with knee OA symptom severity, although not all patients develop detectable  
40 sensitization (40). Pujol et al.(27) studied pain sensitization in knee OA and analysed cortical  
41 activation to punctate and cold stimuli and found a significant difference between sensitized  
42 and non-sensitized patients. Punctate stimulation on the interarticular line showed similar  
43 activation of the neural elements involved in pain perception across groups, whereas the  
44 punctate stimulation on the distant site (anterior surface of the tibial region) showed greater  
45 activation in sensitized patients. In sensitized patients, a state of hyperexcitability consisting  
46 of enhanced responses to noxious stimulation has been shown (41).  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Also Gwyllim et al.(25) found signs of sensitization in patients with knee OA in  
4 response to punctate stimulation of referred pain areas, compared with the response of  
5 healthy controls: the magnitude of this activation positively correlated with the extent of  
6 neuropathic-like elements to the patient's pain. Sofat et al.(26) in contrast, investigated  
7 sensitization (pain thresholds to pressure stimulus modality) in patients with hand OA  
8 through fMRI data during flexion-extension of the hand and found augmented activation in  
9 the thalamus, cingulate, frontal and somatosensory cortex in the OA group but not in healthy  
10 controls. The work by Sofat et. al. (26) is the only included study that assessed brain  
11 activation during spontaneous pain (without an external stimulation), observing that their  
12 cohort of hand OA patients, who presented relatively high VAS pain scores and lower  
13 pressure pain threshold measured by algometers, showed a greater activation in the areas  
14 that involve emotion, learning, memory and central pain processing. In a previous study,  
15 Parks et al.(42) addressed the differences in brain activity during induced pain (with a  
16 pressure-evoked pain stimulus) and spontaneous OA knee pain. They suggested that, for  
17 spontaneous pain, the engagement of brain regions involved in emotional assessment is  
18 mediated sensitization, while evoked pain presents similar mechanisms in patients with  
19 knee OA and healthy subjects (42). Hyperalgesia and signs of sensitization have already  
20 been reported in a previous clinical study that analysed the nerve sensitivity of patients with  
21 hand OA and investigated pressure pain thresholds on local and distant anatomical sites  
22 (43, 44).

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49 Understanding both structural and functional brain adaptations to persistent pain may  
50 be useful to define specific OA phenotypes of pain and, subsequently, to assess the impact  
51 of different therapeutic/preventive modalities on brain changes. Russel et al.(45) evaluated  
52 volumetric changes in brain regions of patients with hand OA treated with centrally acting  
53 analgesics, showing no structural differences in the insular cortex or thalamus at baseline  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 and after treatment. Therapeutic exercise has also been investigated for its possible  
4 influence on brain changes in patients with knee OA, highlighting the potential use of  
5 neuroimaging biomarkers to predict the intervention effect on pain (46). Exercise has the  
6 potential to regulate the motor control system by directly modulating the supplementary  
7 motor area function, as well as the dorsolateral prefrontal cortex by shifting attention to  
8 movements. Moreover, exercise can influence strengthening cognition and self-regulation  
9 through the descending pain modulation system and the brain areas associated with  
10 emotion (amygdala) and pain (insula) (46).  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

### 23 *Limitations*

24  
25  
26 This systematic review did have some limitations; the majority of the studies that has  
27 been included did not report cohorts with consecutive patients and preliminary sample size  
28 calculations. Three studies had a single assessment at follow-up and 2 studies had a follow  
29 up of 6 or 12 months to evaluate the post-surgical outcomes. The sample size of the included  
30 studies was relatively small, thus studies with larger sample sizes are needed to further  
31 validate the findings. Classification bias might be present as the criteria for the diagnosis of  
32 OA were not well described in the studies. Most of the studies were of low quality.  
33 Specifically, no RCTs testing the effect of interventions were identified for this review. All the  
34 mentioned biases, along with the retrospective nature of studies, may influence the strength  
35 of the results. Additionally, given the heterogeneity in methods we are unable to provide  
36 recommendations regarding the most effective combination of imaging and QST  
37 approaches to best quantify brain-related changes in people with OA and sensitization.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

### 56 *Clinical and Research Implications*

1  
2  
3 Gaining information on neuroplastic changes and neural “rearrangement” may be a new  
4 option to develop multimodal approaches for managing OA, including therapies targeting  
5 the cortex and sensitization. The studies included in this review were generally rated as  
6 lower quality and had heterogeneous methodology.  
7  
8  
9  
10  
11  
12

13  
14 Brain imaging provides a method to underline the patterns of neural activity in the human  
15 brain and spinal cord, providing foundation for research of the afferent and efferent circuitry  
16 in the patient with pain. Future studies should be focused on the thorough definition of the  
17 neuroplastic adaptations related to the several phenotypes of pain including sensitization.  
18 Defining the variation of neuronal patterns during treatments targeting sensitization proxies  
19 should be considered in order to improve the quality of personalized treatment based on  
20 pain patient phenotype. The optimal methodology to determine if sensitization in people with  
21 OA is related to cortical reorganisation would be longitudinal studies of disease/condition  
22 progression and recovery after intervention. Unfortunately these studies are difficult and  
23 expensive to perform; however, such studies will be needed to untangle the extent to which  
24 cortical reorganization occurs in response to prolonged pain, or if differences in structure  
25 and function give rise to prolonged pain and/or sensitization, and to determine if any  
26 identified changes are in fact reversible. Alternatively, cross-sectional case-control studies  
27 with rigorous designs could provide more information about differences in cortical structure  
28 and function. In addition, future studies must consider consistent definitions of sensitization  
29 and optimal QST measures used to define such sensitization.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Conclusion

This systematic review shows conflicting evidence concerning the structural and functional brain changes related to assessed sensitization occurring in patients with OA. The tools used and applied as proxies for sensitization vary across studies making direct comparisons of the sensitization aspects difficult. The imaging findings of five low-quality studies suggest a greater pain matrix activation associated to clinical measures of sensitization in patients with OA compared to pain-free subjects and possible reduced gray matter volume in the cortical areas involved in the nociceptive processing. The conclusions have a high risk of bias. It was not possible to make a quantitative analysis aimed at comparing the brain area involved. Future works should consider larger sample sizes and longitudinal study designs should be contemplated in order to determine the chronology of brain alterations in relation to the course of the painful condition as well as to evaluate the effects of pain treatments on neural reorganization in patients suffering from OA with concurrent specific sensitized phenotypes.

**Acknowledgements:**

The authors thank to Dr Stefano Moschin, for his support with the data extraction.

LAN: Centre for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121).

**Conflict of interest statement:**

The authors have no conflicts of interest to declare.

**Authors contributions:**

All the authors meet the following authorship conditions:

1) Substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; 2) drafting the article or revising it critically for important intellectual content; 3) final approval of the version to be published.

## Bibliography

1. Van Spil WE, Kubassova O, Boesen M, Bay-Jensen AC, Mobasher A. Osteoarthritis phenotypes and novel therapeutic targets. *Biochem Pharmacol.* 2019;165:41-48.
2. Villafane JH, Pedersini P, Bertozzi L, Drago L, Fernandez-Carnero J, Bishop MD, Berjano P. Exploring the relationship between chronic pain and cortisol levels in subjects with osteoarthritis: Results from a systematic review of the literature. *Osteoarthritis Cartilage.* 2020.
3. Disease GBD, Injury I, Prevalence C. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2018;392:1789-1858.
4. Villafane JH, Valdes K, Pedersini P, Berjano P. Osteoarthritis: a call for research on central pain mechanism and personalized prevention strategies. *Clinical rheumatology.* 2019;38:583-584.
5. Ter Heegde F, Luiz AP, Santana-Varela S, Magnusdottir R, Hopkinson M, Chang Y, Poulet B, Fowkes RC, Wood JN, Chenu C. Osteoarthritis-related nociceptive behaviour following mechanical joint loading correlates with cartilage damage. *Osteoarthritis Cartilage.* 2020;28:383-395.
6. Bedson J, Croft PR. The discordance between clinical and radiographic knee osteoarthritis: a systematic search and summary of the literature. *BMC musculoskeletal disorders.* 2008;9:116.
7. Finan PH, Buenaver LF, Bounds SC, Hussain S, Park RJ, Haque UJ, Campbell CM, Haythornthwaite JA, Edwards RR, Smith MT. Discordance between pain and radiographic severity in knee osteoarthritis: findings from quantitative sensory testing of central sensitization. *Arthritis and rheumatism.* 2013;65:363-372.
8. Arendt-Nielsen L, Skou ST, Nielsen TA, Petersen KK. Altered Central Sensitization and Pain Modulation in the CNS in Chronic Joint Pain. *Curr Osteoporos Rep.* 2015;13:225-234.
9. Suokas AK, Walsh DA, McWilliams DF, Condon L, Moreton B, Wylde V, Arendt-Nielsen L, Zhang W. Quantitative sensory testing in painful osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis and cartilage.* 2012;20:1075-1085.
10. Akin-Akinyosoye K, Frowd N, Marshall L, Stocks J, Fernandes GS, Valdes A, McWilliams DF, Zhang W, Doherty M, Ferguson E, Walsh DA. Traits associated with central pain augmentation in the Knee Pain In the Community (KPIC) cohort. *Pain.* 2018;159:1035-1044.
11. Aoyagi K, He J, Nicol AL, Clauw DJ, Kluding PM, Jernigan S, Sharma NK. A Subgroup of Chronic Low Back Pain Patients With Central Sensitization. *Clin J Pain.* 2019;35:869-879.
12. Caamano-Barrios LH, Galan-Del-Rio F, Fernandez-de-Las-Penas C, Plaza-Manzano G, Arendt-Nielsen L, Ortega-Santiago R. Widespread Pressure Pain Sensitivity over Nerve Trunk Areas in Women with Frequent Episodic Tension-Type Headache as a Sign of Central Sensitization. *Pain Med.* 2019.
13. Fernandez-de-Las-Penas C, Arendt-Nielsen L. Myofascial pain and fibromyalgia: two different but overlapping disorders. *Pain Manag.* 2016;6:401-408.
14. Grundstrom H, Gerdle B, Alehagen S, Bertero C, Arendt-Nielsen L, Kjolhede P. Reduced pain thresholds and signs of sensitization in women with persistent pelvic pain and suspected endometriosis. *Acta Obstet Gynecol Scand.* 2019;98:327-336.
15. Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain.* 2011;152:S2-15.



16. Arendt-Nielsen L, Nie H, Laursen MB, Laursen BS, Madeleine P, Simonsen OH, Graven-Nielsen T. Sensitization in patients with painful knee osteoarthritis. *Pain*. 2010;149:573-581.
17. De Oliveira Silva D, Rathleff MS, Petersen K, Azevedo FM, Barton CJ. Manifestations of Pain Sensitization Across Different Painful Knee Disorders: A Systematic Review Including Meta-analysis and Metaregression. *Pain medicine*. 2019;20:335-358.
18. Arendt-Nielsen L, Morlion B, Perrot S, Dahan A, Dickenson A, Kress HG, Wells C, Bouhassira D, Mohr Drewes A. Assessment and manifestation of central sensitisation across different chronic pain conditions. *Eur J Pain*. 2018;22:216-241.
19. Lewis GN, Parker RS, Sharma S, Rice DA, McNair PJ. Structural Brain Alterations Before and After Total Knee Arthroplasty: A Longitudinal Assessment. *Pain Med*. 2018;19:2166-2176.
20. Yang S, Chang MC. Chronic Pain: Structural and Functional Changes in Brain Structures and Associated Negative Affective States. *Int J Mol Sci*. 2019;20.
21. Boissoneault J, Sevel L, Letzen J, Robinson M, Staud R. Biomarkers for Musculoskeletal Pain Conditions: Use of Brain Imaging and Machine Learning. *Curr Rheumatol Rep*. 2017;19:5.
22. Arendt-Nielsen L, Simonsen O, Laursen MB, Roos EM, Rathleff MS, Rasmussen S, Skou ST. Pain and sensitization after total knee replacement or nonsurgical treatment in patients with knee osteoarthritis: Identifying potential predictors of outcome at 12 months. *Eur J Pain*. 2018;22:1088-1102.
23. Cagnie B, Coppieters I, Denecker S, Six J, Danneels L, Meeus M. Central sensitization in fibromyalgia? A systematic review on structural and functional brain MRI. *Semin Arthritis Rheum*. 2014;44:68-75.
24. Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ journal of surgery*. 2003;73:712-716.
25. Gwilym SE, Keltner JR, Warnaby CE, Carr AJ, Chizh B, Chessell I, Tracey I. Psychophysical and functional imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis patients. *Arthritis Rheum*. 2009;61:1226-1234.
26. Sofat N, Smee C, Hermansson M, Howard M, Baker EH, Howe FA, Barrick TR. Functional MRI demonstrates pain perception in hand osteoarthritis has features of central pain processing. *Journal of biomedical graphics and computing*. 2013;3.
27. Pujol J, Martinez-Vilavella G, Llorente-Onaindia J, Harrison BJ, Lopez-Sola M, Lopez-Ruiz M, Blanco-Hinojo L, Benito P, Deus J, Monfort J. Brain imaging of pain sensitization in patients with knee osteoarthritis. *Pain*. 2017;158:1831-1838.
28. Soni A, Wanigasekera V, Mezue M, Cooper C, Javaid MK, Price AJ, Tracey I. Central Sensitization in Knee Osteoarthritis: Relating Presurgical Brainstem Neuroimaging and PainDETECT-Based Patient Stratification to Arthroplasty Outcome. *Arthritis & rheumatology*. 2019;71:550-560.
29. Wajed J, Ejindu V, Heron C, Hermansson M, Kiely P, Sofat N. Quantitative sensory testing in painful hand osteoarthritis demonstrates features of peripheral sensitisation. *Int J Rheumatol*. 2012;2012:703138.
30. Salomons TV, Iannetti GD, Liang M, Wood JN. The "Pain Matrix" in Pain-Free Individuals. *JAMA Neurol*. 2016;73:755-756.
31. Craig AD. Pain mechanisms: labeled lines versus convergence in central processing. *Annu Rev Neurosci*. 2003;26:1-30.
32. Kim W, Kim SK, Nabekura J. Functional and structural plasticity in the primary somatosensory cortex associated with chronic pain. *J Neurochem*. 2017;141:499-506.

- 1  
2  
3 33. Kuner R, Flor H. Structural plasticity and reorganisation in chronic pain. *Nat Rev*  
4 *Neurosci.* 2017;18:113.  
5 34. Mao CP, Bai ZL, Zhang XN, Zhang QJ, Zhang L. Abnormal Subcortical Brain  
6 Morphology in Patients with Knee Osteoarthritis: A Cross-sectional Study. *Frontiers in aging*  
7 *neuroscience.* 2016;8:3.  
8 35. Niddam DM, Lee SH, Su YT, Chan RC. Brain structural changes in patients with chronic  
9 myofascial pain. *Eur J Pain.* 2017;21:148-158.  
10 36. Gwilym SE, Filippini N, Douaud G, Carr AJ, Tracey I. Thalamic atrophy associated with  
11 painful osteoarthritis of the hip is reversible after arthroplasty: a longitudinal voxel-based  
12 morphometric study. *Arthritis and rheumatism.* 2010;62:2930-2940.  
13 37. Ruscheweyh R, Wersching H, Kugel H, Sundermann B, Teuber A. Gray matter  
14 correlates of pressure pain thresholds and self-rated pain sensitivity: a voxel-based  
15 morphometry study. *Pain.* 2018;159:1359-1365.  
16 38. Graven-Nielsen T, Wodehouse T, Langford RM, Arendt-Nielsen L, Kidd BL.  
17 Normalization of widespread hyperesthesia and facilitated spatial summation of deep-tissue  
18 pain in knee osteoarthritis patients after knee replacement. *Arthritis and rheumatism.*  
19 2012;64:2907-2916.  
20 39. Skou ST, Graven-Nielsen T, Rasmussen S, Simonsen OH, Laursen MB, Arendt-Nielsen  
21 L. Widespread sensitization in patients with chronic pain after revision total knee  
22 arthroplasty. *Pain.* 2013;154:1588-1594.  
23 40. Lluch E, Nijs J, Courtney CA, Rebbeck T, Wylde V, Baert I, Wideman TH, Howells N,  
24 Skou ST. Clinical descriptors for the recognition of central sensitization pain in patients with  
25 knee osteoarthritis. *Disabil Rehabil.* 2018;40:2836-2845.  
26 41. den Boer C, Dries L, Terluin B, van der Wouden JC, Blankenstein AH, van Wilgen CP,  
27 Lucassen P, van der Horst HE. Central sensitization in chronic pain and medically  
28 unexplained symptom research: A systematic review of definitions, operationalizations and  
29 measurement instruments. *J Psychosom Res.* 2019;117:32-40.  
30 42. Parks EL, Geha PY, Baliki MN, Katz J, Schnitzer TJ, Apkarian AV. Brain activity for  
31 chronic knee osteoarthritis: dissociating evoked pain from spontaneous pain. *Eur J Pain.*  
32 2011;15:843 e841-814.  
33 43. Pedersini P, Negrini S, Cantero-Tellez R, Bishop MD, Villafane JH. Pressure algometry  
34 and palpation of the upper limb peripheral nervous system in subjects with hand  
35 osteoarthritis are repeatable and suggest central changes. *J Hand Ther.* 2020;33(1):103-  
36 111.  
37 44. Pedersini P, Valdes K, Cantero-Tellez R, Cleland JA, Bishop MD, Villafane JH. The  
38 effects of neurodynamic mobilizations on pain hypersensitivity in patients with hand  
39 osteoarthritis compared to robotic assisted mobilization: A randomized clinical trial. *Arthritis*  
40 *Care Res (Hoboken).* 2021 Feb;73(2):232-239.  
41 45. Russell MD, Barrick TR, Howe FA, Sofat N. Reduced anterior cingulate grey matter  
42 volume in painful hand osteoarthritis. *Rheumatol Int.* 2018;38:1429-1435.  
43 46. Liu J, Chen L, Tu Y, Chen X, Hu K, Tu Y, Lin M, Xie G, Chen S, Huang J, Liu W, Wu J,  
44 Xiao T, Wilson G, Lang C, Park J, Tao J, Kong J. Different exercise modalities relieve pain  
45 syndrome in patients with knee osteoarthritis and modulate the dorsolateral prefrontal  
46 cortex: A multiple mode MRI study. *Brain, behavior, and immunity.* 2019;82:253-263.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Figure Legends**

4 **Figure 1.** Flow diagram of studies through the different phases of the systematic review

5  
6 **Table 1.** Characteristics of included studies.

7  
8 **Table 2.** Methodological items for non-randomised studies (MINORS scale)

9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.** Characteristics of included studies.

Author, yrs	Number of participants	Mean (SD) age	Female Ratio (%)	Affected Joint	Used QST modalities	Type of MRI technique	Country	Design of the study	Length of follow-up	Statistical analysis
Pujol et al. (2017) [27]	84 participants (60 Knee OA and 24 healthy subjects.)	-Knee OA 66.7 (7.8) -Healthy subjects 62.8 (7.7)	-Knee OA 71.7% -Healthy subjects 58.3%	Knee OA	PPT, TS to repeated pressure stimulation	fMRI (BOLD) during: Test 1(T1): pressure stimulation on the articular line. T2: pressure stimulation on the anterior surface of the tibial region. T3: during a painful heat stimulation on the forearm.	Spain	Cohort study	Single screening at baseline.	ANOVA to assess within-group and between-group differences.
Lewis et al. (2018) [19]	47 participants (29 Knee OA and 18 healthy subjects)	-Knee OA 68 (10) -Healthy subjects 71.0 (8.0)	-Knee OA 48.3% -Healthy subjects 61.1%	Knee OA after TKA	PPT, TS to repeated pressure stimulation and CPM	MRI VBM	New Zealand	Case-control	Screening before and 6 months after TKA.	Independent t-test to investigate differences between groups for VBM and QST data.
Sofat et al. (2013) [26]	26 participants (13 hand OA patients and 13 healthy subjects)	-Hand OA 61.0 (13.7) -Healthy subjects 52.8 (5.3)	-Hand OA 100% -Healthy subjects 100%	Hand OA	PPT	Evoked brain response assessed through fMRI (BOLD) during a finger flexion-extension.	United Kingdom	Case-control	Single screening at baseline	t-test was performed
Gwilym et al. (2009) [25]	32 participants (20 hip OA patients, 12 underwent fMRI; 12 healthy subjects)	-Hip OA 63.0 (8.0) -Healthy subjects 64.0 (9.0)	-Hip OA 50% -Healthy subjects 45%	Hip OA	punctate stimulus detection threshold, punctate hyperalgesia, change in thermal perception thresholds and thermal pain threshold level	Evoked brain response assessed through fMRI (BOLD) during a punctate stimuli and cold stimuli.	United Kingdom	Case-control	Single screening at baseline	Mann-Whitney test and unpaired t-test was performed
Soni et al. (2019) [28]	24 participants (10 nociceptive pain group and 14 neuropathic pain group)	-Nociceptive group 70.0 (7.0) -Neuropathic group 67.0 (10)	-Nociceptive group, 30.0% -Neuropathic group, 57.0%	Knee OA	mechanical pain threshold, cold detection threshold and cold pain threshold)	Evoked brain response (BOLD) assessed through fMRI during a punctate stimuli and cold stimuli.	United Kingdom	Cohort study	Presurgical, baseline and 12 months follow-up assessment.	Student's t-test, Wilcoxon-Mann-Whitney, Fisher's exact test were performed


**OA:** Osteoarthritis; **fMRI:** functional magnetic resonance imaging; **VBM:** Voxel based morphometry; **TKA:** total knee arthroplasty; **QST:** quantitative sensory testing, **PPT:** pressure pain threshold; **TS:** temporal summation; **CPM:** conditioned pain modulation; **BOLD:** Blood oxygenation level dependent.

**Table 2.** Methodological items for non-randomised studies (MINORS scale)

Authors, Yrs	A clearly stated aims	Inclusion of consecutive patients	Prospective collection data	Endpoints appropriate to the aim of the study	Unbiased assessment of the study endpoint	Follow-up appropriate	Loss to follow-up less than 5%	Prospective calculation of the study size	An adequate control group	Contemporary groups	Baseline equivalence of groups	Adequate statistical analysis	Score
Pujol et al. (2017) [27]	2	2	0	2	1	2	0	0	-	-	-	-	9/16
Lewis et al. (2018) [19]	2	1	1	2	0	1	2	0	1	2	0	2	14/24
Sofat et al. (2013) [26]	2	0	0	1	0	1	0	1	1	2	1	2	11/24
Gwilym et al. (2009) [25]	2	1	1	2	0	1	0	0	1	2	1	1	12/24
Soni et al. (2019) [28]	2	1	1	0	0	2	0	2	-	-	-	-	8/16

The items are scored 0 (not reported), 1 (reported but inadequate) or 2 (reported and adequate). The global ideal score being 16 for non-comparative studies and 24 for comparative studies

**Figure. 1** Flow diagram based on PRISMA statement ([www.prisma-statement.org](http://www.prisma-statement.org), accessed on: 30/01/2020).



Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit [www.prisma-statement.org](http://www.prisma-statement.org).

Official Journal of the American Academy of Pain Medicine