



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Nonpharmacological Treatment of Persistent Postconcussion Symptoms in Adults

*A Systematic Review and Meta-analysis and Guideline Recommendation*

Rytter, Hana Malá; Graff, Heidi J; Henriksen, Henriette K; Aen, Nicolai; Hartvigsen, Jan; Hoegh, Morten; Nisted, Ivan; Næss-Schmidt, Erhard Trillingsgaard; Pedersen, Lisbeth Lund; Schytz, Henrik Winther; Thastum, Mille Møller; Zerlang, Bente; Callesen, Henriette Edemann

*Published in:*

JAMA NETWORK OPEN

*DOI (link to publication from Publisher):*

[10.1001/jamanetworkopen.2021.32221](https://doi.org/10.1001/jamanetworkopen.2021.32221)

*Creative Commons License*

CC BY 4.0

*Publication date:*

2021

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Rytter, H. M., Graff, H. J., Henriksen, H. K., Aen, N., Hartvigsen, J., Hoegh, M., Nisted, I., Næss-Schmidt, E. T., Pedersen, L. L., Schytz, H. W., Thastum, M. M., Zerlang, B., & Callesen, H. E. (2021). Nonpharmacological Treatment of Persistent Postconcussion Symptoms in Adults: A Systematic Review and Meta-analysis and Guideline Recommendation. *JAMA NETWORK OPEN*, 4(11), [e2132221]. <https://doi.org/10.1001/jamanetworkopen.2021.32221>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Nonpharmacological Treatment of Persistent Postconcussion Symptoms in Adults A Systematic Review and Meta-analysis and Guideline Recommendation

Hana Malá Rytter, PhD; Heidi J. Graff, PhD; Henriette K. Henriksen, PT; Nicolai Aaen, MSc; Jan Hartvigsen, PhD; Morten Hoegh, PhD; Ivan Nisted, MSc; Erhard Trillingsgaard Næss-Schmidt, PhD; Lisbeth Lund Pedersen, MSc, PT; Henrik Winther Schytz, MD, PhD, DMSc; Mille Møller Thastum, MSc, PhD; Bente Zerlang, OT; Henriette Edemann Callesen, PhD

## Abstract

**IMPORTANCE** Persistent (>4 weeks) postconcussion symptoms (PPCS) are challenging for both patients and clinicians. There is uncertainty about the effect of commonly applied nonpharmacological treatments for the management of PPCS.

**OBJECTIVE** To systematically assess and summarize evidence for outcomes related to 7 nonpharmacological interventions for PPCS in adults (aged >18 years) and provide recommendations for clinical practice.

**DATA SOURCES** Systematic literature searches were performed via Embase, MEDLINE, PsycINFO, CINAHL, PEDro, OTseeker, and Cochrane Reviews (via MEDLINE and Embase) from earliest possible publication year to March 3, 2020. The literature was searched for prior systematic reviews and primary studies. To be included, studies had to be intervention studies with a control group and focus on PPCS.

**STUDY SELECTION** A multidisciplinary guideline panel selected interventions based on frequency of use and need for decision support among clinicians, including early information and advice, graded physical exercise, vestibular rehabilitation, manual treatment of neck and back, oculomotor vision treatment, psychological treatment, and interdisciplinary coordinated rehabilitative treatment. To be included, studies had to be intervention studies within the areas of the predefined clinical questions, include a control group, and focus on symptoms after concussion or mild traumatic brain injury.

**DATA EXTRACTION AND SYNTHESIS** Extraction was performed independently by multiple observers. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used for data abstraction and data quality assessment. Included studies were assessed using the Assessment of Multiple Systematic Reviews (AMSTAR) tool and the Cochrane Risk of Bias (randomized clinical trials) tool. Meta-analysis was performed for all interventions where possible. Random-effects models were used to calculate pooled estimates of effects. The level and certainty of evidence was rated and recommendations formulated according to the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) framework.

**MAIN OUTCOMES AND MEASURES** All outcomes were planned before data collection began according to a specified protocol. The primary outcomes were the collective burden of PPCS and another outcome reflecting the focus of a particular intervention (eg, physical functioning after graded exercise intervention).

**RESULTS** Eleven systematic reviews were identified but did not contribute any primary studies; 19 randomized clinical trials comprising 2007 participants (1064 women [53.0%]) were separately

(continued)

## Key Points

**Question** What is the evidence for nonpharmacological interventions to treat persistent postconcussion symptoms?

**Findings** Following a systematic review and meta-analysis of 19 randomized clinical trials comprising 2007 participants, using the Grading of Recommendations, Assessment, Development, and Evaluations method, weak recommendations for the following were assigned: systematic provision of early information and advice, use of graded physical exercise, vestibular rehabilitation, manual treatment of neck and back, psychological treatment, and interdisciplinary rehabilitation. No studies were identified regarding oculomotor vision treatment, resulting in a consensus-based statement.

**Meaning** Based on very low to low certainty of evidence or on consensus, individually tailored nonpharmacological treatment of persistent symptoms was recommended, both through specific disciplines and interdisciplinary rehabilitation.

+ [Invited Commentary](#)

+ [Supplemental content](#)

Author affiliations and article information are listed at the end of this article.

**Open Access.** This is an open access article distributed under the terms of the CC-BY License.

Abstract (continued)

identified and included. Evidence for the 7 interventions ranged from no evidence meeting the inclusion criteria to very low and low levels of evidence. Recommendations were weak for early information and advice, graded physical exercise, vestibular rehabilitation, manual treatment of the neck and back, psychological treatment, and interdisciplinary coordinated rehabilitative treatment. No relevant evidence was identified for oculomotor vision treatment, so the panel provided a good clinical practice recommendation based on consensus.

**CONCLUSIONS AND RELEVANCE** Based on very low to low certainty of evidence or based on consensus, the guideline panel found weak scientific support for commonly applied nonpharmacological interventions to treat PPCS. Results align with recommendations in international guidelines. Intensified research into all types of intervention for PPCS is needed.

JAMA Network Open. 2021;4(11):e2132221. doi:10.1001/jamanetworkopen.2021.32221

## Introduction

Concussion or mild traumatic brain injury (mTBI) accounts for up to 90% of all TBIs.<sup>1</sup> The yearly incidence in Denmark is 457 per 100 000 inhabitants<sup>2</sup>; however, these numbers do not include those who consult general practitioners or do not seek care. The true incidence is estimated to be around 600 per 100 000.<sup>3</sup>

Concussion or mTBI is defined as a transmission of mechanical energy due to a blow to the head, neck, or body that results in disruption of brain function.<sup>4,5</sup> Consensus regarding diagnostic criteria emphasizes at least 1 of the following signs: (1) any alteration of mental state immediately after injury, (2) posttraumatic amnesia for less than 24 hours, (3) loss of consciousness for less than 30 minutes, or (4) other signs of focal and transient neurologic dysfunction.<sup>5-7</sup>

Most people who sustain a concussion or mTBI recover quickly, but a significant proportion experience long-term symptoms.<sup>8-10</sup> According to the *International Statistical Classification of Diseases and Related Health Problems, Tenth Revision (ICD-10)*,<sup>11</sup> recovery from concussion is expected within the first 2 to 3 weeks, and persistent symptoms are defined as those lasting for more than 4 weeks after injury.<sup>12</sup> Studies indicate that up to 34% to 44% of patients with concussion or mTBI experience symptoms at 3 to 6 months after injury, and between 5% and 20% experience symptoms at 12 months after injury.<sup>8,13-15</sup>

Persistent postconcussion symptoms (PPCS) comprise a combination of physical (headache, dizziness, blurred vision, sleep disturbance, neck pain, and fatigue), cognitive (slowed thinking, difficulties with attention, concentration, memory, or executive functions), and emotional or behavioral symptoms (changed emotional responsivity, irritability, quickness to anger, disinhibition, or emotional lability),<sup>16</sup> and might be associated with changes in personality and difficulties regarding personal and professional identity.<sup>17</sup> There is an uncertainty regarding the effectiveness of commonly applied nonpharmacological interventions to treat PPCS, and to our knowledge, there is a scarcity of meta-analyses dedicated to this topic. Furthermore, although systematic approaches are in general applied, the Grades of Recommendation, Assessment, Development, and Evaluation (GRADE)<sup>18</sup> approach has only been used sporadically.

The Danish Health Authority commissioned a set of National Clinical Guidelines with the objective to evaluate and summarize the evidence for the effectiveness of nonpharmacological interventions in adults experiencing PPCS and to provide recommendations for clinical practice. The selected areas of interest were (1) early information and advice, (2) graded physical exercise, (3) vestibular rehabilitation, (4) manual treatment of neck and spine, (5) oculomotor vision treatment, (6) psychological treatment, and (7) interdisciplinary coordinated rehabilitative treatment. This article reports the methodology and results of this guideline based on meta-analyses and the GRADE approach.

## Methods

The content of this guideline is structured around selected clinical questions in accordance with the Population, Intervention, Comparison, and Outcome (PICO) framework.<sup>19</sup> Making recommendations for diagnostic procedures, care pathways, or providing estimates of associated costs were beyond the scope of this guideline. Recommendations are based on systematic reviews and follow the principles described in GRADE.<sup>18</sup> This guideline followed the AGREE reporting checklist. The complete clinical guideline is available in Danish online.<sup>20</sup> A protocol for this systematic review and meta-analysis has been registered in the PROSPERO database.<sup>21</sup> The protocol for this study was reviewed and approved by the Danish Health Authority. This study followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline.

## Organization

The work was performed by a multidisciplinary group forming the guideline panel. This included a chairman, search specialist, methodologist, lead reviewer, and members appointed by professional societies and a patient organization. Members had clinical and academic expertise from neurology, physical therapy, neuropsychology, occupational therapy, optometry, chiropractic medicine, pain science, and epidemiology. The guideline panel was involved in every major step of the process. The work was coordinated by the chairman (H.M.R.) assisted by the lead reviewer (H.J.G.) and the methodologist (H.E.C.). If an included study was authored by a member of the guideline panel, the study was appraised by other members of the group. In addition, a reference group including stakeholders from the Danish health care system (ie, municipalities, hospitals, rehabilitation institutions) gave feedback on the PICO questions and the final recommendations. Any potential conflicts of interests were declared before initiating the work and can be accessed online in Danish.<sup>22</sup> The draft of the clinical guideline was peer reviewed by <sup>2</sup> external reviewers as well as in a public hearing.

## Clinical Questions

Seven clinical questions regarding the outcomes of the following nonpharmacological interventions were selected: (1) early information and advice to prevent PPCS, (2) graded physical exercise, (3) vestibular rehabilitation to treat persistent vestibular dysfunction, (4) manual treatment of the neck and back, (5) oculomotor vision treatment to treat persistent visual symptoms, (6) psychological treatment and (7) interdisciplinary coordinated rehabilitative treatment. For early information and advice, such information had to be provided systematically and start within 4 weeks after injury. In the remaining clinical questions, patients had to be diagnosed with concussion or mTBI and experience symptoms that persisted for more than 4 weeks after injury. The interventions were compared with either no intervention or treatment as usual (eTable 4 in the Supplement).

## Eligibility Criteria

To be included, studies had to (1) be intervention studies within the areas of the predefined clinical questions, (2) include a control group, and (3) focus on symptoms after concussion or mTBI. Studies including both concussion or mTBI and moderate to severe TBI were only included if it was possible to extract separate data for the concussion/mTBI population. Participants had to be aged 18 years or older and be diagnosed with concussion or mTBI. Consent (written or oral) from the participants was not obtained, as the study only used data from previously published studies. Studies including adolescents were included only if adolescent participants represented a minority of the study sample. Data on patient race and ethnicity were not collected because we did not assume that these variables would have a significant effect on the outcomes in the included studies. Collection of data on race and ethnicity was not required by the funding agency.

## Outcomes

For each clinical question, 2 primary outcomes were chosen alongside a number of secondary outcomes. The primary outcome for the majority of clinical questions was the collective burden of postconcussion symptoms supplemented by another primary outcome closely reflecting the focus of the respective interventions (eg, physical functioning after graded physical exercise, pain after manual treatment of neck and back, emotional symptoms after psychological treatment). The secondary outcomes also varied across the clinical questions and included pain, physical functioning, emotional symptoms, behavioral response, quality of life, negative effect on prognosis, and return to daily activities. A time frame for each outcome was chosen a priori. A complete overview of outcomes and time frames is available in eTable 2 in the [Supplement](#).

## Literature Searches and Study Selection

A systematic literature search was conducted from November 22, 2019, to March 3, 2020, in Embase, MEDLINE, PsycINFO, CINAHL, PEDro, OTseeker, and Cochrane Reviews (via MEDLINE and Embase) from the earliest possible publication year and up to March 3, 2020. The search strategy was subdivided into searches for relevant randomized clinical trials (RCTs) or observational studies with a control group included in either the guidelines or the systematic reviews. Finally, a search was performed to identify individual studies not included in existing guidelines or systematic reviews and for interventions for which no reviews were available. When relevant, the search date for individual studies was limited to the dates of the latest search in the included reviews (eTable 1 in the [Supplement](#)).

Titles and abstracts of potential studies were screened by the lead reviewer (H.J.G.). Subsequently, the full text of potential studies was screened by the lead reviewer (H.J.G.) and members of the working group (J.H., M.H., I.N., E.T.N-S., L.L.P., H.W.S., M.M.T., and B.Z.) for eligibility. Disagreement was discussed by consultation with a third reviewer (H.M.R.) and/or the methodologist (H.E.C.). The selection process was aided by using Covidence software, a web-based tool.<sup>23</sup> The review authors were not blinded to the journal titles, study authors, study institutions, or the year of publication. A PRISMA flowchart was created to document the number of included and excluded studies identified during the search process (eFigure 1 in the [Supplement](#)).

## Data Extraction and Quality Assessment

Information extracted included the population demographic and baseline characteristics, details on intervention and control treatments, study design, outcomes, and time of measurement. The lead reviewer and methodologist independently extracted the data and assessed the risk of bias for all RCTs using the Cochrane risk of bias tool.<sup>24</sup> If information was missing or unclear, this was noted without making any assumptions regarding the cause. If clinical guidelines or systematic reviews were identified, their quality was evaluated using the Appraisal of Guidelines for Research and Evaluation (AGREE-II)<sup>25</sup> or Assessment of Multiple Systematic Reviews (AMSTAR) tools.<sup>26</sup> Any discrepancies were identified and resolved through discussion with a third reviewer.

## Statistical Analysis

Meta-analyses were performed to give a summary measure of the outcome if data across trials for a given outcome were comparable. Analyses were performed in the program RevMan5, version 5.3 (Cochrane Collaboration). If data were inadequate for inclusion in a meta-analysis, the study results were narratively described. For dichotomous outcomes, the relative risk including the 95% CI was calculated. For continuous outcomes, effect size was calculated using standardized mean difference including the 95% CI if different measurement scales were used. If the continuous outcomes across studies were measured using the same measurement scale, effect size was assessed as mean difference. Random-effect models were used to calculate pooled estimates of effects. Statistical heterogeneity was quantified using the  $I^2$  statistic,<sup>27</sup> with an  $I^2$  value greater than 50% considered to

be substantial heterogeneity. A forest plot was provided for each meta-analysis (eFigure 2 in the Supplement).

### Certainty of Evidence

The outcome measures were combined into a single summary of findings table. The certainty of evidence was then determined according to the GRADE,<sup>28</sup> resulting in 4 possible ratings: high, moderate, low, and very low (Table 1). If applicable, downgrading was done for each outcome by evaluating the extent of risk of bias, inconsistency, indirectness, imprecision, and publication bias.<sup>28,29</sup> The overall quality of evidence for each clinical question was based on the lowest rating for the primary outcome.

### From Evidence to Recommendation

Either a strong or weak recommendation in favor of or against an intervention was made based on a combined assessment of the strength of the available evidence, assumed patient preferences, and weighting of benefits and harms. A good clinical practice recommendation was made in case there was no available evidence, ie, RCTs or observational studies with a control group (Table 2).<sup>30</sup>

**Table 1. Definitions of the Certainty of Evidence Based on the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) Approach<sup>a</sup>**

Certainty of evidence	Definition
High	We are confident that the estimated effect lies close to the true effect.
Moderate	We are moderately confident that the estimated effect is likely to be close to the true effect. However, there is a possibility that it is substantially different.
Low	We have limited confidence in the estimated effect as it may be substantially different from the true effect.
Very low	We have very limited confidence in the estimated effect, as it is likely to be substantially different from the true effect.

<sup>a</sup> This table was adapted from Balshem et al.<sup>28</sup>

**Table 2. Definitions of Recommendations Based on the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) Approach and by the Danish Health Authority<sup>a</sup>**

Recommendation	Definition
Strong recommendation for	A strong recommendation in favor of an intervention is given when there is high-quality evidence showing that the overall benefits of the intervention are clearly greater than the disadvantages. The majority of the patients would want the intervention.
Weak recommendation for	A weak recommendation in favor of an intervention is given when it is assessed that the advantages of the intervention outweigh the disadvantages or if the available evidence cannot rule out a significant benefit of the intervention while at the same time the harmful effects are few or absent. This recommendation is also given when there are substantial variations in patient preferences.
Strong recommendation against	A strong recommendation against an intervention is given when there is high-quality evidence showing that the overall disadvantages of the intervention are clearly greater than the benefits. The majority of the patients would not want the intervention.
Weak recommendation against	A weak recommendation against an intervention is given when it is assessed that the disadvantages of the intervention outweigh the advantages but where it is not substantiated by high-quality evidence. This recommendation is also used where there is high-quality evidence for both beneficial and harmful effects but where the balance between them is difficult to determine. This recommendation is also given when there are substantial variations in patient preferences.
Good clinical practice statement	A good clinical practice statement is used when there is no relevant evidence to answer the clinical questions and thus the recommendation is based on professional consensus among the members of the working group that drafted the guideline. The recommendation can be either for or against the intervention. Because this is based on professional consensus, this type of recommendation is weaker than any evidence-based recommendation.

<sup>a</sup> This table was adapted from the Danish Health Authority.

## Results

A total of 13 097 articles were identified; of those, 109 were excluded as duplicates. Among the remaining 12 988 articles, titles and abstracts were screened for inclusion and exclusion criteria. A total of 297 articles received a full-text review, and 11 relevant systematic reviews<sup>31-41</sup> were identified for 5 of the clinical questions. Only 1 systematic review<sup>33</sup> matched our clinical question (graded physical exercise) and was of sufficient methodological quality (eTable 3 in the Supplement); however, it did not contribute any primary study matching our inclusion and exclusion criteria. The search date for individual studies for this question was then limited to the date of the latest search in the included systematic review,<sup>33</sup> and here we identified 2 relevant RCTs.<sup>42,43</sup> Furthermore, 1 relevant RCT<sup>44</sup> was identified in a systematic review for manual treatment of neck and back. A total of 19 individual RCTs<sup>42-60</sup> comprising 2007 participants (1064 women [53.0%]; 943 men [47.0%]) were identified for 6 of the clinical questions. Nine<sup>43,45-47,49,52,53,55,60</sup> of these included a minority of adolescents in the sample. No evidence was identified for the effect of oculomotor vision treatment. Thus, all recommendations were based exclusively on primary studies. For characteristics of the included studies, see eTable 5 in the Supplement. For flowcharts of included literature, see eFigure 1 in the Supplement.

The available evidence was limited for all interventions, ranging from low to very low mainly owing to risk of bias, imprecision (few studies identified or few patients included), and indirectness (patients aged <18 years included) (Table 3). For results of meta-analyses, see eFigure 2 in the Supplement. For summary of findings table, including the rating of the evidence and risk of bias assessment, see eTable 4 in the Supplement. For recommendations including the supporting evidence, see Table 4.

## Recommendations

A weak recommendation was given for systematically providing early information and advice, the use of graded physical exercise, the use of vestibular rehabilitation, the use of manual treatment of neck and back, psychological treatment, and interdisciplinary rehabilitative treatment. No relevant evidence was identified for oculomotor vision treatment, so the panel provided a good clinical practice recommendation based on consensus (Tables 3 and 4).

## Discussion

This guideline considered 7 clinical questions regarding nonpharmacological management of PPCS. Based on very low to low certainty of evidence, the guideline panel provided weak recommendations for early information and advice, graded physical exercise, vestibular rehabilitation for persistent vestibular dysfunction, manual treatment of the neck and back, psychological treatment, and interdisciplinary coordinated rehabilitative treatment. A consensus-based recommendation was

Table 3. Overview of Recommendations in the Guideline and the Certainty of Evidence<sup>a</sup>

PICO	Intervention	Certainty of evidence	Recommendation
PICO 1	Systematically offered information and advice	Very low	Weak recommendation for
PICO 2	Graded physical exercise	Very low	Weak recommendation for
PICO 3	Vestibular rehabilitation	Very low	Weak recommendation for
PICO 4	Spinal manual therapy	Very low	Weak recommendation for
PICO 5	Oculomotor vision treatment	No relevant evidence identified	Good clinical practice statement
PICO 6	Psychological treatment	Low	Weak recommendation for
PICO 7	Interdisciplinary coordinated rehabilitative treatment	Low	Weak recommendation for

Abbreviation: PICO, Population, Intervention, Comparison, and Outcome.

<sup>a</sup> Definitions of certainty of evidence and recommendations are noted in Tables 1 and 2.



**Table 4. Overall Population, Intervention, Comparison, and Outcome (PICO) Questions, Recommendations, Definitions of Interventions, and Primary Outcomes, Supporting Evidence, and Rationale**

Recommendation	Definition of intervention	Primary outcomes	Included studies	Rationale
<b>PICO 1. Should patients, early after concussion, be systematically offered early information and advice on preventing persistent postconcussive symptoms?</b>				
Consider systematically offering early information and advice to patients within the first 4 wk after concussion.	Definition of intervention: Systematic education, instructions, advice and guidance regarding postconcussion symptoms, symptom management, restitution, and self-care provided individually or in groups, either in person or as telephone guidance by a health care professional, using oral and/or written information. The intervention must be initiated within the first 4 wk after injury and should be provided by relevant health professionals.	<ol style="list-style-type: none"> <li>1. The collective burden of postconcussion symptoms assessed a minimum of 2 wk after completed intervention</li> <li>2. Emotional symptoms assessed a minimum of 1 mo after completed intervention</li> </ol>	Bell et al, <sup>45</sup> Heskestad et al, <sup>46</sup> Matuseviciene et al, <sup>47</sup> Mittenberg et al, <sup>48</sup> Ponsford et al, <sup>49</sup> Suffoletto et al, <sup>50</sup> and Varner et al <sup>51</sup>	The intervention was associated with a positive effect on the overall symptom burden 2 wk after completion. Furthermore, intervention reduced the number of patients who subsequently experienced memory problems and the number of patients in which leisure and working life was affected. There were no reported serious adverse effects; however, this was not systematically assessed. The certainty of evidence was very low due to risk of bias, indirectness, and imprecision. It was assumed that there was no substantial variability in terms of patient preferences and that the majority of patients would want the intervention. Based on a collective assessment of these findings, a weak recommendation was given for the use of systematic information and advice.
<b>PICO 2. Should patients with persistent postconcussive symptoms be offered graded physical exercise?</b>				
Consider offering graded physical exercise in addition to other treatment to patients with persistent postconcussion symptoms.	Definition of intervention: Graded physical exercise, ie, physical exercise with a gradual increase in intensity and/or complexity over time, such as general physiotherapy, general physical activity, sensorimotor training, aerobic and anaerobic training, performed minimally 1 time/wk for 4 wk.	<ol style="list-style-type: none"> <li>1. The collective burden of postconcussion symptoms assessed at the end of completed intervention</li> <li>2. Physical functioning assessed at the end of completed intervention</li> </ol>	Rytter et al <sup>42</sup> and Thastum et al <sup>43</sup>	The intervention in addition to other treatment was associated with a positive effect on both the overall burden of symptoms, the level of physical functioning, behavioral reactions, emotional symptoms, quality of life, and the general satisfaction with the current work situation. There were no reported serious adverse events; however, this was not systematically assessed. The certainty of the evidence was very low due to serious risk of bias, indirectness, and imprecision. It was assumed that there was no substantial variability in terms of patient preferences and that the majority of patients would want the intervention. Based on a collective assessment of these findings, a weak recommendation was given for the use of graded physical exercise.
<b>PICO 3. Should patients with persistent vestibular dysfunction after concussion be offered vestibular rehabilitation?</b>				
Consider offering vestibular rehabilitation in addition to other treatments to patients who experience persistent vestibular dysfunction after concussion.	Vestibular rehabilitation, including the otolith manipulating procedures, habituation and adaptation exercises, substitution training, and balance training administered minimally 1 time/wk for a period of 4 wk.	<ol style="list-style-type: none"> <li>1. The collective burden of postconcussion symptoms assessed at the end of completed intervention</li> <li>2. Vestibular dysfunction assessed at the end of completed intervention</li> </ol>	Kleffelgaard et al <sup>53</sup> and Schneider et al <sup>52</sup>	The intervention was associated with a positive effect on the level of physical functioning as well as the number of patients considered ready to return to sport after completed intervention. No serious adverse events were reported; however, this was not systematically assessed. The certainty of the evidence was very low due to risk of bias, indirectness, and imprecision. It was assumed that there was no substantial variability in terms of patient preferences and that the majority of patients would want the intervention. Based on a collective assessment of these findings, a weak recommendation was given for the use of vestibular rehabilitation.
<b>PICO 4. Should patients with persistent postconcussive symptoms be offered spinal manual therapy, ie, mobilization and manipulation of the neck and spine?</b>				
Consider offering manual treatment of neck and spine in addition to other treatments to patients with persistent symptoms after concussion.	Manual therapy in the form of hands-on mobilization and/or manipulation of the spine or other joints, typically performed by physiotherapists or chiropractors.	<ol style="list-style-type: none"> <li>1. Physical functioning assessed at the end of completed intervention</li> <li>2. Pain assessed at the end of completed intervention</li> </ol>	Schneider et al <sup>52</sup> and Jensen et al <sup>44</sup>	The intervention was associated with a positive effect on pain as well as the number of patients considered ready to return to sport after completed intervention. No serious adverse events were reported; however, this was not systematically assessed. The certainty of the evidence was very low due to risk of bias, indirectness, and imprecision. It is expected that the intervention will include differences in patient preferences, as some patient would want the treatment whereas others would not. Based on a collective assessment of these findings, a weak recommendation was given for the use of spinal manual therapy.

(continued)



Table 4. Overall Population, Intervention, Comparison, and Outcome (PICO) Questions, Recommendations, Definitions of Interventions, and Primary Outcomes, Supporting Evidence, and Rationale (continued)

Recommendation	Definition of intervention	Primary outcomes	Included studies	Rationale
<b>PICO 5. Should patients with persistent visual symptoms after concussion be offered oculomotor vision treatment?</b>				
It is good clinical practice to consider offering oculomotor vision treatment to patients who experience persistent visual symptoms after concussion.	Oculomotor vision treatment, ie, oculomotor training to treat vergence, accommodative, or eye movement dysfunction after concussion, including computer-based training and optometric instrumental training administered as an optometric session minimally 1 time/wk during a period of 4 wk.	<ol style="list-style-type: none"> <li>Oculomotor dysfunction assessed at the end of completed intervention</li> <li>Visual functioning assessed at the end of completed intervention</li> </ol>	No relevant trials identified	Clinical experience shows that oculomotor visual treatment improves the visual symptoms as well as reduces other symptoms such as headache and tiredness in patients with persistent symptoms after concussion. In addition, there is clinical consensus that oculomotor visual therapy has a positive effect on the number of patients returning to work. Because there were no relevant trials identified, this recommendation is largely based on this clinical experience. There are, however, peer-reviewed studies without a control group, showing a positive effect of oculomotor visual therapy in patients with concussion (Gallaway et al <sup>61</sup> ; Thiagarajan and Cuiffreda <sup>62</sup> ; Thiagarajan et al <sup>63</sup> ; Thiagarajan and Cuiffreda <sup>64</sup> ). Based on a collective assessment of these findings, a good clinical practice statement was given for the use of visual therapy.
<b>PICO 6. Should patients with persistent post-concussive symptoms be offered psychological treatment?</b>				
Consider offering psychological treatment in addition to other treatment to patients with persistent symptoms after concussion.	Psychological treatment by psychologists or clinicians with similar professional background administered minimally 1 h/wk as either individual or group therapy for a period of minimally 4 wk.	<ol style="list-style-type: none"> <li>The collective burden of postconcussion symptoms assessed after completed intervention</li> <li>Emotional symptoms assessed a minimum of 3 mo after completed intervention</li> </ol>	Caplain et al, <sup>54</sup> Elgmark-Andersson et al, <sup>55</sup> Kjeldgaard et al, <sup>56</sup> Potter et al, <sup>57</sup> Rytter et al, <sup>42</sup> Silverberg et al, <sup>58</sup> Thastum et al, <sup>43</sup> Tiersky et al, <sup>59</sup> and Vikane et al <sup>60</sup>	The intervention was associated with a positive effect on the overall burden of symptoms after the completion of the intervention as well as at longest follow-up. A positive effect was also seen with respect to emotional symptoms and quality of life at the longest follow-up. No serious adverse events were reported; however, this was not systematically assessed. The overall certainty of evidence was low due to risk of bias and indirectness. It is expected that the intervention will include differences in patient preferences, as some patient would want the treatment whereas others would not. Based on a collective assessment of these findings, a weak recommendation was given for the use of psychological treatment.
<b>PICO 7. Should patients with persistent postconcussive symptoms be offered an interdisciplinary coordinated rehabilitative treatment?</b>				
Consider offering interdisciplinary coordinated rehabilitative treatment to patients with persistent symptoms after concussion.	An interdisciplinary coordinated rehabilitative treatment is a treatment provided by health professionals from at least 2 different disciplines, who collaborate on the rehabilitation of the patient. The treatment includes at least 2 interventions, eg, vestibular rehabilitation, graded physical exercise, oculomotor vision therapy, manual treatment, (neuro)psychological and psychotherapeutic intervention, advice on managing everyday activities, and vocational rehabilitation and appears as a comprehensive interdisciplinary approach. The treatment is administered minimally 1 time/wk during a period of 4 wk by clinicians with relevant background, eg, physiotherapists, occupational therapist, nurses, rehabilitation therapists, (neuro) psychologists, neurologists.	<ol style="list-style-type: none"> <li>The collective burden of postconcussion symptoms assessed after completed intervention</li> <li>Return to daily activities assessed a minimum of 3 month after completed intervention</li> </ol>	Rytter et al, <sup>42</sup> Thastum et al, <sup>43</sup> and Vikane et al <sup>60</sup>	The intervention was associated with a positive effect on the overall burden of symptoms, the level of physical functioning, emotional symptoms, as well as on quality of life and the general satisfaction with work life. No serious adverse events were reported; however, this was not systematically assessed. The overall certainty of evidence was low due to risk of bias and indirectness. It was assumed that there was no substantial variability in terms of patient preferences and that the majority of patients would want the intervention. Based on a collective assessment of these findings, a weak recommendation was given for the use of interdisciplinary coordinated rehabilitative treatment

provided for oculomotor vision treatment to treat persistent visual symptoms. The guideline panel concluded that intensified research into all intervention types is needed.

Given that concussion or mTBI can be associated with persistent symptoms and disability,<sup>65-68</sup> there has been a shift from passively awaiting symptom remission to recommending active management when symptoms persist. Thus, our recommendations align with the Synthesis of

Practice Guidelines from the American Congress of Rehabilitation Medicine's Mild TBI task force<sup>4</sup>; the American Physical Therapy Association's (APTA) guideline for physical therapy evaluation and treatment after concussion or mild TBI<sup>69</sup>; the third edition of the Ontario Neurotrauma Foundation's (ONF) Guideline for concussion or mild TBI and persistent symptoms<sup>5</sup>; and the latest update on the consensus statement published by the Concussion in Sports Group (CISG).<sup>70</sup> However, compared with our guideline, these guidelines have different scopes and include pediatric and adolescent populations. Furthermore, although these guidelines apply a systematic approach to guideline generation, they do not apply the GRADE approach.

Early information and advice was recommended based on 7 RCTs,<sup>45-51</sup> of which 6 were included in the meta-analysis. The study by Ponsford et al<sup>49</sup> was not included because it was presented as a short report. Collectively, the studies concluded that information should be tailored to individual patient needs and that information and advice are particularly beneficial if provided over a longer period of time. This recommendation aligns with the Synthesis of Practice Guidelines<sup>4</sup> as well as with the ONF<sup>5</sup> and APTA<sup>69</sup> guidelines that give a grade A, or strong, recommendation. The CISG guideline<sup>70</sup> does not address this question. Our guideline provides a weak recommendation based on GRADE owing to possible risks of bias in the included studies as well as indirectness.

Graded physical exercise was recommended based on 2 RCTs.<sup>42,43</sup> In both, the intervention was part of interdisciplinary rehabilitation programs with several interventions applied in parallel. Critical outcomes included physical functioning, which presumably reflected the effect of graded physical exercise. Both the APTA<sup>69</sup> and the CISG guideline<sup>70</sup> recommend graded physical exercise as a treatment of autonomic instability or physical deconditioning, with the APTA<sup>69</sup> indicating level A, or strong evidence. The ONF guideline<sup>5</sup> recommends exercise below symptom threshold in management of posttraumatic fatigue (grade C). The Synthesis of Practice Guidelines<sup>4</sup> advocates exercise in general. Our guideline arrived at a weak recommendation based on GRADE.

Vestibular rehabilitation was recommended based on 2 RCTs.<sup>52,53</sup> In both, participants also received exercise and manual treatment of the cervical and thoracic spine. The ONF<sup>5</sup> and APTA guidelines<sup>69</sup> give a grade A, or strong, recommendation for dealing with benign paroxysmal positional vertigo. Furthermore, the ONF guideline gives a grade A recommendation for treatment of unilateral peripheral vestibular dysfunction, whereas the APTA guideline indicates level B, or moderate evidence, for vestibular rehabilitation in conjunction with oculomotor rehabilitation. The ONF guideline gives a grade C consensus-based recommendation, and APTA indicates level C, or weak evidence, for treatment of functional balance impairment. The CISG guideline<sup>70</sup> recommends a targeted physical therapy program for patients with vestibular dysfunction without further specification.

Manual treatment of the neck and back, ie, mobilization and/or manipulation, is recommended based on 2 small RCTs,<sup>44,52</sup> of which only one<sup>52</sup> could be included in the meta-analysis owing to poor reporting in the other.<sup>44</sup> The 2 studies indicate possible beneficial outcomes regarding pain and return to sport, resulting in a weak recommendation. The Synthesis of Practice Guidelines<sup>4</sup> and the ONF guideline<sup>5</sup> do not include manual treatment of the neck and back. The APTA guideline recommends that practitioners address cervical and thoracic spine dysfunction, including manual treatment based on level B, or moderate, evidence.<sup>69</sup> The CISG guideline recommends targeted physical therapy in patients with cervical dysfunction without further details.<sup>70</sup>

Oculomotor vision treatment is recommended based on consensus in the guideline panel. Low-quality studies without appropriate control groups, however, do exist and report positive effects.<sup>61-64</sup> The ONF guideline<sup>5</sup> gives grade C, or a consensus-based recommendation, for vision assessment and treatment, whereas the APTA guideline<sup>69</sup> indicates level B, or moderate, evidence. However, this guideline deals with oculomotor rehabilitation in combination with vestibular treatment, which may explain the discrepancy. The CISG guideline<sup>70</sup> does not address this question.

Psychological treatment is recommended based on 9 RCTs,<sup>42,43,54-60</sup> of which 8 were included in the meta-analysis.<sup>42,43,54-60</sup> Tiersky et al<sup>59</sup> was not included owing to insufficient reporting. Psychological interventions may be applied for multiple reasons, eg, to treat mental health disorders

(primarily anxiety and depression), to address cognitive symptoms, and/or to change maladaptive behavioral strategies. Accordingly, the RCTs applied a combination of different approaches, such as psychoeducation, counseling, cognitive behavioral therapy, computer-based cognitive remediation training, and energy management. The Synthesis of Practice Guidelines<sup>4</sup> emphasizes primarily an active approach to mental health disorders, with fast initiation of cognitive behavioral therapy and/or pharmacotherapy. The ONF guideline<sup>5</sup> gives a grade B recommendation regarding a routine screen for depression and anxiety followed by a specialist treatment, and a grade B recommendation is given for cognitive behavioral therapy as a supplementary intervention for patients with psychological risk factors. The ONF guideline<sup>5</sup> gives a grade A recommendation for neuropsychological assessment, grade B recommendation for treatment of persistent cognitive difficulties, and grade C recommendation for stress management techniques. The CISG guideline<sup>70</sup> recommends cognitive behavioral therapy for persistent mood disorders or behavioral issues.

Interdisciplinary coordinated rehabilitation was recommended based on 3 RCTs.<sup>42,43,60</sup> Their results indicate its usefulness regarding the primary outcome (collective burden of symptoms) as well as several secondary outcomes (Table 4). The Synthesis of Practice Guidelines<sup>4</sup> highlights the need for individually tailored interdisciplinary treatment. The ONF guideline<sup>5</sup> gives a grade A recommendation for evidence-based neurorehabilitation for patients with cognitive impairment and a grade B recommendation for interdisciplinary vocational assessment for those who do not resume preinjury work duties. The CISG guideline<sup>70</sup> recommends a psychological, cervical, and vestibular rehabilitation performed in a collaborative manner based on detailed assessment of physical and psychosocial factors.

Even though the development of the guideline followed a rigorous methodology, our confidence in the recommendations is low or very low. This low confidence is due to scarcity and inadequacies of the underlying literature wherein studies have risk of bias due to investigating the effects of experiential interventions with difficulties related to blinding of participants and therapists. Moreover, in clinic, interventions were often used in combination as part of a multimodal approach with multiple interventions given in parallel.<sup>10,42</sup>

The weak recommendations call for shared decision-making, involving a discussion with the patient. People with PPCS represent a heterogeneous group regarding both their trauma-induced consequences and their premorbid functioning levels, coping strategies, and life situation in general.<sup>71</sup>

Expert groups have used the lack of evidence for benefit or harm for a particular intervention as an argument for not putting forward a recommendation.<sup>72</sup> Such positions have, however, been met with frustration by health care professionals who look to expert groups for guidance.<sup>73</sup> Fortunately, the GRADE approach accommodates these circumstances because it provides interpretations for patients, clinicians, and policy makers.<sup>30</sup> Faced with either no or weak evidence, it is important that patients know that their particular preference among the various treatments should guide choice of intervention; clinicians must recognize that different interventions may be appropriate for different patients and help each patient to arrive at a management decision consistent with their values. Policy makers must involve relevant professional groups and stakeholders when determining design-of-care pathways.<sup>30</sup> Importantly, guideline panels should not refrain from making recommendations because individual patients and clinicians will make different choices when faced with a weak recommendation. In fact, this is to be expected. Consequently, the GRADE Working Group encourages panels to make recommendations whenever possible whether they are based on solid evidence or not.<sup>30</sup>

### Strengths and Limitations

Strengths of this guideline include the broad scope informed by a need for direction in clinical practice. The guideline panel adhered rigorously to the GRADE methodology. The group was composed of experienced academics and clinicians within this research area, an expert librarian, and a skilled methodologist. The process was overseen by the Danish Health Authority, and the final

product was reviewed by<sup>2</sup> external experts who provided valuable input to the final draft. Finally, the interdisciplinary reference group and the public hearing strengthened the outreach to clinical practice and health administration.

Study limitations include the fact that the guideline panel itself selected the interventions in focus. Also, we focused on adults, so the recommendations do not cover adolescents, leading to exclusion of several studies.<sup>74,75</sup> Last, RCTs are not well suited to address potential harms associated with treatments; however, the guideline panel considered the included interventions to be associated with low frequency of harmful consequences.

---

## Conclusions

There is an urgent need for more methodologically robust research evaluating the outcomes of nonpharmacological treatments for persistent symptoms after concussion or mTBI. Given the best available evidence to date, and based on the findings of this systematic review and meta-analysis, active management and treatment of PPCS is recommended, both through individual disciplines targeting specific problems and through interdisciplinary rehabilitation. There was agreement on this recommendation across the available guidelines, including the one presented here, regardless of their applied methodology.

---

## ARTICLE INFORMATION

**Accepted for Publication:** August 31, 2021.

**Published:** November 9, 2021. doi:10.1001/jamanetworkopen.2021.32221

**Open Access:** This is an open access article distributed under the terms of the [CC-BY License](#). © 2021 Rytter HM et al. *JAMA Network Open*.

**Corresponding Author:** Hana Malá Rytter, PhD, Danish Concussion Center, Amagerfaelledvej 56A, 2300 Copenhagen S, Denmark ([hana.mala@psy.ku.dk](mailto:hana.mala@psy.ku.dk)).

**Author Affiliations:** Danish Concussion Center, Copenhagen, Denmark (Rytter, Graff, Henriksen); Department of Psychology, University of Copenhagen, Copenhagen, Denmark (Rytter); Department of Neurology, Bispebjerg-Frederiksberg University Hospital, Copenhagen, Denmark (Rytter); Center for Rehabilitation of Brain Injury, Copenhagen, Denmark (Henriksen); The Danish Concussion Association, Copenhagen, Denmark (Aaen); Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark (Hartvigsen); Chiropractic Knowledge Hub, Odense, Denmark (Hartvigsen); Musculoskeletal Health and Implementation, Department of Medicine, Aalborg University, Aalborg, Denmark (Hoegh); Pain Science Educator, Aarhus, Denmark (Hoegh); Danish College of Optometry, Dania Academy, Randers, Denmark (Nisted); Department of Clinical Medicine, Aarhus University, Aarhus, Denmark (Nisted); Hammel Neurorehabilitation Centre-University Clinic for Neurorehabilitation, Aarhus University, Aarhus, Denmark (Næss-Schmidt, Thastum); Alleens Physiotherapy and Sports Clinic, Odense, Denmark (Pedersen); Danish Headache Center, Rigshospitalet-Glostrup, University of Copenhagen, Copenhagen, Denmark (Schytz); Exercise and Health Training Center, Roskilde Municipality, Roskilde, Denmark (Zerlang); Callesen Method Consulting, Copenhagen, Denmark (Callesen).

**Author Contributions:** Dr Rytter had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

*Concept and design:* Rytter, Graff, Hoegh, Næss-Schmidt, Thastum, Callesen.

*Acquisition, analysis, or interpretation of data:* All authors.

*Drafting of the manuscript:* Rytter, Pedersen, Callesen.

*Critical revision of the manuscript for important intellectual content:* All authors.

*Statistical analysis:* Graff, Hartvigsen, Callesen.

*Obtained funding:* Rytter.

*Administrative, technical, or material support:* Rytter, Graff, Henriksen, Pedersen, Schytz, Callesen.

*Supervision:* Rytter, Aaen, Hartvigsen, Nisted, Næss-Schmidt, Pedersen, Zerlang, Callesen.

Review and grading of included articles: Hartvigsen.

Lead reviewer, formulation of the PICO questions, selection of studies, validating the data extraction and risk of bias evaluation, and rating the certainty of the evidence and formulating the final recommendations: Graff.

**Conflict of Interest Disclosures:** Dr Rytter reported receiving grants from the Danish Ministry of Health where her employer received financial compensation during the conduct of the study. Dr Graff reported receiving grants from the Danish Ministry of Health during the conduct of the study and outside the submitted work. Mr Aaen reported receiving fees from the Danish Health Authority for membership in guideline development during the conduct of the study. Dr Hartvigsen reported receiving fees from the Danish Health Authority for membership in guideline development during the conduct of the study. Dr Hoegh reported receiving personal fees from the Danish Health Authority during the conduct of the study; personal fees from Acatalepsia ApS MH; receiving fees for lectures related to pain and rehabilitation; and receiving royalties for books and book chapters he has written outside the submitted work. Mr Nisted reported receiving fees from the Danish Health Authority for membership in guideline development during the conduct of the study. Dr Næss-Schmidt reported receiving fees from the Danish Health Authority for membership in guideline development during the conduct of the study. Ms Pedersen reported receiving personal fees from the Danish Concussion Center, Copenhagen, Denmark, and from the Danish Health Authority for membership in guideline development during the conduct of the study. Dr Callesen reported receiving personal fees from the Danish Concussion Center, Copenhagen, Denmark, and the Danish Health Authority for consulting during the conduct of the study. No other disclosures were reported.

**Funding/Support:** Support for this study was provided in part by the Danish Ministry of Health based on The Danish Finance Act in 2017, and the work was approved and overseen by the Danish Health Authority. Members of the project group (Drs Rytter, Graff, Ms Henriksen, Dr Callesen) received financial compensation. Likewise, members of the working group (Mr Aaen, Dr Hoegh, Dr Hartvigsen, Mr Nisted, Dr Næss-Schmidt, Ms Pedersen) received a salary if they were not employed by the Danish Regions or Danish Municipalities. The work of members employed by the Danish Regions or Danish Municipalities (Drs Schytz, Thastum, Ms Zerlang) was funded by their employer. The members of the reference group did not receive any salary. Additional funding was provided by the Danish Concussion Center.

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Additional Contributions:** We thank information specialist Kirsten Birkefoss, MSc, for assistance with systematic literature searches and Ida Laenholm for assistance with preparation of the material for the manuscript. Kirsten Birkefoss was financially compensated for her contribution.

## REFERENCES

1. Skandsen T, Einarsen CE, Normann I, et al. The epidemiology of mild traumatic brain injury: the Trondheim MTBI follow-up study. *Scand J Trauma Resusc Emerg Med*. 2018;26(1):34. doi:10.1186/s13049-018-0495-0
2. Pinner M, Børgensen SE, Jensen R, Birket-Smith M, Gade A, Riis JO. *Consensus Report on Commotio Cerebri (Concussion). Konsensusrapport Om Commotio Cerebri (Hjernerystelse)*. Videnscenter for Hjerneskade; 2003:1-82.
3. Cassidy JD, Carroll LJ, Peloso PM, et al; WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*. 2004;43(suppl):28-60. doi:10.1080/16501960410023732
4. Silverberg ND, Iaccarino MA, Panenka WJ, et al; American Congress of Rehabilitation Medicine Brain Injury Interdisciplinary Special Interest Group Mild TBI Task Force. Management of concussion and mild traumatic brain injury: a synthesis of practice guidelines. *Arch Phys Med Rehabil*. 2020;101(2):382-393. doi:10.1016/j.apmr.2019.10.179
5. Marshall S, Bayley MT, McCullagh S, et al. *Guideline for Concussion/Mild Traumatic Brain Injury & Prolonged Symptoms*. 3rd ed. Ontario Neurotrauma Foundation; 2018.
6. Sussman ES, Pendharkar AV, Ho AL, Ghajar J. Mild traumatic brain injury and concussion: terminology and classification. *Handb Clin Neurol*. 2018;158:21-24. doi:10.1016/B978-0-444-63954-7.00003-3
7. Carroll LJ, Cassidy JD, Holm L, Kraus J, Coronado VG; WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. Methodological issues and research recommendations for mild traumatic brain injury: the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*. 2004;43(suppl):113-125. doi:10.1080/16501960410023877
8. van der Naalt J, Timmerman ME, de Koning ME, et al. Early predictors of outcome after mild traumatic brain injury (UPFRONT): an observational cohort study. *Lancet Neurol*. 2017;16(7):532-540. doi:10.1016/S1474-4422(17)30117-5

9. Carroll LJ, Cassidy JD, Cancelliere C, et al. Systematic review of the prognosis after mild traumatic brain injury in adults: cognitive, psychiatric, and mortality outcomes: results of the International Collaboration on Mild Traumatic Brain Injury Prognosis. *Arch Phys Med Rehabil*. 2014;95(3)(suppl):S152-S173. doi:10.1016/j.apmr.2013.08.300
10. Hartvigsen J, Boyle E, Cassidy JD, Carroll LJ. Mild traumatic brain injury after motor vehicle collisions: what are the symptoms and who treats them? a population-based 1-year inception cohort study. *Arch Phys Med Rehabil*. 2014;95(3)(suppl):S286-S294. doi:10.1016/j.apmr.2013.07.029
11. World Health Organization. *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. WHO; 1992.
12. Bellner J, Jensen SM, Lexell J, Romner B. Diagnostic criteria and the use of ICD-10 codes to define and classify minor head injury. *J Neurol Neurosurg Psychiatry*. 2003;74(3):351-352. doi:10.1136/jnnp.74.3.351
13. Clossen MC, van der Naalt J, Spikman JM, et al. Prediction of persistent post-concussion symptoms after mild traumatic brain injury. *J Neurotrauma*. 2018;35(22):2691-2698. doi:10.1089/neu.2017.5486
14. Ponsford J, Cameron P, Fitzgerald M, Grant M, Mikocka-Walus A, Schönberger M. Predictors of postconcussive symptoms 3 months after mild traumatic brain injury. *Neuropsychology*. 2012;26(3):304-313. doi:10.1037/a0027888
15. Voormolen DC, Haagsma JA, Polinder S, et al. Post-concussion symptoms in complicated vs. uncomplicated mild traumatic brain injury patients at three and six months post-injury: results from the CENTER-TBI Study. *J Clin Med*. 2019;8(11):E1921. doi:10.3390/jcm8111921
16. Belanger HG, Barwick FH, Kip KE, Kretzmer T, Vanderploeg RD. Postconcussive symptom complaints and potentially malleable positive predictors. *Clin Neuropsychol*. 2013;27(3):343-355. doi:10.1080/13854046.2013.774438
17. Snell DL, Martin R, Surgenor LJ, Siegert RJ, Hay-Smith EJC. What's wrong with me? seeking a coherent understanding of recovery after mild traumatic brain injury. *Disabil Rehabil*. 2017;39(19):1968-1975. doi:10.1080/09638288.2016.1213895
18. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-394. doi:10.1016/j.jclinepi.2010.04.026
19. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol*. 2011;64(4):395-400. doi:10.1016/j.jclinepi.2010.09.012
20. Rytter HM, Graff HJ, Henriksen HK, et al. National clinical guideline for nonpharmacological treatment of prolonged symptoms after concussion. National klinisk retningslinje for non-farmakologisk behandling af længerevarende symptomer efter hjernerystelse. Accessed August 31, 2021. [https://files.magicapp.org/guideline/ad4ee090-13ea-4f32-bcd6-c1ddb8bfe064/published\\_guideline\\_4774-0\\_5.pdf](https://files.magicapp.org/guideline/ad4ee090-13ea-4f32-bcd6-c1ddb8bfe064/published_guideline_4774-0_5.pdf)
21. Rytter HM, Graff HJ, Henriksen H, et al. Non-pharmacological treatments for persistent symptoms after concussion. April 28, 2020. Accessed June 16, 2021. [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020165194](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020165194)
22. Dansk Center for Hjernerystelse. The national clinical guidelines for concussion. National klinisk retningslinje for hjernerystelse. Accessed August 31, 2021. <https://dcfh.dk/national-klinisk-retningslinje/>
23. Covidence. Better systematic review management. Accessed August 31, 2021. <https://www.covidence.org/home>
24. Cumpston M, Li T, Page MJ, et al. Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for Systematic Reviews of Interventions. *Cochrane Database Syst Rev*. 2019;10:ED000142. doi:10.1002/14651858.ED000142
25. Brouwers MC, Kho ME, Browman GP, et al; AGREE Next Steps Consortium. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ*. 2010;182(18):E839-E842. doi:10.1503/cmaj.090449
26. Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol*. 2007;7:10. doi:10.1186/1471-2288-7-10
27. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558. doi:10.1002/sim.1186
28. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. 2011;64(4):401-406. doi:10.1016/j.jclinepi.2010.07.015
29. Guyatt GH, Oxman AD, Vist G, et al. GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *J Clin Epidemiol*. 2011;64(4):407-415. doi:10.1016/j.jclinepi.2010.07.017



30. Guyatt GH, Oxman AD, Kunz R, et al; GRADE Working Group. Going from evidence to recommendations. *BMJ*. 2008;336(7652):1049-1051. doi:10.1136/bmj.39493.646875.AE
31. Eliyahu L, Kirkland S, Campbell S, Rowe BH. The effectiveness of early educational interventions in the emergency department to reduce incidence or severity of postconcussion syndrome following a concussion: a systematic review. *Acad Emerg Med*. 2016;23(5):531-542. doi:10.1111/acem.12924
32. Gravel J, D'Angelo A, Carrière B, et al. Interventions provided in the acute phase for mild traumatic brain injury: a systematic review. *Syst Rev*. 2013;2:63. doi:10.1186/2046-4053-2-63
33. Lal A, Kolakowsky-Hayner SA, Ghajar J, Balamane M. The effect of physical exercise after a concussion: a systematic review and meta-analysis. *Am J Sports Med*. 2018;46(3):743-752. doi:10.1177/0363546517706137
34. Quatman-Yates C, Cupp A, Gunsch C, Haley T, Vaculik S, Kujawa D. Physical rehabilitation interventions for post-mTBI symptoms lasting greater than 2 weeks: systematic review. *Phys Ther*. 2016;96(11):1753-1763. doi:10.2522/ptj.20150557
35. Kinne BL, Bott JL, Cron NM, Iaquaniello RL. Effectiveness of vestibular rehabilitation on concussion-induced vertigo: a systematic review. *Phys Ther Rev*. 2018;23(6):338-347. doi:10.1080/10833196.2018.1517032
36. Murray DA, Meldrum D, Lennon O. Can vestibular rehabilitation exercises help patients with concussion? a systematic review of efficacy, prescription and progression patterns. *Br J Sports Med*. 2017;51(5):442-451. doi:10.1136/bjsports-2016-096081
37. Schneider KJ, Leddy JJ, Guskiewicz KM, et al. Rest and treatment/rehabilitation following sport-related concussion: a systematic review. *Br J Sports Med*. 2017;51(12):930-934. doi:10.1136/bjsports-2016-097475
38. Al Sayegh A, Sandford D, Carson AJ. Psychological approaches to treatment of postconcussion syndrome: a systematic review. *J Neurol Neurosurg Psychiatry*. 2010;81(10):1128-1134. doi:10.1136/jnnp.2008.170092
39. Bergersen K, Halvorsen JO, Tryti EA, Taylor SI, Olsen A. A systematic literature review of psychotherapeutic treatment of prolonged symptoms after mild traumatic brain injury. *Brain Inj*. 2017;31(3):279-289. doi:10.1080/02699052.2016.1255779
40. Snell DL, Surgenor LJ, Hay-Smith EJC, Siegert RJ. A systematic review of psychological treatments for mild traumatic brain injury: an update on the evidence. *J Clin Exp Neuropsychol*. 2009;31(1):20-38. doi:10.1080/13803390801978849
41. Sullivan KA, Blaine H, Kaye SA, Theadom A, Haden C, Smith SS. A systematic review of psychological interventions for sleep and fatigue after mild traumatic brain injury. *J Neurotrauma*. 2018;35(2):195-209. doi:10.1089/neu.2016.4958
42. Rytter HM, Westenbaek K, Henriksen H, Christiansen P, Humle F. Specialized interdisciplinary rehabilitation reduces persistent post-concussive symptoms: a randomized clinical trial. *Brain Inj*. 2019;33(3):266-281. doi:10.1080/02699052.2018.1552022
43. Thastum MM, Rask CU, Næss-Schmidt ET, et al. Novel interdisciplinary intervention, GAIN, vs. enhanced usual care to reduce high levels of post-concussion symptoms in adolescents and young adults 2-6 months post-injury: a randomised trial. *EclinicalMedicine*. 2019;17:100214. doi:10.1016/j.eclinm.2019.11.007
44. Jensen OK, Nielsen FF, Vosmar L. An open study comparing manual therapy with the use of cold packs in the treatment of post-traumatic headache. *Cephalalgia*. 1990;10(5):241-250. doi:10.1046/j.1468-2982.1990.1005241.x
45. Bell KR, Hoffman JM, Temkin NR, et al. The effect of telephone counselling on reducing post-traumatic symptoms after mild traumatic brain injury: a randomised trial. *J Neurol Neurosurg Psychiatry*. 2008;79(11):1275-1281. doi:10.1136/jnnp.2007.141762
46. Heskestad B, Waterloo K, Baardsen R, Helseth E, Romner B, Ingebrigtsen T. No impact of early intervention on late outcome after minimal, mild and moderate head injury. *Scand J Trauma Resusc Emerg Med*. 2010;18:10. doi:10.1186/1757-7241-18-10
47. Matuskeviciene G, Eriksson G, DeBoussard CN. No effect of an early intervention after mild traumatic brain injury on activity and participation: a randomized controlled trial. *J Rehabil Med*. 2016;48(1):19-26. doi:10.2340/16501977-2025
48. Mittenberg W, Tremont G, Zielinski RE, Fichera S, Rayls KR. Cognitive-behavioral prevention of postconcussion syndrome. *Arch Clin Neuropsychol*. 1996;11(2):139-145. doi:10.1093/arclin/11.2.139
49. Ponsford J, Willmott C, Rothwell A, et al. Impact of early intervention on outcome following mild head injury in adults. *J Neurol Neurosurg Psychiatry*. 2002;73(3):330-332. doi:10.1136/jnnp.73.3.330
50. Suffoletto B, Wagner AK, Arenth PM, et al. Mobile phone text messaging to assess symptoms after mild traumatic brain injury and provide self-care support: a pilot study. *J Head Trauma Rehabil*. 2013;28(4):302-312. doi:10.1097/HTR.0b013e3182847468



51. Varner CE, McLeod S, Nahiddi N, Lougheed RE, Dear TE, Borgundvaag B. Cognitive rest and graduated return to usual activities versus usual care for mild traumatic brain injury: a randomized controlled trial of emergency department discharge instructions. *Acad Emerg Med*. 2017;24(1):75-82. doi:10.1111/acem.13073
52. Schneider KJ, Meeuwisse WH, Nettel-Aguirre A, et al. Cervicovestibular rehabilitation in sport-related concussion: a randomised controlled trial. *Br J Sports Med*. 2014;48(17):1294-1298. doi:10.1136/bjsports-2013-093267
53. Kleffelgaard I, Soberg HL, Tamber AL, et al. The effects of vestibular rehabilitation on dizziness and balance problems in patients after traumatic brain injury: a randomized controlled trial. *Clin Rehabil*. 2019;33(1):74-84. doi:10.1177/0269215518791274
54. Caplain S, Chenuc G, Blanco S, Marque S, Aghakhani N. Efficacy of psychoeducation and cognitive rehabilitation after mild traumatic brain injury for preventing post-concussional syndrome in individuals with high risk of poor prognosis: a randomized clinical trial. *Front Neurol*. 2019;10:929. doi:10.3389/fneur.2019.00929
55. Elgmark Andersson E, Emanuelson I, Björklund R, Stålhammar DA. Mild traumatic brain injuries: the impact of early intervention on late sequelae. a randomized controlled trial. *Acta Neurochir (Wien)*. 2007;149(2):151-159. doi:10.1007/s00701-006-1082-0
56. Kjeldgaard D, Forchhammer HB, Teasdale TW, Jensen RH. Cognitive behavioural treatment for the chronic post-traumatic headache patient: a randomized controlled trial. *J Headache Pain*. 2014;15:81. doi:10.1186/1129-2377-15-81
57. Potter SD, Brown RG, Fleminger S. Randomised, waiting list controlled trial of cognitive-behavioural therapy for persistent postconcussional symptoms after predominantly mild-moderate traumatic brain injury. *J Neurol Neurosurg Psychiatry*. 2016;87(10):1075-1083. doi:10.1136/jnnp-2015-312838
58. Silverberg ND, Hallam BJ, Rose A, et al. Cognitive-behavioral prevention of postconcussion syndrome in at-risk patients: a pilot randomized controlled trial. *J Head Trauma Rehabil*. 2013;28(4):313-322. doi:10.1097/HTR.0b013e3182915cb5
59. Tiersky LA, Anselmi V, Johnston MV, et al. A trial of neuropsychologic rehabilitation in mild-spectrum traumatic brain injury. *Arch Phys Med Rehabil*. 2005;86(8):1565-1574. doi:10.1016/j.apmr.2005.03.013
60. Vikane E, Hellström T, Røe C, Bautz-Holter E, Aßmus J, Skouen JS. Multidisciplinary outpatient treatment in patients with mild traumatic brain injury: a randomised controlled intervention study. *Brain Inj*. 2017;31(4):475-484. doi:10.1080/02699052.2017.1280852
61. Gallaway M, Scheiman M, Mitchell GL. Vision therapy for post-concussion vision disorders. *Optom Vis Sci*. 2017;94(1):68-73. doi:10.1097/OPX.0000000000000935
62. Thiagarajan P, Ciuffreda KJ. Effect of oculomotor rehabilitation on accommodative responsivity in mild traumatic brain injury. *J Rehabil Res Dev*. 2014;51(2):175-191. doi:10.1682/JRRD.2013.01.0027
63. Thiagarajan P, Ciuffreda KJ, Capo-Aponte JE, Ludlam DP, Kapoor N. Oculomotor neurorehabilitation for reading in mild traumatic brain injury (mTBI): an integrative approach. *NeuroRehabilitation*. 2014;34(1):129-146. doi:10.3233/NRE-131025
64. Thiagarajan P, Ciuffreda KJ. Versional eye tracking in mild traumatic brain injury (mTBI): effects of oculomotor training (OMT). *Brain Inj*. 2014;28(7):930-943. doi:10.3109/02699052.2014.888761
65. Graff HJ, Siersma V, Møller A, Egerod I, Rytter HM. Five-year trends in marital stability, academic achievement, and socioeconomic indicators after concussion: a national register study. *J Head Trauma Rehabil*. 2020;35(2):E86-E94. doi:10.1097/HTR.0000000000000501
66. Graff HJ, Siersma V, Møller A, et al. Labour market attachment after mild traumatic brain injury: nationwide cohort study with 5-year register follow-up in Denmark. *BMJ Open*. 2019;9(4):e026104. doi:10.1136/bmjopen-2018-026104
67. de Koning ME, Scheenen ME, van der Horn HJ, et al. Non-hospitalized patients with mild traumatic brain injury: the forgotten minority. *J Neurotrauma*. 2017;34(1):257-261. doi:10.1089/neu.2015.4377
68. McMahon P, Hricik A, Yue JK, et al; TRACK-TBI Investigators. Symptomatology and functional outcome in mild traumatic brain injury: results from the prospective TRACK-TBI study. *J Neurotrauma*. 2014;31(1):26-33. doi:10.1089/neu.2013.2984
69. Quatman-Yates CC, Hunter-Giordano A, Shimamura KK, et al. Physical therapy evaluation and treatment after concussion/mild traumatic brain injury. *J Orthop Sports Phys Ther*. 2020;50(4):CPG1-CPG73. doi:10.2519/jospt.2020.0301
70. McCrory P, Meeuwisse W, Dvořák J, et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med*. 2017;51(11):838-847.

71. Polinder S, Cnossen MC, Real RGL, et al. A multidimensional approach to post-concussion symptoms in mild traumatic brain injury. *Front Neurol*. 2018;9:1113. doi:10.3389/fneur.2018.01113
72. US Preventive Services Task Force. Screening for skin cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2009;150(3):188-193. doi:10.7326/0003-4819-150-3-200902030-00008
73. Petitti DB, Teutsch SM, Barton MB, Sawaya GF, Ockene JK, DeWitt T; US Preventive Services Task Force. Update on the methods of the US Preventive Services Task Force: insufficient evidence. *Ann Intern Med*. 2009;150(3):199-205. doi:10.7326/0003-4819-150-3-200902030-00010
74. Kurowski BG, Hugentobler J, Quatman-Yates C, et al. Aerobic exercise for adolescents with prolonged symptoms after mild traumatic brain injury: an exploratory randomized clinical trial. *J Head Trauma Rehabil*. 2017;32(2):79-89. doi:10.1097/HTR.0000000000000238
75. Leddy JJ, Haider MN, Ellis MJ, et al. Early subthreshold aerobic exercise for sport-related concussion: a randomized clinical trial. *JAMA Pediatr*. 2019;173(4):319-325. doi:10.1001/jamapediatrics.2018.4397

#### SUPPLEMENT.

**eTable 1.** Reproduction of Search Strings by Search Engine for PICO 1-7

**eTable 2.** Clinical Questions in PICO Form, Including Specified Outcome Measures

**eFigure 1.** Flow Charts With Results From the Systematic Review Process for PICO 1-7

**eTable 3.** Evaluation of Systematic Reviews Based on AMSTAR Tool

**eFigure 2.** Meta-Analyses for PICO 1, 2, 3, 4, 6, 7 per Specified Outcome Measures

**eTable 4.** Summary of Findings Table, Including the Rating of the Evidence and Risk of Bias Assessment for PICO 1-7

**eTable 5.** Characteristics of the Included Studies in the Systematic Review