

Aortic dissection diagnosed on stroke computed tomography protocol: a case report

著者名	USUI Takami , SUZUKI Kazufumi , NIINAMI Hiroshi , SAKAI Shuji
journal or publication title	Journal of medical case reports
volume	15
number	1
page range	299
year	2021
URL	http://hdl.handle.net/10470/00033073

doi: 10.1186/s13256-021-02850-1

CASE REPORT

Open Access



Aortic dissection diagnosed on stroke computed tomography protocol: a case report

Takami Usui^{1*}, Kazufumi Suzuki¹ , Hiroshi Niinami² and Shuji Sakai¹

Abstract

Background: Aortic dissection is one of the causes of stroke. Because cerebral infarction with aortic dissection is a contraindication to intravenous recombinant tissue plasminogen activator (rt-PA) therapy, exclusion of aortic dissection is necessary prior to its administration. However, imaging takes time to provide a diagnosis, possibly causing delays in surgical treatment.

Case presentation: A 65-year-old Japanese female patient was transported to the hospital for a suspected stroke, with back pain and left upper and lower extremity palsy which occurred while eating. Upon arrival at the hospital, the left lower limb paralysis had improved, but the left upper limb paralysis remained. Right back pain had also developed. A plain head computed tomography (CT) scan performed 110 minutes after onset showed no acute bleeding or infarction. Subsequent CT perfusion (CTP) showed acute perfusion disturbance in the right hemisphere without infarction, known as ischemic penumbra. The four-dimensional maximum-intensity projection image reconstructed from CTP showed a delayed enhancement at the right internal carotid and right middle cerebral arteries compared to the contralateral side, suggesting a proximal vascular lesion. Contrast helical CT from the neck to abdomen revealed an acute aortic dissection of Stanford type A with false lumen patency. The dissection extended to the proximal right common carotid artery. The patient underwent an emergency total arch replacement and open stent graft. After recovering well, the patient was ambulatory upon discharge from the hospital. The combination of plain head CT, CTP, and helical CT scan from the neck to abdomen enabled us to evaluate for stroke and aortic dissection within a short amount of time, allowing for early therapeutic intervention.

Conclusions: When acute stroke is suspected due to neurological deficits, plain head CT is the first choice for imaging diagnosis. The addition of cervical CT angiography can reliably exclude stroke due to aortic dissection. CTP can identify ischemic penumbra, which cannot be diagnosed by plain head CT or diffusion-weighted magnetic resonance imaging. These combined stroke CT protocols helped us avoid missing an aortic dissection.

Keywords: Computed tomography perfusion, Acute cerebral infarction, Stroke, Acute aortic dissection

Background

Aortic dissection is one of the causes of stroke, with a reported frequency of 0.3–10% [1–7]. Performing an imaging diagnosis may take some time, thus delaying

treatment. Nevertheless, time must be spent on ruling out aortic dissection before administering intravenous recombinant tissue plasminogen activator (rt-PA) therapy. Cerebral infarction with aortic dissection is a contraindication to intravenous rt-PA therapy, and it has been reported to have a lethal course in such patients [2, 8]. Thus, aortic dissection should be ruled out prior to administration. Rapid and reliable diagnostic methods are essential.

*Correspondence: ota.takami@twmu.ac.jp

¹ Department of Diagnostic Imaging and Nuclear Medicine, Tokyo Women's Medical University, 8-1, Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Case presentation

A 65-year-old Japanese female with a history of hyperlipidemia experienced palpitations and uncertain chest discomfort while eating, before developing left upper and lower limb paralysis. During emergency transport, her blood pressure dropped to 62 mmHg, and right joint paralysis developed. She was transported to our hospital 35 minutes after onset with an additional presentation of right back pain. Neurological severity was rated as National Institutes of Health Stroke Scale (NIHSS) 2 (left upper limb paralysis and right conjugate deviation). There was no heart murmur. Left–right difference in blood pressure at the emergency department was less than 20 mmHg (right upper limb 77/54 mmHg, left upper limb 87/45 mmHg, right lower limb 102/92 mmHg, left lower limb 94/51 mmHg). No elevation in white blood cells or C-reactive protein was noted, and there were no findings suggestive of acidosis. D-dimer was mildly elevated at 6.3 $\mu\text{g/mL}$. Serum creatinine and estimated glomerular filtration rate (eGFR) were 1.11 mg/dL and 38.6 mL/minute/1.73 m^2 , respectively, and renal function was not impaired to the extent that contrast media could not be used. An echocardiogram was not performed. Rehydration was started, and there was no hemodynamic deterioration. A chest radiograph taken 110 minutes after onset showed no enlargement of the aorta or mediastinum (Fig. 1). A plain head CT scan did not suggest acute infarction or bleeding (Fig. 2). However, on CT perfusion (CTP), the right middle cerebral artery region showed an elongation of delay (DLY) and increases in time to peak (TTP) and mean transit time (MTT). Cerebral blood




Fig. 1 The patient's chest radiograph shows no enlargement of the mediastinum. There is no calcification of the aortic wall or pleural effusion



Fig. 2 A plain head computed tomography scan does not reveal any sign of acute infarction or bleeding

flow (CBF) was decreased, but cerebral blood volume (CBV) was preserved (Fig. 3). Due to the absence of early CT signs in the plain head CT, we determined that the patient had ischemic penumbra. On the CT angiography (CTA) reconstructed from the arterial phase of the CTP scan, the right middle cerebral artery was poorly depicted (Fig. 4). However, the four-dimensional maximum-intensity projection (4D MIP) image reconstructed from CTP showed a delayed enhancement of both the right internal carotid artery and the right middle cerebral artery. In the late phase, the distal portion of the right middle cerebral artery was well depicted, and no stenosis or obstruction was observed at the intracranial artery, suggesting a proximal vascular lesion (Fig. 5). A contrasted helical CT scan from the neck to abdomen was performed using contrast medium residue soon after CTP imaging. This revealed an acute aortic dissection of Stanford type A with false lumen patency, from the base of the aorta to the level of the renal artery bifurcation. The dissection extended to the proximal right common carotid artery, and the true lumen of the right internal carotid artery was narrowed. The left renal artery was supplied by the false lumen (Fig. 6). The brain tissue status on CTP was reversible ischemia, not irreversible infarction, and the patient was deemed to be eligible for emergency surgery, which included total arch replacement and open stent graft insertion approximately 3.5 hours after the onset of illness. The left upper limb paralysis had recovered to a manual muscle test (MMT) grade of about 4–5 after the surgery. The patient was ambulatory upon discharge from the hospital 23 days after the onset.




Fig. 3 On computed tomography perfusion scan, delay (DLY), time to peak (TTP), and mean transit time (MTT) are elongated. Cerebral blood flow is decreased, but cerebral blood volume (CBV) is preserved in the region of the right middle cerebral artery. The lesions with elongated DLY, TTP, and MTT may indicate major vessel occlusion. An area with decreased CBV is diagnosed as an infarction, which is irreversible ischemic tissue damage. Because CBV remained almost normal, we determined that the patient had a large ischemic penumbra and no infarction yet in this case




Fig. 4 Computed tomography angiography reconstructed from computed tomography perfusion scan shows the right middle cerebral artery poorly depicted. The right internal carotid artery appears to be intact

Discussion and conclusions

Intravenous rt-PA therapy is contraindicated in cases of cerebral infarction with aortic dissection. When rt-PA is considered in acute ischemic stroke, aortic dissection should be excluded. Amr *et al.* noted that brain hypoperfusion alone should not be a contraindication for urgent surgical treatment, regardless of initial clinical neurological severity [8]. Ultrasonography such as point-of-care ultrasound is usually performed to arrive at an early diagnosis [9], but unfortunately it was not performed in this case. Although ultrasonography can be used for an early diagnosis, it is first necessary to understand the exact

type and condition of aortic dissection before surgery in order to determine whether emergency surgery should be performed. In this case, we were misled by the importance of neurological findings despite the presence of symptoms suggestive of acute aortic dissection. However, by using the stroke CT protocol, we arrived at the correct diagnosis of acute aortic dissection with a single examination modality. We believe that contrast CT is useful in all cases when stroke is suspected.

When acute stroke is suspected due to neurological deficits, plain head CT is the first choice of imaging for diagnosis [10]. CTP can evaluate ischemic penumbra, which cannot be diagnosed using plain head CT or diffusion-weighted magnetic resonance imaging (DW-MRI). Cerebral circulatory disturbance in acute aortic dissection may cause impaired consciousness and paralysis, and surgery is indicated when the disease complications can be alleviated or its progression can be controlled by surgery [11, 12]. In this case, CTP revealed that the condition of the brain tissue was reversible ischemia, not irreversible infarction.

Herein, we present our stroke CT protocols as they are currently employed. First, plain head CT is used to determine the presence of cerebral hemorrhage and evaluate early CT signs. Next, a CTP is taken using 50 mL of contrast agent to check for ischemic penumbra. The addition of cervical CTA using 40 mL of contrast agent can reliably exclude stroke due to aortic dissection. An aorta study may be taken instead with helical CT from the neck to the abdomen or mid-thigh level to prepare for catheterization. This usually takes only 10–15 minutes, and in this way, maximum imaging can be achieved in a short amount of time. In this case, the time from scouting to contrast CT imaging was 24 minutes. The delay may have been because the case came in at night and was received by a technician unfamiliar with our stroke CT protocols.




Fig. 5 Four-dimensional maximum-intensity projection image reconstructed from computed tomography perfusion scan. **a, b** The right internal carotid artery and the right middle cerebral artery indicate delayed enhancement compared to the contralateral side. **c** In the late phase, the distal middle cerebral artery also exhibits delayed enhancement, and no stenosis or obstruction is observed, suggesting a proximal vascular lesion. **d** No venous abnormality was seen in venous phase

Nevertheless, although the examination took longer than usual, the patient was still diagnosed within 90 minutes after the onset of symptoms and was able to undergo emergency surgery in about 3.5 hours.

This case may seem like a normal course of action to some experts, but it highlights some important issues. Stroke guidelines recommend the use of CTP between 6 and 24 hours after onset to determine indications for revascularization in ischemic stroke [10]. However, aortic dissection complications and ischemic penumbra can be assessed simultaneously using contrast CT as indicated by the appropriate stroke protocols, as in the present case. This approach can be of benefit in patients up to 6 hours from the onset of stroke.

However, some problems still need to be addressed before CTP can become the standard. There is a risk of developing an allergy from the use of iodine contrast agents in some patients. In addition, an optimizing protocol is required, such as the use of iterative reconstruction methods and the optimization of intermittent scans to reduce radiation exposure with a multiphase scan protocol.

Aortic dissection is associated with a high mortality rate and poor prognosis, but if it is diagnosed and treated quickly, a positive turnaround may be possible [8, 13]. The stroke CT protocol may have advantages in diagnosing stroke and aortic dissection due to its short examination time [14–17].




Fig. 6 Contrast computed tomography from the neck to abdomen was taken using a contrast medium. **a** A cross section at the level of the thoracic aorta revealed an acute aortic dissection of Stanford type A with false lumen patency. **b** A cross section of the aorta to the level of the renal artery bifurcation. The left renal artery is perfused by the false lumen. **c** Coronal reconstruction section of the neck. The dissection extends to the proximal right common carotid artery, with narrowing of the lumen of the right internal carotid artery

Abbreviations

CBF: Cerebral blood flow; CBV: Cerebral blood volume; CTA: Computed tomography angiography; CTP: Computed tomography perfusion; DLY: Delay; NIHSS: National Institutes of Health Stroke Scale; MMT: Manual muscle test; MTT: Mean transit time; TTP: Time to peak; 4D MIP: Four-dimensional maximum-intensity projection.

Acknowledgements

None.

Authors' contributions

TU and KS prepared the manuscript. SS and HN obtained written informed consent from the patient. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article.

Declarations

Ethics approval and consent to participate

This case report was reviewed and approved by the institutional review board.

Consent for publication

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Diagnostic Imaging and Nuclear Medicine, Tokyo Women's Medical University, 8-1, Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan.

²Department of Cardiovascular Surgery, Tokyo Women's Medical University, 8-1, Kawada-cho, Shinjuku-ku, Tokyo 162-8666, Japan.

Received: 23 July 2020 Accepted: 4 April 2021

Published online: 26 May 2021

References

- Hagan PG, Nienaber CA, Isselbacher EM, *et al.* The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA*. 2000;283(7):897–903.
- Fessler AJ, Alberts MJ. Stroke treatment with tissue plasminogen activator in the setting of aortic dissection. *Neurology*. 2000;54(4):1010.
- Iguchi Y, Kimura K, Sakai K, *et al.* Hyper-acute stroke patients associated with aortic dissection. *Intern Med*. 2010;49(6):543–7.
- Bossone E, Corteville DC, Harris KM, *et al.* Stroke and outcomes in patients with acute type A aortic dissection. *Circulation*. 2013;128(11 Suppl 1):S175–179.
- Guglielmi V, Groeneveld NS, Posthuma L, *et al.* Aortic dissection masquerading as a code stroke: a single-centre cohort study. *Eur Stroke J*. 2020;5(1):56–62.
- Sakamoto Y, Koga M, Ohara T, *et al.* Frequency and detection of Stanford type A aortic dissection in hyperacute stroke management. *Cerebrovasc Dis*. 2016;42(1–2):110–6.
- Ohara T, Koga M, Tokuda N, *et al.* Rapid identification of type A aortic dissection as a cause of acute ischemic stroke. *J Stroke Cerebrovasc Dis*. 2016;25(8):1901–6.
- Amr G, Boulouis G, Bricout N, *et al.* Stroke presentation of acute type A Aortic dissection with 100% perfusion-weighted imaging-diffusion-weighted imaging mismatch: a call for urgent action. *J Stroke Cerebrovasc Dis*. 2016;25(5):1280–3.

9. Nishigami K. Update on cardiovascular echo in aortic aneurysm and dissection. *Ann Vasc Dis.* 2018;11(4):437–42.
10. Powers WJ, Rabinstein AA, Ackerson T, *et al.* Guidelines for the Early Management of Patients with Acute Ischemic Stroke: 2019 update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke: A Guideline for Healthcare Professionals from the American Heart Association/American Stroke Association. *Stroke.* 2019;50(12):e344–418.
11. Estrera AL, Garami Z, *et al.* Acute type A aortic dissection complicated by stroke: can immediate repair be performed safely? *J Thorac Cardiovasc Surg.* 2006;132(6):1404–8.
12. Fann JI, Sarris GE, Mitchell RS, *et al.* Treatment of patients with aortic dissection presenting with peripheral vascular complications. *Ann Surg.* 1990;212(6):705–13.
13. Alonso JV, Martin D, Kinderman H, Farhad I, Swallow P, Siggers A. Acute ischemic stroke what is hidden behind? *J Cardiol Cases.* 2017;16(5):174–7.
14. Mayer TE, Hamann GF, Baranczyk J, *et al.* Dynamic CT perfusion imaging of acute stroke. *AJNR Am J Neuroradiol.* 2000;21(8):1441–9.
15. Suzuki K, Morita S, Masukawa A, Machida H, Ueno E. Utility of CT perfusion with 64-row multi-detector CT for acute ischemic brain stroke. *Emerg Radiol.* 2011;18(2):95–101.
16. JCS Joint Working Group. Guidelines for diagnosis and treatment of aortic aneurysm and aortic dissection (JCS 2011): digest version. *Circ J.* 2013;77(3):789–828.
17. Sotoudeh H, Bag AK, Brooks MD. “Code-Stroke” CT perfusion; challenges and pitfalls. *Acad Radiol.* 2019;26(11):1565–79.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

