

University of Groningen

Modulation of yaw by the caudal fin in the yellow boxfish (*Ostracion cubicus*)

Boute, Pim G.; Van Wassenbergh, Sam; Stamhuis, Eize J.

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Boute, P. G., Van Wassenbergh, S., & Stamhuis, E. J. (2021). *Modulation of yaw by the caudal fin in the yellow boxfish (*Ostracion cubicus*)*. 115. Abstract from SEB 2021 Annual Conference.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ABSTRACT BOOK

SEB VIRTUAL CONFERENCE 2021

29 JUNE - 8 JULY 2021

SEBIOLOGY.ORG

#SEBCONFERENCE




SEB 2021 ANNUAL CONFERENCE

CONTENTS

1. SCIENCE ACROSS BOUNDARIES ABSTRACTS (ANIMAL, CELL AND PLANT BIOLOGY)	4
2. ANIMAL BIOLOGY ABSTRACTS	56
3. CELL BIOLOGY ABSTRACTS	194
4. PLANT BIOLOGY ABSTRACTS	216
5. AUTHOR INDEX	268

ANIMAL BIOLOGY ABSTRACTS





James Kempton (University of Oxford, United Kingdom), Caroline Brighton (University of Oxford, United Kingdom), Sofia Minano Gonzalez (University of Oxford, United Kingdom), Marco Klein Heerenbrink (University of Oxford, United Kingdom), Lydia Paris (University of Oxford, United Kingdom), Graham Taylor (University of Oxford, United Kingdom)

@ james.kempton@zoo.ox.ac.uk

Guidance and control of aerial attack behaviour is expected to be intimately related to its ecological setting. Here we analyse a very large dataset of state-of-the-art motion capture data, confirming that Harris' Hawk guidance is best modelled by a mixed guidance law, proportional navigation pursuit (PNP), in contrast to the proportional navigation (PN) guidance best modelling Peregrine Falcons. Furthermore, with PN fitted to both species, it is parameterised very differently between them, likely reflecting disparate ecologies: hawks operating in cluttered environments, hunting agile prey against a close background; falcons operating in open environments against distant backgrounds and potentially less agile prey. We extend our understanding by showing that PNP pursuit in hawks is just as effective when sensory input is modelled in a local background reference frame (LRF) as when in an inertial frame (IRF). This is a priori surprising as effective PN guidance is impossible using an LRF, and engineered missile systems absolutely rely on IRF measurements. The possibility only arises in hawks because of their use of mixed PNP guidance. This is interesting, because the motion vision system naturally makes measurements in the LRF. This result implies input from a single sensory modality – vision – could successfully implement PNP, with no necessary requirement for fusion with inertial measurements from the vestibular system. This may explain why hawks use PNP, while falcons use PN given target motion measured against a distant background approximates being measured in an IRF.

A8.35 HIERARCHY OF GRAVITY-RECEPTIVE ORGANS IN CRICKETS

Tuesday 6 July 2021 12:40

Christoph Bruns (Dept. of Biomimetics Hochschule Bremen - City University of Applied Sciences, Germany), Susanna Labisch (Biomimetics-Innovation-Centre Hochschule Bremen - City University of Applied Sciences Bremen, Germany), Jan-Henning Dirks (Biomimetics-Innovation-Centre Hochschule Bremen - City University of Applied Sciences Bremen, Germany)

@ jan-henning.dirks@hs-bremen.de

For almost all organisms a reliable detection of the gravitational vector is an important fitness criterion. This detection is based on relatively simple physical principles, however physiologically "implemented" a variety of different sensor types. Arthropods and in particular insects show a great variety of different exoskeletal sensors sensitive to the gravitational vector. Previous studies show that in many cases the gravity reception of insects most likely relies on some sort of sensor fusion, resulting in an astonishing overall accuracy and precision. Within these complex sensor systems, it however seems likely that some gravity receptive organs might play a more important role in determining the overall gravitational vector than other sensors. To answer this question, we performed behavioural experiments on house crickets (*Acheta domestica*) with selectively deactivated gravity perceiving structures. Animals with different treatments were

buried inside a transparent box filled with granular material. The box was then randomly rotated to disorient the insect. We then recorded the directions in which the crickets tried to escape the box and the respective time. Our results show that ablation of the cerci resulted in a decrease in sensing performance of approximately 39.59%. Additional ablation of the antennae further decreased the overall sensing performance. A possible participation of the leg proprioceptors in gravity perception could not be shown experimentally. In general, the failure of individual sensory structures could be fully compensated, whereas the failure of the club-shaped sensilla on the cerci could only be compensated with an overall loss in sensing performance.

A8.36 MODULATION OF YAW BY THE CAUDAL FIN IN THE YELLOW BOXFISH (*OSTRACION CUBICUS*)

Tuesday 6 July 2021 14:00

Pim G. Boute (Experimental Zoology Group - Wageningen University and Research, Netherlands), Sam Van Wassenbergh (Department of Biology - University of Antwerp, Belgium), Eize J. Stamhuis (Department of Ocean Ecosystems - University of Groningen, Netherlands)

@ pim.boute@wur.nl

Boxfishes (Ostraciidae; Tetraodontiformes) have a rigid carapace which restricts body undulation. Swimming movements can only be generated by the fins which protrude from the carapace. Nevertheless, these fishes are highly manoeuvrable and manage to swim with remarkably dynamic stability. However, the rigid carapace of boxfishes shows an inherently unstable response in yaw caused by course-disturbing flows. Hence, any net stabilising effect should come from the fishes' fins. Here, we aim to determine the effect of the surface area and orientation of the caudal fin on the yaw torque exerted on the square cross-sectional shaped yellow boxfish (*Ostracion cubicus*). Yaw torques were quantified in a flow tank using a 3D printed physical yellow boxfish model with an attachable closed or open caudal fin. The model was positioned at different body and tail angles and exposed to different water flow speeds. We show that the caudal fin is crucial for yaw control. These flow tank results were confirmed by computational fluid dynamics simulations. The caudal fin acts as both a course-stabiliser and rudder for the naturally unstable rigid carapace with regard to yaw. By using physical models and computer simulations, we quantitatively show that actively changing the shape and orientation of the caudal fin plays an important role in controlling yaw torque in yellow boxfish. Further study is needed to unravel how all components of the boxfishes' locomotor apparatus function together, from a dynamic perspective, during lateral gust flows and turning.

A8.37 SNAKES AND SNAKE ROBOTS TRAVERSING LARGE, SMOOTH OBSTACLES

Tuesday 6 July 2021 14:35

Qiyuan Fu (Johns Hopkins University, United States), Sean W Gart (Johns Hopkins University, United States), Chen Li (Johns Hopkins University, United States)

@ fqiyuan1@jhu.edu