

| | | |
|--|---|---|
| | <p style="text-align: center;">CONTEMPORARY AGRICULTURE SAVREMENA POLJOPRIVREDA</p> <p style="text-align: center;">The Serbian Journal of Agricultural Sciences <i>Srpski časopis za poljoprivredne nauke</i></p> <p style="text-align: center;">Vol. 64, No. 3 - 4, Pp. 235 - 240, 2015.</p> <p style="text-align: center;">www.contagri.info</p> <p style="text-align: center;">ISSN: 0350-1205 UDC: 63(497.1)(051)-"540.2"</p> | <p style="text-align: center;">UNIVERSITAS STUDIORUM NEOPLANTENSIS</p> <p style="text-align: center;">University of Novi Sad, Serbia</p> <p style="text-align: center;">Published by Faculty of Agriculture, Novi Sad</p> |
|--|---|---|

Original scientific paper

UDC: 581.142:641.12

EFFECTS OF COMPOST PRODUCTS ON SEED GERMINATION OF VEGETABLES*

Mira MILINKOVIĆ, Blažo LALEVIĆ, Snežana Oljača, Vlado LIČINA,
Jelena JOVIČIĆ PETROVIĆ, Vera RAIČEVIĆ¹*

Summary: The aim of this work is determination of influence of different compost leachates and teas types on vegetables seed germination. Composts used for leachate and tea production were produced of municipal waste (MSW) and waste from tobacco industry (TW). Results achieved with MSW products were comparable to control. Compost products derived from TW showed significant phytotoxicity, which can be correlated with their chemical composition. Leachates from MSW compost lead to the lower germination index in comparison to MSW compost teas, which indicates the possibilities their application.

Key words: Compost products, germination index, vegetables seed germination, waste

INTRODUCTION

Growth media, mainly comprised of organic matter, fertilizer and soil, is often used for horticultural plants (Boldrin et al., 2010) and vegetables (Arenas et al., 2002). One of the most important highly valuable, cost-effective and environmental friendly organic components for vegetables growth medium is peat (Herrera et al., 2009; Ingelmo et al., 1998). However, the excessive use of peat leads to the release of CO₂ counted as a greenhouse gas (Boldrin et al., 2010). Several studies have addressed the use of compost from different waste materials as an alternative to peat for growth of vegetables (Sterrett, 2001). The application of compost improves the characteristics of growth substrates (Keserović et al., 2012) and increases organic matter and nutrients content for plant growth (Darby et al., 2006). Problems of waste disposal and low content of organic matter in soils can be solved by application of composted municipal solid waste (MSW), which is suitable for agriculture due to its positive effects on soil properties (Wolkowski, 2003). Similar effects, due to high nutrients content (Chaturvedi et al., 2008) and low content of toxic elements (Okur et al., 2008), shows tobacco waste (TW) and its composting is useful for agricultural purpose (Saithep et al., 2009).

¹PhD Mira Milinković, research associate, Fruit research institute Čačak; dr Blažo Lalević, assistant professor, dr Snežana Oljača, full professor, dr Vlado Ličina, full professor, dr Jelena Jovičić Petrović, assistant professor, dr Vera Raičević, full professor, University of Belgrade, Faculty of agriculture

*Corresponding author: Blažo Lalević, e-mail: blazol@agrif.bg.ac.rs; phone: +381 11 2615315/162

**This research was partially supported by the Ministry of education and science of Republic of Serbia (grant number TR 31080) and EU Commission project AREA, contract No. 316004.

As a new approach, compost leachates and teas have a vital role in providing of nutrients for plant growth (Mesbahzadeh and Astarai, 2008). For the phytotoxicity evaluation of these products, one of the suggested methods is germination index test, based on seed germination and initial growth of plants (Sanchez-Monedero et al., 2002). Because initial phases of plant growth are fundamental, the aim of this paper was to determinate the influence of different compost leachates and teas on vegetables seed germination.

MATERIAL AND METHODS

For preparation of compost products, two composts were used: (i) MSW compost from Čačak municipality (Serbia) and (ii) TW compost, obtained from tobacco waste of „Philip Morris International“ company (Niš, Serbia). In order to obtain compost, waste material was periodically turned and aerated. These 1-year-old composted materials were passed (< 7mm) and 5 hours after addition of water until the complete saturation, extraction of leachates was performed. The compost teas were prepared from the same compost materials and water (1:3, w/w). Extraction period lasted for 10 days with continuous aeration by aquarium pumps.

The electrical conductivity (EC) of compost products was measured by Conductivity Meter LF 538 (Windaus Labortechnik WTW, Germany), and pH by pH-meter CyberScan pH 510, Eutech Instruments (Singapore). Humic and fulvic acids were determined according to Jakovljevic et al. (1995). The total carbon content (C_{tot}) was determined by the dichromate oxidation method. The excess of the dichromate was determined according to Yeomans and Bremner (1989). The total nitrogen (N_{tot}) was estimated by Kjeldahl method (Bremner and Mulvaney, 1982). For determination of metal content, 1 g of air dried samples were 4-hours-shaked with 20 mL 0.5% NaEDTA extractants and after filtration samples were subjected to microwave acid (5 mL HNO_3 +2 mL H_2O_2) digestion (Speedwave MWS3, Berghof, Germany) using 5 mL HNO_3 conc. (70%)+2 mL H_2O_2 (33%) and measured by ICP-OES (Spectro Genesis FEE, Germany).

The tested seeds of plants (green beans-*Phaseolus vulgaris* L., cabbage-*Brassica oleracea* L., pepper-*Capsicum annum* L. and lettuce-*Lactuca sativa* L.) originated from Institute for vegetable crops ltd. (Smederevska Palanka, Serbia), while tomato seeds (*Solanum lycopersicum* L.) were obtained from “Superior seeds” (Velika Plana, Serbia). Ten seeds of green beans, 20 seeds of tomato and pepper, 30 seeds of cabbage and 50 seeds of lettuce) were sterilized with H_2O_2 for 30 seconds, washed with distilled water and placed on filter paper (previously impregnated with 1 mL of compost products, separately) in each Petri dish. In control variant filter paper was impregnated with distilled water. All experiments were performed in triplicate, in darkness at the temperature of 25°C for 7 days. The germination rate was determined at the end of experiment.

The potential phytotoxicity of compost products was determined using the seed germination test. For this test, compost products were diluted with distilled water to yield 0, 25, 50, 75, and 100% extracts. According to ISO 11269-2 (1995) directions, ten seeds of lettuce and barley (*Hordeum vulgare* L.) were placed in Petri dishes on filter paper previously treated with 5 mL of adequate treatment extract, while in control with 5 mL of distilled water, with three replicate per treatment. Seeds were germinated at 25°C in darkness, and after 5 days the number of germinated seeds was counted and root length was measured in each Petri dish.

The germination index (GI) was determined according to the following formula: $(G \times L \times 100\%) / (G_c \times L_c)$ where G and L are respectively the percentage of germination and mean root length of treatments, while G_c and L_c are respectively the mean germination rate and root length of control plates.

The data was subjected to ANOVA using MSTAT-C statistical computer package (Michigan State University, East Lansing, MI, USA). The LSD test ($p = 0.05$) was used for treatment means comparison.

RESULTS

Chemical analysis showed high pH values of compost products (Table 1) and low EC values. The values of chemical parameters of TW products were higher than in MSW products.

Table 1. Chemical properties of compost leachates and teas

| Parameter | leachates | | teas | |
|--|-----------|--------|--------|--------|
| | MSW | TW | MSW | TW |
| pH | 7.10 | 7.70 | 7.75 | 7.20 |
| Dry matter (%) | 0.401 | 1.632 | 0.207 | 0.263 |
| ash (%) | 0.374 | 0.893 | 0.186 | 1.100 |
| organic matter (%) | 0.027 | 0.740 | 0.021 | 1.531 |
| N _{tot} (mg/L) | 22.4 | 426.0 | 10.5 | 704.5 |
| C _{tot} (mg/L) | 81.5 | 5217.0 | 37.0 | 7825.0 |
| C _{org} in humic acids/C _{org} in fulvic acids | 1.2:1 | 1.0:1 | 1.8:1 | 0.9:1 |
| C/N | 3.7:1 | 12.2:1 | 3.5:1 | 11.1:1 |
| EC (mS/cm) | 5.19 | 9.78 | 2.61 | 13.81 |
| salts (mg/L) | 3321.5 | 6259.0 | 1668.5 | 8838.0 |
| K (mg/L) | 705 | 2812 | 356.5 | 3750 |
| Na (mg/L) | 182 | 9 | 85 | 14 |
| Ca (mg/L) | 482.5 | 255 | 352 | 972 |
| Mg (mg/L) | 95.5 | 160 | 104 | 436 |
| P (mg/L) | 0.4 | 3.4 | 0.4 | 15.6 |
| S (mg/L) | 334 | 201 | 150.5 | 222.0 |
| Fe (mg/L) | 1.9 | 11.7 | 5.2 | 17 |
| Mn (mg/L) | 0.47 | 1.01 | 0.25 | 2.50 |
| Cu (mg/L) | 0.11 | 1.36 | 0.11 | 2.25 |
| Zn (mg/L) | 0.11 | 3.75 | 0.09 | 4.75 |
| B (mg/L) | 0.98 | 2.21 | 0.84 | 3.51 |
| Co (mg/L) | 0.005 | 0.099 | 0.005 | 0.127 |
| Mo (mg/L) | 0.06 | 0.03 | 0.02 | 0.03 |
| Cd (mg/L) | 0.002 | 0.043 | 0.004 | 0.057 |
| Cr (mg/L) | 0.01 | 0.13 | 0.04 | 0.23 |
| Ni (mg/L) | 0.05 | 0.20 | 0.07 | 0.25 |
| Pb (mg/L) | 0.04 | 0.24 | 0.04 | 0.29 |

The germination rate of plant seeds was affected by compost product types and its origin (Table 2). Statistically significant influence of vegetables seeds on germination rate was noticed.

Table 2. Influence of plants and compost products on germination rate (%)

| Factors | Parameters | Control | MSW | TW |
|------------------|----------------------------------|--------------|--------------|--------------|
| A Plant seeds | Green beans | 93.3±2.108ab | 90.0±5.164ab | 88.3±4.014a |
| | Cabbage | 91.1±1.391ab | 72.2±4.439b | 0.0±0.00c |
| | Tomato | 90.0±0.00b | 80.8±8.795ab | 40.8±16.094b |
| | Pepper | 55.0±3.651c | 30.0±8.062c | 0.8±0.833c |
| | Lettuce | 96.7±0.843a | 97.7±0.954a | 0.00±0.00c |
| B Products | Leachates | 85.2±4.258a | 81.4±5.389a | 20.0±9.844b |
| | Teas | 85.2±4.259a | 66.9±8.399b | 32.0±10.553a |
| AXB | Plant seeds x leachates | 93.3±3.33a | 80.0±5.773b | 93.3±3.33a |
| | | 91.1±2.20a | 81.1±2.967b | 0.0±0.00c |
| | | 90.0±0.00a | 100.0±0.00a | 5.0±2.887c |
| | | 55.0±5.773b | 46.7±6.009d | 1.7±1.67c |
| | | 96.7±1.33a | 99.3±0.667a | 0.0±0.00c |
| | Plant seeds x compost teas | 93.3±3.33a | 100.0±0.00a | 83.3±6.67b |
| | | 91.1±2.20a | 63.4±3.33c | 0.00±0.0c |
| | | 90.0±0.0a | 61.7±4.409c | 76.7±1.66b |
| | | 55.0±5.773b | 13.3±3.33e | 0.0±0.00c |
| | | 96.7±1.33a | 96.0±1.154a | 0.0±0.0c |
| A | | * | * | * |
| B | | NS | * | * |
| AXB | | NS | * | * |

Values within each column labeled with the same letter are insignificantly different at the $p \leq 0.05$ by LSD test

In general, the compost products showed lower seeds' germination rate compared to the control (Table 2). The germination rate using MSW leachate was highest at lettuce and lowest at pepper, while both TW products led to the lowest germination rate at cabbage, pepper and lettuce seeds. However, these TW products induced the highest germination rate at green beans and tomato seeds. MSW leachate has significant influence on pepper, cabbage and lettuce seed germination compared to MSW compost tea, especially at tomato seed.

As can be seen from table 3, in most of treatments using the TW products, germination index was 0. On the other hand, only 100% and 75% dilution of MSW products had GI values close to 50.

Table 3. Germination index of lettuce and barley seeds treated with leachates and teas

| Plants | Compost products' dilution | Leachates | | Teas | |
|---------|----------------------------|-----------------------|-----|-------|------|
| | | Germination index (%) | | | |
| | | MSW | TW | MSW | TW |
| Lettuce | 100 | 25.4 | 0 | 52.8 | 0 |
| | 75 | 30.7 | 0 | 58.3 | 0 |
| | 50 | 52.1 | 0 | 68.7 | 0 |
| | 25 | 73.4 | 0 | 132.1 | 0 |
| Barley | 100 | 44,0 | 0 | 95.3 | 0 |
| | 75 | 58,0 | 0 | 83.4 | 0 |
| | 50 | 86,0 | 0 | 93.2 | 0 |
| | 25 | 98,0 | 2,8 | 92.0 | 37.7 |

DISCUSSION

Different types of waste are disposed off in landfills (Chatterjee et al., 2013). As showed previously, more than 50% of wastes are potentially biodegradable (Engineering Solutions and Design, Inc., 2004). By composting as a promised activity, waste is turned into a soil conditioner and can be used in agricultural production (Hart and Plumiers, 1996).

Lower EC values comparing to previous research (Mokhtarani et al., 2012) were obtained. Also, the concentrations of K, Mg, Zn and Cu in TW products resulted in higher EC, which has been also confirmed (Pant et al., 2012). Heavy metal content of the products was lower than proposed values (Jones-Lee and Lee, 1993), thus avoiding possible metals precipitation at high pH values (Esakku et al., 2003).

Germination rate of examined seeds in control treatment was higher compared to compost products treatments. These results are similar with the findings of Ingelmo et al. (1998) who studied different organic amendments added to soil.

The lowest germination rate at cabbage, pepper and lettuce seeds on tobacco products can be correlated with salinity of these products, which inhibit the seed germination (Marchiol et al., 1999). The similar influence of TW on tomato seeds was obtained in previous research (Okur et al., 2008). On the other side, results of some other authors confirm the negative influence of TW treatment on some properties of tomato (Chaturvedi et al., 2008).

Germination index (GI) of selected plants (lettuce and barley) has been determined because of their high sensitivity to salt concentrations and phytotoxic substances (Dolgen et al., 2004). According to criteria proposed by Zucconi et al. (1981), used leachates showed rather high phytotoxicity (Table 3). The tea production where aeration system used had less phytotoxicity compared with nonaerated production processes, which is confirmed in previous research (Carballo et al., 2009).

CONCLUSION

Our results suggest that influence of MSW compost tea and leachate on germination of vegetables seeds' germination that their phytotoxicity is excluded as a barrier to their application. Probably, the increased salinity appear to be major factor of limiting use of TW compost products. Further investigation will be directed to the possible application of these products as soil amendments with the aim of improvement the agricultural production.

REFERENCES

- ARENAS M, VAVRINA CS, CORNELL JA, HANLON EA, HOCHMUTH GJ: Coir as an alternative to peat in media for tomato transplant production. *Hort Science*, 37(2)309-312, 2002.
- BOLDRIN A, HARTLING KR, LAUGEN M, CHRISTENSEN TH: Environmental inventory modelling of the use of compost and peat in growth media preparation. *Resour. Conserv. Recy.*, 54(12)1250-1260, 2010.
- BREMNER JM, MULVANEY CS: Nitrogen-area. Methods of soil analysis. Pt- 2. Chemical and microbiological properties. In: *Agronomy Monograph 9* (A.L. Page, R.H. Miller and D.R. Keeney, eds.). Am. Soc. Agron., Madison, WI; pp. 699-709, 1982.
- CARBALLO T, GIL MV, CALVO LF, MORAN A: The influence of aeration system, temperature and compost origin on the phytotoxicity of compost tea. *Compost Sci. Util.*, 17(2)127-139, 2009.
- CHATTERJEE N, FLURY M, HINMAN C, COGGER CG: Chemical and physical characteristics of compost leachates - a review. Report prepared for the Washington State Department of Transportation, pp. 1-54, 2013.
- CHATURVEDI S, UPRETI DK, TANDON DK, SHARMA A, DIXIT A: Bio-waste from tobacco industry as tailored organic fertilizer for improving yields and nutritional values of tomato crop. *J. Environ. Biol.*, 29(5)759-763, 2008.
- DARBY HM, STONE AG, DICK RP: Compost and manure mediated impacts on soilborne pathogens and soil quality. *Soil Sci. Soc. Am. J.*, 70(2)347-358, 2006.
- DOLGEN D, ALPASLAN MN, DENLE N: Use of an agro-industry treatment plant sludge on iceberg lettuce growth. *Ecol.Eng.*, 23(2)117-125, 2004.
- ENGINEERING SOLUTIONS AND DESIGN, INC.: State of Ohio waste characterization study. Technical report, Ohio Department of Natural Resources, Columbus, OH, 2004.
- ESAKKU S, PALANIVELU K, JOSEPH K: Assessment of heavy metals in a municipal solid waste dumpsite. Workshop on Sustainable Landfill Management, Chennai, India, 139-145, 2003.
- HART D, PLUIMERS J: Composting of organic household waste. UWEP working document. Niuwehaven 201, pp. 1-91, 1996.
- HERRERA F, CASTILLO JE, LOPEZ-BELLIDO RJ, BELLIDO LL: Replacement of a peat-lite medium with municipal solid waste compost for growing melon (*Cucumis melo* L.) transplant seedlings. *Compost Sci. Util.*, 17(1)31-39, 2009.
- INGELMO F, CANET R, IBAFIEZ MA, POMARES F, GARCIA J: Use of MSW compost, dried sewage sludge and other wastes as partial substitute for peat and soil. *Bioresource Technol.*, 63(2)123-129, 1998.
- ISO 11269-2: Soil Quality – Determination of the effects of pollutants on soil flora – Effects of chemicals on the emergence and growth of higher plants, 1995.
- JAKOVLJEVIĆ M, PANTOVIĆ M, BLAGOJEVIĆ S: Handbook of soil and water chemistry. University of Belgrade, Faculty of Agriculture, pp. 1-117, 1995.
- JONES-LEE A, LEE GF: Ground water pollution by municipal landfills: leachate composition, detection and water quality significance. Proceedings of Sardinia'93, IV International landfill symposium, Sardinia, Italy, 11-15, October, 1993, pp. 1093-1103.
- KESEROVIĆ Z, INJAC M, MAGAZIN N, MILIĆ B, DORIĆ M: Integrated and organic apple production. *Savremenapoljop.*, 61(3-4)286-295, 2012.
- MARCHIOL L, MONDINI C, LEITA L, ZERBI G: Effects of municipal waste leachate on seed germination in soil-compost mixtures. *Restor. Ecol.*, 7(2)155-161, 1999.
- MESBAHZADEH T, ASTARAEI AR: Comparison of urban solid waste leachate and vermi-compost as foliar spray and root media on nutritional performance of *Phaseolus vulgaris* L. *Desert*, 13(1)19-23, 2008.
- MOKHTARANI N, BAYATFARD A, MOKHTARANI B: Full scale performance of compost's leachate treatment by biological anaerobic reactors. *Waste Manage. Res.*, 30(5)524-529, 2012.
- OKUR N, KAYIKCIOGLU HH, OKUR B, DELIBACAK S: Organic amendment based on tobacco waste compost and farmyard manure: influence on soil biological properties and butter-head lettuce yield. *Turk. J. Agric. For.*, 32(2)91-99, 2008.
- PANT AP, RADOVICH TJK, HUE NV, PAULL RE: Biochemical properties of compost tea associated with compost quality and effects on pakchoi growth. *Sci. Hortic.*, 148:138-146, 2012.
- SAITHEP N, DHEERANUPATANA S, SUMRIT P, JEERAT S, BOONCHALEARMKIT S, WONGSANON J, JATISATIENR C: Composting of tobacco plant waste by manual turning and forced aeration system. *Maejo Int. J. Sci. Tech.*, 3(2)248-260, 2009.
- SANCHEZ-MONEDERO MA, URPIAINEN ST, CABANAS-VARGAS DD, KAMILAKI A, STENTIFORD EI: Assessing the stability and maturity of compost at large-scale plants. Proceedings of XXVIII Interamerican Congress of Sanitary and Environmental Engineering, Cancun, Mexico, 27-31. October, 2002, pp. 1-6.
- STERRETT SB: Composts as horticultural substrates for vegetable transplant production. In: *Compost Utilization in Horticultural Cropping Systems* (P.J. Stoffella and B.A. Kahn, eds.). Lewis Publication, Boca Raton, FL, pp. 227-240, 2001.
- WOLKOWSKI R: Nitrogen management considerations for landspreading municipal solid waste compost. *J. Environ. Qual.*, 32(5)1844-1850, 2003.
- YEOMANS J, BREMNER JM: A rapid and precise method for routine determination of organic carbon in soil. *Commun. Soil Sci. Plan.*, 19(13)1467-1476, 1989.
- ZUCCONI F, PERA A, FORTE M, BERTOLDI M: Evaluating toxicity of immature compost. *Biocycle*, 22:54-57, 1981.

UTICAJ PROIZVODA KOMPOSTA NA KLIJANJE SEMENA POVRĆA

*Mira MILINKOVIĆ, Blažo LALEVIĆ, Snežana OLJAČA,
Vlado LIČINA, Jelena JOVIČIĆ PETROVIĆ, Vera RAIČEVIĆ*

Izvod: Cilj ovog rada je određivanje uticaja različitih kompostnih čajeva i ekstrakata na klijanje semena povrća. Kompost korišćen za dobijanje ekstrakata i čajeva potiče od komunalnog otpada (MSW) i otpada iz duvanske industrije (TW). Rezultati postignuti sa komposnim produktima MSW su uporedivi sa kontrolom. Kompostni produkti dobijeni od TW pokazuju značajnu fitotoksičnost koja se može dovesti u vezu sa njihovim hemijskim sastavom. Ekstrakti od MSW komposta doveli su do nižeg germinacionog indeksa u poređenju sa čajevima, što ukazuje na mogućnosti njihove primene.

Ključne reči: kompostni proizvodi, germinacioni indeks, klijanje semena povrća, otpad

Received / Priljen: 20.04.2015.

Accepted / Prihvaćen: .16.11.2015.