



## UvA-DARE (Digital Academic Repository)

### Effects of eHealth interventions on medication adherence: a systematic review of the literature

Linn, A.J.; Vervloet, M.; van Dijk, L.; Smit, E.G.; van Weert, J.C.M.

**DOI**

[10.2196/jmir.1738](https://doi.org/10.2196/jmir.1738)

**Publication date**

2011

**Document Version**

Final published version

**Published in**

Journal of Medical Internet Research

[Link to publication](#)

**Citation for published version (APA):**

Linn, A. J., Vervloet, M., van Dijk, L., Smit, E. G., & van Weert, J. C. M. (2011). Effects of eHealth interventions on medication adherence: a systematic review of the literature. *Journal of Medical Internet Research*, 13(4), [e103]. <https://doi.org/10.2196/jmir.1738>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

Journal List > J Med Internet Res > v.13(4); Oct-Dec 2011 > PMC3278089



J Med Internet Res. 2011 Oct-Dec; 13(4): e103.

PMCID: PMC3278089

Published online 2011 December 5. doi: [10.2196/jmir.1738](https://doi.org/10.2196/jmir.1738)

## Effects of eHealth Interventions on Medication Adherence: A Systematic Review of the Literature

Reviewed by Pythia Nieuwkerk and Carl Stepnowsky

[Annemiek J Linn](#), MSc,<sup>1</sup> [Marcia Vervloet](#), MSc,<sup>1,2</sup> [Liset van Dijk](#), PhD,<sup>2</sup> [Edith G Smit](#), PhD,<sup>1</sup> and [Julia CM Van Weert](#), PhD<sup>1</sup>

[Author information](#) ▶ [Article notes](#) ▶ [Copyright and License information](#) ▶

### Abstract

Go to:

#### Background

Since medication nonadherence is considered to be an important health risk, numerous interventions to improve adherence have been developed. During the past decade, the use of Internet-based interventions to improve medication adherence has increased rapidly. Internet interventions have the potential advantage of tailoring the interventions to the needs and situation of the patient.

#### Objective

The main aim of this systematic review was to investigate which tailored Internet interventions are effective in improving medication adherence.

#### Methods

We undertook comprehensive literature searches in PubMed, PsycINFO, EMBASE, CINAHL, and Communication Abstracts, following the guidelines of the Cochrane Collaboration. The methodological quality of the randomized controlled trials and clinical controlled trials and methods for measuring adherence were independently reviewed by two researchers.

#### Results

A total of 13 studies met the inclusion criteria. All included Internet interventions clearly used moderately or highly sophisticated computer-tailored methods. Data synthesis revealed that there is evidence for the effectiveness of Internet interventions in improving medication adherence: 5 studies

### Formats:

[Abstract](#) | [Full Text](#) | [Printer Friendly](#)

### Related citations in PubMed

Reminder packaging for improving adherence to self-administered long-term m [Cochrane Database Syst Rev. 2011]

Interventions to improve adherence to antiretroviral therapy in children with HIV infectior [Cochrane Database Syst Rev. 2011]

Is immediate imaging important in managing low back pain? [J Athl Train. 2011]

Interventions to improve adherence to lipid lowering medication. [Cochrane Database Syst Rev. 2010]

Interventions for enhancing medication adherence. [Cochrane Database Syst Rev. 2008]

[See reviews...](#)


[See all...](#)

### Links

[PubMed](#)

### Recent activity

[Turn Off](#) [Clear](#)

 Effects of eHealth Interventions on Medication Adherence: A Systematic Review of... PMC

[See more...](#)

(3 high-quality studies and 2 low-quality studies) showed a significant effect on adherence; 6 other studies (4 high-quality studies and 2 low-quality studies) reported a moderate effect on adherence; and 2 studies (1 high-quality study and 1 low-quality study) showed no effect on patients' adherence. However, most studies used self-reported measurements to assess adherence, which is generally perceived as a low-quality measurement. In addition, we did not find a clear relationship between the quality of the studies or the level of sophistication of message tailoring and the effectiveness of the intervention. This might be explained by the great difference in study designs and the way of measuring adherence, which makes results difficult to compare. There was also large variation in the measured interval between baseline and follow-up measurements.

### Conclusion

This review shows promising results on the effectiveness of Internet interventions to enhance patients' adherence to prescribed long-term medications. Although there is evidence according to the data synthesis, the results must be interpreted with caution due to low-quality adherence measurements. Future studies using high-quality measurements to assess medication adherence are recommended to establish more robust evidence for the effectiveness of eHealth interventions on medication adherence.

**Keywords:** Internet interventions, medication adherence, compliance, systematic review, tailoring, eHealth, effects, RCT

### Introduction

[Go to:](#)

Recent reports of the World Health Organization and the National Institute for Health and Clinical Excellence reveal that 30%–50% of patients with chronic illnesses do not adhere to prescribed medication [1]. Other studies also show that rates of nonadherence are very high and depend on the type of disease. The highest adherence rates are found for patients with human immunodeficiency virus infection, while diabetes patients have the lowest rate [2]. As such, medication nonadherence can be considered an important health care problem. This is especially true for patients with a chronic illness because medication adherence is a crucial factor in the effectiveness of a therapy [2]. Consequently, many patient-centered interventions are developed to improve adherence, and the impact of the Internet in the development of these interventions is increasing. It is therefore important to understand how these interventions work and to know whether they are effective in improving adherence. To our knowledge, no recent review has studied the effectiveness of patient-centered Internet interventions on patients' medication adherence. Therefore, we conducted a systematic literature study in which we reviewed evidence from studies on Internet interventions that were developed to assist patients in their medication management. The purpose of our study was fourfold: first, to gain insight into the current stage of development of these interventions; second, to assess the included studies for their effectiveness on medication adherence; third, to investigate to what degree adherence is determined by the characteristics of the intervention; and fourth, to investigate whether there is a relationship between the characteristics of the study and the reported effectiveness of the interventions.

Different terms are used in the literature to describe the concept of adherence—for example, compliance, adherence, and persistence. They have all been used to indicate that the patient is using the medication following the prescribed regimen. These terms differ in exact meaning. In this paper, we use the term adherence. *Adherence* is defined as the extent to which the patient's behavior matches

---

Variations in patients' adherence to medical recommendations: a quantitative review of † [Med Care. 2004]

---

**Review** Review article: understanding adherence to medication in ulcerative colitis: [Aliment Pharmacol Ther. 2010]

Intentional and unintentional nonadherence: a study of decision making. [J Behav Med. 2002]

the agreed recommendations of the prescriber [3]. According to this definition, nonadherence is a wide concept that varies from missing an occasional dose to never taking the prescribed medications [3]. Patients have different reasons for being nonadherent. These different reasons have something in common: the patient does not execute the treatment plan and does not persist. Execution is a continuous process where the actual dosing history corresponds to the ideal doses [4,5].

To improve adherence and develop target interventions, it is important to address the specific reasons why a patient is not able or willing to execute the treatment plan. From this perspective, interventions should be personalized or tailored to address individual needs and beliefs. The definition of tailoring describes the features that make tailored health messages different from other approaches: "It is assessment-based and as a result the message can be individual-focused" [6]. In other words, tailoring is based on gathering and assessing personal data related to health outcomes or several determinants in order to determine the most effective strategy to meet that person's needs [6]. With these characteristics, a tailored message is able to provide personal feedback, commands greater attention, is processed more deeply, and is perceived as more likable by patients than a general message [7,8]. Because of these possibilities, tailored health messages are also more likely than generic information to be read, remembered, and viewed as personally relevant [6,9].

Computer technologies can be used to tailor health messages to the personal situation of the patient and might therefore contribute significantly to the development of tailored message strategies. The Internet is potentially a powerful medium for delivering those tailored messages. The management of a chronic disease should be personalized to an individual because the person is ultimately responsible for the success of the intervention [10]. The technology provides an opportunity to tailor the information in several formats and modalities, which enhances the user's experience of the material and will result in a better understanding [7,8]. Moreover, Internet interventions have the advantage that they can provide interactive and responsive programs [10]. These interventions can provide effective data and information provision and retrieval. The advantages of tailored message strategies can contribute to the incorporation of interactive and continued self-monitoring, feedback, and information exchange, which play an increasingly important role in changing patients' behavior.

## Methods

[Go to:](#)

For this review, we used the guidelines of the Cochrane Collaboration to assess the studies on their internal validity and to summarize the existing evidence about Internet interventions to improve medication adherence in patients. The Cochrane Collaboration method is described in more detail in the Cochrane Handbook for Systematic Reviews of Interventions [11].

### Inclusion Criteria

We included a study when the following inclusion criteria were met: (1) the study described a patient-centered Internet intervention, (2) the study described an intervention for patients who use prescribed medication for a chronic condition, (3) at least one of the outcome measures was adherence, (4) the study was quantitative, and (5) the study was published in either the English or Dutch language.

### Search Methods

We conducted a systematic literature search to identify articles containing information about the effect of Internet interventions on medication adherence. Comprehensive literature searches were

---

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

Does tailoring matter? Meta-analytic review of tailored print health behavior change interventions. [Psychol Bull. 2007]

**Review** Tailored and targeted health communication: strategies for enhancing informati [Am J Health Behav. 2003]

---

**Review** The effectiveness of Web-based vs. non-Web-based interventions: a meta-analy [J Med Internet Res. 2004]

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

Does tailoring matter? Meta-analytic review of tailored print health behavior change interventions. [Psychol Bull. 2007]

undertaken in the databases PubMed, PsycINFO, EMBASE, CINAHL, and Communication Abstracts.

The search strategies used the following keywords: (medication therapy management OR medication adherence OR patient compliance OR self-care) AND (internet) AND (intervention study OR randomized controlled trial OR clinical controlled trial).

We then continued with the snowball method by looking for references in publications, especially those of the included studies and reviews on interventions to promote adherence. The search was conducted in June 2010. Since Internet intervention is a relatively new topic, no time limits were applied.

Application of the search strategy to the specified databases resulted in a total of 620 hits ([Table 1](#)). In total, we selected 13 studies from these results.

Database	Number of hits	Percentage of total hits
PubMed	120	19.35%
PsycINFO	100	16.13%
EMBASE	150	24.19%
CINAHL	100	16.13%
Communication Abstracts	50	8.06%
Total	620	100%

**Table 1**

Results of database searches.

Reference Manager version 11.0 (Thomson Reuters, New York, NY, USA) was used to manage the citations. Duplications were logged, leaving 520 unique results (see [Figure 1](#)). On the basis of title and abstract, two researchers (pairs of AL, MV, LvD, JvW) independently selected studies for inclusion. If the study seemed to meet the inclusion criteria or if there were doubts about the inclusion, the full text of the article was obtained. Based on the full articles two reviewers independently reviewed whether these studies fit all the inclusion criteria. Disagreements were solved by discussions between the two researchers. For a more detailed description of the excluded studies see [Multimedia Appendix 1](#).



**Figure 1**

Flow diagram of study search and selection.

### Assessment of Methodological Quality

The methodological quality of included randomized controlled trials (RCTs) and clinical controlled trials (CCTs) was independently reviewed by two researchers (AL and JvW) using the list from the Cochrane Collaboration Back Review Group [[12](#)]. The list consists of 11 criteria for internal validity, namely:

- 3 criteria regarding selection bias: whether (a) randomization was adequate, (b) treatment allocation was concealed, and (c) groups were similar at baseline regarding the most important indicators,
- 4 criteria for performance bias: whether (d) patients were blinded to the intervention, (e) care provider was blinded to the intervention, (g) co-interventions were avoided, and (h) compliance with the intervention was acceptable,
- 2 criteria regarding attrition bias: whether (i) the dropout rate after baseline was acceptable, and (k) the analysis included an intention-to-treat analysis, and

- 2 criteria for detection bias: whether (f) the outcome assessor was blinded to the intervention, and (j) outcome assessments in all groups were similar.

For each included study, all criteria were scored as “yes,” “no,” or “unclear.” All unclear scores were later rated as “no.” Studies were rated as high quality (HQ) when at least 6 of the 11 criteria for internal validity were met. Otherwise, studies were considered of low quality (LQ). Disagreements were discussed until consensus was reached. If disagreement or indistinctness persisted a third reviewer (LvD) was consulted.

In addition, two researchers (AL and LvD) independently assessed the quality of the methods for measuring adherence to a medical regimen. A standard method to assess adherence does not exist and every method has its limits [13,14]. In clinical trials, adherence can be measured based on, for example, interviews, diary, questionnaire-based self-reporting, prescription refills, pill counts, or electronic monitoring [14,15]. We categorized the measurements in high- and low-quality adherence assessments based on previous findings concerning the objectivity of these measurements [13-15]. In this review electronic monitoring and physiological/biomedical measures are defined as high-quality adherence assessment. These measurements are considered the most objective standard [15]. Previous research has shown that data from pill counts and electronic monitoring are strongly correlated [16]. Yet others consider pill counts not to be accurate [17,18]. In addition, meta-analyses have shown that self-reported adherence is also strongly correlated with electronic monitoring [19,20]. Like pill counts, the accuracy of self-reported measurements is debatable. Some argue that self-reports may be an accurate measurement for measuring adherence [21,22], while others state that the use of self-reported measurements is not an accurate method [14,15,23,24]. Taking all arguments into account, we considered self-measurements, such as questionnaires, pill counts, prescription refills, interviews, and diaries, to be most subjective for measuring adherence [13]. We therefore considered these measurements low-quality adherence assessment. However, if two or more different low-quality adherence measurements were used in the same study, such as a combination of questionnaires and prescription refills, the method was considered high-quality adherence assessment.

### Data Extraction

One researcher (AL) documented the following characteristics of the included studies: (1) method (type of study), (2) participants (total number of participants, sex per group, mean age per group, type of disease), (3) intervention (name of experimental intervention, name of control condition, period, number of times/minutes per week), (4) outcome measures (type of outcome measures, time of measurement), (5) results (short description), and (6) author’s conclusion.

### Data Synthesis

Due to diversity in the features of the interventions and the methods used to measure adherence, it was not possible to pool the data. Therefore, we conducted a best-evidence synthesis (see [Textbox 1](#)) based on [12] and adapted by a Dutch study [25].

#### Principles of Best-Evidence Synthesis.

Evidence:

Provided by consistent, statistically significant findings in outcome measures in at least 2 high-

---

All that glitters is not gold: a comparison of electronic monitoring versus filled prescrip [BMC Health Serv Res. 2006]

**Review** Methods for measuring and monitoring medication regimen adherence in clinical trials and clini [Clin Ther. 1999]

Relationships among subjective and objective measures of adherence to oral antipsychotic medic [Psychiatr Serv. 2007]

Time to stop counting the tablets? [Clin Pharmacol Ther. 1989]

The natural history of medication compliance in a drug trial: limitations of pill counts. [Clin Pharmacol Ther. 1989]

Correlation between adherence rates measured by MEMS and self-reported question [Health Qual Life Outcomes. 2010]

**Review** Concordance of adherence measurement using self-reported adherence questior [Pharmacoeconomics. 2010]

Pill count, self-report, and pharmacy claims data to measure medication adherence in the elderl [Ann Pharmacother. 1998]

Patient adherence and medical treatment outcomes: a meta-analysis. [Med Care. 2002]

[See more ...](#)

---

Updated method guidelines for systematic reviews in the cochrane collaboration back rev [Spine (Phila Pa 1976). 2003]



quality randomized controlled trials (RCTs).

**Moderate evidence:**

Provided by consistent, statistically significant findings in outcome measures in at least 1 high-quality RCT and at least 1 low-quality RCT or high-quality CCT.

**Limited evidence:**

Provided by statistically significant findings in outcome measures in at least 1 high-quality RCT

Or

Provided by consistent, statistically significant findings in outcome measures in at least 2 high-quality CCTs (in the absence of high-quality RCTs).

**Indicative findings:**

Provided by statistically significant findings in outcome measures in at least 1 high-quality CCT or low-quality RCT (in the absence of high-quality RCTs).

**No/insufficient evidence:**

If the number of studies that have significant findings is <50% of the total number of studies found within the same category of methodological quality and study design

Or

In case the results of eligible studies do not meet the criteria for one of the above stated levels of evidence

Or

In case of conflicting (statistically significantly positive and statistically significantly negative) results between RCTs and CCTs

Or

In case of no eligible studies.

The best-evidence synthesis was conducted by attributing various levels of evidence to the effectiveness of the interventions. The synthesis takes into account the design, methodological quality, and outcomes of the studies. [Textbox 1](#) shows that at least 1 HQ RCT or 2 HQ CCTs were needed to establish robust evidence for the effectiveness of an intervention.

### **Sensitivity Analysis**

We conducted a sensitivity analysis to identify how sensitive the results of the best-evidence syntheses were to changes in the way the study quality was assessed. For the sensitivity analysis, the best-evidence synthesis was repeated in two different ways, using the following principles: (1) LQ studies were excluded, (2) studies were rated as HQ if they met at least 4 of the 11 criteria of internal validity instead of 6.

We then compared the results of the sensitivity analysis with the results of the best-evidence synthesis

and described the sensitivity of the results [25,26].

**Effectiveness**

Study effectiveness was categorized as significant effect on adherence, moderate effect on adherence, and no effect on adherence. We defined a study effect as moderate if the authors reported a positive effect of the intervention on adherence but there were limitations, such as the following: improvement of adherence was found only in a subgroup of the intervention group; adherence was measured indirectly (eg, the study drew conclusions about the use of beta-agonist indicating that adherence was improved); or the significance of the results to medication adherence was not tested, but the authors used convincing arguments to explain the effectiveness of the intervention (see results section for explanation per study).

**Intervention: Tailoring Level of Sophistication of the Website**

Tailored Internet interventions differ in how they deliver their message [7]. The difference is based on the sophistication of the way the message is tailored. We categorized the interventions in being low, moderate, or high in sophistication. Some interventions involve a form of online assessments (low sophistication), and others use online assessments, tailored feedback, and content matching (moderate sophistication). The third group of interventions provides instant feedback and a complex tailored health program with several tools and activities that would enable patients to achieve their health goals (high sophistication) (see Figure 2) [7].



**Figure 2**

Continuum level of sophistication of tailored intervention.

**Results**

[Go to:](#)

The main characteristics of the included studies are presented in Table 2 [27-39] and further described below (for a more detailed description of the included studies see Multimedia Appendix 2).

**Table 2**

Characteristics of included studies.

[See links ...](#)

**Methodological Quality: Assessment of Internal Validity**

For this review 10 RCTs were included, and 9 of them were assessed on their internal validity (Table 3 [27-39]). We included 1 RCT with no data on our primary outcome variable (ie, medication adherence) for the control group [27]. This means that for this tenth study, we could not assess validity criteria. Moreover, we reviewed 2 prospective cohort designs and 1 survey. A total of 7 RCTs met 6 or more of the 11 validity criteria and therefore qualified as HQ studies.

**Table 3**

Results of methodological quality.

---

**Review** The effects of psychosocial methods on depressed, aggressive and apathetic beha [Int J Geriatr Psychiatry. 2005]

---

**Review** Computer-tailored health interventions delivered over the Web: review and analy [Patient Educ Couns. 2009]

---

Pilot study of a Web-based compliance monitoring device for patients with congestive heart failure. [Heart Lung. 2003]

---

An internet-based intervention to improve psychosocial outcomes in heart transplant [J Heart Lung Transplant. 2004]



**Intervention: Tailoring Level of Sophistication of the Website**

All Internet interventions reported computer-tailoring methods. Interventions were categorized as having low sophistication (online assessments), moderate sophistication (online assessments, tailored feedback, and content matching), and high sophistication (a more complex tailored health program) (see [Figure 2](#)) [7].

**Online Assessment and Feedback** Online assessment and feedback are used in interventions with single-incident computer-assisted risk or health assessments. For example, feedback is emailed to the patients or provided online [7]. In addition, these interventions are brief and usually done once at the beginning of the intervention. None of the reviewed studies used online assessment and feedback.

**Tailored Content** With tailored content a program provides (1) tailored text messages composed in a unique way according to how patients respond to certain questions, or (2) restricted access to content sections per patient [7]. Tailored content was used by 9 of the studies we reviewed; 1 study [28], Blue Angel for Asthma Kids, conducted an Internet-based interactive asthma educational and monitoring program in which patients were able to complete an electronic diary, record symptoms and need for rescue medication, and upload their videos when they were using their inhaler. Based on these outcomes, the program comprised both an action plan with a warning system and a written treatment plan. A similar customized educational and monitoring Website for patients with asthma was conducted in 2 studies [29,30], and 1 study [31] tested the asthma management program Puff City. The program used tailoring to alter behavior through individualized health messages based on the patients' beliefs, attitudes, and personal barriers to change or maintain the behavior. Another form of a tailored website, System Providing Access to Records Online (SPARO), was examined in 1 study [32]. This website provided the medical record, an educational guide, and a message system for patients. Moreover, patients could contact the health provider by email. The telemedicine diabetes disease management program Health Buddy was tested in 2 studies [33,34]. However, the modality used in the diabetes program was different from the previously mentioned monitoring programs. Patients answered personalized questions that enabled them to monitor their disease symptoms, medication adherence, and disease knowledge by pressing buttons for response. The 2 studies using the Health Buddy differed in the intensity of the feedback. One study [27] tested a medication compliance device. Data and answers to questions were recorded by the device and uploaded daily to a central server. Based on these answers health providers were able to monitor the patients, provide advice, and update the treatment regimens in the Med-eMonitor devices. One study [35] also tested a handheld device, Pocket Personal Assistant for Tracking Health (PATH), developed for patients after lung transplantation to record health data, review data trends, and report their condition changes to the transplant team. The device included decision-support programs to promote self-care behaviors.

**Customized Health Programs** Interventions that provide not only tailored content but also individualized instructions for meeting certain health goals, self-management goals, or goal-setting activities are so-called customized health programs [7], used by 4 of the included studies. Of these, 2 studies [36,37] tested the effects of an Internet-based self-management program for asthma patients. This website allowed monitoring through the website, text messages, use of an Internet-based treatment plan, online education, and the possibility to communicate with the health provider. The intervention WebEase [38] consisted of three modules that were designed to assess an individual's status related to self-management practices and create a plan for change or to maintain the behavior. The modules in WebEase required the patient to answer questions related to these topics. Feedback was provided

---

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

---

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

---

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

An internet-based interactive telemonitoring system for improving childhood asthma outcomes [Telemed J E Health. 2007]

Internet-based home monitoring and education of children with asthma is comparable to ideal office-based care [Pediatrics. 2007]

An Internet-based store-and-forward video home telehealth system for improving asthma management [Am J Health Syst Pharm. 2003]

A web-based, tailored asthma management program for urban African-American children [Am J Respir Crit Care Med. 2007]

Providing a web-based online medical record with electronic communication capabilities to patients [J Med Internet Res. 2004]

Diabetes disease management program for an indigent population empowered by telemedicine [Diabetes Technol Ther. 2002]

Improving asthma outcomes and self-management behaviors of inner-city children: a randomized trial [Arch Pediatr Adolesc Med. 2002]

Pilot study of a Web-based compliance monitoring device for patients with congestive heart failure. [Heart Lung. 2003]

Evaluation of a hand-held, computer-based intervention to promote early self-care behaviors after lung transplantation [Clin Transplant. 2009]

---

**Review** Computer-tailored health interventions delivered over the Web: review and analysis [Patient Educ Couns. 2009]

Internet-based self-management plus education compared with usual care in asthma: a randomized trial [Ann Intern Med. 2009]

based on these responses. Patients entered data into MyLog, which is a screen for recording data about medication-taking behavior, stress, etc. In addition, the intervention included a knowledge component and a discussion board. This means that each patient was directed to another path [38]. Another study [39] tested a customized health program where patients chose which components of the website they wanted to use. The website included a home page, posttransplant skills workshops, discussion groups, “ask an expert,” question-and-answer possibility, healthy-living tips, resources, and reference library. The way the patients used the website was based more on voluntary participation than in the study that used MyLog [38].

Table 4 and Table 5 show for each study which method for delivering the tailored message was used (see column 2).

Study	Study quality	Substitution of tailoring	Year
Chen et al. (2004)	High	Medication	2004
Chen et al. (2005)	High	Medication	2005
Chen et al. (2006)	High	Medication	2006
Chen et al. (2007)	High	Medication	2007
Chen et al. (2008)	High	Medication	2008
Chen et al. (2009)	High	Medication	2009
Chen et al. (2010)	High	Medication	2010
Chen et al. (2011)	High	Medication	2011
Chen et al. (2012)	High	Medication	2012
Chen et al. (2013)	High	Medication	2013
Chen et al. (2014)	High	Medication	2014
Chen et al. (2015)	High	Medication	2015
Chen et al. (2016)	High	Medication	2016
Chen et al. (2017)	High	Medication	2017
Chen et al. (2018)	High	Medication	2018
Chen et al. (2019)	High	Medication	2019
Chen et al. (2020)	High	Medication	2020
Chen et al. (2021)	High	Medication	2021
Chen et al. (2022)	High	Medication	2022
Chen et al. (2023)	High	Medication	2023
Chen et al. (2024)	High	Medication	2024
Chen et al. (2025)	High	Medication	2025

Table 4

Effectiveness of short-term interventions (<6 months).

Study	Study quality	Substitution of tailoring	Year
Van der Meer (2002)	High	High	2002
Van der Meer (2003)	High	High	2003
Van der Meer (2004)	High	High	2004
Van der Meer (2005)	High	High	2005
Van der Meer (2006)	High	High	2006
Van der Meer (2007)	High	High	2007
Van der Meer (2008)	High	High	2008
Van der Meer (2009)	High	High	2009
Van der Meer (2010)	High	High	2010
Van der Meer (2011)	High	High	2011
Van der Meer (2012)	High	High	2012
Van der Meer (2013)	High	High	2013
Van der Meer (2014)	High	High	2014
Van der Meer (2015)	High	High	2015
Van der Meer (2016)	High	High	2016
Van der Meer (2017)	High	High	2017
Van der Meer (2018)	High	High	2018
Van der Meer (2019)	High	High	2019
Van der Meer (2020)	High	High	2020
Van der Meer (2021)	High	High	2021
Van der Meer (2022)	High	High	2022
Van der Meer (2023)	High	High	2023
Van der Meer (2024)	High	High	2024
Van der Meer (2025)	High	High	2025

Table 5

Effectiveness of long-term interventions (>6 months).

**Role of Health Providers** Interventions also differ in the type and extent of health provider involvement. User control allows individuals to take a major role in managing their own care, whereas in expert control an expert or therapist takes a more directive role [7].

Only 4 of the interventions were based on user control and 9 interventions used contact with the health provider. A Web-based asthma management program was developed in 1 study [31]. The program used tailoring to alter behavior through individualized health messages based on the user’s beliefs, attitudes, and personal barriers to change. The health provider did not interfere. Three studies were also user based with treatment algorithms to give feedback [36-38].

In contrast, in the Blue Angel for Asthma Kids, a customized educational and monitoring website site providing secure email contact between patients and their therapist, the therapist had a more directive role [28]. Like the Blue Angel for Asthma Kids, SPPARO included a messaging system that made it possible to exchange secure messages with the health provider [32]. The intervention developed in 2 studies [29,30] consisted of a case manager who reviewed the data, sent emails about the peak flow, inhaler technique, and symptoms, and forwarded them the website. Patients (the virtual group as well as the office-based group) had access to their case manager 24 hours a day, 7 days a week. One study [39] conducted a website including skills workshops, discussion group, ask an expert, question and answer, health tips, recourses, and references. In this case, the role of health providers was to provide the possibility for a patient to ask an expert.

Several interventions used a special device to exchange information between patients and health care providers. The Med-eMonitor recorded data and answers to questions that could be uploaded by the health provider. Based on these outcomes, the health provider provided advice and updated the treatment regimens [27]. The PATH [35] recorded data and provided a tailored decision-support program and email contact with the health provider. The Health Buddy device in 2 studies [33,34] is like the Med-eMonitor and PATH based on the feedback and participation of the health provider, but

Weekly self-monitoring and treatment adjustment benefit patients with partly controlled and uncontr [Respir Res. 2010]

Evaluation of WebEase: an epilepsy self-management Web site. [Health Educ Res. 2009]

An internet-based intervention to improve psychosocial outcomes in heart transplant [J Heart Lung Transplant. 2004]

there is a difference in the intensity of participation of the health provider. In 1 study [33] the health provider contacted the patient only when necessary, while in the other 2 studies [34,35] the patient received feedback instantly after sending a question.

Our analysis to examine the extent to which medication adherence is determined by different tailoring levels revealed no clear relationship between the intervention's level of sophisticated tailoring and the extent to which the intervention was effective (Table 4).

### Summary of Effects on Adherence

We found 5 studies with a significant effect on adherence (3 HQ studies and 2 LQ studies) [28,31,34,35,38]. The first study [28] concluded that the intervention had a positive and significant effect on use of the inhaled corticosteroid and that this effect significantly differed from the baseline. In addition, the second study [31] found positive changes in controller medication adherence. The third study [38] tested the WebEase intervention and found a significant effect on adherence. The fourth study [34] found that patients were more likely to take their asthma medication when they used the Health Buddy. In the fifth study [35], patients who received the PATH intervention were more likely to adhere to their medical regimen.

A moderate effect on adherence was reported in 6 studies (4 HQ studies and 2 LQ studies). The SPPARO proved to be feasible and improved general adherence. Adherence to medication showed a similar trend but these results did not reach significance [32]. The second study [30] concluded that, after the intervention, the use of beta-agonist decreased, which is an indication of better adherence. The third study [33] reported that medication compliance improved from 65% at pretest to 94% at posttest, but the difference was not statistically tested. The medication compliance rate in the fourth study [27] was 94% for the monitor group as measured by the monitor system. However, because there was no pretest and the data of the control group were not available, it is unknown whether the results were significant as compared with the pretest or the control group. The intervention of the last 2 studies [36,37] improved adherence in patients with partly and uncontrolled asthma at baseline. The authors concluded that the intervention was most effective in improving adherence for patients with partly or uncontrolled asthma at baseline.

No significant results on patients' adherence were found in 2 studies (1 HQ and 1 LQ) [29,39].

### Assessment of Adherence Measurements

Regarding the measurement of adherence, the 13 studies we reviewed showed a large variability of methods: 12 studies used a low-quality measurement to assess adherence and 1 used a combination of these methods (ie, a high-quality measurement to assess adherence) (Table 3).

**Low Quality of Adherence Measurement** In 10 studies, self-reported scales were used to obtain the adherence rate. Although 5 studies [28,31,34,36,37] used self-reports to measure adherence, they did not describe what kind of instrument they used. One study [39] used self-reported data by asking questions regarding adherence during the initial interview. Reports of therapist and patients were compared. In addition, 1 study [33] used a self-developed medication adherence survey on the Health Buddy appliance. The other 3 studies chose existing, valid, self-reported adherence scales. One study [32] used a combination of the Morisky scale and the General Adherence Scale from the Medical Outcomes Study, and 1 study [35] used the Health Habits Assessment, a self-reported scale to measure adherence. One study [38] used the self-report USCF Adherence Questionnaire and the

Antiretroviral General Adherence Scale. Finally, 2 studies used measurements such as counting pills [27], and 1 study [29] used a computerized prescription refill record (after evaluation of the pilot study in which completing the diary turned out to be time consuming and inconvenient; see [30]).

**High Quality of Adherence Measurement** One study used a combination of methods. This study [30] used a diary in combination with a computerized prescription refill record.

Table 3 shows the results of the assessment of the internal validity and the quality of adherence measurement.

#### Relation Between Quality of Adherence Measurement and Effectiveness

Our investigation of the relationship between the quality of the adherence measurement and the effectiveness of the interventions revealed no clear relationship (there was only 1 study using a high-quality method to assess adherence), although self-reported adherence measurements seemed to result more often in significant effects than did pill counts and pharmacist adherence measurements (Table 4 and Table 5). Of the 10 studies using self-reports (low-quality adherence measurement), 5 reported a significant effect of the intervention on adherence [28,31,34,35,38], 4 a moderate effect [32,33,36,37], and 1 no effect [39]. From the 2 studies in which pharmacist data or pill counting was used, 1 reported a moderate effect [27] and 1 no effect [29]. The 1 study that used a combination of methods to measure adherence [30] found a moderate effect on adherence.

#### Relation Between Interval of Adherence Measurement and Effectiveness

There was no clear relationship between the timing of the adherence measurements and the effectiveness of the intervention. The intervals between baseline and follow-up measurements differed between projects. Short-term adherence (ie, within 6 months) was measured in 6 studies. The first study [38] showed that WebEase improved adherence 6 weeks after baseline. Patients who used PATH were more likely to show higher adherence than the control group after an interval of 8 weeks after baseline [35]. The third study [28] found a significant effect on adherence after 12 weeks, and the fourth study [34] reported an improvement in adherence after 12 weeks. The fifth study [27] found an adherence rate of 94% in the experimental group after 12 weeks. Because of the lack of adherence data for the control group and the lack of a pretest, the effects on adherence could not be established. The sixth study [39] examined the proportion of nonadherent patients in both an intervention and a control group after 16 weeks. That study's authors found uniformly small and nonsignificant differences between the control and intervention groups. However, they found an important difference within the intervention group. Subgroup differences appeared when the intensity of using parts of the intervention was related to the effectiveness. For example, patients who used the Managing Medical Regimen Workshop more often or intensely appeared to be more adherent than those using the intervention less often or intensely [39].

Long-term adherence was measured in 7 studies—that is, adherence with an interval of 6 months or longer, mostly of 1 year or more. Two studies [30,36] reported a moderate effect in their pilot on adherence after 6 months. In 2 studies [31,32], they found a moderate [32] and significant [31] improvement in adherence after 1 year. Two studies [33,37] found a moderate effect on adherence after 1 year. This means that all of the included studies using an interval of 6 months or longer showed an effect (significant or moderate) on long-term adherence. One study [29] did not find an effect on adherence after 1 year.

[Table 4](#) and [Table 5](#) give an overview of the methodological quality of the studies, the level of sophistication of each intervention, the quality of measurement of adherence, and an overview of the short-term and long-term effects.

### Data Synthesis

Using the principles of the best-evidence synthesis (see [Textbox 1](#)), taking into account the design, methodological quality, and outcomes of the studies, the following conclusions can be drawn. In total, 7 studies were considered HQ. We found 3 HQ studies [[28,31,35](#)] and 2 LQ studies [[34,38](#)] that had a significant effect on enhancing medication adherence and that met 6 of 11 criteria. This means that there is evidence that tailored Internet interventions are successful in improving medication adherence.

### Sensitivity Analysis

The sensitivity analysis showed the same results as the best-evidence synthesis. The results remained the same when the analysis was repeated with the 6 LQ studies excluded (ie, taking only the 7 HQ studies into account). Moreover, when studies were rated to be HQ if 4 instead of 6 criteria of interval validity were met, results stayed the same.

## Discussion

[Go to:](#)

### Principal Results

First, our objective was to gain insight into the current state of the use of Internet interventions to improve medication adherence. Results of this review indicate that this is still a new field. This is visible in the differences in interventions with respect to crucial aspects such as the level of sophisticated tailoring and the role of health care providers. Despite the differences, it is remarkable that none of the interventions used a low level of tailoring and the majority (9 of 13) provided the opportunity to contact a health provider.

Second, the studies were assessed on their effectiveness on medication adherence. There is evidence that Internet interventions can improve medication adherence. This evidence comes from 3 HQ studies and 2 LQ studies, finding significant results on medication adherence.

Third, we wanted to investigate to what degree adherence is determined by the characteristics of the intervention. All interventions discussed in this review used tailored methods and used a moderately or highly sophisticated tailored intervention. These types of health programs, especially customized health programs, are more complex, generally long-term, allowing the patients to access the programs several times [[7](#)], and are considered appropriate for difficult-to-influence behaviors. We did not find a clear relationship between how sophisticated the tailoring of the intervention was and the extent to which the intervention appeared to be effective, possibly due to the various methods that were used.

Last, we wanted to investigate whether there is a relationship between the characteristics of a study and the reported effectiveness of the interventions. We found that there was variation not only in the level of tailoring, but also in the measurement of adherence, the timing of measuring adherence, and the intensity of the intervention. The included studies used self-reporting measurements (ie, interviews, diary, self-reporting via questionnaires) or pill counts, or prescription refills, or a combination. No study used electronic monitoring, which is perceived as a high-quality method for

assessing adherence [15]. Of the 13 studies we reviewed, 7 measured long-term adherence, using an interval of 6 months or longer. There is no clear evidence that the duration of the intervention is related to the effectiveness of the intervention. Nevertheless, of the 7 studies measuring long-term adherence, 1 HQ study showed positive effects and 4 HQ studies and 1 LQ study showed moderate effects on adherence. This indicates that long-term interventions are promising. However, more research in this field is needed.

There is evidence that Internet interventions can be effective in improving medication adherence. The evidence comes from 3 HQ studies. However, the results should be interpreted with caution. Self-reported scales were used in 10 studies, which is considered a low-quality adherence measurement: 5 reported a significant effect of the intervention on adherence, 4 a moderate effect, and 1 no effect. Self-measurements can contribute to overestimating of the effects of interventions [40]. This could be explained by the possibility that patients may forget that they missed a dose. Biases that appear most prominent in estimating adherence by the patient from structured questionnaires are social desirability and social approval [5,13]. In other words, studies relying on self-reporting may have a tendency to err on the optimistic side when it comes to adherence, certainly compared with more objective pill-counting studies. However, it must be noted that anonymous self-report questionnaires are found to be significantly correlated with electronic monitoring [41] and virologic response [40], considered more objective methods. Research also shows that using specific strategies, such as ensuring patients that their responses will be kept confidential [40] or stratifying patients according to their socially desirable response [42], improves the prediction of adherence by self-reports. This indicates that self-reports are not useless, but future research should examine more strategies to reinforce accurate reporting by patients [40]. On the other hand, we included RCTs and, consequently, self-reported adherence can be expected to be overestimated in both treatment arms. Thus, the intervention effect (ie, difference between intervention and control group) was not necessarily overestimated. Additionally, a distinction can be made between valid self-report measurements and measurements that are not. If self-reported measurements are used, for instance because this is a cost-effective method, using validated measurements is recommended.

Electronic monitoring or observation is considered to be more accurate. It electronically records the time and date of the actual dosing events. Because every single method has its limitations, the best approach is to use multiple assessment techniques concurrently, as a way to improve the accuracy of adherence assessment [14]. One study in our review [30] used self-reported diary and prescription refills to measure adherence and made a distinction in interpreting the results. They used a self-reported diary to assess how the patients used their inhaler, based on the idea that when patients are not using their inhaler according to the health providers' advice, they can be considered nonadherent. In addition, they used prescription refills to measure how many refills the patient was obtaining. In line with this method of measuring adherence, the optimal approach could be suggested to be a combination of self-reports and more objective measurements [43]. In addition, every single measurement needs a different interpretive approach because it has different relationships to clinical outcomes [13].

Of the studies we reviewed, 6 measured short-term adherence that varied from 6 weeks to 4 months: 4 of them were found highly effective and 1 moderately effective regarding adherence. The question is whether adherence improved in the long term, because the period was too short to measure persistence. According to an international expert forum on adherence [13], it is not easy to identify



adherent and nonadherent patients beforehand. There is a large body of evidence dominated by reports identifying factors that are predictive or associated with nonadherence. Adherence could be seen as a dynamic behavior that is determined or influenced by unrelated factors that fluctuate and change over time [44]. As an adherent patient can become nonadherent over time, the importance of time (ie, persistence) has been emphasized [5]. The quality of execution of the treatment plan can influence persistence. Factors such as perceptions of treatment outcomes, beneficial effects, and adverse effects can influence the quality of execution over time. Therefore, conducting interventions that address long-term adherence and overcome reasons why a patient is not able or not willing to adhere are recommended.

### Methodological Limitations

The search method was top-down in that we relied on existing databases and search terms. This approach has the possibility of missing important articles due to miscoding of search terms. A bottom-up strategy is more time consuming but has the advantage of being more comprehensive.

### Clinical Implications

Monitoring adherence to optimize effects and minimize nonadherence could be time consuming. Computers, however, are very good at collecting data concerning the monitoring of adherence. Internet interventions can be tailored, collect data, and monitor adherence. In addition, based on this systematic literature review, there is evidence that tailored Internet interventions can be an effective method to improve medication adherence. This means that Web-based interventions can be effective at increasing medication adherence among chronically ill patients. Health providers, who want to enhance patients' adherence, are encouraged to use tailored websites or reminder systems. They could use these interventions in addition to their everyday work.

### Implications for Research

Because we did not find a clear relationship between the effectiveness and the degree of tailoring, we recommend that future studies should be conducted with variation in the level of sophistication of tailoring to further test which characteristics of the tailored messages have the most positive effects on adherence. Moreover, website compliance is often not completely reported. While the frequency in which the patient used the website is often reported, studies do not describe how exactly patients used the website. Therefore, it is difficult to compare the results of different interventions, because the way patients use the website can have implications for the effectiveness of the website.

### Conclusion

With more than 40 million people using the Internet for a variety of purposes, health communication programs in the future are more likely to be delivered online [6]. These types of interventions especially have the potential to address difficult-to-change behaviors such as adherence. This review shows promising results on the effectiveness of tailored Internet interventions to enhance medication adherence of chronically ill patients. There is evidence that these interventions can enhance adherence. But it remains a relatively new field, and studies using more objective measurements to assess adherence are recommended.

### Acknowledgments

Go to:



The authors acknowledge the support of 2Comply. 2Comply had no role in the design and the collection, analysis and interpretation of the data, nor in the writing of this manuscript.

## Abbreviations

[Go to:](#)

CCT clinical controlled trial

HQ high quality (study)

LQ low quality (study)

PATH Pocket Personal Assistant for Tracking Health

RCT randomized controlled trial

SPPARO System Providing Access to Records Online

## Multimedia Appendix 1

[Go to:](#)

Detailed description of the excluded studies.

[Click here to view.](#) (128K, pdf)

## Multimedia Appendix 2

[Go to:](#)

Detailed description of the characteristics of the included studies.

[Click here to view.](#) (111K, pdf)

## Footnotes

[Go to:](#)

Conflicts of Interest:

None declared

Contributed by

JvW and LvD designed the study and acquired funding. MV conducted comprehensive literature searches through the databases. Two researchers pairs (AL/JvW and MV/LvD) performed a first selection for inclusion based on the abstracts. After having obtained full text of articles that seemed to meet the inclusion criteria, the same researchers pairs independently reviewed whether these studies fulfilled all the inclusion criteria. The methodological quality of included RCT's and CCT's were independently reviewed by AL and JvW. In addition, AL and LvD independently assessed the quality of the methods for measuring adherence to a medical regime. AL conducted the 'Best Evidence Synthesis' and drafted the manuscript. MV, LvD, ES and JvW have been involved in critically revising the manuscript. All authors have read and approved the final manuscript.

## References

[Go to:](#)

1. Sabaté E. Adherence to Long-Term Therapies: Evidence for Action. Geneva: World Health Organization; 2003.
2. DiMatteo MR. Variations in patients' adherence to medical recommendations: a quantitative review of 50 years of research. *Med Care*. 2004 Mar;42(3):200–9.00005650-200403000-00002 [[PubMed](#)]
3. Kane SV, Robinson A. Review article: understanding adherence to medication in ulcerative colitis - innovative thinking and evolving concepts. *Aliment Pharmacol Ther*. 2010 Nov;32(9):1051–8. doi: 10.1111/j.1365-2036.2010.04445.x. [[PubMed](#)] [[Cross Ref](#)]

4. Wroe AL. Intentional and unintentional nonadherence: a study of decision making. *J Behav Med.* 2002 Aug;25(4):355–72. [PubMed]
5. Urquhart J, Vrijens B. New findings about patient adherence to prescribed drug dosing regimes: an introduction to pharmionics. *Eur J Hosp Pharm Sci.* 2005;11(5):103–6.
6. Kreuter MW, Farrell D, Olevitch L, Brenna L. *Tailoring Health Messages: Customizing Communication With Computer Technology.* Mahwah, NJ: Lawrence Erlbaum Associates; 2000.
7. Lustria ML, Cortese J, Noar SM, Glueckauf RL. Computer-tailored health interventions delivered over the Web: review and analysis of key components. *Patient Educ Couns.* 2009 Feb;74(2):156–73. doi: 10.1016/j.pec.2008.08.023.S0738-3991(08)00469-2 [PubMed] [Cross Ref]
8. Noar SM, Benac CN, Harris MS. Does tailoring matter? Meta-analytic review of tailored print health behavior change interventions. *Psychol Bull.* 2007 Jul;133(4):673–93. doi: 10.1037/0033-2909.133.4.673.2007-09203-006 [PubMed] [Cross Ref]
9. Kreuter MW, Wray RJ. Tailored and targeted health communication: strategies for enhancing information relevance. *Am J Health Behav.* 2003;27 Suppl 3:S227–32. [PubMed]
10. Wantland DJ, Portillo CJ, Holzemer WL, Slaughter R, McGhee EM. The effectiveness of Web-based vs. non-Web-based interventions: a meta-analysis of behavioral change outcomes. *J Med Internet Res.* 2004 Nov 10;6(4):e40. doi: 10.2196/jmir.6.4.e40. <http://www.jmir.org/2004/4/e40/v6e40> [PMC free article] [PubMed] [Cross Ref]
11. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions, version 5.1.0.* The Cochrane Collaboration; 2011. Mar, [2011-10-07]. [webcite http://www.cochrane-handbook.org/](http://www.cochrane-handbook.org/)
12. van Tulder M, Furlan A, Bombardier C, Bouter L., Editorial Board of the Cochrane Collaboration Back Review Group Updated method guidelines for systematic reviews in the cochrane collaboration back review group. *Spine (Phila Pa 1976)* 2003 Jun 15;28(12):1290–9. doi: 10.1097/01.BRS.0000065484.95996.AF. [PubMed] [Cross Ref]
13. Sluijs E, van Dulmen S, van Dijk L, Ridder D, Heerink R, Bensing J. *Patient Adherence to Medical Treatment: A Meta Review.* Utrecht: NIVEL; 2006.
14. Wetzels GE, Nelemans PJ, Schouten JS, van Wijk BL, Prins MH. All that glisters is not gold: a comparison of electronic monitoring versus filled prescriptions--an observational study. *BMC Health Serv Res.* 2006;6:8. doi: 10.1186/1472-6963-6-8. <http://www.biomedcentral.com/1472-6963/6/8.1472-6963-6-8> [PMC free article] [PubMed] [Cross Ref]
15. Farmer KC. Methods for measuring and monitoring medication regimen adherence in clinical trials and clinical practice. *Clin Ther.* 1999 Jun;21(6):1074–90; discussion 1073. doi: 10.1016/S0149-2918(99)80026-5.S0149-2918(99)80026-5 [PubMed] [Cross Ref]
16. Velligan DI, Wang M, Diamond P, Glahn DC, Castillo D, Bendle S, Lam YW, Ereshefsky L, Miller AL. Relationships among subjective and objective measures of adherence to oral antipsychotic medications. *Psychiatr Serv.* 2007 Sep;58(9):1187–92. doi: 10.1176/appi.ps.58.9.1187. <http://ps.psychiatryonline.org/cgi/pmidlookup?view=long&pmid=17766564.58/9/1187> [PubMed] [Cross Ref]

17. Pullar T, Kumar S, Tindall H, Feely M. Time to stop counting the tablets? *Clin Pharmacol Ther.* 1989 Aug;46(2):163–8. 0009-9236(89)90355-X [[PubMed](#)]
18. Rudd P, Byyny RL, Zachary V, LoVerde ME, Titus C, Mitchell WD, Marshall G. The natural history of medication compliance in a drug trial: limitations of pill counts. *Clin Pharmacol Ther.* 1989 Aug;46(2):169–76. 0009-9236(89)90356-1 [[PubMed](#)]
19. Shi L, Liu J, Fonseca V, Walker P, Kalsekar A, Pawaskar M. Correlation between adherence rates measured by MEMS and self-reported questionnaires: a meta-analysis. *Health Qual Life Outcomes.* 2010;8:99. doi: 10.1186/1477-7525-8-99. <http://www.hqlo.com/content/8/99.1477-7525-8-99> [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
20. Shi L, Liu J, Koleva Y, Fonseca V, Kalsekar A, Pawaskar M. Concordance of adherence measurement using self-reported adherence questionnaires and medication monitoring devices. *Pharmacoeconomics.* 2010;28(12):1097–107. doi: 10.2165/11537400-000000000-00000.5 [[PubMed](#)] [[Cross Ref](#)]
21. Grymonpre RE, Didur CD, Montgomery PR, Sitar DS. Pill count, self-report, and pharmacy claims data to measure medication adherence in the elderly. *Ann Pharmacother.* 1998;32(7-8):749–54. [[PubMed](#)]
22. DiMatteo MR, Giordani PJ, Lepper HS, Croghan TW. Patient adherence and medical treatment outcomes: a meta-analysis. *Med Care.* 2002 Sep;40(9):794–811. doi: 10.1097/01.MLR.0000024612.61915.2D. [[PubMed](#)] [[Cross Ref](#)]
23. Sajatovic M, Velligan DI, Weiden PJ, Valenstein MA, Ogedegbe G. Measurement of psychiatric treatment adherence. *J Psychosom Res.* 2010 Dec;69(6):591–9. doi: 10.1016/j.jpsychores.2009.05.007.S0022-3999(09)00186-X [[PubMed](#)] [[Cross Ref](#)]
24. McMahon JH, Jordan MR, Kelley K, Bertagnolio S, Hong SY, Wanke CA, Lewin SR, Elliott JH. Pharmacy adherence measures to assess adherence to antiretroviral therapy: review of the literature and implications for treatment monitoring. *Clin Infect Dis.* 2011 Feb 15;52(4):493–506. doi: 10.1093/cid/ciq167.ciq167 [[PMC free article](#)] [[PubMed](#)] [[Cross Ref](#)]
25. Steultjens EMJ, Dekker J, Bouter LM, van de Nes JCM, Cardol M, van den Ende CHM. *Occupational Therapy for Multiple Sclerosis (Protocol for a Cochrane Review)* Oxford: Update Software; 2002.
26. Verkaik R, van Weert JC, Francke AL. The effects of psychosocial methods on depressed, aggressive and apathetic behaviors of people with dementia: a systematic review. *Int J Geriatr Psychiatry.* 2005 Apr;20(4):301–14. doi: 10.1002/gps.1279. [[PubMed](#)] [[Cross Ref](#)]
27. Artinian NT, Harden JK, Kronenberg MW, Vander Wal JS, Daher E, Stephens Q, Bazzi RI. Pilot study of a Web-based compliance monitoring device for patients with congestive heart failure. *Heart Lung.* 2003 Aug;32(4):226–33.S0147956303000268 [[PubMed](#)]
28. Jan RL, Wang JY, Huang MC, Tseng SM, Su HJ, Liu LF. An internet-based interactive telemonitoring system for improving childhood asthma outcomes in Taiwan. *Telemed J E Health.* 2007 Jun;13(3):257–68. doi: 10.1089/tmj.2006.0053. [[PubMed](#)] [[Cross Ref](#)]
29. Chan DS, Callahan CW, Hatch-Pigott VB, Lawless A, Proffitt HL, Manning NE, Schweikert M,

- Malone FJ. Internet-based home monitoring and education of children with asthma is comparable to ideal office-based care: results of a 1-year asthma in-home monitoring trial. *Pediatrics*. 2007 Mar;119(3):569–78. doi: 10.1542/peds.2006-1884. <http://pediatrics.aappublications.org/cgi/pmidlookup?view=long&pmid=17332210>.119/3/569 [PubMed] [Cross Ref]
30. Chan DS, Callahan CW, Sheets SJ, Moreno CN, Malone FJ. An Internet-based store-and-forward video home telehealth system for improving asthma outcomes in children. *Am J Health Syst Pharm*. 2003 Oct 1;60(19):1976–81. [PubMed]
31. Joseph CL, Peterson E, Havstad S, Johnson CC, Hoerauf S, Stringer S, Gibson-Scipio W, Ownby DR, Elston-Lafata J, Pallonen U, Strecher V., Asthma in Adolescents Research Team A web-based, tailored asthma management program for urban African-American high school students. *Am J Respir Crit Care Med*. 2007 May 1;175(9):888–95. doi: 10.1164/rccm.200608-1244OC. <http://ajrccm.atsjournals.org/cgi/pmidlookup?view=long&pmid=17290041>.200608-1244OC [PMC free article] [PubMed] [Cross Ref]
32. Ross SE, Moore LA, Earnest MA, Wittevrongel L, Lin CT. Providing a web-based online medical record with electronic communication capabilities to patients with congestive heart failure: randomized trial. *J Med Internet Res*. 2004 May 14;6(2):e12. doi: 10.2196/jmir.6.2.e12. <http://www.jmir.org/2004/2/e12/v6e12> [PMC free article] [PubMed] [Cross Ref]
33. Cherry JC, Moffatt TP, Rodriguez C, Dryden K. Diabetes disease management program for an indigent population empowered by telemedicine technology. *Diabetes Technol Ther*. 2002;4(6):783–91. [PubMed]
34. Guendelman S, Meade K, Benson M, Chen YQ, Samuels S. Improving asthma outcomes and self-management behaviors of inner-city children: a randomized trial of the Health Buddy interactive device and an asthma diary. *Arch Pediatr Adolesc Med*. 2002 Feb;156(2):114–20. <http://archpedi.ama-assn.org/cgi/pmidlookup?view=long&pmid=11814370>.poa10257 [PubMed]
35. DeVito Dabbs A, Dew MA, Myers B, Begey A, Hawkins R, Ren D, Dunbar-Jacob J, Oconnell E, McCurry KR. Evaluation of a hand-held, computer-based intervention to promote early self-care behaviors after lung transplant. *Clin Transplant*. 2009;23(4):537–45. doi: 10.1111/j.1399-0012.2009.00992.x.CTR992 [PMC free article] [PubMed] [Cross Ref]
36. van der Meer V, Bakker MJ, van den Hout WB, Rabe KF, Sterk PJ, Kievit J, Assendelft WJ, Sont JK., SMASHING (Self-Management in Asthma Supported by Hospitals&sbquo; ICT&sbquo; Nurses and General Practitioners) Study Group Internet-based self-management plus education compared with usual care in asthma: a randomized trial. *Ann Intern Med*. 2009 Jul 21;151(2):110–20.151/2/110 [PubMed]
37. van der Meer V, van Stel HF, Bakker MJ, Roldaan AC, Assendelft WJ, Sterk PJ, Rabe KF, Sont JK., SMASHING (Self-Management of Asthma Supported by Hospitals&sbquo; ICT&sbquo; Nurses and General practitioners) Study Group Weekly self-monitoring and treatment adjustment benefit patients with partly controlled and uncontrolled asthma: an analysis of the SMASHING study. *Respir Res*. 2010;11:74. doi: 10.1186/1465-9921-11-74. <http://respiratory-research.com/content/11//74>.1465-9921-11-74 [PMC free article] [PubMed] [Cross Ref]
38. Dilorio C, Escoffery C, McCarty F, Yeager KA, Henry TR, Koganti A, Reisinger EL, Wexler B.

Evaluation of WebEase: an epilepsy self-management Web site. *Health Educ Res.* 2009 Apr;24(2):185–97. doi: 10.1093/her/cyn012. <http://her.oxfordjournals.org/cgi/pmidlookup?view=long&pmid=18682382.cyn012> [[PubMed](#)] [[Cross Ref](#)]

39. Dew MA, Goycoolea JM, Harris RC, Lee A, Zomak R, Dunbar-Jacob J, Rotondi A, Griffith BP, Kormos RL. An internet-based intervention to improve psychosocial outcomes in heart transplant recipients and family caregivers: development and evaluation. *J Heart Lung Transplant.* 2004 Jun;23(6):745–58. [[PubMed](#)]

40. Nieuwkerk PT, Oort FJ. Self-reported adherence to antiretroviral therapy for HIV-1 infection and virologic treatment response: a meta-analysis. *J Acquir Immune Defic Syndr.* 2005 Apr 1;38(4):445–8.00126334-200504010-00010 [[PubMed](#)]

41. Hugen PW, Langebeek N, Burger DM, Zomer B, van Leusen R, Schuurman R, Koopmans PP, Hekster YA. Assessment of adherence to HIV protease inhibitors: comparison and combination of various methods, including MEMS (electronic monitoring), patient and nurse report, and therapeutic drug monitoring. *J Acquir Immune Defic Syndr.* 2002 Jul 1;30(3):324–34. [[PubMed](#)]

42. Nieuwkerk PT, de Boer-van der Kolk IM, Prins JM, Locadia M, Sprangers MA. Self-reported adherence is more predictive of virological treatment response among patients with a lower tendency towards socially desirable responding. *Antivir Ther.* 2010;15(6):913–6. doi: 10.3851/IMP1644. [[PubMed](#)] [[Cross Ref](#)]

43. Drotar D, Richard KA, Burgess E, Levi R, Nobile C, Seti A, Walders N. Adherence to treatment in childhood chronic illness: issues and recommendations to enhance pediatric research and training. In: Drotar D, editor. *Promoting Adherence to Treatment in Childhood Chronic Illness: Concepts, Methods and Interventions.* Mahwah, NJ: Erlbaum; 2002.

44. Reynolds NR. Adherence to antiretroviral therapies: state of the science. *Curr HIV Res.* 2004 Jul;2(3):207–14. [[PubMed](#)]

Articles from Journal of Medical Internet Research are provided here courtesy of **Gunther Eysenbach**

You are here: [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#)

[Write to the Help Desk](#)

#### GETTING STARTED

[NCBI Education](#)  
[NCBI Help Manual](#)  
[NCBI Handbook](#)  
[Training & Tutorials](#)

#### RESOURCES

[Chemicals & Bioassays](#)  
[Data & Software](#)  
[DNA & RNA](#)  
[Domains & Structures](#)  
[Genes & Expression](#)  
[Genetics & Medicine](#)  
[Genomes & Maps](#)  
[Homology](#)  
[Literature](#)  
[Proteins](#)  
[Sequence Analysis](#)  
[Taxonomy](#)  
[Training & Tutorials](#)

#### POPULAR

[PubMed](#)  
[Nucleotide](#)  
[BLAST](#)  
[PubMed Central](#)  
[Gene](#)  
[Bookshelf](#)  
[Protein](#)  
[OMIM](#)  
[Genome](#)  
[SNP](#)  
[Structure](#)

#### FEATURED

[Genetic Testing Registry](#)  
[PubMed Health](#)  
[GenBank](#)  
[Reference Sequences](#)  
[Map Viewer](#)  
[Human Genome](#)  
[Mouse Genome](#)  
[Influenza Virus](#)  
[Primer-BLAST](#)  
[Sequence Read Archive](#)

#### NCBI INFORMATION

[About NCBI](#)  
[Research at NCBI](#)  
[NCBI Newsletter](#)  
[NCBI FTP Site](#)  
[NCBI on Facebook](#)  
[NCBI on Twitter](#)  
[NCBI on YouTube](#)