



## UvA-DARE (Digital Academic Repository)

### The optical and X-ray decline of V 0332 + 53 (X 0331 + 53, BQ Cam)

Corbet, R.H.D.; Charles, P.A.; van der Klis, M.

**Publication date**  
1986

**Published in**  
Astronomy & Astrophysics

[Link to publication](#)

**Citation for published version (APA):**

Corbet, R. H. D., Charles, P. A., & van der Klis, M. (1986). The optical and X-ray decline of V 0332 + 53 (X 0331 + 53, BQ Cam). *Astronomy & Astrophysics*, 162(1-2), 117-120.

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# The optical and X-ray decline of V 0332 + 53 (X 0331 + 53, BQ Cam)

R.H.D. Corbet<sup>1</sup>, P.A. Charles<sup>1</sup>, and M. van der Klis<sup>2</sup>

<sup>1</sup> Department of Astrophysics, University of Oxford, South Parks Road, Oxford OX1 3RQ, UK

<sup>2</sup> Space Science Department/ESA, ESTEC, Postbus 299, 2900 AG Noordwijk, The Netherlands

Received October, 29, 1985; accepted February 14, 1986

**Summary.** We present optical spectra of the X-ray transient V0332 + 53 (BQ Cam). The equivalent width of the H $\alpha$  emission line in the spectrum has declined by a factor  $\sim 2$  compared with previous observations. This is interpreted as showing that V0332 + 53 is surrounded by an evolving circumstellar envelope. EXOSAT observations failed to detect X-ray emission from V0332 + 53 showing that the H $\alpha$  emission decline has been accompanied by a cessation of the X-ray flaring previously observed. From measurements of optical I.S. bands we derive a reddening of  $E(B-V) = 1.88 \pm 0.10$ . We argue that the optical counterpart of V0332 + 53 is a Be star viewed at a low inclination angle.

**Key words:** X-ray binaries – Be stars

## 1. Introduction

The transient X-ray source V0332 + 53 was discovered by Terrell and Priedhorsky (1984) with the Vela 5B satellite. It was rediscovered by Tanaka et al. (1983) and subsequent EXOSAT observations (Stella et al., 1985a, hereafter SEA) showed V0332 + 53 to be a 4.4s X-ray pulsator with an eccentric ( $e = 0.31$ ) 34.25 day orbit. X-ray outbursts occur close to periastron passage. The pulsations show the presence of an accreting rotating magnetic neutron star in the system. An optical counterpart was found (Argyle, 1983; Kodaira, 1983; Honeycutt and Schlegel, 1983) which was thought to be a reddened B star. Bernacca et al. (1984, hereafter BIS) obtained low dispersion (250 Å/mm) optical spectroscopy of this star and reported H $\alpha$  emission with, at least at times, a P-Cygni profile. Further, higher resolution, spectroscopy was obtained by Stocke et al. (1985) which showed broad H and He emission lines but without P-Cygni profiles. The lack of photospheric absorption and radial velocity variations led Stocke et al. to conclude that the optical continuum does not arise close to the primary star. Spectrophotometry of V0332 + 53 (2.5 Å resolution) has been obtained by Honeycutt and Schlegel (1985, hereafter HS). Kodaira et al. (1985) also report spectroscopic and photometric observations.

In this paper we present medium dispersion (50–130 Å/mm) optical spectroscopy of the counterpart of V0332 + 53. We compare our spectra with previous data and relate optical changes to the cessation of the previously observed X-ray flaring behaviour.

Send offprint requests to: R.H.D. Corbet

We use our measurements of optical I.S. bands to estimate the reddening to V0332 + 53.

## 2. Observations

### 2.1. Optical observations

Spectra of V0332 + 53 were obtained at the 2.5 m Isaac Newton Telescope using the Intermediate Dispersion Spectrograph on the three occasions detailed in Table 1. Details of the spectra obtained are shown in Figs. 1 and 2. The detectors employed were the Imaging Photon Counting System (IPCS) and a GEC CCD. Channel to channel variability in detector response was corrected using data on a calibration lamp obtained at the end of each night. Wavelength calibration was from arc lamp exposures. The overall instrumental response of the IPCS was removed through observations of the flux standard HZ15 (Stone, 1977).

Emission is present at H $\alpha$  in the two spectra of V0332 + 53 covering this region. In no case do we see P-Cygni features or absorption components. Very weak emission may be present at H $\beta$  but the signal-to-noise level prevents a definite identification. Interstellar lines due to Na I plus diffuse I.S. bands are also present. Measurements of the H $\alpha$  line were made by fitting a Gaussian profile plus a quadratic continuum. This procedure produced good fits to the data. We measure a heliocentric velocity ( $v_H$ ) of  $-68 \pm 8 \text{ km s}^{-1}$ , E. W. =  $5.9 \pm 0.6 \text{ \AA}$  and a FWHM of  $250 \pm 20 \text{ km s}^{-1}$  for the July spectrum (resolution =  $97 \text{ km s}^{-1}$ ). For the lower resolution ( $228 \text{ km s}^{-1}$ ) Nov. spectrum we measure  $v_H = -81 \pm 17 \text{ km s}^{-1}$ , E. W. =  $4.6 \pm 0.4 \text{ \AA}$  and a FWHM of  $302 \pm 22 \text{ km s}^{-1}$ . The quoted FWHM have been corrected for the instrumental resolution using measurements of the calibration arc lines. These widths are consistent with the BIS  $300 \text{ km s}^{-1}$  upper limit. The blue end of the IPCS spectrum is relatively noisy but H $\gamma$  absorption may be present (E. W. =  $1.0 \pm 0.37 \text{ \AA}$ ,  $v_H = -25 \pm 40 \text{ km s}^{-1}$ ). He I  $\lambda 4387$  may be weakly present (E. W. =  $0.64 \pm 0.26 \text{ \AA}$ ), but the lack of any visible line at He I  $\lambda 4471$  argues against this identification.


### 2.2. X-ray observations

X-ray observations of V0332 + 53 were made using the ME experiment on the EXOSAT observatory (Taylor et al., 1981) on the three occasions detailed in Table 2. Although all observations were made close to times of predicted X-ray maxima no emission

**Table 1.** Optical spectroscopy of V0332+53

Date	J.D. (-2440000)	Phase	Detector	Exposure time (s)	Spectral range (Å)	Dispersion (Å/mm)	Channel Size (Å)
1984 July 8.2	5889.7	6.94	CCD	1000	6300–7000	50	1.1
1984 Nov. 29.0	6033.5	11.14	IPCS	6150	4200–6100	67	1.0
1984 Nov. 30.1	6034.6	11.17	CCD	1800	5800–7400	131	2.9

Notes: Phases are taken from the ephemeris of Stella et al. (1985a) and are all  $\pm \sim 0.05$



**Fig. 1.** 1984 July 8 spectrum of V0332+53 obtained with the INT Intermediate Dispersion Spectrograph with CCD showing detail of H $\alpha$ . The horizontal bar indicates the FWHM instrumental resolution

was seen on any pointing with an upper limit of  $\sim 1 \mu\text{Jy}$  in all cases (2–10 keV,  $> 3\sigma$  confidence).



### 3. Interstellar reddening

The I.S. bands seen in our spectra and their implied values for  $E(B-V)$  are listed in Table 3. Excluding the 6284 Å band, to avoid problems due to blending with an O<sub>2</sub> band, we adopt  $E(B-V) = 1.88 \pm 0.10$  which implies  $A_v = 6.2$  (for  $R = 3.3$ , Herbst 1975).

This reddening implies an X-ray H column density of  $1.3 \pm 0.3 \cdot 10^{22} \text{ cm}^{-2}$  (Ryter et al., 1975). SEA measured  $N_{\text{H}}$  to be between  $0.6 \pm 0.2$  and  $1.5 \pm 0.1 \cdot 10^{22} \text{ cm}^{-2}$  which corresponds to  $E(B-V) = 0.9$  and  $2.2$  respectively (Ryter et al.). The absorption seen in the X-ray band is due to a combination of I.S. and variable material local to the X-ray source. The *lowest*  $N_{\text{H}}$  seen is thus an *upper limit* to the I.S. absorption. The lowest X-ray  $N_{\text{H}}$  value is thus lower than expected at  $\sim 2\sigma$  significance which may indicate that there is an additional soft component in the X-ray spectrum. Alternatively there may be an unusual gas to dust ratio in the direction of V0332+53. This might indicate the presence of dust local to V0332+53.

### 4. Is the optical counterpart of V0332+53 a Be star?

If the primary is an early type star H $\alpha$  emission implies a supergiant or Be classification. The orbital parameters imply that an OB star



**Fig. 2a and b.** Details of 1984 November 28 spectrum of V0332+53 using the IPCS. The flux calibration is less accurate beyond  $\sim 5950 \text{ Å}$

would underfill its Roche lobe and the X-ray pulsation period is much shorter than those of neutron stars accreting from the winds of supergiants (see e.g. Corbet, 1985). The orbital and pulsation periods of V0332+53 are typical, however, of those of Be star systems. The other X-ray properties of V0332+53 are also similar to those of other Be star systems: periodic flaring, long term on/off states and a very hard spectrum (Terrell and Priedhorsky, 1984). The optical counterpart of V0332+53 is thus almost certainly a Be star.

SEA derive an X-ray mass function of  $\sim 0.1 M_{\odot}$ . Assuming a neutron star mass of  $1.4 M_{\odot}$  and a Be star mass of between 10 and  $20 M_{\odot}$  an inclination angle ( $i$ ) of  $10$  to  $14^{\circ}$  is implied. In Struve's (1931) Be star model emission lines originate in an equatorial envelope about a rapidly rotating B star. Be stars viewed at high  $i$

**Table 2.** X-ray observations of V0332+53

Date	J.D. (−2440000)	Phase	Detector	Exposure time (s)
1984 Sep. 14.2	5957.5	8.92	EXOSAT ME	12000
1984 Sep. 17.4	5961.4	9.03	EXOSAT ME	10800
1984 Oct. 18.6	5992.1	9.93	EXOSAT ME	9000

**Table 3.** Interstellar bands in V0332+53 spectrum

I.S. band	E. W. (Å)	$E(B-V)$	
4430	3.90	1.8	
5780	1.23	2.0	
		1.6	(S and VB)
5797	0.50	2.1	
6284	3.2	2.75	(B and N)

*Notes:* Calibrations between E. W. and  $E(B-V)$  are from Herbig (1975), except those marked (S and VB) and (B and N) which are from Snell and Vanden Bout (1981) and Bromage and Nandy (1973), respectively. For the 6284 band an approximate correction for blending with a telluric O<sub>2</sub> band has been applied

show double peaked emission lines whilst those viewed at low  $i$  have only a single peak. In a binary system we may expect that the envelope and orbital planes will coincide. The single-peaked H $\alpha$  line in the V0332+53 spectrum is thus consistent with a low  $i$ .

Dachs et al. (1981) show that for Be stars with single H $\alpha$  emission there is a strong correlation between line width and projected stellar rotational velocity. Hence we may use the width of the H $\alpha$  line in the V0332+53 spectrum to constrain  $i$ . We adopt:  $v \cdot \sin(i) = (2(\log_e 2)^{1/2})^{-1} W(\alpha)$  from Dachs et al. (1981) where  $W(\alpha)$  is the FWHM of H $\alpha$ . Taking  $W(\alpha) = 250 \text{ km s}^{-1}$ , from our higher resolution July spectrum, we obtain  $i > 15^\circ$  for  $v < 600 \text{ km/s}$  (early B stars break-up speed, Allen, 1973).

## 5. Variability

Our spectra show several differences from those of Stocke et al. (1985). In 1984 Jan. at  $\phi \sim 0.0$  Stocke et al. measured H $\alpha$  to have E. W. =  $9.0 \pm 0.1 \text{ \AA}$  and a velocity of  $-10 \text{ km s}^{-1}$ . Our H $\alpha$  emission lines are weaker by a factor  $\sim 2$  and are blue-shifted by  $\sim 65 \text{ km s}^{-1}$ . Since all the spectra were obtained at similar phase ( $\sim 0$ ) and Stocke et al. find no evidence for velocity changes on a time scale of days, orbital motion cannot account for the velocity shifts. Kodaira et al. (1985) also report that the velocity of H $\alpha$  decreased by  $\sim 70 \text{ km s}^{-1}$  between 1983 Nov./Dec. and 1984 Feb. The decline in strength of H $\alpha$  is almost certainly connected with the cessation of X-ray flaring. The last date SEA detected X-rays from V0332+53 was 1984 Jan. 23. The probable presence of H $\gamma$  in our spectra is consistent with declining Balmer emission. The apparent radial velocity changes in H $\alpha$  may be related to the “ $V/R$ ” variability seen in those Be stars with double emission peaks in which the ratio of the strengths of the two peaks is variable (see e.g. Underhill and Doazan, 1982, Chap. 11).

The connection between optical line emission and X-ray production probably arises in two ways: a) X-rays ionize

hydrogen which then produces emission lines by radiative recombination and b) the emission lines imply the presence of a circumstellar envelope from which the neutron star accretes to produce X-rays. H $\alpha$  is still in emission when the X-ray flaring has ceased, indicating that an envelope is present. The envelope has presumably decayed, however, such that the neutron star’s Alfvén radius is always greater than its corotation radius. Accretion thus cannot take place (cf. Corbet, 1984; Stella et al., 1985b).

HS give  $B = 17.04$  on 1983 Nov. 27 and 28 ( $\phi \sim 0.4$ ) whereas BIS report  $B = 17.44$  on 1983 Dec. 4 ( $\phi \sim 0.6$ ). This either indicates photometric changes on a timescale of days or calibration differences between the observations. The second explanation appears more likely as HS also give  $B = 17.04$  on 1984 Feb. 21 ( $\phi \sim 0.9$ ). HS measure  $B-V = 1.62 \pm 0.06$  whereas BIS give  $B-V = 2.31 \pm 0.04$ . Assuming the I.S. reddening derived in Sect. 3 the first  $B-V$  value implies a spectral class earlier than B3 whereas the BIS  $B-V$  value implies an F type star, casting doubt upon their measurements (although the presence of a circumstellar shell might redden the colours beyond the interstellar value).

## 6. Distance

V0332+53 lies within the galactic reddening layer for distances  $< 4 \text{ kpc}$  (Allen, 1973) since  $b = -2^\circ$ . The NaD lines may thus be used as an approximate distance indicator. The E. W. of the sum of these lines is  $2.04 \text{ \AA}$  implying a distance  $\sim 2 \text{ kpc}$  (Allen, 1973). The velocity variations of the H $\alpha$  emission line mean that it cannot be used to calculate the distance. The low E. W. of H $\gamma$  implies contamination by an emission component and so it cannot be used either. Most Be stars in X-ray binaries have spectral types between B0 and B2 and (by definition) luminosity classes of III to V. A star within this range is expected to have  $M_v = -2.5$  to  $-4.6$  (Balona and Crampton, 1974). Assuming the value of  $A_v$  derived in Sect. 3 and  $V = 15.4$  (HS) a distance of 2.2–5.8 kpc is implied.

## 7. Conclusion

V0332+53 is very likely a member of the Be/neutron star binary class in which a neutron star accretes from the extended envelope of an early type primary. This envelope is probably evolving on a time-scale of years. If the optical emission further declines a determination of the primary’s spectral type should be possible.

*Acknowledgements.* The INT support astronomers were T. Sniijders and E. Valentijn. The Isaac Newton Telescope on the island of La Palma is operated by the Royal Greenwich Observatory at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. MvdK acknowledges receipt of an ESA fellowship and support by ZWO (ASTRON) during part of this work.

## References

- Allen, C.W.: 1973, *Astrophysical Quantities*, 3rd edn., The Athlone Press, London
- Argyle, R.W.: 1983, IAUC 3897
- Balona, L., Crampton, D.: 1974, *Monthly Notices Roy. Astron. Soc.* **166**, 203
- Bernacca, P.L., Iijima, T., Stagni, R.: 1984 (BIS), *Astron. Astrophys.* **132**, L8
- Bromage, G.E., Nandy, K.: 1973, *Astron. Astrophys.* **26**, 17
- Corbet, R.H.D.: 1984, *Astron. Astrophys.* **141**, 91
- Corbet, R.H.D.: 1985, NATO Advanced Research Workshop on the Evolution of Galactic X-ray Binaries, Reidel, Dordrecht
- Dachs, J., Eichendorf, H., Schleicher, H., Schmidt-Kaler, Th., Stift, M., Tug, H.: 1981, *Astron. Astrophys. Suppl.* **43**, 427
- Herbig, G.H.: 1975, *Astrophys. J.* **196**, 129
- Herbst, W.: 1975, *Astron. J.* **80**, 498
- Honeycutt, R.K., Schlegel, E.M.: 1983, IAUC 3899
- Honeycutt, R.K., Schlegel, E.M.: 1985 (HS), *Publ. Astron. Soc. Pacific*, **97**, 300
- Kodaira, K.: 1983, IAUC 3897
- Kodaira, K., et al.: 1985, *Publ. Astron. Soc. Pacific* **37**, 97
- Ryter, C., Cesarsky, C.J., Audouze, J.: 1975, *Astrophys. J.* **198**, 103
- Snell, R.L., Vanden Bout, P.A.: 1981, *Astrophys. J.* **244**, 844
- Stella, L., White, N.E., Davelaar, J., Parmar, A.N., Blissett, R.J., van der Klis, M.: 1985a (SEA), *Astrophys. J.* **288**, L45
- Stella, L., White, N.E., Rosner, R.: 1985b, *Space Sci. Rev.* **40**, 399
- Stoeckle, J., Silva, D., Black, J.H., Kodaira, K.: 1985, *Publ. Astron. Soc. Pacific* **97**, 126
- Stone, R.P.S.: 1977, *Astrophys. J.* **218**, 767
- Struve, O.: 1931, *Astrophys. J.* **73**, 94
- Tanaka, Y. et al.: 1983, IAUC 3891
- Taylor, B.G., Andresen, R.D., Peacock, A., Zobl, R.: 1981, *Space Sci. Rev.* **30**, 479
- Terrell, J., Priedhorsky, W.C.: 1984, *Astrophys. J.* **285**, L15
- Underhill, A., Doazan, V.: 1982, "B Stars With and Without Emission Lines", NASA, SP-456

**Note added in proof:** After the acceptance of this paper we received a preprint from Iye and Kodaira (P.A.S.P., 1985, **97**, 1186) which also discusses H $\alpha$  variability in V 0332 + 53. They measure an H $\alpha$  E.W. of  $\sim 1\text{\AA}$  in 1984 October and  $\sim 3.5\text{\AA}$  in December. Combined with our measurements this shows that there is variability of a factor  $\sim 5$  in the strength of H $\alpha$  on time scales at least as short as one month.