



UvA-DARE (Digital Academic Repository)

Conformal Symmetry for Black Holes in Four Dimensions and Irrelevant Deformations

Baggio, M.; de Boer, J.; Jottar, J.I.; Mayerson, D.R.

DOI

[10.1007/JHEP04\(2013\)084](https://doi.org/10.1007/JHEP04(2013)084)

Publication date

2013

Document Version

Final published version

Published in

The Journal of High Energy Physics

[Link to publication](#)

Citation for published version (APA):

Baggio, M., de Boer, J., Jottar, J. I., & Mayerson, D. R. (2013). Conformal Symmetry for Black Holes in Four Dimensions and Irrelevant Deformations. *The Journal of High Energy Physics*, 2013(4), 084. [https://doi.org/10.1007/JHEP04\(2013\)084](https://doi.org/10.1007/JHEP04(2013)084)

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Conformal symmetry for black holes in four dimensions and irrelevant deformations

Marco Baggio, Jan de Boer, Juan I. Jottar and Daniel R. Mayerson

*Institute for Theoretical Physics, University of Amsterdam,
Science Park 904, Postbus 94485, 1090 GL Amsterdam, The Netherlands*

E-mail: m.baggio@uva.nl, J.deBoer@uva.nl, J.I.Jottar@uva.nl,
d.r.mayerson@uva.nl

ABSTRACT: It has been argued several times in the past that the structure of the entropy formula for general non-extremal asymptotically flat black holes in four dimensions can be understood in terms of an underlying conformal symmetry. A recent implementation of this idea, carried out by Cvetič and Larsen, involves the replacement of a conformal factor in the original geometry by an alternative conformal factor in such a way that the near-horizon behavior and thermodynamic properties of the black hole remain unchanged, while only the asymptotics or “environment” of the geometry are modified. The solution thus obtained, dubbed “subtracted geometry”, uplifts to an asymptotically $\text{AdS}_3 \times S^2$ black hole in five dimensions, and an AdS/CFT interpretation is then possible. Building on this intuition we show that, at least in the static case, the replacement of the conformal factor can be implemented dynamically by means of an interpolating flow which we construct explicitly. Furthermore, we show that this flow can be understood as the effect of irrelevant perturbations from the point of view of the dual two-dimensional CFT, and we identify the quantum numbers of the operators responsible for the flow. This allows us to address quantitatively the validity of CFT computations for these asymptotically flat black holes and provides a framework to systematically compute corrections to the CFT results.

KEYWORDS: Gauge-gravity correspondence, Black Holes in String Theory, AdS-CFT Correspondence

ARXIV EPRINT: [1210.7695](https://arxiv.org/abs/1210.7695)

Contents

1	Introduction	1
2	The STU model	3
2.1	Static ansatz	4
2.2	Diagonalizing the non-linear system: decoupled modes and a family of static black hole solutions	6
2.3	The “original” and “subtracted” geometries	7
3	Interpreting the flow between the original and subtracted geometries	9
3.1	Linearized analysis	10
3.2	Perturbation theory and determination of the sources	11
3.3	Range of validity of the linear approximation	12
4	Uplifting and AdS/CFT interpretation	14
4.1	The $5d$ Lagrangian and equations of motion	14
4.2	Consistent Kaluza-Klein reduction	15
4.3	Asymptotically AdS ₃ solutions and dual operators	17
4.4	Irrelevant deformation of the CFT	18
4.5	Irrelevant mass scale and range of validity of the CFT description	21
5	Discussion	22
A	Conventions and useful formulae	23
A.1	Hodge duality	23
A.2	Details of the Kaluza-Klein reduction	24
A.3	The general solution in terms of $5d$ and $3d$ fields	26

1 Introduction

Black holes and especially their entropy have always been a source of mystery in physics and, despite decades of effort and remarkable results on this topic, a full understanding of their properties is still lacking. The first microscopic understanding of black hole entropy was achieved for supersymmetric three-charge black holes, where the entropy formula was reproduced using an effective CFT description of the low-energy degrees of freedom [1]. Later developments have expanded upon this description for other extremal and near-extremal black holes [2–4]. The Kerr/CFT program [5–7] has further provided indications for a CFT description of the low-energy physics for (near-)extremal rotating black holes.

Even when one identifies some sort of conformal symmetry in black hole related backgrounds, this does not need to have immediate physical implications, nor does it have to

imply the existence of a CFT whose physics is relevant for the black hole. The connection between the CFT and the black hole can be purely numerical, purely kinematical, approximately dynamical, or exactly dynamical. An example of the latter is the BTZ black hole, whose physics is completely captured by that of a $2d$ CFT. When there is an approximate dynamical equivalence the physical quantities of interest may or may not, depending on the question, be approximately reproduced using CFT computations.¹ In this paper we will mostly be interested in this case.

Unfortunately, a systematic approach to find the approximate dynamical CFT description for the low-energy physics of black holes is still lacking. In general, the low energy near-horizon modes of general black holes (as opposed to (near-)extremal ones) do not seem to decouple from the asymptotics of the geometry, thus effectively preventing one from being able to take a decoupling limit and find an effective CFT description of these modes. Even so, there has been much interest in the fact that the massless wave equation in a (general) black hole background admits a $SL(2, \mathbb{R})$ symmetry when certain terms are removed. The offending terms are indeed small and can be neglected in certain limits (near-extremal, near extreme rotating, low energy [5, 8–11]). However reminiscent of conformal symmetry this approximate $SL(2, \mathbb{R})$ symmetry may be, the terms “breaking” this symmetry are not small for general black holes and thus can not justifiably be ignored. The program of “hidden conformal symmetry” asserts that the conformal symmetry is there after all, but it is spontaneously broken [11].

Despite these qualifications, it is intriguing that the entropy of fairly general asymptotically flat black holes can be cast in a form suggestive of Cardy’s asymptotic growth of states $S = 2\pi\sqrt{c\Delta_R/6} + 2\pi\sqrt{c\Delta_L/6}$, even away from extremality [12, 13]. A recent development by Cvetič and Larsen [14, 15], inspired by the approximate $SL(2, \mathbb{R})$ symmetry of the massless scalar wave equation, provides further evidence for an approximate CFT description of general black holes in four and five dimensions far from extremality. They construct a so-called “subtracted” geometry, where the warp factor of the geometry is modified. Thus, the asymptotics of the black hole are changed from asymptotically flat to asymptotically conical [16], but its thermodynamic properties are left untouched. This subtracted geometry can then intuitively be thought of as “putting the black hole in a box”. In addition to providing an exact $SL(2, \mathbb{R})$ symmetry of the wave equation, the subtracted geometry can be uplifted one dimension higher to a geometry that is locally a product of AdS_3 and a two-sphere. Thus, a $2d$ CFT description of the subtracted black hole is immediately obvious.

What is less obvious is the relation between the subtracted geometry and the original, asymptotically flat one, and in particular how this relation would be visible in the CFT description. Further developments [16] have made some progress in this direction by showing that the subtracted geometry can be obtained as a scaling limit of the original geometry. In this paper, we wish to address this problem and provide further evidence of the $2d$ CFT description of the general asymptotically flat, non-rotating, four-charge black holes in four

¹In the purely kinematical case, it is only the near horizon AdS_2 geometry of an extremal black hole which agrees with the near horizon geometry of a BTZ black hole, and in the numerical case one only has a match between the CFT entropy and the black hole entropy.

dimensions first constructed in [17, 18] (and later extended to the rotating case in [19]). If the $2d$ CFT description of the subtracted geometry is related to some IR limit of the asymptotically flat original geometry, then it is natural to expect this original geometry to be described by the CFT plus some irrelevant deformation.

We first construct a family of black hole solutions with various asymptotics in the STU model in four dimensions, of which both the subtracted geometry and the original geometry are members. With this explicit family of solutions, it is easy to find the linear perturbations of the subtracted geometry which must be turned on to flow to the original geometry. These perturbations can then be uplifted to perturbations of $\text{AdS}_3 \times S^2$, where the standard AdS/CFT dictionary allows us to interpret and translate them as irrelevant deformations of the dual $2d$ CFT. We also explicitly determine the sources for these dimension $(2, 2)$ operators in the CFT: in addition to providing further insight into the CFT description of asymptotically flat black holes, the explicit construction of these irrelevant operators and their sources allows us to quantitatively determine the limit of the validity of the CFT description.

The rest of the paper is organized as follows. In section 2 we review the STU model and construct a four-parameter family of four-charged, non-rotating black holes with different asymptotics but the same thermodynamics, and explicitly identify the original (asymptotically flat) and subtracted geometry as members of this family. Then, in section 3, we perform a linear analysis to determine the perturbations needed to flow from the subtracted to the original geometry. In section 4 we uplift the subtracted geometry and the linear perturbations thereof to $5d$; we then consistently reduce to an effective three-dimensional description to easily identify the irrelevant operators and sources through the standard AdS/CFT dictionary. The determination of the sources gives us a clear criterion for the window in which the effective IR CFT description is valid. Finally, in section 5, we summarize and discuss our findings. Various details of our calculations and useful formulae are collected in the appendix.

2 The STU model

The STU model is a four-dimensional $\mathcal{N} = 2$ supergravity theory coupled to three vector multiplets [20–22]. Its Lagrangian is given by²

$$\begin{aligned} \mathcal{L}_4 = & R \star_4 \mathbb{1} - \frac{1}{2} H_{ij} \star_4 dh^i \wedge dh^j - \frac{3}{2f^2} \star_4 df \wedge df - \frac{f^3}{2} \star_4 F^0 \wedge F^0 \\ & - \frac{1}{2f^2} H_{ij} \star_4 d\chi^i \wedge d\chi^j - \frac{f}{2} H_{ij} \star_4 (F^i + \chi^i F^0) \wedge (F^j + \chi^j F^0) \\ & + \frac{1}{2} C_{ijk} \chi^i F^j \wedge F^k + \frac{1}{2} C_{ijk} \chi^i \chi^j F^0 \wedge F^k + \frac{1}{6} C_{ijk} \chi^i \chi^j \chi^k F^0 \wedge F^0, \end{aligned} \quad (2.1)$$

where the fields f and h^i ($i = 1, 2, 3$) are scalars, χ^i are pseudoscalars, and F^0 and F^i are U(1) gauge field strengths. The metric H_{ij} on the scalar moduli space is diagonal with entries $H_{ii} = (h^i)^{-2}$, and the symbol C_{ijk} is pairwise-symmetric in its indices with $C_{123} = 1$

²We mostly follow the notation and conventions of reference [23], which we found particularly useful.

and zero otherwise. The h^i fields are constrained by the relation $h^1 h^2 h^3 = 1$, which must be solved before taking variations of the action. Our conventions for Hodge duality as well as some useful expressions can be found in appendix A.1.

In the following we shall be concerned with solutions where the pseudoscalars χ^i are set to zero. This is not in general a consistent truncation, inasmuch as the pseudoscalar equations of motion then imply the constraints

$$-f H_{ij} \star_4 F^0 \wedge F^j + \frac{1}{2} C_{ijk} F^j \wedge F^k = 0. \quad (2.2)$$

In order to fulfill these conditions we will consider solutions where F^0 is purely electric and the F^i are purely magnetic. If we restrict to this case, we can write a simpler action from which we can derive the equations of motion, namely

$$S = -\frac{1}{2\kappa^2} \int d^4x \sqrt{|g|} \left[R - \frac{e^{-\eta_0}}{4} F_{\mu\nu}^0 F^{0\mu\nu} - \frac{1}{2} \sum_{i=1}^3 \left(\nabla_\mu \eta_i \nabla^\mu \eta_i + \frac{e^{2\eta_i - \eta_0}}{2} F_{\mu\nu}^i F^{i\mu\nu} \right) \right], \quad (2.3)$$

where $\kappa^2 = 8\pi G_4$ (κ has units of length), and we have introduced the shorthand notation

$$\eta_0 \equiv \eta_1 + \eta_2 + \eta_3. \quad (2.4)$$

The scalar fields η_i ($i = 1, 2, 3$) are related to the scalars in (2.1) through

$$h^i = e^{\frac{1}{3}\eta_0 - \eta_i}, \quad (2.5)$$

$$f = e^{-\frac{1}{3}\eta_0}. \quad (2.6)$$

The corresponding equations of motion read³

$$0 = \nabla_\mu \nabla^\mu \eta_i + \frac{1}{4} \left[e^{-\eta_0} F_{\mu\nu}^0 F^{0\mu\nu} + e^{-\eta_0} \sum_{j=1}^3 (1 - 2\delta_{ij}) e^{2\eta_j} F_{\mu\nu}^j F^{j\mu\nu} \right], \quad (2.7)$$

$$0 = \nabla_\mu \left(e^{-\eta_0} F^{0\mu\nu} \right), \quad (2.8)$$

$$0 = \nabla_\mu \left(e^{-\eta_0 + 2\eta_i} F^{i\mu\nu} \right), \quad (2.9)$$

$$G_{\mu\nu} = \frac{1}{2} \left[\sum_{i=1}^3 \left(\nabla_\mu \eta_i \nabla_\nu \eta_i - \frac{g_{\mu\nu}}{2} \nabla_\lambda \eta_i \nabla^\lambda \eta_i \right) + e^{-\eta_0} \left(F_\mu^\rho F_{\nu\rho}^0 - \frac{g_{\mu\nu}}{4} F_{\lambda\rho}^0 F^{0\lambda\rho} \right) + e^{-\eta_0} \sum_{i=1}^3 e^{2\eta_i} \left(F_\mu^{i\rho} F_{\nu\rho}^i - \frac{g_{\mu\nu}}{4} F_{\lambda\rho}^i F^{i\lambda\rho} \right) \right]. \quad (2.10)$$

2.1 Static ansatz

In the present context we will be interested in static, spherically symmetric black hole backgrounds. As discussed above, in order to fulfill the constraint (2.2) we furthermore

³We note that the purely electric configurations of [15, 16] solve the equations of motion following from the action obtained from (2.3) by dualizing the fields F^i as $F^i \rightarrow -e^{\eta_0 - 2\eta_i} (\star_4 F^i)$.

consider an electric ansatz for F^0 and a magnetic ansatz for the F^i . Explicitly, our ansatz for the metric and matter fields reads

$$ds_4^2 = -\frac{G(r)}{\sqrt{\Delta(r)}} dt^2 + \sqrt{\Delta(r)} \left(\frac{dr^2}{X(r)} + d\theta^2 + \frac{X(r)}{G(r)} \sin^2 \theta d\phi^2 \right) \quad (2.11)$$

$$A^0 = A_t^0(r) dt \quad (2.12)$$

$$A^i = B_i \cos \theta d\phi \quad (2.13)$$

$$\eta_i = \eta_i(r), \quad (2.14)$$

where the constants B^i ($i = 1, 2, 3$) are the magnetic charges. Einstein's equations are easily seen to imply $G(r) = \gamma X(r)$, where $\gamma = \text{const}$, and also $X''(r) = 2$. Hence, without loss of generality we set

$$X(r) = G(r) = r^2 - 2mr. \quad (2.15)$$

Given this ansatz, we first notice that the equation for F^0 implies

$$F_{rt}^0 = q_0 \frac{e^{\eta_0}}{\sqrt{\Delta}}, \quad (2.16)$$

where the constant q_0 is the electric charge (up to normalization). The scalar equations then reduce to

$$0 = \left(r(r-2m)\eta_i' \right)' - \frac{e^{\eta_0}}{2\sqrt{\Delta}} \left[q_0^2 + \sum_{j=1}^3 (2\delta_{ij} - 1) B_j^2 e^{2(\eta_j - \eta_0)} \right]. \quad (2.17)$$

Finally, one notices that the independent information contained in Einstein's equations amounts to one second order and one first order equation. These can be taken to be

$$0 = \frac{\Delta''}{\Delta} - \frac{3}{4} \left(\frac{\Delta'}{\Delta} \right)^2 + (\eta_1')^2 + (\eta_2')^2 + (\eta_3')^2 \quad (2.18)$$

$$0 = \left(\frac{\Delta'}{2\Delta} \right)^2 - \frac{2(r-m)}{r(r-2m)} \frac{\Delta'}{\Delta} + \frac{4}{r(r-2m)} + (\eta_1')^2 + (\eta_2')^2 + (\eta_3')^2 - \frac{e^{\eta_0}}{r(r-2m)\sqrt{\Delta}} \left[q_0^2 + \sum_{i=1}^3 e^{-2(\eta_0 - \eta_i)} B_i^2 \right]. \quad (2.19)$$

The first of these equations is a linear combination of the (t, t) and (ϕ, ϕ) components of Einstein's equations, while the first order constraint is the (r, r) component.

2.2 Diagonalizing the non-linear system: decoupled modes and a family of static black hole solutions

Quite remarkably, it is possible to diagonalize the full non-linear system of equations. To this end we introduce new fields ϕ_0, ϕ_i defined as

$$\phi_0(r) = \frac{1}{2} \log \left(\frac{\Delta(r)}{m^4} \right) - \eta_1(r) - \eta_2(r) - \eta_3(r) \quad (2.20)$$

$$\phi_1(r) = \frac{1}{2} \log \left(\frac{\Delta(r)}{m^4} \right) - \eta_1(r) + \eta_2(r) + \eta_3(r) \quad (2.21)$$

$$\phi_2(r) = \frac{1}{2} \log \left(\frac{\Delta(r)}{m^4} \right) + \eta_1(r) - \eta_2(r) + \eta_3(r) \quad (2.22)$$

$$\phi_3(r) = \frac{1}{2} \log \left(\frac{\Delta(r)}{m^4} \right) + \eta_1(r) + \eta_2(r) - \eta_3(r). \quad (2.23)$$

Taking suitable linear combinations of the scalar and Einstein's equations one finds

$$0 = \left(r(r-2m) \phi_0'(r) \right)' + 2 \left(\frac{q_0^2}{m^2} e^{-\phi_0(r)} - 1 \right) \quad (2.24)$$

$$0 = \left(r(r-2m) \phi_i'(r) \right)' + 2 \left(\frac{B_i^2}{m^2} e^{-\phi_i(r)} - 1 \right). \quad (2.25)$$

Upon solving these decoupled equations one has the solution for the original fields $\Delta(r)$ and $\eta_i(r)$, and the solution for F^0 is then given by (2.16). Hence, we have effectively diagonalized the full non-linear system.

We have obtained general solutions to the decoupled equations (2.24)–(2.25),⁴ each of which depends on two arbitrary integration constants. These generic solutions are not regular at the horizon $r = 2m$, but upon imposing regularity they reduce to

$$\phi_0^{\text{reg}}(r) = \log \left[\frac{q_0^2}{4m^4} \frac{(a_0^2 r + 2m)^2}{1 + a_0^2} \right] \quad (2.26)$$

$$\phi_j^{\text{reg}}(r) = \log \left[\frac{B_j^2}{4m^4} \frac{(a_j^2 r + 2m)^2}{1 + a_j^2} \right], \quad (2.27)$$

where the four independent constants a_0, a_i parameterize a family of static black hole solutions. Close to the horizon, one finds

$$\phi_0^{\text{reg}}(r \rightarrow 2m) = \log \left(\frac{q_0^2}{m^2} (1 + a_0^2) \right) + \mathcal{O}(r - 2m), \quad (2.28)$$

$$\phi_j^{\text{reg}}(r \rightarrow 2m) = \log \left(\frac{B_j^2}{m^2} (1 + a_j^2) \right) + \mathcal{O}(r - 2m). \quad (2.29)$$

⁴The general static solutions of the STU model have been constructed in [24] using different techniques. The hidden conformal symmetry of the Klein-Gordon equation on these backgrounds is studied in [25].

Similarly, in the asymptotic region $r \rightarrow \infty$

$$\phi_0^{\text{reg}}(r \rightarrow \infty) = \begin{cases} \log \frac{r^2}{m^2} + \mathcal{O}(1), & a_0 \neq 0 \\ \log \frac{q_0^2}{m^2}, & a_0 = 0 \end{cases} \quad (2.30)$$

$$\phi_j^{\text{reg}}(r \rightarrow \infty) = \begin{cases} \log \frac{r^2}{m^2} + \mathcal{O}(1), & a_j \neq 0 \\ \log \frac{B_j^2}{m^2}, & a_j = 0 \end{cases} \quad (2.31)$$

Going back to the original fields η_i and Δ , the solution reads

$$\Delta(r) = \frac{\sqrt{q_0^2 B_1^2 B_2^2 B_3^2}}{16m^4} \prod_{I=0}^3 \frac{a_I^2 r + 2m}{\sqrt{1 + a_I^2}} \quad (2.32)$$

$$e^{2\eta_1(r)} = \left| \frac{B_2 B_3}{q_0 B_1} \right| \sqrt{\frac{(1 + a_0^2)(1 + a_1^2)}{(1 + a_2^2)(1 + a_3^2)}} \frac{(a_2^2 r + 2m)(a_3^2 r + 2m)}{(a_0^2 r + 2m)(a_1^2 r + 2m)} \quad (2.33)$$

$$e^{2\eta_2(r)} = \left| \frac{B_1 B_3}{q_0 B_2} \right| \sqrt{\frac{(1 + a_0^2)(1 + a_2^2)}{(1 + a_1^2)(1 + a_3^2)}} \frac{(a_1^2 r + 2m)(a_3^2 r + 2m)}{(a_0^2 r + 2m)(a_2^2 r + 2m)} \quad (2.34)$$

$$e^{2\eta_3(r)} = \left| \frac{B_1 B_2}{q_0 B_3} \right| \sqrt{\frac{(1 + a_0^2)(1 + a_3^2)}{(1 + a_1^2)(1 + a_2^2)}} \frac{(a_1^2 r + 2m)(a_2^2 r + 2m)}{(a_0^2 r + 2m)(a_3^2 r + 2m)}. \quad (2.35)$$

It is worth emphasizing that, depending on how many of the constants a_0, a_i are non-zero, the asymptotic behavior of $\Delta(r)$ in our family of solutions can be of the form $\Delta(r \rightarrow \infty) \sim r^\gamma$, with $\gamma = 0, 1, \dots, 4$. In particular, when $a_0 = a_i = 0$ (i.e. $\gamma = 0$) we obtain an asymptotically $\text{AdS}_2 \times S^2$ black hole solution. When $\gamma \neq 0$, the metric displays a ‘‘Lifshitz-covariance’’ of the form $t \rightarrow \lambda^z t$, $r \rightarrow \lambda^{2\theta/\gamma} r$, $ds^2 \rightarrow \lambda^\theta ds^2$ in the $r \gg 2m$ region, where the dynamical exponent z and the hyperscaling violation exponent⁵ θ are related by $\theta = \left(\frac{\gamma}{\gamma-2}\right)z$. We have checked explicitly that our family of solutions satisfies all the coupled equations of motion. In particular, the first order constraint (2.19) is satisfied identically, and places no restriction on the values of the constants a_0, a_i .

2.3 The ‘‘original’’ and ‘‘subtracted’’ geometries

The family of solutions found in section 2.2 contains as a particular case the solutions dubbed ‘‘original’’ and ‘‘subtracted’’ in [14, 15].⁶ The original solution is given in terms of functions

$$p_I(r) = r + 2m \sinh^2 \delta_I, \quad (2.36)$$

⁵These metrics are in a sense a ‘‘global’’ version of the planar black brane solutions that have been used to model condensed matter systems displaying hyperscaling violation; for some representative works, see [26–28] and references therein.

⁶As we have mentioned our solutions are related to the purely electric solutions of [14, 15] by the duality transformation $F^i \rightarrow -e^{\eta_0 - 2\eta_i} (\star_4 F^i)$. In particular the conformal factor $\Delta(r)$ and the scalars $\eta(r)$ are unaffected by this transformation, and it is in this sense that we use the same terminology to refer to the full solutions.

($I = 0, 1, 2, 3$) and it reads

$$\Delta(r) = \prod_{I=0}^3 p_I(r) \tag{2.37}$$

$$e^{-\eta_i(r)} = p_i(r) \sqrt{\frac{p_0(r)}{p_1(r)p_2(r)p_3(r)}} \tag{2.38}$$

$$F_{rt}^0 = -m \frac{\sinh(2\delta_0)}{p_0(r)^2}. \tag{2.39}$$

We then see that the original geometry is asymptotically flat in the $r \rightarrow \infty$ (i.e. $r \gg 2m$) region. Comparing with our general solution we can easily read off the electric and magnetic charges and the parameters a_I in terms of the δ_I :

$$q_0^{\text{orig}} = -m \sinh(2\delta_0), \quad B_i^{\text{orig}} = m \sinh(2\delta_i), \tag{2.40}$$

$$a_I^{\text{orig}} = \frac{1}{\sinh(\delta_I)}. \tag{2.41}$$

Similarly, the so-called subtracted geometry is given by

$$\Delta(r) = (2m)^3 \left[(\Pi_c^2 - \Pi_s^2) r + 2m \Pi_s^2 \right] \tag{2.42}$$

$$e^{\eta_i(r)} = \frac{1}{\sqrt{\Delta(r)}} \prod_{j \neq i} B_j \tag{2.43}$$

$$F_{tr}^0 = -\frac{16m^4}{\Delta^2(r)} \Pi_c \Pi_s B_1 B_2 B_3, \tag{2.44}$$

where the B_i are the magnetic charges as before, and we can read off the electric charge as $q_0 = -16m^4 (B_1 B_2 B_3)^{-1} \Pi_c \Pi_s$.⁷ This solution is asymptotically conical for $r \rightarrow \infty$ [16]. Comparing with our general solution, we learn that the subtracted geometry has

$$a_0^{\text{subt}} = \sqrt{\frac{\Pi_c^2 - \Pi_s^2}{\Pi_s^2}}, \quad a_i^{\text{subt}} = 0. \tag{2.45}$$

As shown in [14], the thermodynamics of the original and subtracted solutions matches if the parameters Π_c and Π_s are given as follows:

$$\Pi_c = \prod_{I=0}^3 \cosh \delta_I, \quad \Pi_s = \prod_{I=0}^3 \sinh \delta_I. \tag{2.46}$$

As we indicated above, depending on how many of the parameters a_I are zero the (large- r) asymptotic behavior of the conformal factor $\Delta(r)$ changes. While the asymptotically flat original geometry has all $a_I \neq 0$ and $\Delta_{\text{orig}} \sim r^4$ for large r (i.e. $r \gg 2m$), we have shown that the subtracted solution has $a_1 = a_2 = a_3 = 0$ and therefore the conformal

⁷Upon dualizing, we obtain a generalization of the solution presented in [14, 15] where we allow for a set of four independent U(1) charges.




Figure 1. Log plot of $\gamma(r) \equiv \frac{d \log(\Delta)}{d \log r}$ for the general solution (2.32). The bottom red curve with $\gamma(r \gg 2m) = 1$ corresponds to the subtracted geometry ($a_1 = a_2 = a_3 = 0$), while the various curves with $\gamma(r \gg 2m) = 4$ correspond to the original geometry with different values for $a_1 = a_2 = a_3 \equiv 1/\sinh(\delta)$. The different curves have increasingly larger values of δ towards the right, so we see that the original and subtracted geometries agree over a broader range in r as the magnetic charges $B \sim \sinh(2\delta)$ increase.

factor scales linearly $\Delta_{\text{subt}} \sim r$. Figure 1 illustrates how we can smoothly interpolate between the subtracted and original geometries by dialing the parameters a_I . In particular notice that when all $\delta_i \gg 1$ a region emerges where the two solutions match to a very good approximation. It is in this sense that we refer to our family of solutions as an interpolating flow, with the different curves in figure 1 corresponding to different points in the space of couplings on a putative dual field theory. In fact, in this limit (also known as dilute-gas approximation) one can think of the subtracted solution as coming from a decoupling limit of the original solution, as we will discuss in the next sections.

3 Interpreting the flow between the original and subtracted geometries

In the previous section we described a four-parameter family of *exact* static solutions of the STU model that interpolates between the original and subtracted geometries; depending on the choice of parameters, this family also includes geometries with different asymptotic behavior from that of the original geometry. We can thus view the subtracted geometry loosely as an IR endpoint of an RG flow starting from the original geometry.

It is noteworthy that our solution implements explicitly the scaling limit discussed in [16] that extracts the subtracted solution from the original one. In the present section we will interpret this scaling limit as a flow between the original and subtracted geometries, while setting the stage for the AdS/CFT discussion to follow in section 4.

Even though (2.32)–(2.35) (with F^0 given by (2.16)) is an exact solution of the full nonlinear equations, we find it instructive to discuss its linearized version. On the one hand the linearized analysis makes the discussion of regularity at the horizon cleaner, since this is related to a choice of state in the holographic context. Secondly, the sources that one gets by linearizing our family of exact solutions do not necessarily correspond to the sources of irrelevant perturbation theory, as we will discuss in detail below. Lastly, while in generic situations exact solutions to the nonlinear equations are not available, the principles behind the linearized analysis still apply. In particular, we will exhibit the existence of linearized modes of the subtracted geometry that start the flow to the original geometry when their sources are chosen correctly. As we will explicitly show in section 4, upon uplifting the solutions to $5d$, these modes will turn out to be dual to irrelevant operators that deform the conformal field theory dual to the subtracted geometry.

3.1 Linearized analysis

We start our analysis by linearizing the field equations around the subtracted solution. Since the full non-linear equations of motion are diagonalized by the fields ϕ_I in (2.20)–(2.23), we can simply consider

$$\phi_I = \phi_I^{\text{subt}} + \delta\phi_I, \tag{3.1}$$

where the $\delta\phi_I$ will be our linearized perturbations. The linearized equations are then

$$r(r - 2m)\delta\phi_0'' + 2(r - m)\delta\phi_0' - 2\frac{q_0^2 B_1^2 B_2^2 B_3^2}{\Delta^2}\delta\phi_0 = 0, \tag{3.2}$$

$$r(r - 2m)\delta\phi_i'' + 2(r - m)\delta\phi_i' - 2\delta\phi_i = 0. \tag{3.3}$$

The equations for the $\delta\phi_i$'s are particularly simple, and we focus on those first. Changing to a new radial variable $x = \frac{r}{m} - 1$, these equations take the form

$$(1 - x^2)\delta\phi_i''(x) - 2x\delta\phi_i'(x) + 2\delta\phi_i(x) = 0, \tag{3.4}$$

which is a Legendre equation whose general solution is

$$\delta\phi_i(x) = \alpha_i x + \beta_i \left(\frac{x}{2} \log \left(\frac{x+1}{x-1} \right) - 1 \right). \tag{3.5}$$

Of these two solutions, only one is regular at the horizon (which is located at $x = 1$), therefore we must set $\beta_i = 0$, or

$$\delta\phi_i = \alpha_i \left(\frac{r}{m} - 1 \right). \tag{3.6}$$

Using the same variable x , and defining the parameters b and c as

$$b = \frac{\Pi_c^2 - \Pi_s^2}{2\sqrt{2}\Pi_c\Pi_s}, \quad c = \frac{\Pi_c^2 + \Pi_s^2}{2\sqrt{2}\Pi_c\Pi_s}, \tag{3.7}$$

the equation for $\delta\phi_0$ becomes

$$(1 - x^2)\delta\phi_0''(x) - 2x\delta\phi_0'(x) + \frac{1}{(bx + c)^2}\delta\phi_0(x) = 0. \tag{3.8}$$

The solution that is regular at the horizon in this case is given by

$$\delta\phi_0 = \alpha_0 \frac{cx + b}{bx + c} = \alpha_0 \frac{(\Pi_c^2 + \Pi_s^2)r - 2m\Pi_s^2}{(\Pi_c^2 - \Pi_s^2)r + 2m\Pi_s^2}. \quad (3.9)$$

Notice that the condition of regularity will translate into a functional relation between the normalizable and non-normalizable modes in the standard holographic setting.

3.2 Perturbation theory and determination of the sources

In the previous section we showed that the solutions dubbed “original” and “subtracted” fit in a four-parameter family of solutions parametrized by a_I . In particular, recall that we have

$$a_i^{\text{orig}} = \frac{1}{\sinh \delta_i}, \quad a_i^{\text{subt}} = 0, \quad (3.10)$$

while the two a_0 's are both different from zero. In order to go from the subtracted to the original geometry, we need to “turn on” the parameters a_i and change the parameter a_0 . We would like to understand this in terms of a flow that is started by linearized fluctuations around the subtracted background. This suggests that the sources α_I should be directly related to the parameters a_I . However, since we are turning on an irrelevant mode, at each order in perturbation theory higher powers of r will be generated, therefore we need to treat the sources as infinitesimal quantities. It is easy to see that linearizing the general solution ϕ_i around $a_i^2 = 0$, at first order in a_i^2 one gets:

$$\phi_i = \phi_i^{\text{subt}} + a_i^2 \left(\frac{r}{m} - 1 \right) + \dots, \quad (3.11)$$

and we recognize the second term on the right-hand side as being the linearized perturbation of the previous subsection. Notice that the higher order terms do not contain terms linear in r , so the sources obtained by linearizing in a_i^2 are equivalent to the sources that one would obtain by extracting the coefficient of order r in a power-series expansion. Therefore we should identify

$$\alpha_i = (a_i^{\text{orig}})^2 = \frac{1}{\sinh^2 \delta_i}. \quad (3.12)$$

Analogously, we have

$$\phi_0 = \phi_0^{\text{subt}} + (a_0^2 - (a_0^{\text{subt}})^2) \frac{\Pi_s^2 (\Pi_c^2 + \Pi_s^2)r - 2m\Pi_s^2}{\Pi_c^2 (\Pi_c^2 - \Pi_s^2)r + 2m\Pi_s^2}, \quad (3.13)$$

and as a consequence

$$\alpha_0 = \left((a_0^{\text{orig}})^2 - (a_0^{\text{subt}})^2 \right) \frac{\Pi_s^2}{\Pi_c^2}. \quad (3.14)$$

Notice however that the sources are in general *not* infinitesimal. They do become infinitesimal in the limit where the three parameters δ_i become very large. In fact in this limit we obtain particularly simple expressions for the sources:

$$\alpha_i \approx 4 e^{-2\delta_i}, \quad (3.15)$$

$$\alpha_0 \approx -4 \sum_i e^{-2\delta_i}. \quad (3.16)$$

Therefore, to leading order we have the relation:

$$\frac{\delta\Delta}{\Delta} = \frac{1}{2} \sum_I \delta\phi_I = 0, \tag{3.17}$$

that is, the metric is not changed to leading order in the parameters $e^{-2\delta_i}$.

Looking at the behavior of the linearized modes for very large r , one is led to the suspicion that the $\delta\phi_i$'s correspond to irrelevant perturbations while $\delta\phi_0$ seems to be associated to a marginal perturbation. In section 4 we will show that this suspicion is correct (after a suitable change of basis), and we will compute the quantum numbers of the operators in the dual CFT₂ that we need to turn on to start the flow to the asymptotically flat original black hole. It is important to notice that these irrelevant perturbations do change the value of the matter fields in the interior, and in particular they are finite (i.e. non zero) at the horizon. This is in contrast to the extremal case, where irrelevant perturbations die off quickly in the interior and do not change the value of the fields at the horizon.

Notice also that since we are turning on irrelevant deformations, there is no intrinsic (i.e. coordinate invariant) way to extract the sources for the dual operators. Their precise definitions must be supplemented with a perturbation scheme to compute higher order corrections. For example, it is easy to see that our choice for the α_i 's is compatible with a scheme where the linear term in r does not receive higher order corrections; however if we used a different radial coordinate, for example $r' = r + c$ where c is a constant, this would not be true anymore. This ambiguity has an analog in quantum field theory, where the question of whether a source of an operator receives quantum corrections or not depends on the renormalization scheme. This ambiguity is obviously not present to leading order in perturbation theory. We will revisit this issue at the end of section 4.3, where we will describe other possible choices for the sources.

3.3 Range of validity of the linear approximation

In many applications, one does not have the exact solutions, and often it is even impossible to solve the linearized equations exactly. In fact, in many interesting situations only the linearized modes in the asymptotic region are available, and a numerical treatment becomes necessary. It is therefore useful to investigate how one could approach this problem from a numerical perspective; we will then be able to compare the numerical results with the analytic results of the previous sections. The first step is to find a region that can be identified with the asymptotic region of the subtracted geometry (that is $\frac{r}{m} \gg 1$) but where the modes that start the flow to the original geometry are still small, so that they can be treated perturbatively. From the discussion in the previous sections, it is clear that this region should be

$$1 < \frac{r}{m} \ll \frac{1}{\alpha}. \tag{3.18}$$

where α is the smallest of the α_i 's. Furthermore, we argued that the α_i 's are related to the parameters δ_i , so that when the latter are large, $\alpha_i \approx 4e^{-2\delta_i}$. As anticipated in the previous section, this is when the three charges B_1 , B_2 , and B_3 are large compared to the fourth, q_0 .




Figure 2. The dashed red line represents the difference between the original and subtracted fields ϕ_1 . The solid black line is a linear function with slope $4e^{-2\delta_1}$. For this plot, we chose $\delta_0 = \delta_1 = \delta_2 = \delta_3 = 15$, $m = 1$, and the domain is $r \in [100, 10^{-4}e^{2\delta_1}]$.




Figure 3. The dashed red line represents the difference between the original and subtracted fields ϕ_0 . The solid black line is the function $4(e^{-2\delta_1} + e^{-2\delta_2} + e^{-2\delta_3}) \frac{(\Pi_c^2 + \Pi_s^2)r - 2m\Pi_s^2}{(\Pi_c^2 - \Pi_s^2)r + 2m\Pi_s^2}$. For this plot, we chose $\delta_0 = \delta_1 = \delta_2 = \delta_3 = 15$, $m = 1$, and the domain is $r \in [100, 10^{-4}e^{2\delta_1}]$.

Since we expect the difference between the original and subtracted solutions for the ϕ_i 's to be linear in this intermediate region, it is possible to determine the sources for these three modes by means of a linear interpolation, as shown in figure 2. The slope of the linear function turns out to be $4e^{-2\delta_i}$, perfectly matching the results of the previous section.

We can also plot ϕ_0 and the function $\alpha_0 \frac{(\Pi_c^2 + \Pi_s^2)r - 2m\Pi_s^2}{(\Pi_c^2 - \Pi_s^2)r + 2m\Pi_s^2}$. We see in figure 3 that the correct source for this mode is $\alpha_0 = -\alpha_1 - \alpha_2 - \alpha_3 = -4(e^{-2\delta_1} + e^{-2\delta_2} + e^{-2\delta_3})$, confirming once again the analysis of the previous section. Before closing this section it is worth emphasizing that, by turning on different combinations of the sources, we can also flow to the various geometries with Lifshitz-like scaling discussed in section 2.2.

4 Uplifting and AdS/CFT interpretation

In this section we uplift our $4d$ solutions to five dimensions, where an AdS/CFT interpretation of the flow is possible. As shown in [14], the subtracted geometry uplifts to a BTZ black hole, which is asymptotically $\text{AdS}_3 \times S^2$. The linearized perturbations of the previous section uplift to linearized perturbations of the BTZ black hole, and we will explicitly show them to be dual to three irrelevant scalar operators with conformal weights $(h, \bar{h}) = (2, 2)$. This allows us to give a more precise description of the dynamical realization of the conformal symmetry for the charged $4d$ black holes under study, while clarifying at the same time the limitations of this program.

4.1 The $5d$ Lagrangian and equations of motion

As shown in [23], the STU model (2.1) can be obtained by dimensional reduction of the following $5d$ Lagrangian:

$$\mathcal{L}_5 = R_5 \star_5 \mathbb{1} - \frac{1}{2} H_{ij} \star_5 dh^i \wedge dh^j - \frac{1}{2} H_{ij} \star_5 \tilde{F}^i \wedge \tilde{F}^j + \frac{1}{6} C_{ijk} \tilde{F}^i \wedge \tilde{F}^j \wedge \tilde{A}^k, \quad (4.1)$$

where H_{ij} and C_{ijk} are defined as in (2.1). The $4d$ and $5d$ line elements are related by

$$ds_5^2 = f^{-1} ds_4^2 + f^2 (dz + A^0)^2, \quad (4.2)$$

and the vector fields by

$$\tilde{A}^i = \chi^i (dz + A^0) + A^i. \quad (4.3)$$

The form of the h^i scalars in our general four-parameter family of solutions is given in (A.47)–(A.49). In particular, uplifting the subtracted solution (2.42)–(2.44) we discover that the $5d$ scalar fields are constant in this case:⁸

$$h_{\text{subt}}^1 = \left(\frac{B_1^2}{B_2 B_3} \right)^{1/3}, \quad h_{\text{subt}}^2 = \left(\frac{B_2^2}{B_1 B_3} \right)^{1/3}, \quad h_{\text{subt}}^3 = \left(\frac{B_3^2}{B_1 B_2} \right)^{1/3}. \quad (4.4)$$

As anticipated, the $5d$ subtracted geometry asymptotes to $\text{AdS}_3 \times S^2$, and this will allow us to interpret the flow we found in the four-dimensional STU theory in terms of deformations of the CFT living on the boundary of the AdS_3 factor. The strategy we will follow consists of performing a consistent Kaluza-Klein reduction of the $5d$ theory on the two-sphere to obtain an effective $(2 + 1)$ -dimensional theory. We will discover that the solutions of this effective theory with constant scalars correspond locally to AdS_3 , and one of them uplifts precisely to the subtracted geometry in five dimensions. Linearizing the theory around this solution will then allow us to identify the dual operators associated with the flow between the original and subtracted geometries.

Before proceeding further we note that the model (4.1) is slightly inconvenient in that the scalar fields satisfy the constraint $h^1 h^2 h^3 = 1$ which must be solved before taking

⁸Without loss of generality, in order to simplify the notation we assume that the magnetic charges satisfy $B_i > 0$ from now on. In the general case, the absolute value of various expressions involving products of the B_i should be considered when appropriate.

variations of the action. Hence, we choose to work instead with unconstrained scalars Ψ and Φ defined through [23]

$$h^1 = e^{\sqrt{\frac{2}{3}}\Psi}, \quad h^2 = e^{-\frac{\Psi}{\sqrt{6}} - \frac{\Phi}{\sqrt{2}}}, \quad h^3 = e^{-\frac{\Psi}{\sqrt{6}} + \frac{\Phi}{\sqrt{2}}}, \quad (4.5)$$

in terms of which

$$\mathcal{L}_5 = R_5 \star_5 \mathbb{1} - \frac{1}{2} \star_5 d\Psi \wedge d\Psi - \frac{1}{2} \star_5 d\Phi \wedge d\Phi - \frac{1}{2} H_{ij}(\Psi, \Phi) \star_5 \tilde{F}^i \wedge \tilde{F}^j + \frac{1}{6} C_{ijk} \tilde{F}^i \wedge \tilde{F}^j \wedge \tilde{A}^k. \quad (4.6)$$

The equations of motion for the matter fields are then

$$0 = d \left(H_{ij} \star_5 \tilde{F}^j \right) - \frac{C_{ijk}}{2} \tilde{F}^j \wedge \tilde{F}^k \quad (4.7)$$

$$0 = d(\star_5 d\Psi) - \frac{1}{2} \frac{\delta H_{ij}}{\delta \Psi} \star_5 \tilde{F}^i \wedge \tilde{F}^j \quad (4.8)$$

$$0 = d(\star_5 d\Phi) - \frac{1}{2} \frac{\delta H_{ij}}{\delta \Phi} \star_5 \tilde{F}^i \wedge \tilde{F}^j. \quad (4.9)$$

Similarly, Einstein's equations read

$$G_{\mu\nu} = \frac{1}{2} \left[\nabla_\mu \Psi \nabla_\nu \Psi - \frac{g_{\mu\nu}}{2} \nabla_\lambda \Psi \nabla^\lambda \Psi + \nabla_\mu \Phi \nabla_\nu \Phi - \frac{g_{\mu\nu}}{2} \nabla_\lambda \Phi \nabla^\lambda \Phi + H_{ij} \left(\tilde{F}_\mu{}^\rho \tilde{F}_{\nu\rho}^j - \frac{g_{\mu\nu}}{4} \tilde{F}_{\lambda\rho}^i \tilde{F}^{j\lambda\rho} \right) \right] \quad (4.10)$$

and we recall that the only non-vanishing components of $H_{ij}(\Psi, \Phi)$ are given by $H_{ii}(\Psi, \Phi) = (h^i(\Psi, \Phi))^{-2}$.

4.2 Consistent Kaluza-Klein reduction

The general structure of the uplifted line element (4.2) is

$$ds_5^2 = e^{\frac{\eta_0}{3}} ds_4^2 + e^{-\frac{2\eta_0}{3}} (dz + A^0)^2 \quad (4.11)$$

$$= e^{\frac{\eta_0}{3}} \sqrt{\Delta(r)} \left(\frac{dr^2}{X(r)} - \frac{G(r)}{\Delta(r)} dt^2 + \frac{e^{-\eta_0}}{\sqrt{\Delta(r)}} (dz + A^0)^2 \right) + e^{\frac{\eta_0}{3}} \sqrt{\Delta(r)} ds^2(S^2) \quad (4.12)$$

where we assume that A^0 has no legs on the sphere directions (i.e. it is purely electric). It is easy to show that the subtracted geometry uplifts to a BTZ \times S^2 black hole. Nevertheless, it is more convenient to take a more general route that will allow us to characterize the general linear perturbations around the uplifted geometry. A Kaluza-Klein (KK) Ansatz that includes all our uplifted solutions is

$$ds_5^2 = ds_{\text{string}}^2(M) + e^{2U(x)} ds^2(Y) \quad (4.13)$$

$$\tilde{F}^i = -B_i \sin \theta d\theta \wedge d\phi \quad (4.14)$$

$$\Psi = \Psi(x) \quad (4.15)$$

$$\Phi = \Phi(x). \quad (4.16)$$

Here, M is the $(2+1)$ -dimensional “external” manifold with coordinates $x = \{r, t, z\}$, and some metric that we keep arbitrary, and Y is the “internal” (compact) manifold, namely the two-sphere with radius ℓ_S and coordinates $y = \{\theta, \phi\}$. We pick the orientation such that the volume form on Y is $\text{vol}_2 = \ell_S^2 \sin \theta d\theta \wedge d\phi$. The subscript “string” in $ds_{\text{string}}^2(M)$ is meant to remind us that the theory that will come out of the reduction will not be immediately in the $(2+1)$ -dimensional Einstein frame, but rather in what could be called string frame. After performing the reduction we will translate the effective theory to Einstein frame before performing the AdS/CFT analysis. The radius of the sphere ℓ_S is set by the equations of motion to be:

$$\ell_S = (B_1 B_2 B_3)^{1/3}. \tag{4.17}$$

Notice however that we have the freedom to rescale U in the reduction. The choice above guarantees that the radius of the reduced $3d$ (locally) AdS₃ metric (in the final $3d$ Einstein frame) is equal to the radius of the (locally) AdS₃ factor in the $5d$ geometry.

The details of the (consistent) KK reduction can be found in appendix A.2. Reducing the $5d$ equations of motion one finds that all reference to the two-sphere drops out, and the resulting three-dimensional equations of motion (A.17), (A.18), (A.41) and (A.42) follow from the effective (string frame) action

$$S_{\text{string}} = -\frac{1}{16\pi G_3} \int d^3x \sqrt{|g|} e^{2U} \left[R + \frac{2}{\ell_S^2} e^{-2U} - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi) + 2(\nabla U)^2 - \frac{1}{2}(\nabla \Psi)^2 - \frac{1}{2}(\nabla \Phi)^2 \right]. \tag{4.18}$$

The three dimensional Newton’s constant G_3 is fixed in terms of the normalization of the $5d$ action and the volume of the internal manifold, which are in turn related to the $4d$ Newton’s constant G_4 :

$$G_3 = \frac{1}{4\pi \ell_S^2} G_5 = \frac{R_z}{2\ell_S^2} G_4. \tag{4.19}$$

Here, R_z is the radius of the circle on which we reduce to go from the $5d$ theory (4.1) to the $4d$ STU model, and is in principle arbitrary. The next step consists in passing to three-dimensional Einstein frame by performing a Weyl rescaling of the metric on M . Denoting with a subscript (E) the quantities in Einstein frame, the transformation we need is

$$ds_{\text{string}}^2(M) = e^{-4U} ds_{(E)}^2(M), \tag{4.20}$$

which in particular implies

$$R = e^{4U} \left[R_{(E)} + 8\Box_{(E)}U - 8(\nabla_{(E)}U)^2 \right]. \tag{4.21}$$

It follows that the Einstein frame effective action is (after dropping a surface term)

$$S_{(E)} = -\frac{1}{16\pi G_3} \int d^3x \sqrt{|g_{(E)}|} \left[R_{(E)} - 6(\nabla_{(E)}U)^2 - \frac{1}{2}(\nabla_{(E)}\Psi)^2 - \frac{1}{2}(\nabla_{(E)}\Phi)^2 + \frac{2}{\ell_S^2} e^{-6U} - \frac{e^{-8U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi) \right]. \tag{4.22}$$

In a slight abuse of notation, we will drop the subscript (E) from now on because we will be working exclusively in Einstein frame. The equations of motion are then

$$G_{\mu\nu} = -\frac{g_{\mu\nu}}{\ell_S^2} \left[-e^{-6U} + \frac{e^{-8U}}{4} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^2} H_{ii}(\Psi, \Phi) \right] + \frac{1}{2} \tilde{T}_{\mu\nu} \quad (4.23)$$

$$0 = \nabla_\mu \nabla^\mu \Psi - \frac{e^{-8U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Psi} \quad (4.24)$$

$$0 = \nabla_\mu \nabla^\mu \Phi - \frac{e^{-8U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Phi} \quad (4.25)$$

$$0 = \nabla_\mu \nabla^\mu U - \frac{e^{-6U}}{\ell_S^2} + \frac{e^{-8U}}{3} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi), \quad (4.26)$$

where we defined the “kinetic” part of the stress tensor as

$$\tilde{T}_{\mu\nu} = \nabla_\mu \Psi \nabla_\nu \Psi - \frac{g_{\mu\nu}}{2} (\nabla \Psi)^2 + \nabla_\mu \Phi \nabla_\nu \Phi - \frac{g_{\mu\nu}}{2} (\nabla \Phi)^2 + 12 \nabla_\mu U \nabla_\nu U - 6 g_{\mu\nu} (\nabla U)^2. \quad (4.27)$$

4.3 Asymptotically AdS₃ solutions and dual operators

We will consider solutions where the scalars take constant values $U = \bar{U}$, $\Psi = \bar{\Psi}$, $\Phi = \bar{\Phi}$, so that $\tilde{T}_{\mu\nu} = 0$. In such a background, the equations (4.23)–(4.26) reduce to

$$G_{\mu\nu} = -\Lambda_{\text{eff}} g_{\mu\nu} \quad (4.28)$$

$$0 = \sum_{i=1}^3 B_i^2 \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Psi} \Big|_{\bar{\Psi}, \bar{\Phi}} \quad (4.29)$$

$$0 = \sum_{i=1}^3 B_i^2 \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Phi} \Big|_{\bar{\Psi}, \bar{\Phi}} \quad (4.30)$$

$$e^{2\bar{U}} = \frac{1}{3} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^2} H_{ii}(\bar{\Psi}, \bar{\Phi}), \quad (4.31)$$

where the effective cosmological constant Λ_{eff} is given by

$$\Lambda_{\text{eff}} = \frac{1}{\ell_S^2} \left[-e^{-6\bar{U}} + \frac{e^{-8\bar{U}}}{4} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^2} H_{ii}(\bar{\Psi}, \bar{\Phi}) \right] = -\frac{e^{-6\bar{U}}}{4\ell_S^2}, \quad (4.32)$$

and we used (4.31) in the last equality. In three dimensions, the only solutions to Einstein’s equations with negative cosmological constant are locally AdS₃; the effective AdS₃ length L in our case is then given by

$$L^2 = -\frac{1}{\Lambda_{\text{eff}}} = 4 e^{6\bar{U}} \ell_S^2. \quad (4.33)$$

There is in fact a unique solution to equations (4.29)–(4.31) for the scalars, given by

$$e^{\bar{U}} = \left(\frac{B_1 B_2 B_3}{\ell_S^3} \right)^{1/3} = 1, \quad e^{\bar{\Psi}} = \left(\frac{B_1^2}{B_2 B_3} \right)^{1/6}, \quad e^{\bar{\Phi}} = \left(\frac{B_3}{B_2} \right)^{1/2}. \quad (4.34)$$

Notice in particular that $\bar{U} = 0$. Comparing with (4.4)–(4.5), we see that these are precisely the values corresponding to the subtracted geometry.⁹ Moreover, as it follows from (4.28), the metric of this three-dimensional solution is locally AdS₃ with radius

$$L = 2 e^{3\bar{U}} \ell_S = 2 \ell_S = 2 (B_1 B_2 B_3)^{1/3}. \quad (4.35)$$

The Brown-Henneaux [29] central charge corresponding to the $2d$ theory on the boundary of this AdS space is

$$c = \frac{3L}{2G_3} = 12\pi \frac{B_1 B_2 B_3}{G_5}. \quad (4.36)$$

From the presence of the Chern-Simons terms in the $5d$ theory, one expects the charges B_i to be quantized in units of $G_5^{1/3}$, which is basically the five-dimensional Planck length. We will describe the global properties of the solution that corresponds to the subtracted geometry in the following subsection.

We can determine the operator content of the dual field theory from the action (4.22): we have the stress tensor coupling to the massless graviton, and three scalar operators that couple to the boundary values of U , Ψ , Φ . Following the standard AdS/CFT dictionary, in order to compute the conformal dimensions of these operators we need to obtain the masses of the linearized bulk fields around the solution corresponding to the subtracted geometry. Linearizing the equations we find that the fluctuations of the three bulk scalars decouple and in fact satisfy the same equation:

$$0 = \nabla_\mu \nabla^\mu \delta F - \frac{8}{L^2} \delta F, \quad (4.37)$$

where δF stands for any of δU , $\delta \Psi$, $\delta \Phi$. Therefore, the masses are given by

$$m_{\delta U}^2 = m_{\delta \Psi}^2 = m_{\delta \Phi}^2 = \frac{8}{L^2}, \quad (4.38)$$

and according to the standard dictionary we conclude that the three scalar operators in the dual theory are irrelevant, with conformal dimension $\Delta = 4$.

4.4 Irrelevant deformation of the CFT

Finally, we relate the $4d$ modes of section 3, parametrized by the α_I 's, to the linearized modes of the $3d$ theory. The scalars are

$$\delta U = \frac{1}{6m} (\alpha_1 + \alpha_2 + \alpha_3) (r - m) \quad (4.39)$$

$$\delta \Psi = \frac{1}{2\sqrt{6}m} (2\alpha_1 - \alpha_2 - \alpha_3) (r - m) \quad (4.40)$$

$$\delta \Phi = \frac{1}{2\sqrt{2}m} (\alpha_3 - \alpha_2) (r - m), \quad (4.41)$$

⁹For completeness, the explicit form of the $3d$ scalars in our general family of solutions is given in (A.53)–(A.55).

corresponding to non-normalizable modes. To identify the marginal mode, a little work is required. As explained in [15], the uplifted subtracted geometry can be cast in the BTZ form with the change of coordinates (we work in a gauge where $A_0 \rightarrow 0$ as $r \rightarrow \infty$):

$$\rho^2 = \frac{R_z^2}{\ell_S^4} \Delta(r) = \frac{R_z^2}{\ell_S^4} (2m)^3 \left((\Pi_c^2 - \Pi_s^2) r + 2m\Pi_s^2 \right) \quad (4.42)$$

$$t = \frac{R_z}{2\ell_S^4} (2m)^3 (\Pi_c^2 - \Pi_s^2) t_3 \quad (4.43)$$

$$z = -R_z \phi_3, \quad (4.44)$$

so that the metric reads

$$ds^2 = -\frac{(\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)}{L^2 \rho^2} dt_3^2 + \frac{L^2 \rho^2}{(\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)} d\rho^2 + \rho^2 \left(d\phi_3 + \frac{\rho_+ \rho_-}{L \rho^2} dt_3 \right)^2, \quad (4.45)$$

with the position of the inner (ρ_-) and outer (ρ_+) horizons given by

$$\rho_+ = \frac{16m^2 R_z}{L^2} \Pi_c, \quad \rho_- = \frac{16m^2 R_z}{L^2} \Pi_s. \quad (4.46)$$

The left- and right-moving temperatures are then

$$T_L = \frac{\rho_+ + \rho_-}{2\pi L^2} = \frac{8m^2 R_z}{\pi L^4} (\Pi_c + \Pi_s), \quad T_R = \frac{\rho_+ - \rho_-}{2\pi L^2} = \frac{8m^2 R_z}{\pi L^4} (\Pi_c - \Pi_s), \quad (4.47)$$

and the black hole mass, angular momentum, entropy density and temperature are

$$M = \frac{1}{8G_3} \left(\frac{\rho_+^2 + \rho_-^2}{L^2} \right) = \frac{32m^4 R_z^2}{L^6 G_3} (\Pi_c^2 + \Pi_s^2) \quad (4.48)$$

$$J = \frac{1}{8G_3} \left(\frac{2\rho_+ \rho_-}{L} \right) = \frac{64m^4 R_z^2}{L^5 G_3} \Pi_c \Pi_s \quad (4.49)$$

$$S = \frac{(4\pi\rho_+)}{8G_3} = \frac{8\pi m^2 R_z}{L^2 G_3} \Pi_c \quad (4.50)$$

$$T = \frac{2T_L T_R}{T_L + T_R} = \frac{8m^2 R_z}{\pi L^4} \left(\frac{\Pi_c^2 - \Pi_s^2}{\Pi_c} \right). \quad (4.51)$$

Notice that, in terms of CFT variables, the entropy of the original 4d black hole, which by construction equals that of the BTZ black hole, can be written as

$$S = \frac{\pi^2 L}{3} (cT_L + cT_R). \quad (4.52)$$

This agrees with Cardy's entropy formula for a CFT with central charge (4.36) on an circle of length $2\pi L$.

When we uplift the perturbations (3.6) and (3.9), we find that they superficially destroy the BTZ asymptotics. This is due to the fact that all the independent perturbations in the 4d theory involve a change in the metric. However, since the 3d linearized Einstein's equations decoupled from the matter fields, the solution must still be locally AdS₃ and

the uplifted perturbations must correspond to a change in the BTZ parameters up to diffeomorphisms. Indeed, we can perform a linearized diffeomorphism

$$\delta g_{\mu\nu} = 2\nabla_{(\mu}\xi_{\nu)}, \tag{4.53}$$

that brings the metric to the original BTZ form, with the change of parameters:

$$\delta\rho_+ = \left(\alpha_0 + \sum_i \alpha_i\right) \frac{4m^2 R_z}{L^2} \Pi_c \tag{4.54}$$

$$\delta\rho_- = -\left(\alpha_0 + \sum_i \alpha_i\right) \frac{4m^2 R_z}{L^2} \Pi_s. \tag{4.55}$$

Incidentally, this shows that the marginal mode is non-normalizable from the AdS₃ perspective, and that there is a non-trivial change of basis between the independent 4d modes and the 3d modes. We can translate this into a change of mass and angular momentum of the BTZ black hole:

$$\delta M = \delta \left(\frac{\rho_+^2 + \rho_-^2}{8G_3 L^2} \right) = \left(\alpha_0 + \sum_i \alpha_i \right) \frac{16m^4 R_z^2}{G_3 L^6} (\Pi_c^2 - \Pi_s^2) \tag{4.56}$$

$$\delta J = 0. \tag{4.57}$$

Notice that the variations of the physical BTZ parameters vanish when

$$\alpha_0 = -\sum_i \alpha_i. \tag{4.58}$$

As we now explain, we can choose our sources so that they satisfy the relation above, and this corresponds to a scheme where the entropy of the black hole does not change order by order in perturbation theory.

Recall that the precise relation between the parameters α_I and the parameters a_I that describe the family of exact black hole solutions depends on the renormalization scheme, as explained at the end of section 3.2. The choice (3.12)–(3.13) is one possibility, but here we will present an alternative that is more natural from the point of view of AdS/CFT. In quantum field theory one has the freedom to redefine the sources at each order in perturbation theory, so that

$$J = J_0 + \lambda J_1 + \dots + \lambda^n J_n + \dots, \tag{4.59}$$

where λ is the coupling constant. It is customary to choose a scheme where

$$J = J_0, \tag{4.60}$$

i.e. where the source is not renormalized. From the point of view of AdS/CFT, this means that the coefficient of $\rho^{\Delta-d}$ that corresponds to the source of the dual operator does not change at higher order in perturbation theory. This corresponds to the choice

$$\alpha_i = \frac{\Pi_c^2 - \Pi_s^2}{\Pi_c^2 \sinh^2 \delta_i - \Pi_s^2 \cosh^2 \delta_i} \approx 4e^{-2\delta_i}, \tag{4.61}$$

showing once again that the leading contribution is independent of the scheme. It is possible to do the same for the metric mode associated to the dual stress tensor, but in this context it seems more natural to choose a scheme where the $4d$ metric does not change at the horizon order by order in perturbation theory. This yields

$$\alpha_0 = - \sum_{i=1}^3 \alpha_i \approx -4 \sum_{i=1}^3 e^{-2\delta_i}, \tag{4.62}$$

which once again agrees with the previous results to leading order in $e^{-2\delta_i}$. Notice that this choice corresponds to keeping the physical parameters of the BTZ black hole fixed, so that at first order the marginal mode associated to the metric is turned off. We conclude that the flow to the original geometry is started by turning on three irrelevant operators in the dual CFT.

4.5 Irrelevant mass scale and range of validity of the CFT description

Finally, we can determine the mass scale set by the irrelevant deformations, which represents the UV cutoff of the dual field theory. Consider the asymptotic behavior of the field δU :

$$\delta U = \frac{\ell_S^4 \sum_{i=1}^3 \alpha_i}{3R_z^2 (2m)^4 (\Pi_c^2 - \Pi_s^2)} \rho^2 + \dots \tag{4.63}$$

As a consequence, the source of the operator dual to U reads

$$J_U = \frac{L^8 \sum_{i=1}^3 \alpha_i}{48R_z^2 (2m)^4 (\Pi_c^2 - \Pi_s^2)} = \frac{1}{12\pi^2 T_L T_R} \sum_i \alpha_i. \tag{4.64}$$

Recall that the temperature in a CFT sets an infrared cutoff, while the mass scale of the irrelevant deformation sets an ultraviolet cutoff. Equation (4.64) shows that when the α_i 's are of order 1, the infrared cutoff and the ultraviolet cutoff are of the same order, so that there is no regime where the conformal field theory description is meaningful. On the other hand, when the α_i 's become small (or $\delta_i \gg 1$), an energy window appears where perturbation theory on the CFT should be a good description of the system:

$$1 < \frac{E^2}{T_L T_R} \ll \frac{1}{\alpha}. \tag{4.65}$$

This is the CFT analog of the condition (3.18) that we have identified in the $4d$ system.

We can phrase the result above in terms of standard effective field theory. The contributions of irrelevant couplings to a process characterized by an energy scale E are typically suppressed by powers of E/M , where M is the UV cutoff set by the irrelevant couplings. In our case we have

$$M^2 \approx \frac{1}{\alpha} T_L T_R, \tag{4.66}$$

and this should be compared to the IR cutoff of the system, that is the temperature. One way to see this is that contributions from the region $E \sim M$ to thermal expectation values are suppressed by a factor $e^{-\beta M}$; since these contributions cannot be reliably computed in

effective field theory, we require $\beta M \gg 1$. In this sense, M is very large when the α 's are small, opening up a range of energies where effective field theory becomes meaningful. It is precisely in this region that CFT (plus perturbation theory) becomes a good description of the system.

5 Discussion

It has been recently argued that certain questions involving the entropy and thermodynamics of four-dimensional asymptotically flat non-extremal black holes can be elucidated by replacing the original geometry by one with a different conformal factor, dubbed subtracted geometry. The replacement modifies the asymptotics while preserving the near-horizon behavior of the original black hole, in such a way that the role of an underlying conformal symmetry becomes manifest, shedding light on the form of the entropy for black holes away from extremality [14–16]. Building on these works, we have shown that four-dimensional, static, asymptotically flat non-extremal black holes with one electric and three magnetic charges can be connected to their corresponding subtracted geometry by a flow which we have constructed explicitly in the form of an interpolating family of solutions. Upon uplifting the construction to five dimensions the subtracted geometry asymptotes to $\text{AdS}_3 \times S^2$, and an AdS/CFT interpretation of the flow is readily available as the effect of irrelevant perturbations in the conformal field theory dual to the AdS_3 factor. In particular, we have identified the quantum numbers of the deformations responsible for the flow and showed that they correspond to three scalar operators with conformal weights $(h, \bar{h}) = (2, 2)$.

As discussed in detail in section 3 and 4, the mass scale associated to such irrelevant perturbations becomes very large compared to the temperature when the magnetic charges are large. In this limit, it is reasonable to expect that some dynamical questions can be approximately answered by means of perturbation theory in the CFT_2 . At least in the static limit, our construction then puts the procedure followed in [14, 15] on a somewhat more concrete footing. On the other hand, away from this limit the ultraviolet cutoff set by the irrelevant deformations becomes of the same order of the infrared cutoff set by the temperature, and the dual CFT captures an increasingly smaller subset of the dynamics, making the usefulness of such an approach doubtful.

It would be of interest to extend our analysis to include rotating four-dimensional black holes. Even though we do not expect any conceptual difficulties, the rotating case is technically more challenging: in the $5d$ uplifted geometry the two-sphere S^2 is fibered non-trivially over AdS_3 , and the modes that start the flow presumably involve non-trivial harmonics on the sphere. This case will be addressed elsewhere. It would also be very interesting to set up the perturbation scheme in the dual CFT_2 and determine what observables of the $4d$ black hole can be reliably computed in terms of perturbation theory in the irrelevant couplings. Effective field theory makes sense only up to the scale set by the irrelevant deformations. However, our perturbations can be resummed geometrically to all orders, allowing us to go beyond the region where perturbation theory is meaningful and reach the asymptotically flat region. From the field theoretic perspective, it is then natural to wonder whether this allows us to say something about the regime where effec-

tive field theory breaks down. Similar questions can be considered for the black holes with Lifshitz-like asymptotics that can be obtained by turning on only a subset of the irrelevant deformations.

Acknowledgments

It is a pleasure to thank Borun Chowdhury, Geoffrey Compère and Balt van Rees for helpful conversations, and especially Finn Larsen for discussions that helped to motivate this project. We also thank Natalia Pinzani-Fokeeva for collaboration on an early stage of this project. This work is part of the research programme of the Foundation for Fundamental Research on Matter (FOM), which is part of the Netherlands Organization for Scientific Research (NWO).

A Conventions and useful formulae

A.1 Hodge duality

Let ω be a p -form in D -dimensions,

$$\omega = \frac{1}{p!} \omega_{\mu_1 \dots \mu_p} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}. \quad (\text{A.1})$$

We define the action of the Hodge star on the basis of forms as

$$\star(dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p}) = \frac{1}{(D-p)!} \epsilon_{\nu_1 \dots \nu_{D-p}}^{\mu_1 \dots \mu_p} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_{D-p}}, \quad (\text{A.2})$$

where $\epsilon_{\mu_1 \dots \mu_D}$ are the components of the Levi-Civita tensor. Equivalently, in components we find

$$(\star\omega)_{\mu_1 \dots \mu_{D-p}} = \frac{1}{p!} \epsilon_{\mu_1 \dots \mu_{D-p} \nu_1 \dots \nu_p} \omega^{\nu_1 \dots \nu_p}. \quad (\text{A.3})$$

If $\varepsilon_{\mu_1 \dots \mu_D}$ denotes the components of the Levi-Civita *symbol* (a tensor density), we have

$$\epsilon_{\mu_1 \dots \mu_D} = \sqrt{|g|} \varepsilon_{\mu_1 \dots \mu_D} \quad \Leftrightarrow \quad \epsilon^{\mu_1 \dots \mu_D} = \frac{(-1)^t}{\sqrt{|g|}} \varepsilon^{\mu_1 \dots \mu_D} \quad (\text{A.4})$$

where t denotes the number of timelike directions, and we have adopted the convention that the Levi-Civita symbol ε with up or down indices is the same. The volume element is given by

$$\star \mathbb{1} = \sqrt{|g|} d^D x \equiv \text{vol}_D \quad \Rightarrow \quad \star \text{vol}_D = (-1)^t \mathbb{1}. \quad (\text{A.5})$$

A useful observation is that, for any two p -forms A and B ,

$$\star A \wedge B = \star B \wedge A = \frac{1}{p!} A^{\mu_1 \dots \mu_p} B_{\mu_1 \dots \mu_p} \text{vol}_D. \quad (\text{A.6})$$

Similarly, if ϕ is a scalar it follows

$$d\star d\phi = (-1)^{D-1} \nabla_\mu \nabla^\mu \phi \text{vol}_D \quad \Rightarrow \quad \star d\star d\phi = (-1)^{t+D-1} \nabla^\mu \nabla_\mu \phi, \quad (\text{A.7})$$

while a one-form A with field strength $F = dA$ satisfies

$$\star d\star dA = \star d\star F = (-1)^{t+1} \nabla_\nu F^\nu_\lambda dx^\lambda. \quad (\text{A.8})$$

A.2 Details of the Kaluza-Klein reduction

Here we provide further details on the reduction of the 5d theory (4.6) on the two-sphere. We found the techniques outlined in [30] particularly useful for this purpose. As described in the main text, our KK ansatz is

$$ds_5^2 = ds_{\text{string}}^2(M) + e^{2U(x)} ds^2(Y) \quad (\text{A.9})$$

$$\tilde{F}^i = -B_i \sin\theta d\theta \wedge d\phi \quad (\text{A.10})$$

$$\Psi = \Psi(x) \quad (\text{A.11})$$

$$\Phi = \Phi(x), \quad (\text{A.12})$$

where M is the $(2+1)$ -dimensional external manifold with coordinates $x = \{r, t, z\}$ and Y is the two-sphere with radius ℓ_S and coordinates $y = \{\theta, \phi\}$. We pick the orientation such that the volume form on Y is $\text{vol}_2 = \ell_S^2 \sin\theta d\theta \wedge d\phi$.

Because the field strengths \tilde{F}^i are purely magnetic, and proportional to the volume form of the two-sphere, the vector equations (4.7) are satisfied trivially in our ansatz and do not yield lower-dimensional equations of motion. Let us now consider the reduction of the scalar equations (4.8)–(4.9). In order to reduce the coupling of the scalars to the $U(1)$ field strengths it is useful to notice that via (A.6) our ansatz implies

$$\star_5 \tilde{F}^i \wedge \tilde{F}^i = \frac{1}{2!} \tilde{F}^{i\mu\nu} \tilde{F}_{\mu\nu}^i \text{vol}_5 = \frac{B_i^2}{\ell_S^4} e^{-4U(x)} \text{vol}_5 = \frac{B_i^2}{\ell_S^4} e^{-2U(x)} \text{vol}_3 \wedge \text{vol}_2. \quad (\text{A.13})$$

Next, we note that for any one-form A with support in M

$$\star_5 A = e^{2U(x)} \star_3 A \wedge \text{vol}_2. \quad (\text{A.14})$$

In particular, if Ψ is a scalar in M , applying this result to $d\Psi$ we find

$$\star_5 d\Psi = e^{2U(x)} \star_3 d\Psi \wedge \text{vol}_2. \quad (\text{A.15})$$

The decomposition of the scalar Laplacian then follows:

$$d\star_5 d\Psi = e^{2U(x)} \left[d(\star_3 d\Psi) + 2dU(x) \wedge \star_3 d\Psi \right] \wedge \text{vol}_2. \quad (\text{A.16})$$

Plugging this result together with (A.13) into (4.8)–(4.9) we find the effective 3d equations for the scalar fields on M :

$$0 = d(\star_3 d\Psi) + 2dU \wedge \star_3 d\Psi - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}}{\delta \Psi} \text{vol}_3 \quad (\text{A.17})$$

$$0 = d(\star_3 d\Phi) + 2dU \wedge \star_3 d\Phi - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}}{\delta \Phi} \text{vol}_3. \quad (\text{A.18})$$

Equivalently, in component notation we have

$$0 = \nabla_\mu \nabla^\mu \Psi + 2\nabla_\mu U \nabla^\mu \Psi - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Psi} \quad (\text{A.19})$$

$$0 = \nabla_\mu \nabla^\mu \Phi + 2\nabla_\mu U \nabla^\mu \Phi - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} \frac{\delta H_{ii}(\Psi, \Phi)}{\delta \Phi}. \quad (\text{A.20})$$

We now turn our attention to the reduction of the 5d Einstein's equations (4.10). In order to reduce the Ricci tensor, we first study the decomposition of the spin connection and the curvature two-form. Let \hat{e}^M denote the 5d local Lorentz frame, and M, N, \dots denote the flat indices on the 5d manifold. Denoting by a, b, \dots the flat indices on M , and by α, β, \dots the flat indices on the compact manifold Y , our choice of vielbein reads

$$\hat{e}^a = e^a \tag{A.21}$$

$$\hat{e}^\alpha = e^U e^\alpha, \tag{A.22}$$

where e^a and e^α are orthonormal frames for M and Y , respectively. Denoting by ω^a_b the spin connection associated with M and by ω^α_β the spin connection appropriate to Y , solving the torsionless condition for the 5d spin connection $\hat{\omega}$ we find

$$\hat{\omega}^a_b = \omega^a_b \tag{A.23}$$

$$\hat{\omega}^\alpha_\beta = \omega^\alpha_\beta \tag{A.24}$$

$$\hat{\omega}^\alpha_a = P_a e^\alpha, \tag{A.25}$$

where we introduced the shorthand

$$P_a \equiv e^U (\partial_a U). \tag{A.26}$$

It is useful to notice that $\hat{\omega}^a_\alpha \wedge \hat{\omega}^\alpha_b = P^a P_b \eta_{\alpha\beta} e^\alpha \wedge e^\beta = 0$. Next, let Θ denote the curvature two-form. Then, on the 5d manifold we have $\hat{\Theta}^M_N = d\hat{\omega}^M_N + \hat{\omega}^M_P \wedge \hat{\omega}^P_N$. Computing the different components we find

$$\hat{\Theta}^a_b = \Theta^a_b \tag{A.27}$$

$$\hat{\Theta}^\alpha_\beta = \Theta^\alpha_\beta - P_a P^a \eta_{\beta[\gamma} \delta^\alpha_{\sigma]} e^\sigma \wedge e^\gamma \tag{A.28}$$

$$\hat{\Theta}^\alpha_a = \delta^\alpha_\gamma (\nabla_c P_a) e^c \wedge e^\gamma. \tag{A.29}$$

The antisymmetrization symbol $[\dots]$ used above includes a factor of $1/2!$, and ∇_a denotes the connection on M . From these expressions we can identify the non-vanishing components of the Riemann tensor, defined as $\hat{\Theta}^M_N = \frac{1}{2!} \hat{R}^M_{NPQ} \hat{e}^P \wedge \hat{e}^Q$:

$$\hat{R}^a_{bcd} = R^a_{bcd} \tag{A.30}$$

$$\hat{R}^\alpha_{\beta\gamma\delta} = e^{-2U} \mathcal{R}^\alpha_{\beta\gamma\delta} - 2e^{-2U} P_a P^a \delta^\alpha_{[\gamma} \eta_{\delta]\beta} \tag{A.31}$$

$$\hat{R}^\alpha_{a\beta b} = -\delta^\alpha_\beta e^{-U} \nabla_b P_a \tag{A.32}$$

$$\hat{R}^\alpha_{\alpha b\beta} = -\eta_{\alpha\beta} e^{-U} \nabla_b P^\alpha. \tag{A.33}$$

In the above notation R^a_{bcd} are the components of the Riemann tensor of the external manifold M , and $\mathcal{R}^\alpha_{\beta\gamma\delta}$ those of the Riemann tensor of the compact manifold Y . Finally, for the decomposition of the Ricci tensor $\hat{R}_{MN} = \hat{R}^P_{MPN}$ we find

$$\begin{aligned} \hat{R}_{ab} &= R_{ab} - d_Y e^{-U} \nabla_b P_a \\ &= R_{ab} - d_Y (\nabla_b \nabla_a U + \nabla_a U \nabla_b U) \end{aligned} \tag{A.34}$$

$$\begin{aligned} \hat{R}_{\alpha\beta} &= e^{-2U} \mathcal{R}_{\alpha\beta} - (d_Y - 1) e^{-2U} P_a P^a \eta_{\alpha\beta} - e^{-U} \nabla_c P^c \eta_{\alpha\beta} \\ &= e^{-2U} \mathcal{R}_{\alpha\beta} - d_Y (\nabla_a U \nabla^a U) \eta_{\alpha\beta} - (\nabla^a \nabla_a U) \eta_{\alpha\beta} \end{aligned} \tag{A.35}$$

$$\hat{R}_{a\alpha} = 0, \tag{A.36}$$

where d_Y is the dimension of the compact manifold ($d_Y = 2$ in our case). In particular, for the Ricci scalar $\hat{R} = \eta^{MN} \hat{R}_{MN}$ it follows that

$$\hat{R} = R + e^{-2U} \mathcal{R} - 2d_Y \nabla^a \nabla_a U - d_Y (1 + d_Y) \nabla^a U \nabla_a U, \quad (\text{A.37})$$

where R is the scalar curvature on M , and \mathcal{R} that of Y . Since the two-sphere has radius ℓ_S we have $\mathcal{R}_{\alpha\beta} = \eta_{\alpha\beta}/\ell_S^2$. Setting $d_Y = 2$ in the above expressions we find that the only non-vanishing components in our reduction are

$$\hat{R}_{ab} = R_{ab} - 2(\nabla_b \nabla_a U + \nabla_a U \nabla_b U) \quad (\text{A.38})$$

$$\hat{R}_{\alpha\beta} = \left(\frac{e^{-2U}}{\ell_S^2} - \nabla^a \nabla_a U - 2\nabla_a U \nabla^a U \right) \eta_{\alpha\beta}. \quad (\text{A.39})$$

The Ricci scalar is then given by

$$\hat{R} = R + \frac{2}{\ell_S^2} e^{-2U} - 4\nabla^a \nabla_a U - 6\nabla^a U \nabla_a U. \quad (\text{A.40})$$

Using the decomposition of the Ricci tensor, from the components of the 5d Einstein's equations in the directions of the external manifold M we get (using flat indices on M)

$$R_{ab} = 2(\nabla_b \nabla_a U + \nabla_a U \nabla_b U) + \frac{1}{2} \left[\nabla_a \Psi \nabla_b \Psi + \nabla_a \Phi \nabla_b \Phi - \frac{\eta_{ab}}{3} e^{-4U} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi) \right]. \quad (\text{A.41})$$

Similarly, noting that with flat indices $\tilde{F}_\alpha{}^P \tilde{F}_{\beta P}^i = (e^{-4U} B_i^2/\ell_S^4) \eta_{\alpha\beta}$, from the components of the 5d Einstein's equations in the directions of Y we find

$$\nabla^a \nabla_a U + 2\nabla_a U \nabla^a U - \frac{e^{-2U}}{\ell_S^2} + \frac{e^{-4U}}{3} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi) = 0. \quad (\text{A.42})$$

Since all reference to the two-sphere dropped out from the equations of motion, the proposed truncation is consistent. Finally, we point out that the resulting three-dimensional equations of motion (A.17), (A.18), (A.41) and (A.42) can be obtained from the following effective action (in string frame):

$$S_{\text{string}} = -\frac{1}{16\pi G_3} \int d^3x \sqrt{|g|} e^{2U} \left[R + \frac{2}{\ell_S^2} e^{-2U} - \frac{e^{-4U}}{2} \sum_{i=1}^3 \frac{B_i^2}{\ell_S^4} H_{ii}(\Psi, \Phi) + 2(\nabla U)^2 - \frac{1}{2}(\nabla \Psi)^2 - \frac{1}{2}(\nabla \Phi)^2 \right]. \quad (\text{A.43})$$

A.3 The general solution in terms of 5d and 3d fields

In terms of the 4d fields (2.11)–(2.14), our four-parameter family of solutions was given in (2.32)–(2.35) (with F^0 given by (2.16)). The h^i fields appearing in the 5d theory (and

also in the 4d STU model) are related to the diagonal fields (2.20)–(2.23) through

$$h^1 = \exp \left[\frac{1}{6} (2\phi_1 - \phi_2 - \phi_3) \right] \quad (\text{A.44})$$

$$h^2 = \exp \left[\frac{1}{6} (2\phi_2 - \phi_1 - \phi_3) \right] \quad (\text{A.45})$$

$$h^3 = \exp \left[\frac{1}{6} (2\phi_3 - \phi_1 - \phi_2) \right]. \quad (\text{A.46})$$

Hence, in our general family of solutions (2.26)–(2.27) they read

$$h^1(r) = \left(\frac{B_1^2}{|B_2 B_3|} \frac{\sqrt{(1+a_2^2)(1+a_3^2)}}{1+a_1^2} \frac{(a_1^2 r + 2m)^2}{(a_2^2 r + 2m)(a_3^2 r + 2m)} \right)^{1/3} \quad (\text{A.47})$$

$$h^2(r) = \left(\frac{B_2^2}{|B_1 B_3|} \frac{\sqrt{(1+a_1^2)(1+a_3^2)}}{1+a_2^2} \frac{(a_2^2 r + 2m)^2}{(a_1^2 r + 2m)(a_3^2 r + 2m)} \right)^{1/3} \quad (\text{A.48})$$

$$h^3(r) = \left(\frac{B_3^2}{|B_1 B_2|} \frac{\sqrt{(1+a_1^2)(1+a_2^2)}}{1+a_3^2} \frac{(a_3^2 r + 2m)^2}{(a_1^2 r + 2m)(a_2^2 r + 2m)} \right)^{1/3}. \quad (\text{A.49})$$

We recall that the 5d line element is given in terms of the 4d one by (4.11). Similarly, in terms of the decoupled fields the 3d scalars are given by

$$U = \frac{\phi_1 + \phi_2 + \phi_3}{6} + \log \frac{m}{\ell_S} \quad (\text{A.50})$$

$$\Psi = \frac{1}{2\sqrt{6}} (2\phi_1 - \phi_2 - \phi_3) \quad (\text{A.51})$$

$$\Phi = \frac{1}{2\sqrt{2}} (\phi_3 - \phi_2), \quad (\text{A.52})$$

so in our general solution we obtain

$$U(r) = \frac{1}{3} \log \left[\frac{1}{\sqrt{(1+a_1^2)(1+a_2^2)(1+a_3^2)}} \left(\frac{a_1^2}{2m} r + 1 \right) \left(\frac{a_2^2}{2m} r + 1 \right) \left(\frac{a_3^2}{2m} r + 1 \right) \right] \quad (\text{A.53})$$

$$\Psi(r) = \frac{1}{\sqrt{6}} \log \left[\frac{B_1^2}{|B_2 B_3|} \frac{\sqrt{(1+a_2^2)(1+a_3^2)}}{1+a_1^2} \frac{(a_1^2 r + 2m)^2}{(a_2^2 r + 2m)(a_3^2 r + 2m)} \right] \quad (\text{A.54})$$

$$\Phi(r) = \frac{1}{\sqrt{2}} \log \left[\left| \frac{B_3}{B_2} \right| \sqrt{\frac{1+a_2^2}{1+a_3^2}} \left(\frac{a_3^2 r + 2m}{a_2^2 r + 2m} \right) \right]. \quad (\text{A.55})$$

References

- [1] A. Strominger and C. Vafa, *Microscopic origin of the Bekenstein-Hawking entropy*, *Phys. Lett. B* **379** (1996) 99 [[hep-th/9601029](#)] [[INSPIRE](#)].

- [2] J.R. David, G. Mandal and S.R. Wadia, *Microscopic formulation of black holes in string theory*, *Phys. Rept.* **369** (2002) 549 [[hep-th/0203048](#)] [[INSPIRE](#)].
- [3] P. Kraus, *Lectures on black holes and the AdS₃/CFT₂ correspondence*, *Lect. Notes Phys.* **755** (2008) 193 [[hep-th/0609074](#)] [[INSPIRE](#)].
- [4] A. Sen, *Black Hole Entropy Function, Attractors and Precision Counting of Microstates*, *Gen. Rel. Grav.* **40** (2008) 2249 [[arXiv:0708.1270](#)] [[INSPIRE](#)].
- [5] M. Guica, T. Hartman, W. Song and A. Strominger, *The Kerr/CFT Correspondence*, *Phys. Rev. D* **80** (2009) 124008 [[arXiv:0809.4266](#)] [[INSPIRE](#)].
- [6] T. Hartman, K. Murata, T. Nishioka and A. Strominger, *CFT Duals for Extreme Black Holes*, *JHEP* **04** (2009) 019 [[arXiv:0811.4393](#)] [[INSPIRE](#)].
- [7] G. Compere, *The Kerr/CFT correspondence and its extensions: a comprehensive review*, *Living Rev. Rel.* **15** (2012) 11 [[arXiv:1203.3561](#)] [[INSPIRE](#)].
- [8] M. Cvetič and F. Larsen, *General rotating black holes in string theory: Grey body factors and event horizons*, *Phys. Rev. D* **56** (1997) 4994 [[hep-th/9705192](#)] [[INSPIRE](#)].
- [9] J.M. Maldacena and A. Strominger, *Black hole grey body factors and D-brane spectroscopy*, *Phys. Rev. D* **55** (1997) 861 [[hep-th/9609026](#)] [[INSPIRE](#)].
- [10] M. Cvetič and F. Larsen, *Greybody Factors and Charges in Kerr/CFT*, *JHEP* **09** (2009) 088 [[arXiv:0908.1136](#)] [[INSPIRE](#)].
- [11] A. Castro, A. Maloney and A. Strominger, *Hidden Conformal Symmetry of the Kerr Black Hole*, *Phys. Rev. D* **82** (2010) 024008 [[arXiv:1004.0996](#)] [[INSPIRE](#)].
- [12] M. Cvetič and D. Youm, *All the static spherically symmetric black holes of heterotic string on a six torus*, *Nucl. Phys. B* **472** (1996) 249 [[hep-th/9512127](#)] [[INSPIRE](#)].
- [13] F. Larsen, *A String model of black hole microstates*, *Phys. Rev. D* **56** (1997) 1005 [[hep-th/9702153](#)] [[INSPIRE](#)].
- [14] M. Cvetič and F. Larsen, *Conformal Symmetry for General Black Holes*, *JHEP* **02** (2012) 122 [[arXiv:1106.3341](#)] [[INSPIRE](#)].
- [15] M. Cvetič and F. Larsen, *Conformal Symmetry for Black Holes in Four Dimensions*, *JHEP* **09** (2012) 076 [[arXiv:1112.4846](#)] [[INSPIRE](#)].
- [16] M. Cvetič and G. Gibbons, *Conformal Symmetry of a Black Hole as a Scaling Limit: A Black Hole in an Asymptotically Conical Box*, *JHEP* **07** (2012) 014 [[arXiv:1201.0601](#)] [[INSPIRE](#)].
- [17] M. Cvetič and D. Youm, *BPS saturated and nonextreme states in Abelian Kaluza-Klein theory and effective N = 4 supersymmetric string vacua*, [hep-th/9508058](#) [[INSPIRE](#)].
- [18] M. Cvetič and D. Youm, *Dyonic BPS saturated black holes of heterotic string on a six torus*, *Phys. Rev. D* **53** (1996) 584 [[hep-th/9507090](#)] [[INSPIRE](#)].
- [19] M. Cvetič and D. Youm, *Entropy of nonextreme charged rotating black holes in string theory*, *Phys. Rev. D* **54** (1996) 2612 [[hep-th/9603147](#)].
- [20] E. Cremmer et al., *Vector Multiplets Coupled to N = 2 Supergravity: SuperHiggs Effect, Flat Potentials and Geometric Structure*, *Nucl. Phys. B* **250** (1985) 385 [[INSPIRE](#)].
- [21] M. Duff, J.T. Liu and J. Rahmfeld, *Four-dimensional string-string-string triality*, *Nucl. Phys. B* **459** (1996) 125 [[hep-th/9508094](#)] [[INSPIRE](#)].

- [22] K. Behrndt, R. Kallosh, J. Rahmfeld, M. Shmakova and W.K. Wong, *STU black holes and string triality*, *Phys. Rev. D* **54** (1996) 6293 [[hep-th/9608059](#)] [[INSPIRE](#)].
- [23] A. Virmani, *Subtracted Geometry From Harrison Transformations*, *JHEP* **07** (2012) 086 [[arXiv:1203.5088](#)] [[INSPIRE](#)].
- [24] P. Galli, T. Ortín, J. Perz and C.S. Shahbazi, *Non-extremal black holes of $N = 2$, $D = 4$ supergravity*, *JHEP* **07** (2011) 041 [[arXiv:1105.3311](#)] [[INSPIRE](#)].
- [25] T. Ortín and C. Shahbazi, *A Note on the hidden conformal structure of non-extremal black holes*, *Phys. Lett. B* **716** (2012) 231 [[arXiv:1204.5910](#)] [[INSPIRE](#)].
- [26] L. Huijse, S. Sachdev and B. Swingle, *Hidden Fermi surfaces in compressible states of gauge-gravity duality*, *Phys. Rev. B* **85** (2012) 035121 [[arXiv:1112.0573](#)] [[INSPIRE](#)].
- [27] X. Dong, S. Harrison, S. Kachru, G. Torroba and H. Wang, *Aspects of holography for theories with hyperscaling violation*, *JHEP* **06** (2012) 041 [[arXiv:1201.1905](#)] [[INSPIRE](#)].
- [28] S.A. Hartnoll and E. Shaghoulian, *Spectral weight in holographic scaling geometries*, *JHEP* **07** (2012) 078 [[arXiv:1203.4236](#)] [[INSPIRE](#)].
- [29] J.D. Brown and M. Henneaux, *Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity*, *Commun. Math. Phys.* **104** (1986) 207.
- [30] J.P. Gauntlett and O. Varela, *Universal Kaluza-Klein reductions of type IIB to $N = 4$ supergravity in five dimensions*, *JHEP* **06** (2010) 081 [[arXiv:1003.5642](#)] [[INSPIRE](#)].