



UvA-DARE (Digital Academic Repository)

Exactly soluble model of a quantum spin glass

Nieuwenhuizen, T.M.

DOI

[10.1103/PhysRevLett.74.4289](https://doi.org/10.1103/PhysRevLett.74.4289)

Publication date

1995

Published in

Physical Review Letters

[Link to publication](#)

Citation for published version (APA):

Nieuwenhuizen, T. M. (1995). Exactly soluble model of a quantum spin glass. *Physical Review Letters*, 74, 4289-4292. <https://doi.org/10.1103/PhysRevLett.74.4289>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Exactly Solvable Model of a Quantum Spin Glass

Th. M. Nieuwenhuizen

van der Waals–Zeeman Laboratorium, Universiteit van Amsterdam Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands
(Received 20 July 1994)

A mean-field spherical model with random couplings between pairs, quartets, and possibly higher multiplets of spins is considered. It has the same critical behavior as the Sherrington-Kirkpatrick model. It thus exhibits replica symmetry breaking. The order parameter function is solved exactly in the whole low temperature phase. The zero-field-cooled susceptibility remains finite at low T . Next a quantum version of the system is considered. Whereas the magnetic properties are not altered qualitatively, the thermodynamics is now regular at small temperatures.

PACS numbers: 75.10.Nr, 75.10.Jm, 75.40.Cx, 75.50.Lk

The phenomenon of spin glasses has generated a major new field of research. The basic phenomenon is breaking of ergodicity. This leads to highly nontrivial properties as observed recently, e.g., in aging experiments [1]. The best known spin glass model was formulated by Sherrington and Kirkpatrick (SK) [2]. Its solution was presented by Parisi; see Ref. [3] for a review. It gave a description of ergodicity breaking in terms of overlaps. For more recent reviews on spin glasses, see Refs. [4–6].

The spherical model has always played a special role in understanding the basic phenomena of phase transitions. It was introduced by the late Marc Kac and solved by Berlin and Kac [7]. It constituted a simple model for studying critical behavior; see Ref. [8] for a review.

The spherical random bond spin glass was studied in Ref. [9]. It shows freezing but no replica symmetry breaking. This model was extended by the present author to include short-range ferromagnetic interactions [10]. The spherical model with random couplings between p spins was studied recently by Crisanti and Sommers (CS). For $p > 2$ it was found that there occurs a one-step replica symmetry breaking [11]. Dynamical aspects of this transition were studied in detail [12].

For Ising spin glasses the Parisi solution is known explicitly near the spin glass transition. In the frozen phase the solution has been put in the form of a stochastic differential equation, for which no explicit solution is known. As a model solvable in the whole low temperature phase, we propose here a mean-field spherical model with random couplings between pairs, quartets, and, possibly, higher multiplets of spins. In contrast to previous spherical models, where only one type of these couplings occurs, the model belongs to the universality class of the Ising spin glass. We shall derive the explicit form of the order parameter in the low temperature phase. Then we shall

consider the recent quantum formulation of the spherical model [13] and study its low temperature behavior.

Consider a system with N spins with random couplings between sets of p spins. The Hamiltonian reads

$$\mathcal{H} = - \sum_{p=2}^{\infty} \sum_{i_1 < i_2 < \dots < i_p} J_{i_1 i_2 \dots i_p} S_{i_1} S_{i_2} \dots S_{i_p} - H \sum_i S_i. \quad (1)$$

The J 's are independent Gaussian random variables with average zero and variances $\langle J_{i_1 i_2 \dots i_p}^2 \rangle = (p-1)! J_p^2 N^{1-p}$. The spins are subject to the spherical constraint $\sum_i S_i^2 = N\sigma$. The classical partition sum reads

$$Z_{\text{cl}} = \int DS e^{-\beta \mathcal{H}}, \quad (2)$$

where $DS = \delta(\sum_{i=1}^N S_i^2 - N\sigma) \prod_i (dS_i / \sqrt{\pi})$. The thermodynamics of this model with only one of the p terms was worked out by CS. In a replica formulation of the average partition sum, it was found that the free energy only depends on the overlap $q_{\alpha\beta} = (1/N) \sum_{i=1}^N S_i^\alpha S_i^\beta$. This result has to be summed over p . We introduce

$$f(q) = \sum_{p=2}^{\infty} \frac{1}{p} J_p^2 q^p, \quad (3)$$

and we obtain a replicated free energy

$$2\beta F_n = -\beta^2 \sum_{\alpha,\beta=1}^n [f(q_{\alpha\beta}) + H^2 q_{\alpha\beta}] - \sum_{\alpha=1}^n \{\ln q\}_{\alpha\alpha} - n. \quad (4)$$

If $J_2 = 0$ the model has the one-step replica symmetry breaking solution studied by CS. When $J_2 > J_2^*(J_3, J_4, \dots)$ is large enough, a continuous transition will take place. From now on we assume that this is the case. In Eq. (4) the diagonal terms $q_{\alpha\alpha} \equiv q_d$ also occur; in the classical situation they are equal to $q_d = \sigma$. Expanding in off-diagonal elements $q_{\alpha\beta}$ one obtains

$$\begin{aligned} 2\beta F_n = & -n \{1 + \ln q_d + \beta^2 f(q_d) + \beta^2 H^2 q_d\} - \beta^2 H^2 \sum' q_{\alpha\beta} - \frac{1}{2} \left(\beta^2 J_2^2 - \frac{1}{q_d^2} \right) \sum' q_{\alpha\beta}^2 \\ & - \frac{1}{3q_d^3} \sum'_{\alpha\beta\gamma} q_{\alpha\beta} q_{\beta\gamma} q_{\gamma\alpha} - \frac{1}{3} \beta^2 J_3^2 \sum' q_{\alpha\beta}^3 - \frac{1}{4} \beta^2 J_4^2 \sum' q_{\alpha\beta}^4 + \dots, \end{aligned} \quad (5)$$

where the primes indicate exclusion of diagonal terms. In the case $J_3 = 0$, but $J_4 > 0$, this is exactly the relevant part of the free energy functional of the SK model [14,15]. The model thus belongs to the same universality class, and exhibits replica symmetry breaking. This continuous transition sets in at $T_G = J_2 q_d = J_2 \sigma$. For $J_3 > 0$ the model belongs to the universality class of the random bond Potts model. When the $J_2 > J_2^*$ and the next nonzero term is J_p for some $p > 4$, the model belongs to a new universality class. We shall not consider such situations here and assume $J_2 > J_2^*$, $J_3 = 0$, and $J_4 > 0$. The higher couplings are only relevant well below T_G .

We express the $q_{\alpha\beta}$ in the Parisi function $q(x)$. It takes the plateau value q_1 for $x_1 < x < 1$. In an external field it also has a plateau value $q(x) = q_0$ for $0 < x < x_0$. The inverse function is $x(q)$. Using the expressions of an appendix of Ref. [11], we obtain the explicit expression for the classical free energy

$$2\beta F_{cl} = -\beta^2 \int_0^1 dx \{f(q_d) - f(q(x)) + H^2 q_d - H^2 q(x)\} - \int_0^{q_1} \frac{dq}{I(q)} - \ln(q_d - q_1) - 1, \quad (6)$$

where

$$I(q) = q_d - q_1 + \int_q^{q_1} x(q') dq'. \quad (7)$$

The saddle point equation for $q(x)$ reads

$$\beta^2 f'(q(x)) + \beta^2 H^2 = \int_0^{q(x)} dq' \frac{1}{I(q')^2}. \quad (8)$$

In the region where $q'(x) \neq 0$, one has $\beta^2 f''(q) = I(q)^{-2}$. It follows that $x(q)$ has a universal shape at all T

$$x(q) = T \frac{f''(q)}{2\{f''(q)\}^{3/2}}, \quad (9)$$

which gives $q(x; T) = q(\beta x)$ after inversion. In an external field H , the plateau value q_0 follows from

$$H^2 = q_0 f''(q_0) - f'(q_0). \quad (10)$$

This condition is independent of T . q_1 follows from

$$q_1 = q_d - \frac{T}{\sqrt{f''(q_1)}}. \quad (11)$$

Of special interest is the case where the couplings J_p are such that

$$f(q) = J_2^2 \frac{q}{4a} \ln \frac{1+aq}{1-aq} \quad (12)$$

for some a in the range $0 < a < 1/\sigma$. In zero field one finds $q(x) = (\beta J_2 / 2a^2)x$ for $0 < x < x_1$ and

$$q_1 = \frac{2(q_d J_2 - T)}{J_2 + \sqrt{J_2^2 - 4a^2 T(q_d J_2 - T)}}. \quad (13)$$

Note that here the order parameter function $q(x)$ is either linear or constant in the whole low temperature

phase. The breakpoint $x_1 = 2a^2 T q_1$ vanishes both near the critical temperature and near zero temperature.

In an external field the freezing transition is given by the de Almeida–Thouless (AT) line [16]. Here the *replicon* mode [15] becomes massless. We propose to give this mode the more physical name *ergodon*. The critical field where this occurs, $H_{fr}(T)$, is obtained by inserting $q_0 = q_1(T)$ into Eq. (10). Note that H_{fr} remains finite at $T = 0$.

In the high temperature phase the susceptibility follows the Curie law $\chi = \beta q_d$. The zero-field-cooled susceptibility reads

$$\chi_{ZFC} = \beta(q_d - q_1) = \frac{1}{\sqrt{f''[q_1(T)]}}. \quad (14)$$

It remains finite at $T = 0$. This is usually observed experimentally. However, the SK model predicts a vanishing value at $T = 0$ [17]. The field-cooled susceptibility

$$\chi_{FC} = \beta \int_0^1 dx [q_d - q(x)] = \frac{1}{J_2} \quad (15)$$

is constant in the frozen phase, in good agreement with experiments. The internal energy and the entropy read

$$U_{cl} = \beta [f(q_1) - f(q_d)] + \frac{f'(q_1)}{\sqrt{f''(q_1)}} - \int_0^{q_1} dq \sqrt{f''(q)},$$

$$2S_{cl} = \beta^2 [f(q_1) - f(q_d)] + \frac{\beta f'(q_1)}{\sqrt{f''(q_1)}} + 1 + \ln(q_d - q_1). \quad (16)$$

For small T one has $S_{cl} \approx \frac{1}{4} - \frac{1}{4} \ln \beta^2 f''(q_1)$. As usual for classical vector or spherical spins, it goes to $-\infty$ as $T \rightarrow 0$. This also implies that the specific heat goes to a constant, $C \rightarrow 1/2$, at low T . For the case $\sigma = 1/2$ with pair and quartet couplings ($J_2 = 1, J_4 = 2$), the specific heat and the entropy have been plotted in Fig. 1.

We have wondered whether the anomalous low temperature behavior of our model can be cured. In order to do so, one should regularize the low temperature behavior of




FIG. 1. Specific heat and entropy as a function of temperature in the classical model. The arrow marks T_G .

the spherical model in general. This can be done by going to a description of spherical spins in a standard thermal field theory for bosonic *quantum* spins [13]. The situation becomes simplest when the spins are complex valued and couplings are Hermitian. For the case of random pair and quartet couplings we now assume the Hamiltonian

$$\mathcal{H} = -\frac{1}{2} \sum_{i,j} J_{ij} S_i^* S_j - \frac{1}{24} \sum_{i,j,k,l} J_{ijkl} S_i^* S_j S_k^* S_l - H \sum_i (\text{Re} S_i + \text{Im} S_i). \quad (17)$$

For each pair (i, j) there are now two independent random variables, J' and J'' , with average zero and variance $J_2^2/2N$, in terms of which $J_{ij} = J_{ji}^* = J' + iJ''$. For each quartet (i, j, k, l) with $i < k, j < l$ there are four independent random variables, $J'_{1,2}$ and $J''_{1,2}$, each having average zero and variance $9J_4^2/2N^3$. In terms of $J_{1,2} = J'_{1,2} + iJ''_{1,2}$, the couplings in Eq. (17) read $J_{ijkl} = J_{jikl}^* = J_1 + iJ_2$ and $J_{ijlk} = J_{ijlk}^* = J_1 - iJ_2$. The thermal partition sum involves spins on the imaginary-time interval $0 < \tau < \beta$

$$Z = \int \mathcal{D}S \exp \int_0^\beta d\tau \left\{ \frac{-1}{4\alpha} \sum_{i=1}^N S_i^*(\tau) \frac{d}{d\tau} S_i(\tau) - \mathcal{H}(S_i(\tau)) \right\} \quad (18)$$

with boundary conditions $S_i(\beta) = S_i(0)$. The τ axis is discretized in $\mathcal{M} = \beta/d\tau$ steps, with first $N \rightarrow \infty$ and then $\mathcal{M} \rightarrow \infty$, while $dS(\tau) = S(\tau + d\tau) - S(\tau)$. The integration measure is a repetition of the spherical measure at all τ , $\mathcal{D}S = C^N \prod_\tau DS(\tau)$, where $DS = \delta(2N\sigma - \sum_i S_i^* S_i) \prod_i (dS_i^* dS_i/2\pi)$ is the spherical measure for complex spins and $C = (1/2\alpha) \prod_{n \neq 0} (\pi|n|/2\alpha)$.

For a ferromagnet Eq. (18) leads to similar critical behavior as the classical spherical model. However, the entropy is non-negative and vanishes at $T = 0$. The zero point magnetization, $M_0 = \sqrt{\sigma - \alpha}$, shows a quantum reduction from the classical value $M_0 = \sqrt{\sigma}$ [13].

Here we wish to see whether the low temperature behavior is also cured for our spin glass model. We therefore extend the Crisanti-Sommers approach to thermal fields. We introduce a "Fermi level" $\mu_\alpha(\tau)$ related to the spherical constraint at imaginary time τ and overlaps $q_{\alpha\beta}(\tau, \tau') = (1/N) \sum_i S_i^*(\tau) S_i(\tau')$. We look for time-invariant solutions where $\mu_\alpha(\tau) = \mu$ and where $q_{\alpha\beta}(\tau, \tau') = \sum_\epsilon q_{\alpha\beta}(\epsilon) \exp\{2\pi i n T(\tau - \tau')\}$ with bosonic Matsubara frequencies $\epsilon \equiv \pi n T/2\alpha$. In the high temperature phase at zero field only the $q_{\alpha\alpha}(\epsilon)$ are nonzero. We shall denote $q_d \equiv q_{\alpha\alpha}(\epsilon = 0)$, which now becomes a smooth, increasing, strictly positive function of temperature. The spin glass temperature T_G is the solution of the relation $T_G = J_2 q_d(T_G)$. Here the off-diagonal elements $q_{\alpha\beta}(\epsilon = 0)$ become nonzero; since $T_G < \sigma$, this occurs below the classical transition point. The $q_{\alpha\beta}(\epsilon)$ with $\epsilon \neq 0$ are always equal to zero [18]. For $\epsilon \neq 0$ we

set $p_\epsilon \equiv \beta q_{\alpha\alpha}(\epsilon)$ and find that the full free energy reads

$$F = 2F_{\text{cl}}[q(x); q_d] + F_{\text{qc}}(q_d; \mu; p_\epsilon), \quad (19)$$

where F_{cl} is the free energy given by Eq. (6) with $f(q) = J_2^2 q^2/2 + J_4^2 q^4/4$. [The factor 2 in Eq. (19) arises since now spins are complex.] The quantum correction equals

$$\begin{aligned} \beta F_{\text{qc}} = & - \sum_{\epsilon \neq 0} \left\{ \frac{J_2^2}{2} p_\epsilon^2 + 1 - (\mu + 2i\epsilon)p_\epsilon + \ln 2i\epsilon p_\epsilon \right\} \\ & - \frac{T^2 J_4^2}{4} \sum_{\epsilon_1 + \epsilon_2 = \epsilon_3 + \epsilon_4} p_{\epsilon_1} p_{\epsilon_2} p_{\epsilon_3} p_{\epsilon_4} \\ & + \beta \mu (q_d - \sigma) + \ln 2\alpha, \end{aligned} \quad (20)$$

where terms with all ϵ_i equal to zero are excluded; they occur already in F_{cl} . In the quantum situation q_d, μ , and all p_ϵ have to be treated as additional variational parameters. For large temperatures one finds $\mu = Ty/\alpha$, $q_d = \alpha/y$, $p_\epsilon = 1/(\mu + 2i\epsilon)$, with $\tanh y = \alpha/\sigma$. The entropy $S_\infty = y\sigma/\alpha - \ln 2 \sinh y$ agrees with the result in the situation without interactions [13]. Only in the classical limit $\alpha \rightarrow 0$ does one recover the previous relation $q_d = \sigma$. Therefore, the susceptibility $\chi = q_d(T)/T$ has been reduced from its classical value at any T .

First consider the situation where only pair couplings occur, $J_4 = 0$. This case, where no replica symmetry breaking occurs, is exactly solvable since only Gaussian integrals occur. The free energy may be expressed as

$$\beta F = -\beta \mu \sigma + \int d\lambda \rho(\lambda) \ln 2 \sinh \alpha \beta (\mu - \lambda). \quad (21)$$

Here $\rho(\lambda)$ denotes the semicircular density of eigenvalues of the coupling matrix J_{ij} , viz. $\rho(\lambda) = \sqrt{4J_2^2 - \lambda^2}/2\pi J_2^2$. Alternatively we can solve the p_ϵ from Eq. (20). They read $p_\epsilon = 1/[\frac{1}{2}\mu + i\epsilon + \sqrt{(\frac{1}{2}\mu + i\epsilon)^2 - J_2^2}]$ with $\mu = 2J_2$ in the condensed phase. When expanding the $\ln \sinh$ in Eq. (21) as an ϵ sum, we find a term-by-term agreement with Eq. (20). It is clear that $S_0 = 0$, showing that our quantum formulation indeed regularizes the low- T anomaly of the entropy of Ref. [9]. The entropy and specific heat now behave as $S \sim C \sim T^{3/2}$, related to the square root singularity of the semicircular law at $\lambda = 2J_2$. This describes gapless excitations. In contrast, the mean-field quantum spherical ferromagnet does have a gap [13].

When also quartet couplings, and possibly higher couplings, are present, replica symmetry is broken. The entropy at low T can be obtained by analyzing the continuum limit of the equations for p_ϵ . Details of this derivation will not be presented here. It turns out that the same behavior occurs, $S \sim T^{3/2}$, $C \sim T^{3/2}$.

Figure 2 presents the specific heat and entropy for $\alpha = 1/4$ in the situation of Fig. 1, $\sigma = 1/2$, $J_2 = 1$, $J_4 = 2$.

In conclusion, we have presented a set of models where the order parameter function can be solved exactly in the whole low temperature phase. The critical behavior is




FIG. 2. Specific heat and entropy in the quantum model.

exactly the one derived by Parisi for the SK model. As compared to that model, the solution is explicit for all T . It is shown that the order parameter function has a universal shape in the whole low temperature region. The breakpoint x_1 is small near the transition and near $T = 0$. Well below T_G states with very different overlaps, and very different nature, occur.

We have also considered a quantum version of the model. It is seen that the shape of the order parameter function is not changed. The low- T divergency of the entropy in the classical model is eliminated. Now the entropy and specific heat vanish as $T^{3/2}$.

It is interesting to point out that the magnetic properties of the model, such as $q(x)$, χ_{ZFC} , and χ_{FC} have not changed qualitatively in the quantum description. The only relevant difference is that the self-overlap q_d has become a smooth function of T . In our model χ_{ZFC} remains nonzero at $T = 0$. This disagrees with the SK model, but is very often observed experimentally. Another new aspect of our model is the boundedness of the critical field below which freezing takes place. For nonzero field one has in the frozen phase $M(H, T) = H/\sqrt{f''(q_0)} = M(H)$ and $S(T, H) = S(T)$. Therefore, the Parisi-Toulouse hypothesis is satisfied [19].

It is hoped that the proposed model can be used as a basis to explain experiments. To this end, inclusion of ferromagnetic couplings and extension to vector spins seems of great interest. In a dynamical study of the

model, a closed set of equations will occur that bear the same physics as the more complicated set of the SK model.

The author is grateful for discussions with J. H. Sommers, H. Horner, and J. A. Mydosh. He thanks J. M. Luck and N. Garcia for hospitality at the CEA, Saclay, and the Universidad Autonoma de Madrid, where part of this work was done. This work was made possible by the Royal Dutch Academy of Arts and Sciences (KNAW).

-
- [1] F. Lefloch, J. Hamman, M. Ocio, and E. Vincent, *Europhys. Lett.* **18**, 647 (1992).
 - [2] D. Sherrington and S. Kirkpatrick, *Phys. Rev. Lett.* **35**, 1972 (1975).
 - [3] M. Mezard, G. Parisi, and M. A. Virasoro, *Spin Glass Theory and Beyond* (World Scientific, Singapore, 1987).
 - [4] K. Binder and A. P. Young, *Rev. Mod. Phys.* **58**, 801 (1986).
 - [5] K. H. Fischer and J. A. Hertz, *Spin Glasses*, Cambridge Studies in Magnetism 1, edited by D. Edwards and D. Melville (Cambridge University Press, Cambridge, 1991).
 - [6] J. A. Mydosh, *Spin Glasses: An Experimental Introduction* (Taylor and Francis, London, 1993).
 - [7] T. H. Berlin and M. Kac, *Phys. Rev.* **86**, 821 (1952).
 - [8] G. S. Joyce, in *Phase Transitions and Critical Phenomena* 2, edited by C. Domb and M. S. Green (Academic, London, 1972).
 - [9] M. Kosterlitz, D. J. Thouless, and R. C. Jones, *Phys. Rev. Lett.* **36**, 1217 (1976).
 - [10] Th. M. Nieuwenhuizen, *Phys. Rev. B* **31**, 7487 (1985).
 - [11] A. Crisanti and H.-J. Sommers, *Z. Phys. B* **87**, 341 (1992).
 - [12] A. Crisanti, H. Horner, and H. J. Sommers, *Z. Phys. B* **92**, 257 (1993).
 - [13] Th. M. Nieuwenhuizen, following Letter, *Phys. Rev. Lett.* **74**, 4293 (1995).
 - [14] E. Pytte and J. Rudnick, *Phys. Rev. B* **19**, 3603 (1979).
 - [15] A. J. Bray and M. A. Moore, *Phys. Rev. Lett.* **41**, 1086 (1978).
 - [16] J. R. L. de Almeida and D. J. Thouless, *J. Phys. A* **11**, 983 (1978).
 - [17] D. J. Thouless, P. W. Anderson, and R. G. Palmer, *Philos. Mag.* **35**, 593 (1977).
 - [18] Y. Y. Goldschmidt and P. Y. Lai, *Phys. Rev. Lett.* **64**, 2467 (1990).
 - [19] G. Parisi and G. Toulouse, *J. Phys. Lett.* **41**, L361 (1980).