



UvA-DARE (Digital Academic Repository)

Lentiviral shRNA silencing of murine bone marrow cell CCR2 leads to persistent knockdown of CCR2 function in vivo

Bot, I.; van Eck, M.; van Santbrink, P.J.; Groot, P.H.E.; Hildebrand, R.B.; Seppen, J.; van Berkel, T.J.C.; Biessen, E.A.L.

DOI

[10.1182/blood-2004-12-4839](https://doi.org/10.1182/blood-2004-12-4839)

Publication date

2005

Document Version

Final published version

Published in

Blood

[Link to publication](#)

Citation for published version (APA):

Bot, I., van Eck, M., van Santbrink, P. J., Groot, P. H. E., Hildebrand, R. B., Seppen, J., van Berkel, T. J. C., & Biessen, E. A. L. (2005). Lentiviral shRNA silencing of murine bone marrow cell CCR2 leads to persistent knockdown of CCR2 function in vivo. *Blood*, *106*(4), 1147-1153. <https://doi.org/10.1182/blood-2004-12-4839>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Lentiviral shRNA silencing of murine bone marrow cell CCR2 leads to persistent knockdown of CCR2 function in vivo

Ilze Bot, Jian Guo, Miranda Van Eck, Peter J. Van Santbrink, Pieter H. E. Groot, Reeni B. Hildebrand, Jurgen Seppen, Theo J. C. Van Berkel, and Erik A. L. Biessen

A major barrier in hematopoietic gene function studies is posed by the laborious and time-consuming generation of knockout mice with an appropriate genetic background. Here we present a novel lentivirus-based strategy for the in situ generation of hematopoietic knockdowns. A short hairpin RNA (shRNA) was designed targeting murine CC-chemokine receptor 2 (CCR2), which was able to

specifically blunt CCR2 expression at the mRNA, protein, and functional levels in vitro. Reconstitution of irradiated recipient mice with autologous bone marrow that had been ex vivo transduced with shRNA lentivirus led to persistent down-regulation of CCR2 expression, which translated into a 70% reduction in CCR2-dependent recruitment of macrophages to an inflamed peritoneal cavity without

noticeable side effects on related chemokine receptors or general inflammation status. These findings clearly demonstrate the potential of shRNA lentivirus-infected bone marrow transplantation as a rapid and effective method to generate hematopoietic knockdowns for leukocyte gene function studies. (Blood. 2005;106:1147-1153)

© 2005 by The American Society of Hematology

Introduction

Bone marrow transplantation studies have greatly facilitated the identification of macrophage genes that play a significant role in inflammatory disorders¹; for example, atherosclerosis.²⁻⁴ However, this experimental setup essentially depends on the availability of transgenic mice or knockouts of the gene of interest for providing bone marrow. Thus, progress is considerably hampered by the laborious construction of, in particular, knockout mice and the subsequent backcrossing to an appropriate disease background for analysis of gene function. Moreover, the development of knockouts is even more difficult when deletion results in embryonic lethality as is the case for tissue factor (TF)⁵ or vascular endothelial growth factor (VEGF).⁶

Efficient and stable gene transfer to hematopoietic stem cells has been a major challenge in gene therapy. For several years, a range of vectors of onco-retroviral origin, which integrate into the host-cell genome, has been developed.⁷ A major drawback of these retroviral systems, mostly based on Moloney leukemia viruses, is their inability to integrate in the genome of nonmitotic cells. Since hematopoietic stem cells are rather quiescent, their retroviral transduction generally is inefficient. While transduction efficiency can be improved by cytokine supplementation,⁸ by reducing cyclin-dependent kinase inhibitor levels⁹ or by fibronectin coating to induce stem-cell proliferation,^{10,11} these manipulations may likely affect stem-cell differentiation. Unlike onco-retroviral vectors, lentivirus offers the advantage of stably transducing not only mitogenic but also quiescent cells.¹²⁻¹⁵ Indeed, efficient lentiviral transduction of bone marrow stem cells using lentiviral vectors has

been reported.¹⁶ Such transduction is not only required to obtain required gene expression levels in vivo, but also obviates the use of time-consuming preselection strategies.

We argued that the high levels of lentiviral gene transfer to bone marrow may be exploited to knockdown hematopoietic genes in vivo via RNA interference (RNAi), defined as sequence-specific posttranscriptional gene silencing.¹⁷⁻²⁴ The advent of RNAi or small interfering RNA (siRNA) has instigated a true revolution in functional genomics and allowed the in situ knockdown of genes. Synthetic short double-strand (ds) DNA sequences of 21 to 25 nucleotides in length²⁵ were reported to be efficient in inducing selective mRNA degradation and to suppress gene expression.^{26,27} Recently, Brummelkamp et al²⁸ have generated a mammalian expression vector, encoding a short hairpin dsRNA transcript under the control of the RNA polymerase III-dependent H1 gene promoter for sustained silencing of a transgene. At present, several groups reported successful lentiviral delivery of short hairpin RNA (shRNA) to eukaryotic cells in vitro. Tiscornia et al²⁹ showed decreased green fluorescent protein (GFP) levels after transduction with a lentiviral shRNA construct. Likewise, successful lentiviral delivery of shRNAs to human cells^{30,31} and even to human hematopoietic stem cells was reported.³²

In this study we have addressed the potential of lentivirus-aided gene silencing of genes in bone marrow for the in situ generation of chimeras with blunted gene expression in the hematopoietic-cell lineage in vivo. As a model gene we chose murine CC-chemokine receptor 2 (*Ccr2*), which was previously shown to play a central

From the Division of Biopharmaceutics, Gorlaeus Laboratories, Leiden University, Leiden, The Netherlands; and the Academic Medical Center, Department of Experimental Hepatology, Amsterdam, The Netherlands.

Submitted December 21, 2004; accepted April 22, 2005. Prepublished online as *Blood* First Edition Paper, May 10, 2005; DOI 10.1182/blood-2004-12-4839.

Supported by grant 016.026.019 from the Netherlands Organization for Scientific Research (I.B.) and grants 2003T201 (E.A.L.B.), 2001T041 (M.V.E.), and M93.001 (P.J.V.S.) from the Netherlands Heart Foundation, and GlaxoSmithKline Pharmaceuticals, United Kingdom (J.G.).

I.B. and J.G. contributed equally to this study.

An Inside *Blood* analysis of this article appears in the front of this issue.

Reprints: Ilze Bot, Division of Biopharmaceutics, Leiden/Amsterdam Center for Drug Research, Gorlaeus Laboratories, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands; e-mail: i.bot@lacdr.leidenuniv.nl.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2005 by The American Society of Hematology

role in the recruitment of monocytes to sites of inflammation and during atherogenesis.³³⁻³⁵ An shRNA construct was developed against CCR2 (shCCR2), which proved highly efficient in down-regulating CCR2 expression both at an mRNA as well as at a protein level in CCR2-transfected HEK 293 cells. Furthermore, we have efficiently transduced bone marrow cells and reconstituted irradiated C57Bl/6 mice with shCCR2-transduced bone marrow cells, which considerably attenuated macrophage recruitment to the site of inflammation to a level comparable with that of mice reconstituted with *Ccr2*^{-/-} bone marrow.

Materials and methods

Cloning and expression of murine CCR2-GFP in HEK 293 cells

Total RNA was extracted from the murine monocytic cell line WEHI, reverse transcribed using M-MuLV reverse transcriptase (RevertAid; MBI Fermentas, Leon-Roth, Germany) and the cDNAs were amplified by polymerase chain reaction (PCR) to obtain full-length cDNA for murine CCR2 using the following primers: forward: 5'-AGCCTCGAGATGGAA-GACAATAATATGTT-3' and reverse: 5'-ATAAGCTTTTACAAC-CCAACCGAGACCTCT-3'. Primers contained extra *XhoI* and *HindIII* restriction sites to facilitate cloning (underlined). Purified PCR products were subsequently cloned into pCR2.1 (Invitrogen, Breda, The Netherlands) and subcloned into pEGFPN-1 (Clontech, Palo Alto, CA). Murine CCR2 and GFP were fused at the GFP N-terminus via removal of the stop codon, affording the pEGFPN-1/CCR2 fusion vector. A CCR2 expression vector was generated by cloning full-length *Ccr2* into pcDNA3.1(-) (Invitrogen), designated as pcDNA3.1(-)/CCR2. All constructs were sequenced to confirm the identity of murine *Ccr2*.

Design and cloning of shRNA directed against murine CCR2

A pair of 64-nucleotide oligonucleotides encoding a 19-nucleotide CCR2 shRNA (5'-GATCCCCCTGTGTGATTGACAAGCACTTCAAGAGAG-TGCTTGTCATCACACAGTTTTTGGAAA-3' and 5'-AGCTTTTCC-AAAACTGTGTGATTGACAAGCACTCTTGAAGTGTCTGTCAA-TCACACAGGGG-3') were designed, which contained extra *BgIII* and *HindIII* restriction sites to facilitate cloning. The position of the core 19-nucleotide sequence (underlined) targeted nucleotides 867 to 885 of the murine *Ccr2*. Blast search was performed using the National Center for Biotechnology Information (NCBI) Expressed Sequence Tags (EST) database to ensure that the shRNA construct was targeting only murine CCR2. The 64-nucleotide oligonucleotides were annealed and cloned into the *BgIII* and *HindIII* sites of the pSUPER vector (kindly provided by Dr Agami, The Netherlands Cancer Institute, Amsterdam, The Netherlands) and the resulting vector, designated pSUPER-H1.shCCR2, was subsequently sequenced to confirm identity.

Silencing effect of pSUPER-H1.shCCR2

HEK 293 cells, grown at 37°C and 5% CO₂ in Dulbecco modified Eagle medium (DMEM) supplemented with 10% fetal calf serum (FCS), 100 µg/mL streptomycin, 100 U/mL penicillin, and 2 mM L-glutamine (all from

Cambrex Bio Science, Verviers, Belgium), were transiently cotransfected with a constant amount of pEGFPN-1/CCR2 (0.5 µg) and 0.1 µg, 0.5 µg, and 1.5 µg pSUPER-H1.shCCR2, supplemented to 1.5 µg with pSUPER-H1.Empty vector using Lipofectamine 2000 (Invitrogen). As controls, the effects of an shRNA designed against luciferase (pSUPER-H1.shLuc, CGTACGCGGAATACTTCGA³⁶) and a nonsense sequence (pSUPER-H1.shInert, AGGCTGCTTGACGATCTA) cloned into pSUPER on CCR2 expression levels were determined. After 48 hours, total RNA was extracted from the HEK 293 cells and reverse transcribed as described under "Cloning and expression of murine CCR2-GFP in HEK 293 cells." Quantitative analysis of CCR2 expression was performed using an ABI PRISM 7700 Taqman apparatus (Applied Biosystems, Foster City, CA) as described previously³⁷ (Table 1). The amplicon encompassed the shCCR2 core sequence (nucleotides 851-991), enabling the analysis of sequence-specific degradation of murine CCR2 mRNA and excluding artifacts due to the presence of shRNA. Human hypoxanthine phosphoribosyltransferase (hHPRT) was used as a standard housekeeping gene and non-reverse transcribed RNA samples were used as controls to determine genomic DNA contamination. Murine CCR2 mRNA expression was calculated relative to that of hHPRT on the basis of ΔCt values.

After transfection with pEGFPN-1/CCR2 and pSUPER-H1.Empty, pSUPER-H1.shLuc, pSUPER-H1.shInert, or different amounts of pSUPER-H1.shCCR2, the cells were harvested, resuspended in phosphate-buffered saline (PBS), and analyzed using fluorescence activated cell sorting (FACS; FACScalibur, BD Biosciences) for GFP expression. For Western blotting, transfected cells were washed gently with PBS, scraped off, and subsequently lysed in sodium dodecyl sulfate (SDS) sample buffer supplemented with a protease inhibitor cocktail (Roche, Mannheim, Germany). Equal amounts of protein were applied to a polyacrylamide gel (12.5%) and transferred to nitrocellulose. Visualization of GFP with a mouse anti-GFP monoclonal antibody and a peroxidase-conjugated rat anti-mouse IgG as a secondary antibody (1:2000 and 1:1000 dilution, respectively; Santa Cruz Biotechnology, Santa Cruz, CA) was performed via the enhanced chemiluminescence (ECL) technique (Amersham Pharmacia Biotechnology, Piscataway, NJ). Blots were simultaneously probed with a mouse anti-β-tubulin antibody (Sigma, Zwijndrecht, The Netherlands) to verify equal loading of the samples.

Functional loss of CCR2 by shCCR2

The effect of the shRNA vector on CCR2 function was determined by analyzing calcium influx in response to the endogenous ligand for murine CCR2, CC-chemokine ligand 2 (CCL2), or JE. HEK 293 cells, expressing murine CCR2, were transfected with 0.1 µg pSUPER-H1.shCCR2 or with pSUPER-H1.Empty control vector. After 48 hours, cells were seeded onto poly-lysine-coated glass coverslips to measure transient intracellular calcium influx. Cells were loaded with 20 µM Fura-2-AM (Invitrogen) for 60 minutes and after washing with Hank balanced salt solution, the coverslips were mounted in an incubation chamber. JE (Santa Cruz Biotechnology) was added to a final concentration of 1 nM and real-time digital fluorescence imaging was performed using a fluorescence microscope (excitation: 340 nm and 380 nm; emission: 495 nm) for 20 minutes. The mean fluorescence intensity was determined to calculate intracellular calcium concentrations.

Table 1. Primer set sequences used for PCR analysis

Gene	Forward (5'-3')	Reverse (5'-3')
<i>hHPRT</i>	TGACACTGGCAAACAATGCA	GGTCCTTTTCACCAGCAAGCT
<i>mHPRT</i>	TTGCTCGAGATGTCTGAAGGA	AGCAGGTGACAAAGAAGCTTATAG
<i>CCR2</i>	CTTGGGAATGAGTAACTGTGTGA	GGAGAGATACCTTCGGAAGCTTCT
<i>CCR3</i>	TGCAGGTGACTGAGGTGATTG	CGGAACCTCTCACAAACAAG
<i>CCR5</i>	GACTGTGACGAGGAAGTGACAT	CTTTGACGCCAGCTGAGCAA
<i>CXCR3</i>	GCTGCTGTCCAGTGGGTTTT	AGTTGATGTTGAACAAGGCGC
<i>IFN_γ</i>	ATAACTATTTTAACTCAAGTGGCATAGATGT	TCTGGCTCTGCAGGATTTTCA
<i>IFN_β</i>	GCTCCTGGAGCAGCTGAATG	TCCGTATCTCCATAGGGATCTT
<i>SRY</i>	GTGTGGTCCCTGGGTGAGA	TGTAAGGCTTTTCCACCTGCAT

Lentivirus vector construction and production

The lentiviral expression vectors pRRI-cPPT-H1.PreSIN (H1.Empty) and pRRI-cPPT-H1.shCCR2-PreSIN (H1.shCCR2) were constructed by removing the cytomegalovirus (CMV) promoter from the expression vector pRRI-cPPT-CMV-PreSIN¹⁵ using *ClaI* and *PstI* digestion, followed by insertion of the H1 promoter or the complete H1.shCCR2 construct. Virus was produced as described³⁸ using transient calcium phosphate cotransfection of 293T cells with the H1.Empty or H1.shCCR2 vector together with pMDL/RRE, pRSV-REV, and pVSV-G. Viral supernatants were passed through a 0.45 μ m filter and concentrated by ultracentrifugation (41 000 g, 2 hours, 4°C). Viral titers were determined essentially as described by Sastry et al.³⁹ In short, HEK 293 cells were transduced with serially diluted viral supernatant, and 48 hours after transduction total genomic DNA was isolated from these cells and the number of vector DNA copies was determined using PCR analysis with pRRI-cPPT-H1.PreSIN vector as the calibration standard (forward primer: GTGCAGCAGCAGAACAATTG; reverse primer: CCCAGACTGTGAGTTGCAA).

Animals

All animal work was approved by the regulatory authority of Leiden and performed in compliance with the Dutch government guidelines. C57Bl/6 mice were obtained from Charles River (Maastricht, the Netherlands). *Ccr2*^{-/-} mice were kindly provided by Dr Maeda (University of North Carolina, Chapel Hill). Mice that underwent bone marrow transplantation were housed in sterile filter-top cages and fed a sterile regular chow diet (RM3; Special Diet Services, Witham, United Kingdom). Drinking water was infused with antibiotics (83 mg/L ciprofloxacin and 67 mg/L polymyxin B sulfate) and 6.5 g/L sugar and was provided ad libitum.

In vitro transduction protocol for bone marrow cells

Bone marrow cell suspensions were isolated from C57Bl/6 mice by flushing the femurs and tibias with PBS. Single-cell suspensions were prepared by passing the cells through a cell strainer and 10⁵ cells/well were plated in a 12-well plate. Viral transductions were performed by incubating the bone marrow cells with different titers of pRRI-cPPT-PGK-GFP-PreSIN in complete DMEM supplemented with 10 μ g/mL diethylaminoethyl (DEAE)-dextran at 37°C. After 24 hours, cells were washed with PBS and fixed in 4% paraformaldehyde for FACS analysis to determine GFP expression levels. Viral copy number inserted in the bone marrow cells after 24 hours was determined by isolation of total genomic DNA (from 10⁶ bone marrow cells) and PCR analysis.

Irradiation and bone marrow transplantation

To induce bone marrow aplasia, female C57Bl/6 mice (12 to 14 weeks of age) were exposed to a single dose of 9 Gy (0.19 Gy/min, 200 kV, 4 mA) total body irradiation, using an Andrex Smart 225 Röntgen source (YXLON International, Copenhagen, Denmark) with a 6-mm aluminium filter, one day before transplantation. Bone marrow cell suspensions were isolated from C57Bl/6 mice as described in the previous section. Subsequently, bone marrow cells (1.0 \times 10⁷) were transduced with either H1.Empty or H1.shCCR2 lentivirus in the presence of 10 μ g/mL DEAE-dextran (multiplicity of infection [moi] = 15). After 24 hours, the cells were injected into the tail vein of the irradiated recipients (n = 4 per group). As a positive control, irradiated C57Bl/6 mice (n = 5) received 1.0 \times 10⁷ bone marrow cells isolated from male *Ccr2*^{-/-} mice. In a similar set-up, C57Bl/6 mice (n = 2) were injected with bone marrow cells transduced with GFP lentivirus. In these mice, GFP expression levels in blood cells were determined 6 weeks after transplantation by preparing cytopspins of white blood cells (9 g, 5 minutes) after lysis of the erythrocytes. The images were taken with a Leica DM-RE microscope and LeicaQwin software (Leica Imaging Systems).

Macrophage recruitment

Six weeks after bone marrow transplantation, the mice receiving H1.Empty or H1.shCCR2 lentivirus-transduced bone marrow and the mice reconsti-

tuted with *Ccr2*^{-/-} bone marrow were injected intraperitoneally with sterile 3% (wt/vol) Brewer thioglycollate solution (Difco, Detroit, MI). Peritoneal macrophages were isolated 5 days later by lavage of the peritoneal cavity with PBS (10 mL) and counted after lysis of erythrocytes. Also, blood smears were prepared, stained with Wright Giemsa (Sigma), and scored manually. Total RNA was isolated from peritoneal macrophages and gene expression was determined by reverse transcription (RT)-PCR compared with murine HPRT (Table 1). Also, bone marrow was isolated from each individual mouse and, after lysis of the bone marrow cells, genotyped for the presence of the *Ccr2* gene in a single reaction (primers: CCR2 forward, 5'-GATGATGGTGAGCCTTGTC-3'; CCR2 reverse, 5'-CACAGCAT-GAACAATAGCCA-3'; and a pgk-*neo* specific primer, Neo 5'-TTA-AGGGCC-AGTCATCTCT-3'). In wild-type mice, a 360 base pair sequence is amplified corresponding with intact *Ccr2*, whereas in the *Ccr2*^{-/-} mice a 290 base pair sequence is generated. Also, viral copy number in the bone marrow was determined. In addition, the percentage repopulation of male bone marrow in the female recipients at that time point was determined by means of Taqman PCR analysis on the *Sry* gene, located on the Y-chromosome. To quantify the percentage of male bone marrow, a calibration curve was used containing increasing amounts of male bone marrow supplemented with female bone marrow to standardize the amount of genomic DNA per well.

Statistical analysis

Data are expressed as mean plus or minus the standard error of the mean (SEM). A 2-tailed Student *t* test was used to compare individual groups. To determine the significance of the relative mRNA expression levels, statistical analysis was performed on Δ Ct values.

Results

Silencing CCR2 expression by shRNA vectors

To determine the silencing capacity of an shRNA against CCR2 (pSUPER-H1.shCCR2) on murine CCR2 expression, HEK 293 cells were cotransfected with pEGFPN-1/CCR2 fusion vector and increasing amounts of pSUPER-H1.shCCR2. As controls, the effects of an shRNA sequence against luciferase and a nonsense shRNA sequence on CCR2 expression were also determined. CCR2 mRNA levels were already found to be significantly reduced after cotransfection with only 0.1 μ g of pSUPER-H1.shCCR2 (Figure 1A, *P* < .001 compared with the shRNA controls at all concentrations) with a maximum reduction of 94% at 1.5 μ g pSUPER-H1.shCCR2, whereas the shRNA controls and pSUPER-H1.Empty had no effect. In agreement, expression analysis by fluorescence microscopy revealed an almost complete quenching of GFP-tagged murine CCR2 protein expression in HEK 293 cells cotransfected with pSUPER-H1.shCCR2 compared with that of the pSUPER-H1.Empty cotransfected control (Figure 1B). Subsequent FACS analysis (Figure 2A) confirmed that CCR2/GFP fusion protein expression was dose-dependently blunted after cotransfection with increasing amounts of pSUPER-H1.shCCR2, whereas the shRNA controls (ie, pSUPER-H1.Empty, pSUPER-H1.shLuc, and the nonsense shRNA sequence) were ineffective. Also, the fluorescence intensity of the GFP-positive cells was reduced to the level of untransfected HEK 293-cell controls (on average 380) after cotransfection with pSUPER-H1.shCCR2 (Figure 2B). These FACS data were confirmed by Western blot analysis (Figure 2C).

Intracellular calcium levels were measured in HEK 293 cells expressing CCR2 and the H1.shCCR2 or H1.Empty sequence in response to the murine CCR2 ligand JE. Upon incubation with JE (1 nM), transient calcium influx was observed after 2 minutes, and peaked at 4 minutes, after which intracellular calcium levels




Figure 1. Silencing of CCR2 expression by pSUPER-H1.shCCR2. (A) Expression of murine CCR2 in HEK 293 cells after cotransfection with pEGFPN-1/CCR2 and pSUPER-H1.Empty (1.5 μg), relative to that of HPRT, was not affected by pSUPER-H1.shLuc or pSUPER-H1.shlncrt, whereas CCR2 mRNA levels were up to 94% reduced after cotransfection with increasing amounts of pSUPER-H1.shCCR2 ($***P < .001$). Error bars represent SEM. (B) HEK 293 cells cotransfected with either pEGFPN-1/CCR2 and pSUPER-H1.Empty (1.5 μg; left panel) or pEGFPN-1/CCR2 and pSUPER-H1.shCCR2 (0.5 μg or 1.5 μg; middle and right panels, respectively), which show increased quenching of GFP protein expression on the cell surface, indicating almost complete loss of CCR2 surface protein after transfection with pSUPER-H1.shCCR2. Cells were viewed with a Biorad 2-photon confocal laser scanning microscope (1000 \times magnification), and images were digitized with Adobe Photoshop 7.0.

gradually declined to base values. Transfection of these cells with pSUPER-H1.shCCR2 completely abolished the JE-induced calcium influx to levels comparable with that of nonstimulated cells (Figure 2D), indicating that the observed reduction in CCR2 mRNA and protein levels establishes a complete loss of CCR2 function in response to JE.

Transduction of CCR2-transfected HEK 293 cells with H1.shCCR2 lentivirus resulted in a significantly reduced expression of CCR2 relative to hHPRT (5.95 ± 0.67 versus 14.4 ± 3.6 in cells transduced with H1.Empty virus; $P = .02$).

Transduction of bone marrow cells

In the next stage, the transduction efficiency of lentivirus in bone marrow cells was optimized. FACS analysis of bone marrow cells showed that transduction was not satisfactory at low multiplicities of infection ($\text{moi } 6.5\% \pm 0.6\%$ and $9.8\% \pm 3.3\%$ at moi 's of 1 and 2, respectively). The use of higher titers and DEAE-dextran (10 μg/mL) led to a strong increase in infection efficiency from $3.1\% \pm 0.5\%$ in control cells via $67.1\% \pm 6.0\%$ ($\text{moi} = 5$) and $76.2\% \pm 2.0\%$ ($\text{moi} = 10$) to $86.1\% \pm 2.3\%$ ($\text{moi} = 20$) 24 hours after infection (Figure 3A). Viral incorporation analysis revealed that bone marrow cells contained on average 2 copies of virus per *Hprt* gene 24 hours after transduction of the cells with GFP, H1.shCCR2, or H1.Empty control lentivirus (Figure 3B). The high transduction, obtained at moi 's of 10 to 20 in the presence of 10 μg/mL DEAE-dextran, suffices for *in vivo* transplantation purposes and circumvents the need of a laborious preselection step. Six weeks after injection of GFP lentivirus-transduced bone marrow into irradiated C57Bl/6 mice, cytospin analysis showed that approximately 80% to 90% of all white blood cells were GFP positive (Figure 3C-F).

Transplantation of H1.shCCR2-transduced bone marrow

To generate chimeras with silenced CCR2 expression, bone marrow from C57Bl/6 mice was transduced with either H1.shCCR2 or H1.Empty control lentivirus in the presence of 10 μg/mL DEAE-dextran and intravenously injected into irradiated C57Bl/6 recipient mice, while a third group received *Ccr2*^{-/-} bone marrow. No differences were observed in either weight gain during the experiment or in mononuclear-cell and neutrophil content of the total white blood cell population at 7 weeks after transplantation (data not shown). At that point, macrophages were recruited to the peritoneal cavity by an intraperitoneal thioglycollate challenge. Assessment of the number of peritoneal macrophages, 5 days after challenge, revealed a significant 70% reduction in macrophage recruitment in mice that had received H1.shCCR2 lentivirus-transduced bone marrow compared with the H1.Empty lentivirus controls ($8 \times 10^6 \pm 4 \times 10^6$ cells versus $27 \times 10^6 \pm 6 \times 10^6$ cells, respectively; $P = .03$, Figure 4A). Interestingly, macrophage recruitment was comparable to that of mice reconstituted with *Ccr2*^{-/-} bone marrow ($8 \times 10^6 \pm 1 \times 10^6$ cells; $P = .9$ versus the H1.shCCR2 lentivirus group). Macrophage recruitment levels of mice reconstituted with H1.Empty-transduced bone marrow




Figure 2. Silencing of CCR2 protein by pSUPER-H1.shCCR2. (A) FACS analysis of HEK 293 cells after cotransfection with pEGFPN-1/CCR2 and pSUPER-H1.Empty (1.5 μg, control), which was not significantly affected by cotransfection with pSUPER-H1.shLuc or pSUPER-H1.shlncrt. Cotransfection with increasing amounts of pSUPER-H1.shCCR2 led to a significant and dose-dependent silencing of CCR2 protein level up to 75%. $*P \leq .05$. (B) The average fluorescence intensity of the GFP-positive cells was even more markedly reduced after cotransfection with pSUPER-H1.shCCR2 to the untransfected HEK 293 control level (approximately 390). The nonspecific controls had no effect on GFP fluorescence intensity compared with pSUPER-H1.Empty control levels. $**P \leq .01$; $***P \leq .001$. (C) Western blot analysis of HEK 293-cell lysates for GFP protein expression using β -tubulin (50 kDa) as internal standard, showing a dose-dependent and specific reduction in GFP protein expression in cells cotransfected with increasing amounts of pSUPER-H1.shCCR2 (lanes D, E, and F) compared with only pEGFPN-1/CCR2-transfected (lane B) or pSUPER-H1.Empty cotransfected cells (lane C). Lane 1 represents a sample of untransfected HEK 293 cells. (D) Measurements of JE-induced calcium influx in HEK 293 cells expressing murine CCR2 and the H1.Empty (\blacktriangle) or the H1.shCCR2 sequence (\square). As a control, unstimulated HEK 293 cells expressing CCR2 were measured (\bullet). In contrast to pSUPER-H1.Empty-transfected cells, no calcium influx was observed in pSUPER-H1.shCCR2-transfected cells after stimulation with JE, indicating that CCR2 signal transduction is silenced by shCCR2 treatment. Error bars represent SEM.




Figure 3. Efficient transduction of bone marrow cells using GFP lentivirus. (A) Transduction of whole bone marrow with GFP lentivirus led to a titer-dependent increase in percentage of GFP-expressing cells. (B) Virus particle incorporation in bone marrow cells compared with HPRT of untransduced control cells and 24 hours after transduction with GFP-, H1.Empty-, or H1.shCCR2-lentivirus. Error bars in panels A and B represent SEM. (C,D) Fluorescence and light microscopic high-power field, respectively, of a white blood cell cytospin from control mice 6 weeks after transplantation of nontransduced bone marrow (original magnification, $\times 200$). (E,F) White blood cells from mice that underwent transplantation with lentiviral GFP-transduced bone marrow show approximately 90% GFP-positive cells (panel E; original magnification, $\times 200$), indicating that transplantation of lentivirus-transduced bone marrow results in sustained transgene expression by blood cells. Images were captured with a Leica DR-ME microscope equipped with a $20\times$ objective lens and a $10\times$ ocular lens, and Leica QWin software (Leica Imaging Systems) was used.

matched with those found in control mice that underwent transplantation (female C57Bl6 recipient mice that had received untransduced male C57Bl6 bone marrow, data not shown), which indicates that lentiviral transduction of bone marrow per se does not elicit an inflammatory response.

Analysis of CCR2 expression by peritoneal macrophages revealed a highly significant reduction in CCR2 mRNA levels in mice that received a transplant of H1.shCCR2- versus H1.Empty lentivirus-transduced bone marrow (expression relative to murine HPRT: 0.4 ± 0.1 and 1.3 ± 0.2 , respectively; $P = .002$). For comparison, mice that had received *Ccr2*^{-/-} bone marrow showed a relative CCR2 expression of 0.09 ± 0.02 (Figure 4B).

As expected, genotyping showed that C57Bl/6 mice that received a transplant of H1.Empty- or H1.shCCR2 lentivirus-transduced bone marrow displayed the 360 base pair amplicon, indicative of wild-type *Ccr2* (Figure 4C). C57Bl/6 mice that had received *Ccr2*^{-/-} bone marrow showed the *Ccr2*^{-/-} phenotype (290 base pair sequence). Reconstitution with male bone marrow in the female recipient mice was quantitative and persistent for at least 7 weeks in all 3 groups, as judged from PCR analysis of the *Sry* gene (Figure 4D). PCR analysis of genomic DNA from bone marrow revealed that unpurified bone marrow from mice that received a transplant of H1.shCCR2 contained on average 2 virus integrants per *Sry* copy in both the H1.shCCR2 (2.30) and the H1.Empty groups (1.98), while as expected in bone marrow from mice that received a transplant of *Ccr2*^{-/-} no virus particles were detected. Furthermore, no differences were observed in peritoneal macrophage expression levels of other chemokine receptors, such as CCR3, CCR5, and CXCR3 (Figure 5A-C, $P = NS$), or of inflammatory mediators like interferon β (IFN β) and IFN γ (data not shown), indicating that the shRNA against CCR2 was highly specific and did not display significant off-target effects on the expression levels of other relevant genes. Intriguingly, macrophages of mice that received a transplant of *Ccr2*^{-/-} showed an almost 50% reduction in relative CCR5 mRNA expression levels ($P = .04$).

Discussion

Transplantation of transgenic or knockout bone marrow to wild-type recipients is a widely used technique to determine the role of a particular leukocyte gene in several inflammatory disorders, such as arthritis¹ and atherosclerosis.²⁻⁴ Unfortunately, the time-consuming generation of transgenic or knockout mice and the subsequent backcrossing to an appropriate disease background for analysis of gene function considerably delays progression in gene function research. In this study, we demonstrate the high potential of a novel lentiviral gene silencing approach on bone marrow cells for the in situ generation of leukocyte knockdowns of the aimed target gene. Targeted gene silencing by means of shRNA has, since its discovery, proven to be a powerful tool in the analysis of gene function in vitro as well as in vivo as it can result in prolonged selective inhibition of target gene expression even in eukaryotic cells.²¹⁻²⁵ To deliver shRNA and warrant prolonged shRNA expression, we made use of third-generation self-inactivating lentiviruses,^{12,13} which have been demonstrated to efficiently infect and integrate their genome into a range of mitotic and nonmitotic cells.^{14,15,32} Although rather efficient retroviral transduction of stem




Figure 4. Lentiviral shCCR2 results in reduced macrophage influx due to reduced CCR2 expression. (A) Macrophage recruitment to the peritoneum in mice that underwent transplantation with H1.shCCR2 lentivirus-transduced bone marrow is significantly reduced compared with those that received H1.Empty lentivirus-transduced bone marrow. Recruitment levels were found to be similar to control mice that underwent transplantation with *Ccr2*^{-/-} bone marrow ($*P < .05$, $**P < .01$). (B) CCR2 mRNA expression level in peritoneal macrophages of mice that underwent transplantation with H1.shCCR2-transduced bone marrow was reduced 7 weeks after transplantation ($**P < .01$, $***P < .001$ compared with H1.Empty; $\#P < .01$ compared with H1.shCCR2). (C) *Ccr2* genotyping of transplanted C57Bl/6 mice, revealing a 360-base pair amplicon of wild-type *Ccr2* in mice that underwent transplantation with H1.Empty- or H1.shCCR2 lentivirus-transduced bone marrow and the 290-base pair amplicon was observed in mice that underwent transplantation with *Ccr2*^{-/-} bone marrow. (D) Percent male bone marrow in bone marrow cell lysates from recipient mice, determined by PCR analysis on the *Sry* gene, located on the Y-chromosome. Error bars in panels A, B, and C represent SEM.




Figure 5. Absence of nonspecific effects on related chemokine receptors. No nonspecific effects were observed on peritoneal macrophage mRNA levels of other chemokine receptors, as CCR3 (A), CCR5 (B), and CXCR3 (C) in mice that underwent transplantation with H1.shCCR2 lentivirus-transduced bone marrow compared with the H1.Empty controls. *Ccr2*^{-/-} bone marrow transplantation led to a significant decrease in macrophage CCR5 expression ($P = .04$). Δ Ct indicates delta cycle threshold.

cells of up to 64% has been reported,⁸⁻¹¹ this required extensive manipulation of bone marrow stem cells to increase their mitotic capacity, possibly compromising the hematopoietic potential.⁴⁰ Efficient lentiviral infection requires only a short in vitro incubation period and preventing undesired differentiation of the hematopoietic stem cells.

In this study, the transduction of bone marrow cells with GFP lentivirus was almost quantitative (80%-90%) when using titers of 10 moi to 20 moi. Transduction levels in our study exceeded reported transduction efficiencies for human bone marrow-derived CD34⁺ stem cells⁴¹⁻⁴³ and even were sufficiently high to perform bone marrow transplantation without prior selection of infected cells. Generally, 2 or more consecutive transduction steps were required to achieve similarly high levels, although Banerjee et al, for example, communicated on efficient gene transfer in a single-step protocol.⁴⁴ Our protocol is a compromise of a limited culture period of bone marrow cells in growth factor-free medium (24 hours) on the one hand, preventing undesired differentiation or maturation of the bone marrow cells, and lowest possible virus titer on the other. Furthermore, a single transduction protocol will also minimize the risk of multiple, potentially harmful viral integrants per cell due to, among others, insertional mutagenesis.⁴⁵ However, it should be noted that in the case of shRNAs, multiple integrations per host cell may even be beneficial, as multiple integrations will be accompanied by higher transcript expression, leading to more efficient silencing of the target gene. In our study, the initial infection levels of the bone marrow cells by the H1.Empty and the H1.shCCR2 lentivirus were shown to be similar to that of GFP lentivirus-transduced cells.

As a model gene to illustrate the potential of lentiviral gene silencing in vivo, we have chosen the CC-chemokine receptor 2 (*Ccr2*). CCR2 is critically involved in monocyte recruitment to sites of inflammation in response to an inflammatory stimulus⁴⁶ during atherosclerosis, renal fibrosis,⁴⁷ or pulmonary inflammation.⁴⁸ An shRNA sequence, targeting nucleotides 867 to 885 of the murine *Ccr2* gene, reduced CCR2 mRNA levels by 94% and protein levels in HEK 293 cells to an almost similar extent. Furthermore, the shCCR2 construct completely prevented CCR2 signal transduction in response to its natural ligand JE, as revealed by intracellular calcium release measurements.

Interestingly, transplantation of bone marrow cells transduced with a lentivirus carrying the shCCR2 construct strongly attenuated thioglycolate-elicited, CCR2-dependent migration of macrophages into the peritoneal cavity. The amount of recruited peritoneal macrophages in mice that underwent transplantation with H1.shCCR2 lentivirus-transduced bone marrow was comparable to that of mice that received *Ccr2*^{-/-} bone marrow transplants. Macrophage recruitment levels of the H1.Empty and H1.shCCR2 lentivirus group were essentially similar to values described by Kurihara et al⁴⁶ for wild-type and *Ccr2*^{-/-} mice, respectively, indicating that local macrophage recruitment and proliferation was not affected by lentiviral transduction of the bone marrow per se. Peritoneal macrophages derived from the H1.shCCR2-transduced

bone marrow cells displayed a 70% reduction in CCR2 mRNA content compared with the H1.Empty group, which apparently translated in the observed *Ccr2* knockout phenotype. In the mice that underwent transplantation with *Ccr2*^{-/-} bone marrow, only approximately 7% of circulating blood cells originated from the recipient mice, which accounted for the residual CCR2 mRNA expression in blood cells.

The bone marrow transplantation led to a quantitative and persistent repopulation of donor bone marrow as determined by genomic DNA analysis of the Y-chromosomal *Sry* gene. Moreover, insertion of the target gene apparently did not result in a shifted differentiation pattern of hematopoietic stem cells. At the endpoint of the study, viral copy number in the bone marrow from recipient mice was essentially similar to that of the initial transduction level, implying that the lentivirus-infected cells are not outcompeted by residual or nontransduced cells and that the expansive capacity of bone marrow cells is not affected by lentiviral transduction. We realize that these data were all acquired at 7 weeks after transplantation and that more long-term studies are required on the effects and persistence of lentivirus-transduced bone marrow; however, we are confident that the results will not be very different.

In line with the in vitro data, the obtained reduction in CCR2 expression in vivo was sufficient for a complete and specific knockdown of CCR2 function. Importantly, IFN β and IFN γ mRNA levels did not differ, indicating that the shRNA lentivirus-transduced bone marrow did not elicit an inflammatory response after transplantation.⁴⁹ While CCR2 silencing by shCCR2 was not accompanied by an altered expression pattern of other chemokine receptors, the peritoneal macrophages isolated from mice that underwent transplantation with *Ccr2*^{-/-} displayed a significant 50% reduction in CCR5 mRNA levels, pointing toward transcriptional modulation of the flanking *Ccr5* gene by insertion of the PGK-*neo* cassette into the *Ccr2* gene on mouse chromosome 9. Down-regulation of genes downstream of the site of PGK-*neo* cassette insertion has been reported previously^{50,51} and can be circumvented when using the shRNA approach.

In conclusion, here we show effective delivery of lentiviral shRNA to bone marrow cells and subsequent transplantation into irradiated recipient mice as a strategy to generate hematopoietic knockdowns with silenced CCR2 expression. shRNA-mediated silencing of gene expression persisted for at least 7 weeks after transplantation and led to a complete loss of CCR2 function without any noticeable side effects. Speed and efficiency render this strategy very valuable for addressing the role of other hematopoietic, and in particular leukocyte, genes in inflammatory disorders.

Acknowledgments

The authors would like to thank Miranda Stitzinger and Saskia de Jager from the Division of Biopharmaceutics and Hans de Bont from the Division of Toxicology (Leiden Amsterdam Center for Drug Research, Leiden University) for technical assistance.

References

- Van Bekkum DW. Experimental basis of hematopoietic stem cell transplantation for treatment of autoimmune diseases. *J Leukoc Biol*. 2002;72:609-620.
- Linton MF, Fazio S. Macrophages, lipoprotein metabolism, and atherosclerosis: insights from murine bone marrow transplantation studies. *Curr Opin Lipidol*. 1999;10:97-105.
- Boisvert WA, Spangenberg J, Curtiss LK. Treatment of severe hypercholesterolemia in apolipoprotein E-deficient mice by bone marrow transplantation. *J Clin Invest*. 1995;96:1118-1124.
- Herijgers N, Van Eck M, Groot PH, Hoogerbrugge PM, Van Berkel TJ. Effect of bone marrow transplantation on lipoprotein metabolism and atherosclerosis in LDL receptor-knockout mice. *Arterioscler Thromb Vasc Biol*. 1997;17:1995-2003.
- Toomey JR, Kratzer KE, Lasky NM, Stanton JJ, Broze GJ. Targeted disruption of the murine tissue factor gene results in embryonic lethality. *Blood*. 1996;88:1583-1587.
- Carmeliet P, Ferreira V, Breier G, et al. Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. *Nature*. 1996;383:73-75.
- Kohn DB, Weinberg KI, Nolte JA, et al. Engraftment of gene-modified umbilical cord blood cells in neonates with adenosine deaminase deficiency. *Nat Med*. 1995;1:1017-1023.
- Bjorgvinsdottir H, Bryder D, Sitnicka E, et al. Efficient oncoretroviral transduction of extended long-term culture-initiating cells and NOD/SCID repopulating cells: enhanced reconstitution with gene-marked cells through an ex vivo expansion approach. *Hum Gene Ther*. 2002;13:1061-1073.
- Dao MA, Taylor N, Nolte JA. Reduction in levels of the cyclin-dependent kinase inhibitor p27(kip-1) coupled with transforming growth factor beta neutralization induces cell-cycle entry and increases retroviral transduction of primitive human hematopoietic cells. *Proc Natl Acad Sci U S A*. 1998;95:13006-130011.
- Dao MA, Hashino K, Kato I, Nolte JA. Adhesion to fibronectin maintains regenerative capacity during ex vivo culture and transduction of human hematopoietic stem and progenitor cells. *Blood*. 1998;92:4612-4621.
- Abonour R, Williams DA, Einhorn L, et al. Efficient retrovirus-mediated transfer of the multidrug resistance 1 gene into autologous human long-term repopulating hematopoietic stem cells. *Nat Med*. 2000;6:652-658.
- Dull T, Zufferey R, Kelly M, et al. A third-generation lentivirus vector with a conditional packaging system. *J Virol*. 1998;72:8463-8471.
- Zufferey R, Dull T, Mandel RJ, et al. Self-inactivating lentivirus vector for safe and efficient in vivo gene delivery. *J Virol*. 1998;72:9873-9889.
- Seppen J, Barry SC, Klinkspoor JH, et al. Apical gene transfer into quiescent human and canine polarized intestinal epithelial cells by lentivirus vectors. *J Virol*. 2000;74:7642-7645.
- Seppen J, Rijnberg M, Cooreman MP, Oude Elferink RPJ. Lentiviral vectors for efficient transduction of isolated primary quiescent hepatocytes. *J Hepatol*. 2002;36:459-465.
- Levasseur DN, Ryan TM, Pawlik KM, Townes TM. Correction of a mouse model of sickle cell disease: lentiviral/antisickling β -globin gene transduction of unmobilized, purified hematopoietic stem cells. *Blood*. 2003;102:4312-4319.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*. 1998;391:806-811.
- Cogoni C, Macino G. Post-transcriptional gene silencing across kingdoms. *Curr Opin Genet Dev*. 2000;10:638-643.
- Plasterk RH, Ketting RF. The silencing of the genes. *Curr Opin Genet Dev*. 2000;10:562-567.
- Bass BL. RNA interference, the short answer. *Nature*. 2001;411:428-429.
- Hammond SM, Caudy AA, Hannon GJ. Post-transcriptional gene silencing by double-stranded RNA. *Nat Rev Genet*. 2001;2:110-119.
- Matzke MA, Matzke AJM, Pruss GJ, Vance VB. RNA-based silencing strategies in plants. *Curr Opin Genet Dev*. 2001;11:221-227.
- Sharp PA. RNA interference. *Genes Dev*. 2001;15:485-490.
- Waterhouse PM, Wang MB, Lough T. Gene silencing as an adaptive defence against viruses. *Nature*. 2001;411:834-842.
- Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T. Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature*. 2001;411:494-498.
- Novina CD, Murray MF, Dykxhoorn DM, et al. siRNA-directed inhibition of HIV-1 infection. *Nat Med*. 2002;8:681-686.
- Jacque JM, Triques K, Stevenson M. Modulation of HIV-1 replication by RNA interference. *Nature*. 2002;418:435-438.
- Brummelkamp TR, Bernards R, Agami R. A system for stable expression of short interfering RNAs in mammalian cells. *Science*. 2002;296:550-553.
- Tiscornia G, Singer O, Ikawa M, Verma IM. A general method for gene knockdown in mice using lentiviral vectors expressing small interfering RNA. *Proc Natl Acad Sci U S A*. 2003;100:1844-1848.
- Qin XF, An DS, Chen IS, Baltimore D. Inhibiting HIV-1 infection in human T cells by lentiviral-mediated delivery of small interfering RNA against CCR5. *Proc Natl Acad Sci U S A*. 2003;100:183-188.
- An DS, Xie Y, Mao SH, Morizono K, Kung SK, Chen IS. Efficient lentiviral vectors for short hairpin RNA delivery into human cells. *Hum Gene Ther*. 2003;14:1207-1212.
- Scherr M, Battmer K, Dallmann I, Ganser A, Eder M. Inhibition of GM-CSF receptor function by stable RNA interference in a NOD/SCID mouse hematopoietic stem cell transplantation model. *Oligonucleotides*. 2003;13:353-363.
- Boring L, Gosling J, Cleary M, Charo IF. Decreased lesion formation in *Ccr2*^{-/-} mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature*. 1998;394:894-897.
- Dawson TC, Kuziel WA, Osahar TA, Maeda N. Absence of CC chemokine receptor-2 reduces atherosclerosis in apolipoprotein E-deficient mice. *Atherosclerosis*. 1999;143:205-211.
- Guo J, Van Eck M, Twisk J, et al. Transplantation of monocyte CC-chemokine receptor 2-deficient bone marrow into ApoE3-Leiden mice inhibits atherogenesis. *Arterioscler Thromb Vasc Biol*. 2003;23:447-453.
- Nishitsuji H, Ikeda T, Miyoshi H, Ohashi T, Kanagaki M, Masuda T. Expression of small hairpin RNA by lentivirus-based vector confers efficient and stable gene-suppression of HIV-1 on human cells including primary non-dividing cells. *Microbes Infect*. 2004;6:76-85.
- 't Hoen PA, Van der Lans CA, Van Eck M, Bijstervosch MK, Van Berkel TJ, Twisk J. Aorta of ApoE-deficient mice responds to atherogenic stimuli by a prelesional increase and subsequent decrease in the expression of antioxidant enzymes. *Circ Res*. 2003;93:262-269.
- Zufferey R, Nagy D, Mandel RJ, Naldini L, Trono D. Multiply attenuated lentiviral vector achieves efficient gene delivery in vivo. *Nat Biotechnol*. 1997;15:871-875.
- Sastry L, Johnson T, Hobson MJ, Smucker B, Cornetta K. Titering lentiviral vectors: comparison of DNA, RNA and marker expression methods. *Gene Ther*. 2002;9:1155-1162.
- Srouf EF, Brandt JE, Briddell RA, Grigsby S, Leemhuis T, Hoffman R. Long-term generation and expansion of human primitive hematopoietic progenitor cells in vitro. *Blood*. 1993;81:661-669.
- Salmon P, Kindler V, Ducey O, Chapuis B, Zuber RH, Trono D. High-level transgene expression in human hematopoietic progenitors and differentiated blood lineages after transduction with improved lentiviral vectors. *Blood*. 2000;96:3392-3398.
- Schomber T, Kalberer CP, Wodnar-Filipowicz A, Skoda RC. Gene silencing by lentivirus-mediated delivery of siRNA in human CD34⁺ cells. *Blood*. 2004;103:4511-4513.
- Case SS, Price MA, Jordan CT, et al. Stable transduction of quiescent CD34⁺CD38⁻ human hematopoietic cells by HIV-1-based lentiviral vectors. *Proc Natl Acad Sci U S A*. 1999;96:2988-2993.
- Banerjee A, Li MJ, Bauer G, et al. Inhibition of HIV-1 by lentiviral vector-transduced siRNAs in T lymphocytes differentiated in SCID-hu mice and CD34⁺ progenitor cell-derived macrophages. *Mol Ther*. 2003;8:62-71.
- Woods N-B, Muessig A, Schmidt M, et al. Lentiviral vector transduction of NOD/SCID repopulating cells results in multiple vector integrations per transduced cell: risk of insertional mutagenesis. *Blood*. 2003;101:1284-1289.
- Kurihara T, Warr G, Loy J, Bravo R. Defects in macrophage recruitment and host defense in mice lacking the CCR2 chemokine receptor. *J Exp Med*. 1997;186:1757-1762.
- Wada T, Furuichi K, Sakai N, et al. Gene therapy via blockade of monocyte chemoattractant protein-1 for renal fibrosis. *J Am Soc Nephrol*. 2004;15:940-948.
- Maus U, Waelsch K, Kuziel WA, et al. Monocytes are potent facilitators of alveolar neutrophil emigration during lung inflammation: role of the CCL2-CCR2 axis. *J Immunol*. 2003;170:3273-3278.
- Bridge AJ, Pebernard S, Ducaux A, Nicolaz, A-L, Iggo R. Induction of an interferon response by RNAi vectors in mammalian cells. *Nat Genet*. 2003;34:263-264.
- Pham CT, MacIvor DM, Hug BA, Heusel JW, Ley TJ. Long-range disruption of gene expression by a selectable marker cassette. *Proc Natl Acad Sci U S A*. 1996;93:13090-13095.
- Seidl KJ, Manis JP, Bottaro A, et al. Position-dependent inhibition of class-switch recombination by PGK-neo cassettes inserted into the immunoglobulin heavy chain constant region locus. *Proc Natl Acad Sci U S A*. 1999;96:3000-3005.