



## UvA-DARE (Digital Academic Repository)

### Chronic exposure of the oligochaete *Lumbriculus variegatus* to Polycyclic Aromatic Compounds (PACs): Bioavailability and effects on reproduction

León Paumen, M.; Stol, P.; ter Laak, T.L.; Kraak, M.H.S.; van Gestel, C.A.M.; Admiraal, W.

**DOI**

[10.1021/es702500t](https://doi.org/10.1021/es702500t)

**Publication date**

2008

**Document Version**

Final published version

**Published in**

Environmental Science and Technology

[Link to publication](#)

**Citation for published version (APA):**

León Paumen, M., Stol, P., ter Laak, T. L., Kraak, M. H. S., van Gestel, C. A. M., & Admiraal, W. (2008). Chronic exposure of the oligochaete *Lumbriculus variegatus* to Polycyclic Aromatic Compounds (PACs): Bioavailability and effects on reproduction. *Environmental Science and Technology*, 42(9), 3434-3440. <https://doi.org/10.1021/es702500t>

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

*UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)*

# Chronic Exposure of the Oligochaete *Lumbriculus variegatus* to Polycyclic Aromatic Compounds (PACs): Bioavailability and Effects on Reproduction

MIRIAM LEÓN PAUMEN,<sup>\*,†</sup> PAUL STOL,<sup>†</sup>  
THOMAS L. TER LAAK,<sup>‡</sup>  
MICHIEL H. S. KRAAK,<sup>†</sup>  
CORNELIUS A. M. VAN GESTEL,<sup>§</sup> AND  
WIM ADMIRAAL<sup>†</sup>

*Aquatic Ecology and Ecotoxicology, University of Amsterdam, Kruislaan 320, Amsterdam, Noord Holland, 1098SM, The Netherlands, University of Utrecht, P.O. Box 80125, 3508 TC Utrecht, The Netherlands, and VU University Amsterdam, De Boelelaan 1105, 1081 HV Amsterdam, The Netherlands*

Received October 08, 2007. Revised manuscript received January 22, 2008. Accepted February 06, 2008.

This study aimed to monitor PAC availability to the oligochaete *Lumbriculus variegatus* during 28 days of exposure to spiked sediments, in order to obtain reliable chronic effect concentrations for reproduction. Sediment toxicity tests were performed using three pairs of PAC isomers: two homocyclic compounds (anthracene and phenanthrene), two azaarenes (acridine and phenanthridine), and the two main transformation products of the azaarenes (acridone and phenanthridone). During the experiment, available PAC concentrations in pore water (estimated using solid phase microextraction) decreased more than total PAC concentrations in the sediment. Relating effect concentrations to PAC concentrations in pore water and in organisms showed that the two homocyclic compounds caused narcotic effects during chronic exposure, but only one of the four tested heterocyclic PACs caused narcotic effects. The transformation product phenanthridone was not toxic at the tested concentrations (up to 4000  $\mu\text{mol/kg}$  dry sediment), whereas EC<sub>50</sub> values for the parent compound phenanthridine and the isomer acridone were below the estimated limit for narcosis, suggesting a specific mode of action. These results demonstrated the unpredictable (isomer) specific toxicity of azaarenes and their transformation products, emphasizing the need of chronic toxicity testing to gain insight into the long-term effects of heterocyclic PACs, which have been overlooked in risk assessment.

## Introduction

Benthic invertebrates inhabiting PAC contaminated sediments are chronically exposed to a variety of homocyclic and heterocyclic compounds (1, 2). Risk assessment for PACs, however, is based on only a limited set of homocyclic

compounds, ignoring the vast number of heterocyclic compounds (with in- or on-ring substitutions) and transformation products (3). Moreover, in the past PAC toxicity has been commonly assessed in short-term high-dose experiments, in which mortality is often the only end point (4). During such acute exposures PACs act mainly by narcosis (5), but during chronic exposure the same compounds may exert sublethal effects (on, e.g., growth, reproduction) (6, 7). Therefore, the present study focused on assessing chronic effects of azaarenes and their transformation products in addition to homocyclic compounds. Azaarenes are mainly formed and released into the environment by incomplete combustion of fossil fuels, in spills or effluents of several industrial activities, and by pesticide use (8). Azaarenes, containing one in-ring nitrogen substitution, can comprise up to 10% of the total PAC concentration of contaminated sites (9), but toxicity data for this group of compounds are scarce, and very little is known about their chronic effects (6, 7).

Sorption to sediment organic matter may decrease availability of PACs to benthic invertebrates, since tissue residues and effects correlate to dissolved pore water concentrations (10). Nevertheless, in chronic toxicity tests PAC bioavailability during exposure is rarely monitored, although changes in bioavailability could alter the outcome of the experiments (11). Recently, several techniques have been developed to facilitate quantification of the freely dissolved (pore) water concentrations of toxicants (12, 13), including solid-phase microextraction (SPME) (14). The poly(dimethylsiloxane) coating on the SPME fiber mimics the structure and sorptive properties of membranes (15), allowing a reliable estimation of the availability of organic compounds.

The aim of the present study was, therefore, to monitor PAC availability to the Oligochaete *Lumbriculus variegatus* during chronic exposure, in order to obtain reliable effect concentrations for reproduction. Asexual reproduction of *L. variegatus* is a sensitive chronic end point that can easily be combined with SPME and PAC accumulation measurements in the worms' body in a single test. This allowed us to express the observed effects as a function of measured PAC concentrations in sediment, estimated PAC concentrations in pore water, and PAC concentrations in the organism. Since it has been demonstrated that slight differences in chemical structure may result in substantial differences in toxicity (16–18), three three-ring PAC isomer pairs were selected: two homocyclic compounds, two azaarenes, and the two main stable transformation products of the azaarenes.

## Materials And Methods

**Test Organism.** *Lumbriculus variegatus* is an endobenthic oligochaete, widely used in bioaccumulation experiments (11, 19–21). In the laboratory it reproduces via fragmentation, and a standardized protocol to determine effects on asexual reproduction of *L. variegatus* has been developed by the Organisation for Economic Co-operation and Development (OECD). This protocol, with slight modifications, was applied in the present study.

**Test Compounds.** Six test compounds were chosen: two three-ring homocyclic compounds, anthracene and phenanthrene; two azaarene analogues, acridine and phenanthridine, and two phase I azaarene transformation products, acridone and phenanthridone. Several properties of the compounds are listed in Table S1 in the Supporting Information. All compounds were provided by Sigma-Aldrich Chemie BV (Zwijndrecht, The Netherlands).

\* Corresponding author fax: 0031205257716; e-mail: mleon@science.uva.nl.

<sup>†</sup> University of Amsterdam.

<sup>‡</sup> University of Utrecht.

<sup>§</sup> VU University Amsterdam.

**Sediment Spiking.** Drontermeer sediment, a Dutch reference sediment (12–14% organic matter, 15.5% clay, and all individual PAH concentrations below 0.01 mg/kg DW), was used for the toxicity tests. Sediment was provided by Grontmij/AquaSense (Amsterdam, The Netherlands). After sampling, it was homogenized and frozen to eliminate indigenous fauna (−20 °C). Three days before the spiking procedure was started the sediment was thawed at 4 °C.

The sediment was spiked to obtain the following nominal PAC concentrations: 67, 140, 280, 561, and 1122  $\mu\text{mol/kg DW}$  for anthracene, 140, 280, 561, 1122, and 2244  $\mu\text{mol/kg DW}$  for phenanthrene, 139, 279, 558, 1116, and 2232  $\mu\text{mol/kg DW}$  for acridine and phenanthridine, and 154, 318, 640, 1281, and 2561  $\mu\text{mol/kg DW}$  for acridone and phenanthridone.

PACs were spiked to dry sediment (10% of the total amount). Acetone was used as carrier solvent, and for each PAC concentration series equal volumes of acetone were added to all treatments. The spiking was carried out in the dark to prevent photodegradation of the compounds. Controls and solvent controls were included. The homocyclic compounds and the azaarenes were dissolved in 50 mL of acetone and added to 38 g of dry sediment, and the mixture was left in a closed 500 mL glass bottle for 24 h to equilibrate. The transformation products, less soluble in acetone, were dissolved in 300 mL of acetone. After the 24 h equilibration period, the bottles were opened and left for 24 h in a fume hood to allow the acetone to evaporate. Next, the spiked dry sediment was mixed with 800 mL of Dutch Standard Water (DSW, deionized water with 200 mg/L  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 180 mg/L  $\text{MgSO}_4 \cdot \text{H}_2\text{O}$ , 100 mg/L  $\text{NaHCO}_3$ , and 20 mg/L  $\text{KHCO}_3$ ; pH  $\approx$  8.2) and 432 g of wet sediment (20% water content) in 3-L glass bottles. A 20:1 wt/wt mixture of Trouvit and TetraPhyll, diluted in distilled water, was used as additional food source (0.25 mg/worm/day). The glass bottles filled with the water–sediment–food mixture were placed on a roller bank (30 rpm) for 24 h to homogenize. Next, the mixture was divided into eight replicate 200 mL glass jars and allowed to settle for 10 days. Each test replicate consisted of approximately 60 g of wet sediment and 100 mL of DSW. Four replicates per concentration were sacrificed during the experiment for PAC measurements in sediment and pore water, leaving four replicates for the toxicity test.

**Toxicity Tests.** Four-week (28-day) toxicity tests were performed in an incubator at a constant temperature of  $20 \pm 1$  °C, with mercury lamps (light intensity approximately 50  $\mu\text{mol quanta/m}^2/\text{s}$ ) providing a light regime of 16 h of light and 8 h of darkness. UV filters were used to minimize photodegradation of the test compounds by UV–B radiation. Glass jars were covered with plastic foil and constantly aerated to keep oxygen concentrations stable. Once a week, deionized water was added to compensate for evaporation of the test solution. pH in the test jars ranged from 7.5 to 8.5 and oxygen concentration was always above 75% air saturation.

At the beginning of the experiment, 20 *L. variegatus* adult individuals from our laboratory culture were introduced into each of the test jars using a plastic Pasteur pipet. The worms were allowed to burrow into the sediment for 4 h before aeration of the jars was restarted. After 28 days, the sediment was sieved to extract the worms from the sediment. The number of worms in each test jar was determined, and the average reproductive output per concentration (%) was calculated using the formula  $Y = ((X - 20)/20) \times 100$ , where *X* is the number of worms per jar at the end of the test. The worms were kept for four hours in clean tap water for gut clearance. This short gut clearance period was applied to prevent depletion of the heterocyclic compounds (22). Following, worm samples were frozen at −20 °C until analysis.

**PAC Availability.** Availability of anthracene, phenanthrene, acridine, and phenanthridine in sediment pore water was measured using solid-phase microextraction (SPME).

Due to their low hydrophobicity, SPME could not be applied to the transformation products. The fibers, obtained from the Institute for Risk Assessment Sciences (IRAS, University of Utrecht, The Netherlands), had a diameter of 110  $\mu\text{m}$  and a 28.5  $\mu\text{m}$  thick poly(dimethylsiloxane) (PDMS) coating, and were manufactured by Poly Micro Industries (Phoenix, AZ). Three-centimeter pieces of fiber were glued to a 0.4 mm  $\varnothing$  nylon fishing thread, washed with 70% methanol and deionized water and carefully introduced into the test sediment. Two replicate fibers were introduced into each test jar. Fiber measurements were performed in two jars per test concentration. The fibers remained in the sediment for three days, sufficient time for the fiber to reach equilibrium with the PCA concentrations in the pore water (23, 24). Next, they were gently cleaned with a tissue, and a 2-cm fiber piece was cut to get rid of the glued tip of the fiber and extracted in 300  $\mu\text{L}$  of acetonitrile. For each toxicity test, fibers were introduced into the sediment at the beginning and at days 3, 6, 9, 13, 17, 21, and 25 of the experiment. PAC concentrations in the fibers were measured by high-performance liquid chromatography (HPLC), using a 1 mL/min 80% acetonitrile/20% water flow rate and an injection volume of 20  $\mu\text{L}$ . For the homocyclic compounds, fiber–water partition coefficients ( $K_f$ ) from Ter Laak et al. (24) were used to estimate concentrations in the pore water using the SPME measurements. For the azaarenes, static system experiments were performed to determine  $K_f$  values, as in ref (25).

**PAC Concentrations in Sediment, Pore Water, and Oligochaetes.** Sediment and pore water samples were collected at the beginning, after 7 and 14 days, and at the end of the experiment. For sediment samples, 40 mL was centrifuged for 15 min at 3000 rpm. To obtain a clear pore water sample, the supernatant was collected and centrifuged again for 15 min at 3000 rpm. Acetonitrile (150  $\mu\text{L}$ ) was added to two replicated 1-mL pore water samples. Sediment and pore water samples were frozen at −20 °C until analysis.

Ten grams of centrifuged sediment (water content approximately 10%) was mixed with equal amounts of anhydrous  $\text{NaSO}_4$  (Merck, Darmstadt, Germany), and Soxhlet extracted for 5 h in cellulose extraction thimbles. Hexane (Chromasolv, Sigma-Aldrich Chemie BV) was used as extraction solvent for the homocyclic compounds and the azaarenes, and acetonitrile (HPLC-grade, Biosolve BV) was used for the transformation products. The hexane-extracted samples were transferred into acetonitrile by blowing off the hexane using a gentle stream of nitrogen. All extracts were analyzed by HPLC. Recovery checks were performed for each of the compounds to validate the Soxhlet extraction procedure. For this purpose, known amounts of spiking solution in the same concentration range as tested in the experiments were added to clean sediment in the extraction thimble. Extraction recoveries were 80% for anthracene, 85% for phenanthrene, 75% for acridine, 77% for phenanthridine, and 88% for the transformation products. PAC concentrations in the sediment were corrected for recovery. Pore water samples were directly injected in the HPLC, to prevent overestimation of freely dissolved pore water concentrations due to exhaustive extraction of the DOM bound fraction of the compounds (25).

*L. variegatus* tissue was sampled at the end of the experiment (28 days). Worms from the 4 replicates per concentration were pooled in order to obtain enough biomass for a reliable PAC measurement. PACs were extracted from the tissue with acetonitrile. One hundred milligrams of wet tissue was mixed with 1.5 mL of a 1%  $\text{CaCl}_2$  solution and homogenized to break down the tissue. A 3-mL portion of acetonitrile was added, and the mixture was sonicated for 45 min in cool water (approximately 12 °C). Next, the PACs were extracted from the tissue at 4 °C for 24 h. After this first extraction, the sample was centrifuged for 15 min at 3000




FIGURE 1. Relationship between polycyclic aromatic compound (PAC) concentrations in pore water estimated using SPME measurements (Y-axis,  $\mu\text{M}$ ) and measured experimentally (X-axis,  $\mu\text{M}$ ) in spiked Drontermeer sediment during 28-day toxicity tests with *Lumbriculus variegatus*. Dashed line represents 1:1 relationship.

rpm and the supernatant was transferred to a HPLC vial. Acetonitrile (3 mL) was added to the pellet, the sample was homogenized, and the extraction procedure was repeated. The two worm tissue extracts were pooled and measured using HPLC. PAC concentrations in the tissue were corrected for recovery. Extraction recoveries were 70% for anthracene, 72% for phenanthrene, 65% for acridine, 67% for phenanthridine, 68% for acridone, and 72% for phenanthridone.

**Statistical Analysis.** The concentration of the test compounds in sediment, pore water, and worm tissue that caused 50% reduction in reproduction compared to the control was calculated according to Haanstra et al. (26). Average measured PAC concentrations in the sediment, average estimated PAC concentrations in the pore water, and PAC concentrations in the worms at the end of the 28-day exposure period were used. *t* Tests ( $p < 0.05$ ) were performed to compare control and acetone control reproductive outputs. Likelihood ratio tests were performed to compare effect concentrations ( $X^2_1 > 3.84$ ,  $p < 0.05$ ). Statistical analysis was performed using SPSS 11.0 for Windows.

## Results

**PAC Concentrations in Sediment, Pore Water, and *L. variegatus*.** Measured PAC concentrations in the sediment are shown in Table S2. Concentrations of anthracene, acridine, and acridone in the sediment did not decrease substantially during the experiment (respectively,  $3 \pm 2$ ,  $3 \pm 3$ , and 0%). In contrast, concentrations of phenanthrene, phenanthridine, and phenanthridone in the sediment decreased during the experiment, mainly due to microbial degradation (27, 28). Average decreases were  $19 \pm 10\%$  for

phenanthrene,  $30 \pm 16\%$  for phenanthridine, and  $24 \pm 12\%$  for phenanthridone.

PAC concentrations in pore water were estimated using the concentrations in SPME fibers and PDMS-water partitioning coefficients ( $K_i$ , Table S1). In Figure 1, estimated concentrations (Y-axis) were plotted against directly measured PAC concentrations in pore water (X-axis), and a linear regression was performed in order to evaluate the performance of the SPME technique. The  $r^2$  values of the linear regression for anthracene, phenanthrene, and phenanthridine were respectively 0.77, 0.92, and 0.97, showing a consistent linear relationship between measured and estimated PAC concentrations in pore water. For acridine, the  $r^2$  value was 0.64, due to variability in the results obtained with SPME. If direct measurements in pore water would be equivalent to concentrations in pore water estimated using SPME, the slope of the regression line should be 1. This was indeed the case for phenanthrene (1.07, 95% C.I. 0.89–1.19), but for anthracene the slope was significantly higher than 1 (2.43, 95% C.I. 1.85–3.02), meaning that the measured anthracene concentrations in the pore water were lower than estimated ones. For the two azaarenes, acridine and phenanthridine, the slope of the regression line was significantly lower than 1 (respectively 0.54 (95% C.I. 0.32–0.75) and 0.72 (95% C.I. 0.60–0.84)), meaning that estimated concentrations were somewhat lower than concentrations measured in pore water.

During exposure, estimated pore water concentrations of all PACs decreased at all concentrations in the sediment, due to increased sediment sorption in time and degradation by bacteria present in the sediment (28, 29) (Tables S3, S4,

**TABLE 1. EC50 and EC10 values (With Corresponding 95% Confidence Intervals) for the Effects of Polycyclic Aromatic Compounds (PACs) on the Reproduction of *Lumbriculus variegatus* in Spiked Drontermeer Sediments, Calculated Using Average Measured PAC Concentrations in the Sediment ( $\mu\text{mol/kg DW}$ ), Pore Water PAC Concentrations Estimated Using SPME ( $\mu\text{M}$ ), and PAC Concentrations in *L. variegatus* at the End of the 28-Day Exposure Period ( $\mu\text{mol/kg ww}$ )**

|                         |                | sediment<br>effect concentration<br>( $\mu\text{mol/kg DW}$ ) with 95% C.I. |                 | SPME porewater<br>effect concentration<br>( $\mu\text{M}$ ) with 95% C.I. |                   | body residue<br>effect concentration<br>( $\mu\text{mol/kg ww}$ ) with 95% C.I. |                    |
|-------------------------|----------------|---|-----------------|---|-------------------|---|--------------------|
|                         |                | EC50  | EC10            | EC50  | EC10              | ER50  | ER10               |
| homocyclic compounds    | anthracene     | 532<br>(0–1075)   | 17<br>(0–529)   | 0.15<br>(0.064–0.246)   | 0.025<br>(0–0.08) | 116<br>(43–210)   | 111<br>(15.16–207) |
|                         | phenanthrene   | 467<br>(285–650)  | 187<br>(17–358) | 0.299<br>(0.15–0.45)  | 0.07<br>(0–0.4)   | 88<br>(72–105)  | 84<br>(69–99)      |
| azaarenes               | acridine       | 1248<br>(315–2182)  | 197<br>(0–1076) | 3.97<br>(0.47–7.46)   | 0.45<br>(0–3.9)   | 137<br>(93–181)   | 132<br>(90–173)    |
|                         | phenanthridine | 315<br>(65–565)   | 37<br>(0–272)   | 0.148<br>(0.08–0.22)  | 0.04<br>(0–0.45)  | 2<br>(0.2–2.7)  | 0.67<br>(0–2.21)   |
| transformation products | acridone       | 301<br>(240–363)  | 138<br>(80–196) | 0.46 <sup>a</sup><br>(0.37–0.56)  |                   | 30<br>(25–36)   | 25<br>(20–30)      |
|                         | phenanthridone | >4014   | >4014           | >5.77   | >5.77             | >331  | >331               |

<sup>a</sup> EC50 for acridone calculated as in Droge et al. (6).

S5, S6 Supporting Information). The decrease in estimated pore water concentrations was larger in proportion than the decrease in measured concentrations in the sediment.

In general, average PAC concentrations in the worms' tissue ( $\mu\text{mol PAC/kg wet weight}$ ) after 28 days of exposure increased with increasing PAC concentrations in the sediment, but differences in accumulation among the six tested compounds were found (Table S7).

**Effect Concentrations.** Control reproductive output after 28 days of exposure was on average  $80 \pm 9\%$ . Control and solvent control average reproductive output did not differ significantly ( $t$  test,  $p < 0.05$ ) and therefore solvent control values were used for further analysis of the results. All tested compounds except phenanthridone affected reproduction of *L. variegatus*. In Table 1, 50 and 10% effect concentrations (EC50 and EC10s) with their 95% confidence intervals are shown. Likelihood ratio tests were performed to compare the EC50 values.

**EC50s Based on Measured PAC Concentrations in Sediment.** From the compounds that exerted an effect, acridone was the most toxic (EC50 301  $\mu\text{mol/kg sediment DW}$ ), while its parent compound acridine was the least toxic (EC50 1248  $\mu\text{mol/kg sediment}$ ). Likelihood ratio tests showed that the sediment EC50 for acridine was significantly higher than EC50s for phenanthridine, the homocyclic compound anthracene, and the transformation product acridone. In contrast, EC50s for anthracene, phenanthrene, phenanthridine, and acridone did not differ from each other.

**EC50s Based on PAC Concentrations in Pore Water.** Pore water EC50s were calculated using estimated PAC concentrations in the pore water for the homocyclic compounds and the azaarenes. For acridone, the pore water EC50 was estimated from the sediment EC50 using its sediment–water partitioning coefficient ( $K_{oc}$ ) as in Droge et al. (6). Pore water EC50s ranged from 0.148  $\mu\text{M}$  for phenanthridine to 3.97  $\mu\text{M}$  for acridine, and again likelihood ratio tests showed that the EC50 for acridine was significantly higher than the EC50s for the other compounds. EC50s for anthracene, phenanthrene, phenanthridine, and acridone did not differ significantly.

**EC50s Based on PAC Concentrations in the Organism.** Fifty percent effect body residues (ER50s) based on wet weight were calculated. Likelihood ratio tests showed that the ER50s for the two homocyclic compounds, anthracene and phenanthrene, did not differ significantly (116 and 88  $\mu\text{mol/kg WW}$ , respectively). The ER50 for the azaarene acridine (137  $\mu\text{mol/kg WW}$ ) was similar to the homocyclic compound ER50s. In contrast, phenanthridine showed the significantly lowest


ER50 (2  $\mu\text{mol/kg WW}$ ). The transformation product acridone showed a significantly lower ER50 than the homocyclic compounds and acridine (30  $\mu\text{mol/kg WW}$ ).

## Discussion

**PAC Availability to *L. variegatus*.** To our knowledge, this was the first attempt to use SPME to monitor the change in PAC availability in pore water during chronic toxicity tests. Indeed, the decrease in freely dissolved PAC concentrations in pore water during exposure time was substantial, according to previous observations on progressive sorption of PACs to particles (29, 30). Even more important, the present study showed that the freely dissolved PAC concentrations in pore water decreased more than total PAC concentrations in the sediment, emphasizing the importance of monitoring availability during chronic exposure.

For the two azaarenes, estimated concentrations using SPME were somewhat lower than directly measured concentrations in the pore water. This may have been caused by a slight underestimation of PDMS–water partitioning coefficients ( $K_f$ ) due to depletion in the static experimental system in which  $K_f$  values were determined, as suggested by Poerschmann et al. (31). For anthracene, in contrast, concentrations measured directly in the pore water were 2.4 times lower than estimated pore water concentrations using SPME measurements. Due to the low water solubility of anthracene, binding of anthracene to particulate and dissolved organic carbon in the sediment pore water probably occurred during the pore water analysis (25). As a result, very low concentrations were measured. Therefore, pore water concentrations estimated using SPME measurements were more reliable, and were used to calculate effect concentrations.

**Effect Concentrations. Effects Based on Measured PAC Concentrations in the Sediment (Figure 2A).** Due to equilibrium partitioning, toxic effects of PACs will occur at lower concentrations in the (pore) water with increasing  $\log K_{ow}$  of the test compounds (32). On the other hand, sorption of the test compounds to sediment organic matter increases with increasing  $\log K_{ow}$  (24). Because of these two counteracting influences, sediment EC50s for our test compounds were expected to be more or less constant (33). This was indeed the case for the homocyclic compounds (anthracene and phenanthrene), the azaarenes (acridine and phenanthridine), and the transformation product acridone. The transformation product phenanthridone was the only compound that did not affect reproduction of *L. variegatus* at the tested concentrations (up to 4000  $\mu\text{mol/kg dry sediment}$ ). Sediment



**FIGURE 2.** EC50 values for the effect of polycyclic aromatic compounds (PACs) on reproduction of *Lumbriculus variegatus* in spiked Drontemeer sediment as a function of log $K_{ow}$  (•); based on measured concentrations in sediment ( $\mu\text{mol}/\text{kg DW}$ , plot A), estimated concentrations in pore water ( $\mu\text{M}$ , plot B) and concentrations in *L. variegatus* (ER50s,  $\mu\text{mol}/\text{g WW}$ , plot C). (Plot B) Solid line: linear log $\text{LC}_{50}$ –log $K_{ow}$  relationship for *Chironomus riparius* exposed for 96 h to similar PACs ( $y = -0.8162x + 3.4936$ ,  $r^2 = 0.986$ , (47)). Dashed line: ratio  $\text{LC}_{50}$  from ref (47)/3, lower limit for narcosis. For acridone, the calculated pore water EC50 is shown. (Plot C) Dashed lines: according to CBR predictions, upper and lower limits of the tissue concentration interval after chronic exposure to narcotic chemicals (0.2–0.8  $\mu\text{mol}/\text{g WW}$ ).

EC50s were in the range of effect concentrations determined for other benthic invertebrates exposed to PAH-contaminated sediments (34–36).

*Effects Based on PAC Concentrations in the Pore Water (Figure 2B).* In previous studies (6, 35), estimated chronic pore water effect concentrations for soil and sediment invertebrates were compared to an acute  $\text{LC}_{50}$ –log $K_{ow}$  relationship determined by Bleeker et al. (5), which described the narcotic effects of several PACs to the midge *Chironomus riparius*. This way, effects of homo- and heterocyclic compounds were compared, and specific effects emerging besides narcosis during chronic exposure were identified. Applying this approach to the present data set revealed that pore water EC50s for the homocyclic compounds anthracene and phenanthrene agreed well with the acute  $\text{LC}_{50}$ –log $K_{ow}$  relationship. From the four tested heterocyclic compounds, however, only the EC50 for acridine was in agreement with the acute  $\text{LC}_{50}$ –log $K_{ow}$  relationship. The transformation product phenanthridone did not affect reproduction of *L. variegatus*. Based on its log $K_{ow}$  value an EC50 of about 20  $\mu\text{M}$  was expected, but this concentration in the pore water was not reached at the highest concentration in the sediment. In contrast, the EC50s for the azaarene phenanthridine and the transformation product acridone were clearly below the

concentration corresponding to narcosis, suggesting a specific mode of action.

The specific effects found for the azaarene phenanthridine compared to the narcotic effects of its isomer acridine were observed previously for reproduction of the soil invertebrates *Folsomia candida* and *Enchytraeus crypticus* (6). Moreover, phenanthridine was found to be more teratogenic than acridine to *Xenopus laevis* (37), and more genotoxic in the GFP assay (38). Isomer-specific toxicity was also found for the transformation product isomers: acridone was more toxic than expected based on log $K_{ow}$ , as found previously in a mutagenicity study (18), while phenanthridone did not affect reproduction of *L. variegatus* at all at the highest concentration tested. In contrast, both transformation products affected emergence of *C. riparius* (35), underlining the species-specific sensitivity to PACs (6).

*Effects Based on PAC Concentrations in the Organism (Figure 2C).* Toxic effects can only take place if the toxicant is taken up by the organism. Therefore, we also expressed effects as a function of PAC concentrations in the organism. This allowed us to evaluate the observed deviations from narcosis in a second independent way, using the critical body residue (CBR) concept (39) predicts that nonpolar narcotics (e.g., PACs, PCBs) cause 50% of mortality

at threshold body concentrations between 2 and 8  $\mu\text{mol/g}$  wet weight. This concept has been successfully applied to interpret toxicity data for sediment- and soil-inhabiting organisms (4, 40–43). Because of the unspecificity of the narcotic mode of action, it has been suggested that this threshold would be applicable to all organisms. Furthermore, analyses of chronic toxicity data suggest that threshold body concentrations for chronic exposure would be 1 order of magnitude lower than lethal CBRs (between 0.2 and 0.8  $\mu\text{mol/g}$  wet weight) (34, 39, 43, 44), and body concentrations far below the threshold would indicate specific modes of action of the tested compounds occurring besides narcosis (45). In our study, ER50s for anthracene, phenanthrene, and acridine were around 1 order of magnitude lower than 2–8  $\mu\text{mol/g}$  wet weight, in agreement with the chronic CBR concept. Also, ER50s for the two homocyclic compounds were close (within a factor 2) to body residue predictions for chronic exposure to homocyclic PAHs (43). In contrast, ER50s for phenanthridine and acridone were 2 orders of magnitude lower than the chronic CBR range, suggesting additional specific modes of action of the two compounds. Thus, deviations from narcosis found for the ER50s were in agreement with deviations from narcosis found for the pore water EC50s.

In this study, only one of the four tested heterocyclic PACs caused narcotic effects during chronic exposure, showing that (isomer) specific toxicity of azaarenes and their transformation products could not be predicted from acute effect concentrations. About two-thirds of the approximately four million known organic compounds are heterocyclic (46), and in our study toxicity of three out of the four tested heterocycles was poorly predicted by  $\log K_{ow}$ . These results emphasize the need of applying chronic toxicity testing to gain insight in the long term effects of heterocyclic PACs, which have been overlooked by risk assessment.

### Acknowledgments

This research was supported by the Technology Foundation (STW), applied science division of The Netherlands Organization for Scientific Research (NWO) (project AEB 06364).

### Supporting Information Available

Tables S1–S7. This information is available free of charge via the Internet at <http://pubs.acs.org>.

### Literature Cited

- Uhler, A. D.; Emsbo-Mattingly, S.; Liu, B.; Hall, L. W.; Burton, D. T. An integrated case study for evaluating the impacts of an oil refinery effluent on aquatic biota in the Delaware River: Advanced chemical fingerprinting of PAHs. *Hum. Ecol. Risk Assess.* **2005**, *11*, 771–836.
- Lahr, J.; Maas-Diepeveen, J. L.; Stuijzand, S. C.; Leonards, P. E. G.; Druke, J. M. L. S.; Espeldoorn, A.; Kerkum, L. C. M.; van Stee, L. L. P.; Hendriks, A. J. Responses in sediment bioassays used in the Netherlands: Can observed toxicity be explained by routinely monitored priority pollutants? *Water Res.* **2003**, *37*, 1691–1710.
- Crommentuijn, T.; Sijm, D.; de Bruijn, J.; van Leeuwen, K.; van de Plassche, E. Maximum permissible and negligible concentrations for some organic substances and pesticides. *J. Environ. Manage.* **2000**, *58*, 297–312.
- Lee, J. H.; Landrum, P. F.; Koh, C. H. Toxicokinetics and time-dependent PAH toxicity in the amphipod *Hyalella Azteca*. *Environ. Sci. Technol.* **2002**, *36*, 3124–3130.
- Bleeker, E. A. J.; Pieters, B. J.; Wiegman, S.; Kraak, M. H. S. Comparative (photoenhanced) toxicity of homocyclic and heterocyclic PACs. *Polycyclic Aromat. Compd.* **2002**, *22*, 601–610.
- Droge, S. T. J.; Leon Paumen, M.; Bleeker, E. A. J.; Kraak, M. H. S.; van Gestel, C. A. M. Chronic toxicity of polycyclic aromatic compounds to the springtail *Folsomia candida* and the enchytraeid *Enchytraeus crypticus*. *Environ. Toxicol. Chem.* **2006**, *25*, 2423–2431.
- Feldmannova, M.; Hilscherova, K.; Marsalek, B.; Blaha, L. Effects of N-heterocyclic polyaromatic hydrocarbons on survival, reproduction, and biochemical parameters in *Daphnia magna*. *Environ. Toxicol.* **2006**, *21*, 425–431.
- Kuhn, E. P.; Sufliata, J. M. Microbial degradation of nitrogen, oxygen and sulfur heterocyclic compounds under anaerobic conditions: studies with aquifer samples. *Environ. Toxicol. Chem.* **1989**, *8*, 1149–1158.
- Neilson, A. H. *The Handbook of Environmental Chemistry: PAHs and Related Compounds*; Springer Verlag: Berlin, 1998; Volume 3.
- Moermond, C. T. A.; Roessink, I.; Jonker, M. T. O.; Meijer, T.; Koelmans, A. A. Impact of polychlorinated biphenyl and polycyclic aromatic hydrocarbon sequestration in sediment on bioaccumulation in aquatic food webs. *Environ. Toxicol. Chem.* **2007**, *26*, 607–615.
- Schuler, L. J.; Wheeler, M.; Bailer, A. J.; Lydy, M. J. Toxicokinetics of sediment-sorbed benzo[a]pyrene and hexachlorobiphenyl using the freshwater invertebrates *Hyalella azteca*, *Chironomus tentans*, and *Lumbriculus variegatus*. *Environ. Toxicol. Chem.* **2003**, *22*, 439–449.
- Mayer, P.; Vaes, W. H. J.; Hermens, J. L. M. Absorption of hydrophobic compounds into the poly(dimethylsiloxane) coating of solid-phase microextraction fibers: High partition coefficients and fluorescence microscopy images. *Anal. Chem.* **2000**, *72*, 459–464.
- Leppanen, M. T.; Kukkonen, J. V. K. Evaluating the role of desorption in bioavailability of sediment-associated contaminants using oligochaetes, semipermeable membrane devices and Tenax extraction. *Environ. Pollut.* **2006**, *140*, 150–163.
- King, A. J.; Readman, J. W.; Zhou, J. L. The application of solid-phase micro-extraction (SPME) to the analysis of polycyclic aromatic hydrocarbons (PAHs). *Environ. Geochem. Health* **2003**, *25*, 69–75.
- Leslie, H. A.; Hermens, J. L. M.; Kraak, M. H. S. Baseline toxicity of a chlorobenzene mixture and total body residues measured and estimated with solid-phase microextraction. *Environ. Toxicol. Chem.* **2004**, *23*, 2017–2021.
- Kraak, M. H. S.; Wijnands, P.; Govers, H. A. J.; Admiraal, W.; de Voogt, P. Structural-based differences in ecotoxicity of benzoquinoline isomers to the zebra mussel (*Dreissena polymorpha*). *Environ. Toxicol. Chem.* **1997**, *16*, 2158–2163.
- Wiegman, S.; van Vlaardingen, P. L. A.; Bleeker, E. A. J.; de Voogt, P.; Kraak, M. H. S. Phototoxicity of azaarene isomers to the marine flagellate *Dunaliella tertiolecta*. *Environ. Toxicol. Chem.* **2001**, *20*, 1544–1550.
- Bleeker, E. A. J.; Van der Geest, H. G.; Klamer, H. J. C.; De Voogt, P.; Wind, E.; Kraak, M. H. S. Toxic and genotoxic effects of azaarenes: Isomers and metabolites. *Polycyclic Aromat. Compd.* **1999**, *13*, 191–203.
- Ingersoll, C. G.; Brunson, E. L.; Wang, N.; Dwyer, E. J.; Ankley, G. T.; Mount, D. R.; Huckins, J.; Petty, J.; Landrum, P. E. Uptake and depuration of nonionic organic contaminants from sediment by the oligochaete *Lumbriculus variegatus*. *Environ. Toxicol. Chem.* **2003**, *22*, 872–885.
- Leppanen, M. T.; Kukkonen, J. V. K. Relative importance of ingested sediment and pore water as bioaccumulation routes for pyrene to oligochaete (*Lumbriculus variegatus*, Muller). *Environ. Sci. Technol.* **1998**, *32*, 1503–1508.
- Landrum, P. E.; Leppanen, M.; Robinson, S. D.; Gossiaux, D. C.; Burton, G. A.; Greenberg, M.; Kukkonen, J. V. K.; Eadie, B. J.; Lansing, M. B. Comparing behavioral and chronic endpoints to evaluate the response of *Lumbriculus variegatus* to 3,4,3',4'-tetrachlorobiphenyl sediment exposures. *Environ. Toxicol. Chem.* **2004**, *23*, 187–194.
- Mount, D. R.; Dawson, T. D.; Burkhard, L. P. Implications of gut purging for tissue residues determined in bioaccumulation testing of sediment with *Lumbriculus variegatus*. *Environ. Toxicol. Chem.* **1999**, *18*, 1244–1249.
- Leslie, H. A.; Ter Laak, T. L.; Busser, F. J. M.; Kraak, M. H. S.; Hermens, J. L. M. Bioconcentration of organic chemicals: Is a solid-phase microextraction fiber a good surrogate for biota? *Environ. Sci. Technol.* **2002**, *36*, 5399–5404.
- Ter Laak, T. L.; Barendregt, A.; Hermens, J. L. M. Freely dissolved pore water concentrations and sorption coefficients of PAHs in spiked aged, and field-contaminated soils. *Environ. Sci. Technol.* **2006**, *40*, 2184–2190.
- Ter Laak, T. L.; Durjava, M.; Struijs, J.; Hermens, J. L. M. Solid phase dosing and sampling technique to determine partition coefficients of hydrophobic chemicals in complex matrices. *Environ. Sci. Technol.* **2005**, *39*, 3736–3742.

- (26) Haanstra, L.; Doelman, P.; Oude Voshaar, J. H. The use of sigmoidal concentration-response curves in soil ecotoxicological research. *Plant Soil* **1985**, 293–297.
- (27) Willumsen, P. A.; Johansen, J. E.; Karlson, U.; Hansen, B. M. Isolation and taxonomic affiliation of N-heterocyclic aromatic hydrocarbon-transforming bacteria. *Appl. Microbiol. Biotechnol.* **2005**, 67, 420–428.
- (28) van Herwijnen, R.; van de Sande, B.; van der Wielen, F. W. M.; Springael, D.; Govers, H. A. J.; Parsons, J. R. Influence of phenanthrene and fluoranthene on the degradation of fluorene and glucose by *Sphingomonas* sp strain LB126 in chemostat cultures. *FEMS Microbiol. Ecol.* **2003**, 46, 105–111.
- (29) Conrad, A. U.; Comber, S. D.; Simkiss, K. Pyrene bioavailability; effect of sediment-chemical contact time on routes of uptake in an oligochaete worm. *Chemosphere* **2002**, 49, 447–454.
- (30) Kukkonen, J.; Landrum, P. F. Toxicokinetics and Toxicity of Sediment-Associated Pyrene to *Lumbriculus-Variegatus* (Oligochaeta). *Environ. Toxicol. Chem.* **1994**, 13, 1457–1468.
- (31) Poerschmann, J.; Gorecki, T.; Kopinke, F. D. Sorption of very hydrophobic organic compounds onto poly(dimethylsiloxane) and dissolved humic organic matter. 1. Adsorption or partitioning of VHOC on PDMS-coated solid-phase microextraction fibers - A never-ending story. *Environ. Sci. Technol.* **2000**, 34, 3824–3830.
- (32) Lee, J. H.; Landrum, P. F.; Field, L. J.; Koh, C. H. Application of a Sigma polycyclic aromatic hydrocarbon model and a logistic regression model to sediment toxicity data based on a species-specific, water-only LC50 toxic unit for *Hyalella azteca*. *Environ. Toxicol. Chem.* **2001**, 20, 2102–2113.
- (33) van Leeuwen, C. J.; Aldenberg, T.; Verhaar, H. J. M.; Hermens, J. L. M. Application of QSARs, extrapolation and equilibrium partitioning in aquatic effects assessment. I. Narcotic industrial pollutants. *Environ. Toxicol. Chem.* **1992**, 11, 267–282.
- (34) Lotufo, G. R. Lethal and sublethal toxicity of sediment-associated fluoranthene to benthic copepods: application of the critical-body-residue approach. *Aquat. Toxicol.* **1998**, 44, 17–30.
- (35) León Paumen, M.; Borgman, E.; Kraak, M. H. S.; van Gestel, C. A. M.; Admiraal, W. Life cycle responses of the midge *Chironomus riparius* to Polycyclic Aromatic Compound exposure. *Environ. Pollut.* **2008**, 152, 225–232.
- (36) Verrhiest, G.; Clement, B.; Blake, G. Single and combined effects of sediment-associated PAHs on three species of freshwater macroinvertebrates. *Ecotoxicology* **2001**, 10, 363–372.
- (37) Buryskova, B.; Hilscherova, K.; Blaha, L.; Marsalek, B.; Holoubek, I. Toxicity and modulations of biomarkers in *Xenopus laevis* embryos exposed to polycyclic aromatic hydrocarbons and their N-heterocyclic derivatives. *Environ. Toxicol.* **2006**, 21, 590–598.
- (38) Bartos, T.; Letzsch, S.; Skarek, M.; Flegrova, Z.; Cupr, P.; Holoubek, I. GFP assay as a sensitive eukaryotic screening model to detect toxic and genotoxic activity of azaarenes. *Environ. Toxicol.* **2006**, 21, 343–348.
- (39) McCarty, L. S.; Mackay, D. Enhancing Ecotoxicological Modeling and Assessment. *Environ. Sci. Technol.* **1993**, 27, 1719–1728.
- (40) Leslie, H. A.; Kraak, M. H. S.; Hermens, J. L. M. Chronic toxicity and body residues of the nonpolar narcotic 1,2,3,4-tetrachlorobenzene in *Chironomus riparius*. *Environ. Toxicol. Chem.* **2004**, 23, 2022–2028.
- (41) Landrum, P. F.; Lotufo, G. R.; Gossiaux, D. C.; Gedeon, M. L.; Lee, J. H. Bioaccumulation and critical body residue of PAHs in the amphipod *Diporeia* spp.: additional evidence to support toxicity additivity for PAH mixtures. *Chemosphere* **2003**, 51, 481–489.
- (42) Fisher, S. W.; Chordas, S. W.; Landrum, P. F. Lethal and sublethal body residues for PCB intoxication in the oligochaete *Lumbriculus variegatus*. *Aquat. Toxicol.* **1999**, 45, 115–126.
- (43) Di Toro, D. M.; McGrath, J. A.; Hansen, D. J. Technical basis for narcotic chemicals and polycyclic aromatic hydrocarbon criteria. I. Water and tissue. *Environ. Toxicol. Chem.* **2000**, 19, 1951–1970.
- (44) Hendriks, A. J. Modeling response of species to microcontaminants - Comparative ecotoxicology by (sub)lethal body burdens as a function of species size and partition ratio of chemicals. *Ecotoxicol. Environ. Saf.* **1995**, 32, 103–130.
- (45) Emery, V. L.; Dillon, T. M. Chronic toxicity of phenanthrene to the marine polychaete worm *Nereis (Neanthes) arenaceodentata*. *Bull. Environ. Contam. Toxicol.* **1996**, 56, 265–270.
- (46) Adrian, N. R.; Sufliata, J. M. Anaerobic biodegradation of halogenated and nonhalogenated N-heterocyclic, S-heterocyclic, and O-heterocyclic compounds in aquifer slurries. *Environ. Toxicol. Chem.* **1994**, 13, 1551–1557.
- (47) Bleeker, E. A. J.; Wiegman, S.; de Voogt, P.; Kraak, M.; Leslie, H. A.; de Haas, E.; Admiraal, W. Toxicity of azaarenes. *Rev. Environ. Contam. Toxicol.* **2002**, 173, 39–83.

ES702500T