



UvA-DARE (Digital Academic Repository)

General expression for the component size distribution in infinite configuration networks

Kryven, I.

DOI

[10.1103/PhysRevE.95.052303](https://doi.org/10.1103/PhysRevE.95.052303)

Publication date

2017

Document Version

Final published version

Published in

Physical Review E

[Link to publication](#)

Citation for published version (APA):

Kryven, I. (2017). General expression for the component size distribution in infinite configuration networks. *Physical Review E*, 95(5), [052303].
<https://doi.org/10.1103/PhysRevE.95.052303>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

General expression for the component size distribution in infinite configuration networks

Ivan Kryven*

University of Amsterdam, P.O. Box 94214, 1090 GE, Amsterdam, The Netherlands

(Received 16 December 2016; published 2 May 2017)

In the infinite configuration network the links between nodes are assigned randomly with the only restriction that the degree distribution has to match a predefined function. This work presents a simple equation that gives for an arbitrary degree distribution the corresponding size distribution of connected components. This equation is suitable for fast and stable numerical computations up to the machine precision. The analytical analysis reveals that the asymptote of the component size distribution is completely defined by only a few parameters of the degree distribution: the first three moments, scale, and exponent (if applicable). When the degree distribution features a heavy tail, multiple asymptotic modes are observed in the component size distribution that, in turn, may or may not feature a heavy tail.

DOI: [10.1103/PhysRevE.95.052303](https://doi.org/10.1103/PhysRevE.95.052303)

I. INTRODUCTION

Random graphs provide models for complex networks, and in many cases, real-world networks has been accurately described by such models [1–4]. Within the scope of random graph models one finds: Erdős-Rényi model, Barabási-Albert model [1], node copying model [5], small-world network [6], configuration network [7,8], and many others. In the configuration network, N nodes are assigned predefined degrees. The edges connecting these nodes are then considered to be random, and every distinct configuration of edges that satisfies the given degree sequence is treated as a new instance of the network in the sense of random graphs. Interesting properties emerge when the number of nodes, N , approaches infinity, or at the so-called thermodynamic limit [9,10]. In this case, the infinite degree sequence, which provides the only input information for the model, is equivalent to the frequency distribution of degrees, $u(k)$, $k = 1, 2, \dots$, i.e., the probability that a randomly chosen node has degree k .

Component-size distribution, $w(n)$, denotes probability that a randomly chosen node is part of a connected component of finite size n . Connected components in the infinite configuration network can be of finite or infinite size. Molloy and Reed [11] showed that if an infinite component exists, then it is the only infinite component with probability 1. Hence, the infinite component is referred to as the giant component.

Depending upon a specific context behind the network, the component size distribution may summarize an important feature of the modeled system. In polymer chemistry, for example, the infinite configuration network is used as a toy model for hyper-branched and cross-linked polymers. In this context, the component size distribution predicts viscoelastic properties of the material while the emergence of the giant component is interpreted as a phase transition from liquid to solid state of the soft matter [4,12]. Since connected components are closely related to clusters in bond percolation processes, the distribution of component sizes can be used to model outbreaks for SIR epidemiological processes [2]. In linguistics, component size distribution of the sentence similarity graph is an important tool when studying structure of natural

languages [13]. This brief list of application cases is far from being exhaustive. Despite the vast applications, the empirical component size distribution is hard to measure precisely unless the whole topology of the network is known. On another hand, empirical observations on the degree distribution, $u(k)$, are much easier to perform.

Reference [11] provides an elegant criterion that connects moments of $u(k)$ to the fact that the network contains the giant component. A somewhat deeper question further in this direction reads: providing $u(k)$ is given, what is the component size distribution, $w(n)$? In Ref. [14], Newman *et al.* showed that the component size distribution can be recovered by a numerical algorithm that involves solving a fixed-point problem followed by an inversion of a generation function. Such algorithm demonstrates that indeed $u(k)$ and $w(n)$ can be put into a correspondence; however, it becomes computationally infeasible for large values of n . This numerical issue arises due to ill-posedness of the numerical generating function inversion.

On another hand, the component size distribution has been analytically resolved only for a limited number of partial cases of $u(k)$ [15]. Within the scope of analytically solvable cases, only the Yule-Simon degree distribution features a heavy tail, that is to say it decays proportionally to an algebraic function, $n^{-\beta}$, $\beta > 0$ at large n . At the same time, the heavy-tailed (or scale-free) distributions are commonly observed in the empirical data collected from many real-world networks [16–19]. Empirically observed exponents vary in a broad range. Some studies report degree exponents that are as small as $\beta = 0.81$ in the case of the Internet topology [20] and $\beta = 1$ in social networks [21]. On the opposite side of this spectrum, one finds exponent $\beta = 5$ in the generalization of preferential attachment model [22].

The only asymptotic analysis available for component size distribution in the configuration network states that for large n , $w(n)$ is either proportional to $n^{-3/2}$ or to e^{-Cn} , where $C > 0$ is a constant [14]. The current paper uncovers new asymptotic modes for $w(n)$ that emerge only when the degree distribution features a heavy tail. The paper shows that for an arbitrary $u(k)$, $w(n)$ can be expressed as a finite sum. In practice, this sum can be stably computed up to the machine precision in the cost of $O(n^2)$ multiplicative operations. Finally, the paper discusses how a finite cutoff introduced in the degree distribution reflects on the distribution of component sizes.

*i.kryven@uva.nl

II. COMPONENT-SIZE DISTRIBUTION BY LAGRANGE INVERSION

It has been noticed that all components in the infinite configuration model are locally tree-like. Using this fact as a departure point, Newman *et al.* [14] showed that the degree distribution can be put into a correspondence to the component size distribution by applying the generation-function (GF) formalism. Here, by a GF of $u(k)$, $\sum_{k=0}^{\infty} u(k) = 1$, we refer to the series,

$$U(x) = \sum_{k=0}^{\infty} u(k)x^k, \quad x \in \mathbb{C}, |x| \leq 1. \quad (1)$$

According to the approach presented in Ref. [14], the generating function for the component size distribution, $W(x)$ is found as a solution of the following system of functional equations:

$$W(x) = xU[W_1(x)], \quad (2)$$

$$W_1(x) = xU_1[W_1(x)], \quad (2')$$

where $U(x)$ is the GF of $u(k)$, and $U_1(x)$ is the GF for the excess degree distribution,

$$u_1(k) = \frac{k+1}{\mu_1}u(k+1), \quad (3)$$

where $\mu_1 = \sum_{k=1}^{\infty} ku(k)$. Similar to combinatorial tree-counting problems, Eq. (2) can be solved by applying the Lagrange inversion formula [23]. The original formulation of the Lagrange inversion principle is as follows. Suppose $A(x), R(x)$ are such formal power series that $A(x) = xR[A(x)]$, then for an arbitrary formal power series $F(x)$, the coefficient of power series $F[A(x)]$ at x^n reads as

$$[x^n]F[A(x)] = \frac{1}{n}[t^{n-1}]F'(t)R^n(t), \quad n > 0. \quad (4)$$

Here $[t^{n-1}]$, as being the inverse operation to the GF transform Eq. (1), refers to the coefficient at t^{n-1} of the corresponding power series. By substituting $A(x) = W_1(x)$, $F(x) = U(x)$, and applying Eq. (2), one transforms the left-hand side of Eq. (4), $[x^n]F[A(x)] = [x^n]U[W_1(x)] = [x^{n+1}]W(x) = w(n+1)$. Further on, the right-hand side of Eq. (4) is transformed by substituting $R(x) = U_1(x)$ and realizing that, according to the definition Eq. (3), $U'(x) = \mu_1 U_1(x)$. Now, we are ready to write an expression for $w(n)$, even though we have no explicit expression for generating function $W(x)$ itself,

$$\begin{aligned} w(n) &= \frac{1}{n-1}[t^{n-2}]U'(x)U_1(x)^{n-1} \\ &= \frac{\mu_1}{n-1}[t^{n-2}]U_1(x)^n, \quad n > 1. \end{aligned} \quad (5)$$

A similar equation was also derived in Ref. [15] by means of different reasoning. In principle, Eq. (5) provides enough information to analytically recover the component size distribution for a few special cases of the degree distribution [15]. In practice, however, the main difficulty when applying Eq. (5) is that the equation employs the inverse GF transform, $[t^n]$, which limits the choices one has when searching for an exact solution or performing numerical computations. With this in

mind, one may rewrite Eq. (5) so that the new expression does not involve the GF concept at all. It turns out that the only reason why Eq. (5) utilizes the GF formalism is that it provides means for *convolution power*.

The convolution of two distributions, $f(k) * g(k)$, $k > 0$ is defined as a binary multiplicative operation,

$$f(k) * g(k) = \sum_{i+j=k} f(i)g(j),$$

where the summation is performed over all nonnegative ordered couples i, j that sum up to k . This sum contains exactly $k+1$ of such couples. In this paper, the order of operations is chosen in such a way that the pointwise multiplication precedes convolution, for instance, $f(k) * kg(k) = f(k) * [kg(k)]$. The convolution can be inductively extended to the n -fold convolution, or *the convolution power*,

$$f(k)^{*n} = f(k)^{*n-1} * f(k), \quad (6)$$

where $f(k)^{*0} \equiv 1$ by definition. It can be shown that the convolution power can be expanded into a sum of products,

$$f(k)^{*n} = \sum_{\substack{k_1+\dots+k_n=k \\ k_i \geq 0}} \prod_{i=1}^n f(k_i). \quad (7)$$

The convolution has a peculiar property in respect to the GF transform. If $F(x), G(x), U(x)$ are GFs for $f(k), g(k)$, and $u(k) = f(k) * g(k)$, then $U(x) = F(x)G(x)$. Furthermore, if $U(x)$ is GF for $u(k)$, then $U(x)^n$ generates $u(k)^{*n}$. By exploiting this relation one immediately reduces Eq. (5) to

$$w(n) = \begin{cases} \frac{\mu_1}{n-1}u_1^{*n}(n-2), & n > 1, \\ u(0) & n = 1. \end{cases} \quad (8)$$

Here, the value of $w(0)$ is derived directly from the formulation of the problem: nodes with degree zero are also components of size one. This simple equation is ready to be used: by combining Eq. (8) and the definition Eqs. (3) and (7), one may directly express the values of the component size distribution in terms of $u(k)$ for $n > 1$,

$$w(n) = \frac{[ku(k)]^{*n}(2n-2)}{(n-1)\mu_1^{n-1}}, \quad n > 1. \quad (9)$$

For example, the first five values of $w(n)$ read as,

$$\begin{aligned} w(1) &= u(0), \\ w(2) &= \frac{1}{\mu_1}u(1)^2, \\ w(3) &= \frac{3}{\mu_1^2}u(1)^2u(2), \\ w(4) &= \frac{4}{\mu_1^3}u(1)^2[2u(2)^2 + u(1)u(3)], \\ w(5) &= \frac{5}{\mu_1^4}u(1)^2[4u(2)^3 + 6u(1)u(2)u(3) + u(1)^2u(4)]. \end{aligned}$$

The number of terms in this expansion increases rapidly with n . That said, Eq. (8) can be easily readjusted for numerical computations. Namely, one can use Fast Fourier

TABLE I. Exact expressions for component size distributions in configuration network as evaluated with Eq. (8) via Z transform.

Degree Distribution	Component-Size Distribution
Exponential distribution $Ce^{-\lambda k}$	$\frac{(1-e^{-\lambda})^{2n-1}}{e^{\lambda(n-1)}} \frac{\Gamma(3n-2)}{\Gamma(n)\Gamma(2n)}$
Geometric distribution $(1-p)^{k-1}p$	$(1-p)^{n-2}p^{2n-1} \frac{\Gamma(3n-2)}{\Gamma(n)\Gamma(2n)}$
Binomial distribution $\binom{k_{\max}}{k}(1-c)^{k_{\max}-k}c^k$	$\frac{1}{n-1} \binom{n k_{\max}-n}{n-2} \times (1-c)^{n k_{\max}-2n-2} c^{n-2}$

Transform (FFT) to compute the convolution powers, $u_1^{*n}(k) = \mathcal{F}^{-1}[\mathcal{F}[u_1(k)]^n]$. In this case, $O(N^2)$ multiplicative operations is sufficient to compute all values of $w(n), n \leq N$. Alternatively, if $w(n-1)$ is known, then $w(n)$ can be found in the cost of $O(n \log n)$. Besides FFT, there are algorithms that are specifically designed for fast approximation of convolution powers, such as projection onto basis functions that are invariant under convolution [24].

Analytic formulas for convolution powers (sometimes also referred to as *compositas* [25]), were covered by literature for many elementary functions [26,27]. Convolution powers of $u_1(k)$ can also be found analytically by applying discrete functional transforms, for instance, Z transform and discrete Fourier transform. A few examples of such results are given in Table I. Focusing on one of them, the first curve in Fig. 1 demonstrates that both analytical and numerical results for the exponential degree distribution coincide.




FIG. 1. Examples of component size distributions (solid lines) that feature fast (1) and slow (2) convergence to their asymptotes (dashed lines). Both asymptotes are covered by Case A, Table II. (1) $u(k) = Ce^{-1.05k}$, all three: the analytical expression (see Table I), numerical values [according to Eq. (8)] and the asymptote practically coincide. (2) $u(k)$ is nonzero in three points $u(1) = 0.97$, $u(2) = 0.015$, $u(10) = 0.015$; the component size distribution features oscillations before it converges to the asymptote.

III. ASYMPTOTIC ANALYSIS

The format of Eq. (8) naturally suggests a straightforward way to perform an asymptotic analysis for $n \rightarrow \infty$. One may view $u(k)$ as a probability mass function PMF (or alternatively discrete probability density function) of some discrete random variables k_i . Recall the following property of convolution powers: if i.i.d. random variables k_i have PMF $u_1(k)$, then $u_1^{*n}(k)$ gives the PMF for the sum $k_1 + k_2 + \dots + k_n$. The central limit theorem (CLT) gives an estimate for this sum as $n \rightarrow \infty$, and the idea is now to obtain the asymptotes of $w(n)$ by applying CLT to the definition Eq. (8).

A. Light-tailed degree distributions

First, let us assume that distribution $u(k)$ decays faster than algebraically, that is

$$u(k) = o(k^{-\beta}), \quad \beta > 2, \quad k \rightarrow \infty, \quad (10)$$

which is also equivalent to $u_1(k) = o(k^{-\beta+1})$. Then, according to CLT, $u_1^{*n}(k)$ approaches the normal distribution, $u_1^{*n}(k) \xrightarrow{d} (\sqrt{n}\sigma)^{-1} \mathcal{N}(\frac{k-nM}{\sqrt{n}\sigma}, 0, 1)$, when $n \rightarrow \infty$, where $M = \sum_{k=1}^{\infty} k u_1(k)$ and $\sigma^2 = \sum_{k=1}^{\infty} k(k-M)^2 u_1(k) < \infty$ denote the mean value and variance of $u_1(k)$. The normal distribution can now replace $u_1^{*n}(k)$ in Eq. (8), which yields the asymptote for the component size distribution,

$$w(n) \sim \frac{\mu_1 e^{-\frac{(n(1-M)-2)^2}{2n\sigma^2}}}{(n-1)\sqrt{2\pi n\sigma^2}}, \quad \text{as } n \rightarrow \infty. \quad (11)$$

Quantities M, σ^2 are directly expressible in terms of moments of degree distribution $u(k)$,

$$\begin{aligned} M &= \sum_{k=1}^{\infty} k u_1(k) = \frac{1}{\mu_1} \sum_{k=1}^{\infty} (k^2 - k) u(k) = \frac{\mu_2 - \mu_1}{\mu_1}, \\ \sigma^2 &= \sum_{k=1}^{\infty} k(k-M)^2 u_1(k) \\ &= \frac{1}{\mu_1} \sum_{k=0}^{\infty} k(k-M-1)^2 u(k) = \frac{\mu_3 \mu_1 - \mu_2^2}{\mu_1^2}, \end{aligned} \quad (12)$$

where

$$\mu_i = \mathbb{E}[k^i] = \sum_{k=1}^{\infty} k^i u(k), \quad i = 1, 2, \dots$$

Finally, substituting Eqs. (12) into Eq. (11) gives the final version of the asymptote,

$$w(n) \sim \frac{\mu_1^2 n^{-3/2} e^{-\frac{(\mu_2 - 2\mu_1)^2}{2(\mu_1\mu_3 - \mu_2^2)} n}}{\sqrt{2\pi(\mu_1\mu_3 - \mu_2^2)}}, \quad \text{as } n \rightarrow \infty. \quad (13)$$

Two examples of component size distributions that converge with various rates to their asymptotes are given in Fig. 1. Particularly, the only information on $u(k)$ that is contained in the asymptote definition Eq. (13) is the first three moments μ_1, μ_2, μ_3 . Furthermore, depending upon the value of $\theta = \mu_2 - 2\mu_1$, the asymptotic expression Eq. (13) switches between the *two modes*: it either decays exponentially as

$O(e^{-An})$, when $\theta \neq 0$, or it decays as an algebraic function, $O(n^{-3/2})$, when $\theta = 0$ (see also Table II, Case A). The last equality is the well-known giant component criterion,

$$\mu_2 - 2\mu_1 = 0. \quad (14)$$

The criterion Eq. (14) was obtained by Molloy and Reed [11] by means of a different reasoning. In Ref. [11], the authors prove that $\theta > 0$ implies existence of the giant component in the configuration network, whereas $\theta < 0$ implies nonexistence of this component. In Ref. [14], it was hypothesized that the $-3/2$ exponent is universal and must hold for all degree distributions at the critical point $\theta = 0$. We will see now that when the condition Eq. (10) fails to hold, distinct from $-3/2$ exponents may also appear in the asymptotic of $w(n)$.

B. Heavy-tailed degree distributions

Suppose that, contrary to the condition Eq. (10), degree distribution $u(k)$ features a heavy tail,

$$u(k) \sim sk^{-\beta}, \quad \beta > 2, \quad k \rightarrow \infty, \quad (15)$$

which is equivalent to $u_1(k) \sim sk^{-\alpha-1}$, $\alpha = \beta - 2 > 0$, $k \rightarrow \infty$. It turns out that exponent α and the scale s , together with the moments μ_1, μ_2, μ_3 , provide enough information to generalize the asymptote Eq. (13) for the case of heavy-tailed degree distributions. Suppose $0 < \alpha \leq 2$. In terms of $u(k)$ moments, this condition casts out as $\mu_3 = \infty$. As follows from Gnedenko and Kolmogorov's generalization of CLT [28], the mass density distribution for $u_1^{*n}(k)$ approaches the stable law,

$$u_1^{*n}(k) \xrightarrow{d} \frac{1}{\gamma(n)} G^A\left(\frac{k - \mu(n)}{\gamma(n)}, \alpha, 1\right), \quad n \rightarrow \infty. \quad (16)$$

Here, we use the notation of Uchaikin and Zolotarev [29], which includes *exponent parameter* α , the *location parameter*,

$$\mu(n) = \begin{cases} n \frac{\mu_2 - \mu_1}{\mu_1}, & \alpha > 1, \\ sn \ln n, & \alpha = 1, \\ 0, & 0 < \alpha < 1, \end{cases} \quad (17)$$

and the *scale parameter*,

$$\gamma(n) = \begin{cases} \sqrt{sn \ln n}, & \alpha = 2, \\ \sqrt{\pi s} [2\Gamma(\alpha) \sin \frac{\alpha\pi}{2}]^{-1/\alpha} n^{1/\alpha}, & \alpha \in (0, 1) \cup (1, 2), \\ \frac{\pi ns}{2}, & \alpha = 1. \end{cases} \quad (18)$$

No general analytical expression is known for $G^A(x, \alpha, 1)$, and the stable law is defined via its Fourier transform,

$$\mathcal{F}[G^A(x, \alpha, 1)] = \begin{cases} e^{-x^\alpha - i \tan \frac{\alpha\pi}{2}}, & \alpha \in (0, 1) \cup (1, 2), \\ e^{-x^\alpha + i \frac{2\alpha}{\pi}}, & \alpha = 1. \end{cases} \quad (19)$$

Consider the case when $1 < \alpha < 2$. According to Eq. (16), the point in which the stable law is evaluated, $x(n) = \frac{n - \mu(n)}{\gamma(n)}$, approaches positive or negative infinities depending upon the sign of $\theta = \mu_2 - 2\mu_1$. Indeed, as $n \rightarrow \infty$,

$$x(n) \rightarrow \begin{cases} +\infty, & \theta < 0, \\ 0, & \theta = 0, \\ -\infty, & \theta > 0. \end{cases} \quad (20)$$

For these values of α , function $G^A(x, \alpha, 1)$ is nonzero on $(-\infty, +\infty)$. If $x(n) \rightarrow \infty$, the function features an algebraic decay, whereas if $x(n) \rightarrow -\infty$, the decay is exponential.




FIG. 2. Component-size distributions (*solid lines*) and their asymptotes (*dashed lines*) as obtained for degree distributions with exponent $\beta = 3.5$ ($\alpha = 1.5$) and various values of scale parameter. Three distinct asymptotic modes are illustrated: (1) $s = 0.066$: $\theta < 0$, $\eta = \frac{5}{2}$; (2) $s = 0.644$: $\theta \approx 0$, $\eta = \frac{5}{3}$; (3) $s = 0.8$: $\theta > 0$.

Therefore, the limiting value switching that takes place in Eq. (20) may reflect on the asymptotic behavior of $u_1^{*n}(n)$. To give a precise answer one has to consider series expansions of $G^A(x, \alpha, 1)$ around the points of interest, $x \in \{-\infty, 0, +\infty\}$. We use here the leading terms of these series [29],

$$G^A(x, \alpha, 1) = \begin{cases} \frac{\Gamma(1+\frac{1}{\alpha}) \sin \frac{\pi}{\alpha}}{\pi} + O(x), & x \rightarrow 0 \\ \frac{\Gamma(\alpha+1)x^{-\alpha-1}}{\Gamma(2-\alpha)\Gamma(\alpha-1)} + O(x^{-2\alpha-1}), & x \rightarrow \infty, \\ \frac{e^{-(\alpha-1)(\frac{x}{\alpha})^{\alpha-1}} (\frac{x}{\alpha})^{\frac{1}{2}(\frac{1}{\alpha-1}-1)}}{\sqrt{2\pi\alpha(\alpha-1)}} [1 + O(x^{-\frac{\alpha-1}{\alpha}})], & x \rightarrow -\infty. \end{cases} \quad (21)$$

By replacing the expression for the limiting distribution Eq. (16) with the leading terms given in Eq. (21), one obtains the asymptotes for Eq. (8). This time, the asymptote has *three modes*: depending upon the value of θ , it either features a heavy tail with exponent $-\alpha - 1$, a heavy tail with exponent $-\frac{1}{\alpha} - 1$, or an exponential decay, as shown in Table II, Case D. A few examples of such asymptotic modes for a heavy-tailed degree distribution,

$$u(k) = \begin{cases} C & k = 1, \\ s(\beta - 2)k^{-\beta} & k > 1, \end{cases} \quad (22)$$

are given in Fig. 2. The degree distribution Eq. (22) is defined by two parameters: exponent β and scale s ; whereas the constant C is such that the total probability is normalized, $\sum_k u(k) = 1$.

When $\alpha = 2$, the behavior of $\frac{n - \mu(n)}{\gamma(n)}$, $n \rightarrow \infty$ is identical to Eq. (20), but the expression for $\gamma(n)$ is different and the series expansions Eq. (21) lead to somewhat different asymptotes; see Table II, Case C.

According to the definition Eq. (17), the location parameter vanishes, $\mu(n) \equiv 0$, when $\alpha < 1$. In this case, $x(n) = \frac{n}{\gamma(n)} \rightarrow 0$ as $n \rightarrow \infty$, and only *one* asymptotic mode is possible for

TABLE II. Asymptotic behavior of component sizes $w(n)$, in terms of degree distribution parameters: the first three moments μ_1, μ_2, μ_3 , scale parameter s and exponent β . Supporting source code available in Ref. [32].

Finite moments of $u(k)$	$u(k), k \rightarrow \infty$	$\theta = \mu_2 - 2\mu_1$	Asymptote of $w(n)$
$\mu_3 < \infty$	A. $o(k^{-\beta}), \beta > 4$	$\theta \neq 0$	$C_1 e^{-C_2 n} n^{-3/2}$
		$\theta = 0$	$C_1 n^{-3/2}$
		$\theta < 0$	$C_3 n^{-\alpha-1}$
	B. $O(k^{-\beta}), \beta > 4$	$\theta = 0$	$C_1 n^{-3/2}$
		$\theta > 0$	$C_1 e^{-C_2 n} n^{-3/2}$
		$\theta < 0$	$C_3 n^{-\alpha-1}$
$\mu_3 = \infty, \mu_2 < \infty$	C. $O(k^{-\beta}), \beta = 4$	$\theta = 0$	$C_1' \frac{n^{-3/2}}{\sqrt{\log n}}$
		$\theta > 0$	$C_1' \frac{n^{-3/2}}{\sqrt{\log n}} e^{-C_2' \frac{n}{\log n}}$
		$\theta < 0$	$C_3 n^{-\alpha-1}$
	D. $O(k^{-\beta}), 3 < \beta < 4$	$\theta = 0$	$C_4 n^{-\frac{1}{\alpha}-1}$
		$\theta > 0$	$C_5 e^{-C_6 n} n^{-3/2}$
		$\theta < 0$	$C_3 n^{-\alpha-1}$
$\mu_2 = \infty$	E. $O(k^{-\beta}), \beta = 3$	$\theta > 0$	$C_7 e^{-C_8 - C_9 n^{\frac{2}{\pi}}} n^{\frac{1}{\pi}-2}$
		$\theta > 0$	$C_{10} e^{-C_{11} n} n^{-3/2}$
	F. $O(k^{-\beta}), 2 < \beta < 3$	$\theta > 0$	$C_7 = \frac{\sqrt{2}\mu_1}{\pi^{3/2}s},$
		$\theta > 0$	$C_8 = \frac{1}{\pi s} + \frac{1}{2},$
		$\theta > 0$	$C_9 = e^{-1 - \frac{2}{\pi s}},$
		$\theta > 0$	$C_{10} = \frac{\mu_1}{\sqrt{2-2\alpha}} \left(\frac{\sqrt{2}\Gamma(\alpha) \sin \frac{\pi\alpha}{2}}{\alpha\pi^\alpha s} \right)^{\frac{1}{2\alpha-2}},$
		$C_{11} = (1-\alpha) \left(\frac{\sqrt{2}\Gamma(\alpha) \sin \frac{\pi\alpha}{2}}{\pi\alpha^\alpha s} \right)^{\frac{1}{\alpha-1}},$	
		$\alpha = \beta - 2$	

$$C_1 = \frac{\mu_1^2}{\sqrt{2\pi(\mu_1\mu_3 - \mu_2^2)}}, \quad C_1' = \frac{\mu_1}{\sqrt{2\pi s}},$$

$$C_2 = \frac{(\mu_2 - 2\mu_1)^2}{2(\mu_1\mu_3 - \mu_2^2)}, \quad C_2' = \frac{(\mu_2 - 2\mu_1)^2}{2s\mu_1^2},$$

$$C_3 = \frac{s\mu_1^{\alpha+2}\Gamma(\alpha+1)}{(2\mu_1 - \mu_2)^{\alpha+1}\Gamma(\alpha)},$$

$$C_4 = \mu_1 \Gamma\left(1 + \frac{1}{\alpha}\right) \sin \frac{\pi}{\alpha} \left(\frac{2\Gamma(\alpha) \sin \frac{\pi\alpha}{2}}{\pi^{\alpha+1}s}\right)^{1/\alpha},$$

$$C_5 = \frac{\mu_1}{\sqrt{\alpha-1}} \left(\frac{2^{2-\alpha} \left(\frac{\mu_2}{\mu_1} - 2\right)^{2-\alpha} \Gamma(\alpha) \sin \frac{\pi\alpha}{2}}{\alpha\pi^\alpha s}\right)^{\frac{1}{2\alpha-2}},$$

$$C_6 = (1-\alpha) \left(\frac{2 \left(\frac{\mu_2}{\mu_1} - 2\right)^\alpha \Gamma(\alpha) \sin \frac{\pi\alpha}{2}}{\alpha^\alpha \pi s}\right)^{\frac{1}{\alpha-1}},$$

$w(n)$. Stable law $G^A(x, \alpha, 1)$ is supported on $(0, \infty)$, and we make use of the series expansion around $x \rightarrow 0^+$,

$$G^A(x, \alpha, 1) = \frac{e^{-(1-\alpha)(\frac{x}{\alpha})^{\frac{1}{1-\alpha}}} \left(\frac{x}{\alpha}\right)^{\frac{1}{2}(1+\frac{1}{1-\alpha})}}{\sqrt{2\pi\alpha(1-\alpha)}} [1 + O(x^{\frac{1-\alpha}{\alpha}})], \quad x \rightarrow 0^+, \quad (23)$$

which when plugged into Eq. (16) yields faster than algebraic decay of the component-size distribution; see Table II, Case F. Due to the parametrization scheme for the stable law, the point $\alpha = 1$ needs to be considered separately. In this case, $\frac{n-\mu(n)}{\gamma(n)} \rightarrow \infty$ when $n \rightarrow \infty$, and we utilize the leading term of the series expansion,

$$G^A(x, \alpha, 1) = \frac{1}{\sqrt{2\pi}} e^{\frac{x-1}{2} - e^{x-1}} [1 + O(e^{1-x})], \quad x \rightarrow \infty, \quad (24)$$

which admits one subalgebraic asymptotic mode for $w(n)$ as shown in Table II, Case E. This case is special in that the stable law $G^A(x, \alpha, 1)$ is supported on $x \in (-\infty, \infty)$, but asymptotically, $\frac{1}{\gamma(n)} G^A\left(\frac{n-\mu(n)}{\gamma(n)}, \alpha, 1\right)$ always tends to $-\infty$ for large n . At the same time, if for small n the point $x(n) = \frac{n-\mu(n)}{\gamma(n)}$ stays on the positive half-axis where Eq. (24) does not provide correct description for $G^A(x, \alpha, 1)$, the convergence to the asymptote will be slow. In other words, there is an intermediate asymptote that the component size distribution can be approximated with, before it eventually switches to

Eq. (24). This switching point is given by such n_0 that $x(n)$ changes the sign from “+” to “-”, i.e., when n becomes greater than n_0 . By solving $x(n_0) = 0$, one obtains $n_0 = e^{\frac{1}{s}}$, which means that, in principle, the switching between the intermediate and the final asymptotes may be indefinitely postponed if s is small enough. The intermediate asymptote itself is deduced from the leading term of the stable law expansion at ∞ , that is, $G^A(x, 1, 1) = \frac{2}{\pi} x^{-2} + O(x^{-3})$. After the substitutions, one obtains

$$w(n) \simeq \frac{\mu_1 s}{(s \log n - 1)^2} n^{-2}, \quad \frac{1}{s} \gg 0, \quad n < e^{\frac{1}{s}}, \quad \alpha = 1.$$

As illustrated in Fig. 3, similar considerations are also valid for the case $0 < \alpha < 1$, where

$$w(n) \simeq \mu_1 s \frac{\Gamma(\alpha+1)}{\Gamma(\alpha)} n^{-\alpha-1}, \quad \frac{1}{s} \gg 0, \quad 0 < \alpha < 1.$$

When this occurs, such switching has a practical importance when dealing with empirically observed component size data. Indeed, it may happen that one observes only the intermediate asymptote and not the final one due to a small number of samples at the tail of the component size distribution. For instance, the second curve in Fig. 3 does feature an exponential decay at infinity, but if one limits the data points to $n < 10^6$, the component size distribution will seem to be a heavy-tailed one.




FIG. 3. Component-size distributions (solid line) corresponding to degree distributions with exponent $\beta = 2.6$ ($\alpha = 0.6$). In this case, the component size distributions cannot feature a heavy tail; however, depending upon the scale parameter s a transient asymptote with exponent -1.6 (dashed line) emerges: (1) $s = 8.3 \times 10^{-2}$, fast convergence to the exponential asymptote. (2) $s = 8.3 \times 10^{-5}$, the distribution transiently follows what seems to be a heavy tail for $n < 10^6$, whereas for larger n the theory predicts no heavy tail.

Finally, we consider the case when the condition Eq. (15) holds for $\beta > 4$: even though $u_1(k)$ has finite mean M and variance σ^2 it also features a heavy tail. Again, as $n \rightarrow \infty$, $x(n) = \frac{n - \mu(n)}{\sigma(n)}$ features the limiting values that are defined by the sign of θ ; see Eq. (20). One would expect that since σ^2 is finite, this case should be also well approximated with Eq. (13). This is indeed the case for $x(n) \approx 0$. However, large deviations from zero $x(n) \gg 0$ do not follow Gaussian statistics [30,31], and we approximate $u_1^{*n}(k)$ with the Pareto stable law $u_1^{*n}(k) \rightarrow \frac{1}{\sigma(n)} G^P(\frac{k - \mu(n)}{\sigma(n)}, \alpha)$, $n \rightarrow \infty$. It turns out that $G^P(x, \alpha)$ behaves as the normal distribution for $x < C$, where C is a finite positive constant but features a heavy tail with the same exponent as $u_1(k)$ when $x \rightarrow \infty$; see Ref. [30]. Thus, when $\theta \geq 0$ the component size distribution features asymptotic modes as in Eq. (13), while when $\theta < 0$ it features a heavy tail with exponent $-\alpha - 1$; see Table II, Case B. Interestingly, when θ is a small negative number, $w(n)$ transiently follows one asymptote and then switches to the other as demonstrated in Fig. 4. If there is a process that continuously changes the degree distribution so that θ progresses from being negative to positive, the exponent of the associated component size distribution will jump from the subcritical branch, at $\theta < 0$, to the critical one at $\theta = 0$. An example of such a transition between two power-law modes is given in Fig. 4, where a component size distribution switches between power laws with exponents $\eta = 3$ and $\eta = 1.5$.

IV. DISCUSSION AND CONCLUSIONS

The broad generality of the results obtained in the previous section is achieved due to the fact that the configuration networks are locally treelike and have vanishing probability of clustering in the thermodynamic limit, which allows one to benefit from the available in analytic combinatorics tools.




FIG. 4. Examples of component size distributions (solid lines) that are associated with heavy-tailed degree distributions with $\beta = 6$ ($\alpha = 4$). The dashed lines represent the asymptotes in accordance with Case B in Table II. Depending on the sign of θ , three asymptotic modes are distinguished: (1) $s = 1.93$: $\theta = -0.8$, (1a) $s = 9.42$: $\theta = -0.027$, (2) $s = 9.69$: $\theta = 1.6 \times 10^{-7}$, (3) $s = 10.05$: $\theta = 0.038$. When θ is a small negative number (curve 1a), $w(n)$ first decays as $n^{-3/2}$ but eventually switches to asymptote $n^{-\alpha-1} = n^{-5}$.

Equation (8), which was analyzed in the previous section, connects the degree distribution in a configuration network to the distribution of sizes for connected components. The main conclusion one may draw from this equation is that the convolution power provides a smoothing effect. This means that all points of $u(k)$, $k = 1, \dots, \infty$ have a significant contribution to the definition of $w(n)$, but as n increases, the system “forgets” the exact shape of the degree distribution and the component size distribution tends to the asymptote, which is defined by only a few parameters. The only information that is still preserved at the limit $n \rightarrow \infty$ is the first three moments of the degree distribution if such does not feature a heavy tail; see, for example, Fig. 1. If $u(k)$ does feature a heavy tail then the information that characterizes the tail becomes also important, which is the scale parameter s and the exponent β . Depending upon the values of these parameters, many asymptotical modes exist.

The expression for the asymptote is framed in terms of small deviation statistics for a sum of random variables and in some cases can be used as a good approximation for the component size distribution. Table II contains the analytical expressions for the asymptotes. Additionally, supporting code computing the component size distribution and the corresponding asymptotes is provided [32]. When using the asymptotical expressions to approximate $w(n)$, one should pay attention to two factors that follow from central limits: first, n should be large; second, the approximation is best for θ close to zero. Finally, small deviations or a cutoff in a heavy-tailed degree distribution can trigger considerable and nontrivial changes in $w(n)$, for instance, the change of the asymptotical mode of the latter.

A. Degree distributions with a cutoff

In practice, no empirical degree distribution is a heavy-tailed one. Most of the “real-world” degree distributions feature a cutoff, $u(k) = 0$, $k > k_{\text{cut}}$, and therefore fail to be




FIG. 5. The effect of a cutoff imposed on a heavy-tailed degree distribution with $\beta = 3.3$ ($\alpha = 1.3$) and $s = 7.73$. The solid curves correspond to component size distributions with: (1) no cutoff, $\theta > 0$; (2) cutoff at $k = 1000$, $\theta = 0$; (3) cutoff at $k = 100$, $\theta < 0$. The asymptote for 1 is covered by Case D, Table II; while due to the cutoffs, the asymptotes for 2 and 3 are covered by Case A.

heavy-tailed in the strict sense of the definition Eq. (15). It turns out that if a cutoff is featured at large enough k_{cut} , the above-provided asymptotic analysis still has a relevant meaning. This situation can be compared to how we commonly attribute the fractal dimension to real-world geometric objects that fail to be fractals on infinitesimal scales.

Suppose one applies a cutoff at k_{cut} to a degree distribution, $u(k)$, that features a heavy tail. Since $u(k)$ has a finite support, the asymptote of associated $w(n)$ is covered by Case A (Table II); however, if k_{cut} is large, $w(n)$ may also transiently follow the original asymptote. Instead of an analytical investigation, we demonstrate the influence of the cutoff with numerical examples obtained by computing Eq. (8). This influence strongly depends on how the sign of θ is affected by the introduction of the cutoff. For example, if $\theta > 0$ even after the cutoff, the cutoff will cause more nodes to appear in finite-size components, and thus the component size distribution will shift toward larger sizes. The opposite case is valid when $\theta \leq 0$ before (and after) the cutoff, then the cutoff causes the component size distribution to shift toward smaller sizes. The third option is when the cutoff changes the sign of θ from “+” to “-”. In this case, both shifts are possible. Figure 5 shows how a component size distribution that corresponds to degree exponent $\beta = 3.3$ is affected by a cutoff with various values of k_{cut} .

B. Excess degree distribution with no mean value

In principle, the excess degree distributions that do not have a mean value, i.e., $\beta < 2$, do not fall within any of the above categories. However, if one introduces a cutoff, $u(k)$ will feature finite moments including, $\mu_3 < \infty$, hence this case should be treated according to Case A of Table II. Figure 6 shows how cutoffs at $k = k_{\text{cut}}$ influence an instance of component size distribution with $\beta = 1$. Unlike as in the previous example, in which $u(k)$ with no cutoff generates a valid $w(n)$, here the increase of k_{cut} results in vanishing




FIG. 6. The effect of a cutoff imposed on a heavy-tailed degree distribution with exponent $\beta = 1$ ($\alpha = -1$) and $s = -2 \times 10^{-4}$. The following values of the cutoff are considered: (1) $k_{\text{cut}} = 80$, corresponds to $\theta < 0$; (2) $k_{\text{cut}} = 100$, corresponds to $\theta \approx 0$; (3) $k_{\text{cut}} = 150$ corresponds to $\theta > 0$, (4) $k_{\text{cut}} = 10^3$; (5) $k_{\text{cut}} = 10^5$.

probability of finding a finite-size component at all: for any n , $w(n) \rightarrow 0$ when $k_{\text{cut}} \rightarrow \infty$. This illustrates the fact that finite-size components do not exist for $\beta \leq 1$, and the whole configuration network is connected almost surely. Nonexistence of finite components for $\beta \leq 1$ also follows from the fact that in this case μ_1 diverges and the point values of $w(n)$, as given below the definition Eq. (9), tend to zero.

Suppose the cutoff in the empirical, heavy-tailed degree distribution is due to the fact that the network sample has a




FIG. 7. The correspondence between the exponent β in a heavy-tailed degree distribution, and the exponent η in the associated heavy-tailed component size distribution. The critical branch corresponds to $\theta = 0$, subcritical branch to $\theta < 0$. Positive θ is not associated with heavy-tailed component size distributions.

finite size, $k_{\text{cut}} = N$, $N \neq \infty$, then one may approximate the expected number of edges in this sample as

$$n_e = \frac{N\mu_1}{2} = \frac{N}{2} \sum_{k=1}^N ku(k) \simeq \frac{N}{2} \sum_{k=1}^N k^{-\beta+1}, \quad N \gg 1,$$

so that

$$n_e \simeq \begin{cases} N(1 - N^{2-\beta}), & \beta \neq 2, \\ N \log N, & \beta = 2. \end{cases}$$

Subsequently, three scenarios are possible here:

(i) *sparse network*, $n_e = CN$, $C > 0$, $\beta > 2$: the asymptotic modes are given in Table II;

(ii) *semi-dense network*, either $n_e = CN \log N$, $\beta = 2$ or $n_e = CN^{3-\beta}$, $1 < \beta < 2$: the mean value of excess distribution diverges; there are finite components but no power law in the distribution of component sizes;

(iii) *dense network*, $n_e = CN^{3-\beta}$, $\beta \leq 1$: the mean value of degree distribution $\mu_1 \rightarrow \infty$, and finite components vanish as $N \rightarrow \infty$.

C. The role of the giant component

All the cases presented in Table II depend in some way on the value of θ . This is not a coincidence as the sign

of θ is the indicator for the giant component existence. If the degree distribution features a heavy tail with exponent $\beta \geq 3$, depending upon the value of θ , there are two possible heavy-tail exponents for the component size distribution: subcritical branch $\eta = \beta - 1$ when $\theta < 0$, and critical branch $\eta = \min\{\frac{3}{2}, \frac{\beta-1}{\beta-2}\}$ when $\theta = 0$. This relation is illustrated in Fig. 7, where the component size distribution exponent η is plotted versus the degree-distribution exponent β . We can see that if the giant component exists, $\theta > 0$, then irrespectively of what is the degree distribution, the component size distribution always decays faster than the power law. Therefore, it can be concluded that the giant component is not compatible with a heavy-tailed component size distribution. Any degree distribution with $\beta < 3$ leads to a giant component since θ can only be positive in this case. Furthermore, if $\beta \leq 1$, then the giant component is also the only component: with probability 1 the configuration network is fully connected.

ACKNOWLEDGMENTS

This work is part of the research program Veni with Project No. 639.071.511, which is financed by the Netherlands Organisation for Scientific Research (NWO).

-
- [1] A.-L. Barabási and R. Albert, *Science* **286**, 509 (1999).
- [2] M. E. Newman, D. J. Watts, and S. H. Strogatz, *Proc. Natl. Acad. Sci. USA* **99**, 2566 (2002).
- [3] A. Vázquez, *Phys. Rev. E* **67**, 056104 (2003).
- [4] I. Kryven, J. Duivenvoorden, J. Hermans, and P. D. Iedema, *Macromol. Theory Simul.* **25**, 449 (2016).
- [5] U. Bhat, P. L. Krapivsky, R. Lambiotte, and S. Redner, *Phys. Rev. E* **94**, 062302 (2016).
- [6] M. E. J. Newman and D. J. Watts, *Phys. Rev. E* **60**, 7332 (1999).
- [7] M. Molloy and B. Reed, *Combinat. Probabil. Comput.* **7**, 295 (1998).
- [8] M. Newman, *Networks: An Introduction* (Oxford University Press, Oxford, 2010).
- [9] F. Chung, L. Lu, and V. Vu, *Proc. Natl. Acad. Sci. USA* **100**, 6313 (2003).
- [10] I. Kryven, *Phys. Rev. E* **94**, 012315 (2016).
- [11] M. Molloy and B. Reed, *Random Struct. Algor.* **6**, 161 (1995).
- [12] I. Kryven and P. D. Iedema, *Chem. Eng. Sci.* **126**, 296 (2014).
- [13] C. Biemann and A. van den Bosch, *Structure Discovery in Natural Language* (Springer Science & Business Media, Berlin, 2011).
- [14] M. E. J. Newman, S. H. Strogatz, and D. J. Watts, *Phys. Rev. E* **64**, 026118 (2001).
- [15] M. E. J. Newman, *Phys. Rev. E* **76**, 045101 (2007).
- [16] S.-H. Yook, H. Jeong, and A.-L. Barabási, *Proc. Natl. Acad. Sci. USA* **99**, 13382 (2002).
- [17] E. Ravasz and A.-L. Barabási, *Phys. Rev. E* **67**, 026112 (2003).
- [18] V. M. Eguiluz, D. R. Chialvo, G. A. Cecchi, M. Baliki, and A. V. Apkarian, *Phys. Rev. Lett.* **94**, 018102 (2005).
- [19] F. Fu, L. Liu, and L. Wang, *Physica A: Stat. Mech. Appl.* **387**, 675 (2008).
- [20] L. A. Adamic and B. A. Huberman, *Glottometrics* **3**, 143 (2002).
- [21] G. Timár, S. N. Dorogovtsev, and J. F. F. Mendes, *Phys. Rev. E* **94**, 022302 (2016).
- [22] S. N. Dorogovtsev, J. F. F. Mendes, and A. N. Samukhin, *Phys. Rev. Lett.* **85**, 4633 (2000).
- [23] F. Bergeron, G. Labelle, and P. Leroux, *Combinatorial Species and Tree-like Structures* (Cambridge University Press, Cambridge, 1998).
- [24] I. Kryven and P. Iedema, *Macromol. Theory Simul.* **22**, 89 (2013).
- [25] D. V. Kruchinin, *Adv. Diff. Equat.* **2015**, 1 (2015).
- [26] N.-Y. Ma and F. Liu, *Appl. Math. Comput.* **158**, 225 (2004).
- [27] N.-Y. Ma and R. King, *Appl. Math. Comput.* **133**, 83 (2002).
- [28] B. V. Gnedenko and A. Kolmogorov, *Limit Distributions for Sums of Independent Variables* (Addison-Wesley, Cambridge, 1954).
- [29] V. V. Uchaikin and V. M. Zolotarev, *Chance and Stability: Stable Distributions and Their Applications* (Walter de Gruyter, Urecht, 1999).
- [30] C. M. Ramsay, *Commun. Stat. Theory Methods* **35**, 395 (2006).
- [31] J. Nacher and T. Akutsu, *Physica A: Stat. Mech. Appl.* **390**, 4636 (2011).
- [32] Matlab/GNU Octave source code for calculating component size distribution in configuration network, https://github.com/ikryven/PhysRevE_2017_GECS.