



UvA-DARE (Digital Academic Repository)

Tomato reproductive success is equally affected by herbivores that induce or that suppress defenses

Liu, J.; Legarrea, S.; Kant, M.R.

DOI

[10.3389/fpls.2017.02128](https://doi.org/10.3389/fpls.2017.02128)

Publication date

2017

Document Version

Other version

Published in

Frontiers in Plant Science

[Link to publication](#)

Citation for published version (APA):

Liu, J., Legarrea, S., & Kant, M. R. (2017). Tomato reproductive success is equally affected by herbivores that induce or that suppress defenses. *Frontiers in Plant Science*, 8, [2128]. <https://doi.org/10.3389/fpls.2017.02128>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.


UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Supplementary Material


Tomato reproductive success is equally affected by herbivores that induce or that suppress defenses

Jie Liu, Saioa Legarrea, Merijn R. Kant*

* **Correspondence:** Corresponding Author: m.kant@uva.nl



Supplementary Figure 1. Seeds germination rates. The figure shows the germination rate of the seeds across the treatments. Bars annotated with different letters were significantly different according to Fisher's LSD test ($P < 0.05$) after ANOVA.



Supplementary Figure 2. The effect of the treatments on plant biomass as obtained from the third replicate experiment. The figure shows (A) total plant fresh weight (shoots plus roots) across the treatments, (B) fruit fresh weight relative to plant fresh weight (shoots plus roots) across the treatments and (C) viable seeds relative to total mass (shoots plus roots plus fruits). FW = fresh weight.