



## UvA-DARE (Digital Academic Repository)

### A bright mid-infrared excess in MAXI J1820+070

Russell, D.M.; Baglio, M.C.; Bright, J.; Fender, R.; Al Qaissieh, T.; Palado, A.; Gabuya, A.; Asmus, D.; Belloni, T.; Bel, M.C.; Casella, P.; Ceccobello, C.; Corbel, S.; Diaz Trigo, M.; Gallo, E.; Gandhi, P.; Homan, J.; Koljonen, K.; Lewis, F.; Markoff, S.; Miller-Jones, J.C.A.; O'Brien, K.; Russell, T.D.; Saikia, P.; Sivakoff, G.; Shahbaz, T.; Soria, R.; Tetarenko, A.; van den Ancker, M.

#### Publication date

2018

#### Document Version

Final published version

#### Published in

The astronomer's telegram

#### License

Unspecified

[Link to publication](#)

#### Citation for published version (APA):

Russell, D. M., Baglio, M. C., Bright, J., Fender, R., Al Qaissieh, T., Palado, A., Gabuya, A., Asmus, D., Belloni, T., Bel, M. C., Casella, P., Ceccobello, C., Corbel, S., Diaz Trigo, M., Gallo, E., Gandhi, P., Homan, J., Koljonen, K., Lewis, F., ... van den Ancker, M. (2018). A bright mid-infrared excess in MAXI J1820+070. *The astronomer's telegram*, 11533. <http://www.astronomerstelegram.org/?read=11533>

#### General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

#### Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: [Library@uva.nl](mailto:Library@uva.nl), P.O. Box 1001, 1000 AA Amsterdam, The Netherlands. You will be contacted as soon as possible.

## Outside

GCN  
IAUCs

## Other

ATel on Twitter and Facebook  
ATELstream  
ATel Community Site

This space for free for your conference.

[ [Previous](#) | [Next](#) | [ADS](#) ]

## A bright mid-infrared excess in MAXI J1820+070

ATel #11533; *David M. Russell, M. Cristina Baglio (NYU Abu Dhabi), Joe Bright, Rob Fender (Univ. Oxford), Thabet Al Qaissieh, Alejandro Palado, Aldrin Gabuya (Al Sadeem Observatory, Abu Dhabi), Daniel Asmus (ESO, Santiago & Univ. Southampton), Tomaso Belloni (INAF-OAB), Marion Cadolle Bel (MPCDF), Piergiorgio Casella (INAF-OAR), Chiara Ceccobello (Chalmers Univ. of Technology, Sweden), Stephane Corbel (DAP-AIM, CEA, Paris Diderot Univ.), Maria Diaz Trigo (ESO, Garching), Elena Gallo (Univ. Michigan), Poshak Gandhi (University of Southampton), Jeroen Homan (Eureka Scientific), Karri Koljonen (FINCA, Univ. Turku), Fraser Lewis (Faulkes Telescope Project & Astrophysics Research Institute, LJMU), Sera Markoff (Univ. Amsterdam), James C. A. Miller-Jones (ICRAR-Curtin Univ.), Kieran O'Brien (Durham Univ.), Thomas D. Russell (Univ. Amsterdam), Payaswini Saikia (NYU Abu Dhabi), Gregory Sivakoff (Univ. Alberta), Tariq Shahbaz (IAC & ULL), Roberto Soria (NAOC, Chinese Academy of Sciences & ICRAR-Curtin Univ.), Alex Tetarenko (Univ. Alberta), Mario van den Ancker (ESO, Garching)*  
on 12 Apr 2018; 20:57 UT  
Credential Certification: *David M. Russell (dave.russell5@gmail.com)*

Subjects: Radio, Far-Infra-Red, Infra-Red, Optical, X-ray, Binary, Black Hole, Transient

Referred to by ATel #: [11539](#), [11540](#), [11723](#), [12128](#), [12534](#)

MAXI J1820+070 (ASASSN-18ey) is a black hole candidate X-ray binary (e.g. ATel #[11399](#), #[11418](#), #[11420](#)). It is currently in a bright, hard spectral state (ATel #[11423](#), #[11427](#), #[11439](#)). Here, we report photometric mid-IR (5-12  $\mu\text{m}$ ) observations of the source with the VLT Imager and Spectrometer for the mid-InfraRed (VISIR; Lagage et al. 2004, The Messenger, 117, 12) mounted on UT3 of ESO's Very Large Telescope. The observations were made on 2018 April 8 (07:21-08:08 UTC) and April 9 (08:54-09:41 UTC) in clear conditions (MJD 58216-7). We also report on contemporaneous radio and optical data. MAXI J1820+070 is clearly detected (S/N > 200) in all four mid-IR filters. The following preliminary observed flux densities ( $F_{\nu}$ ) were calculated using observations of standard stars taken on the same nights:

Filter - Wavelength( $\mu\text{m}$ ) -  $F_{\nu}$ (mJy; Apr 8) -  $F_{\nu}$ (mJy; Apr 9)

B11.7 - 11.52 - 376 - 378

B10.7 - 10.65 - 353 - 368

J8.9 - 8.72 - 303 - 297

M-band - 4.85 - 292 - 271

Photometric errors are small but systematic errors are up to ~5-15% due to the limited number of available standard stars. These mid-IR flux densities are amongst the brightest reported in a transient low-mass X-ray binary.

## Related

- 12608** Optical spectroscopy and photometry of MAXI J1820+070 (ASASSN-18ey) during the large multi-wavelength re-brightening of March 2019
- 12596** Optical observations of MAXI J1820+070 confirm the re-brightening
- 12577** AMI-LA and Swift confirm the multi-wavelength re-brightening of MAXI J1820+070
- 12573** Swift observation of the re-brightening in MAXI J1820+070
- 12567** Re-brightening of ASASSN-18ey = MAXI J1820+070
- 12534** MAXI J1820+070 is close to quiescence
- 12157** MAXI 1820+070 has completed the decline from the recent optical re-brightening following the soft to hard transition
- 12128** Optical brightening of MAXI J1820+070 over the soft to hard transition observed with LCO and the Al Sadeem Observatory
- 12068** MAXI J1820+070 continuing its rapid evolution toward the hard state
- 12064** Swift observes MAXI J1820+070 in transition from the soft to the hard-intermediate state
- 12061** AMI radio detection of the black hole candidate MAXI J1820+070 during the soft to hard transition.
- 12057** MAXI/GSC detection of a rapid increase in the hard X-ray flux of MAXI J1820+070
- 11951** Detection of a 55 Hz high-frequency QPO in MAXI J1820+070 with NICER
- 11936** Optical timing observations of MAXI J1820+070 with IFI+IQUEYE and AQUEYE+ soon after state transition
- 11899** Short-lived episodes of emission line splitting in the candidate black hole X-ray binary MAXI 1820+070
- 11887** LOFAR observations of MAXI J1820+070 (ASASSN-

The Arcminute Microkelvin Imager Large Array (AMI-LA) observed MAXI J1820+070 (ATel #11420) on 2018 April 8 (06:43-10:46 UTC) at 15.5 GHz. We reduced the data using the `reduce_dc` pipeline. We then performed cleaning (with natural weighting) and additional flagging in CASA. Using IMFIT in CASA, we measured a flux density of the unresolved point source of  $42 \pm 2$  mJy.

The **Al Sadeem Observatory** (Owner/Co-founder Thabet Al Qaissieh, Director/Co-founder Alejandro Palado, Resident Astronomer Aldrin B. Gabuya) is located in Al Wathba South, outside the city of Abu Dhabi in the United Arab Emirates. We observed MAXI J1820+070 on 2018 April 7 (22:21-22:34 UTC) with the Meade LX850 16-inch (41-cm) telescope with an SBIG STT-8300 camera. 20 images (each 30 sec.) were taken in the 'green' Baader LRGB CCD-Filter (similar bandpass to Johnson V-band). The images were bias/dark-subtracted and flat fielded. We measured a brightness of  $V = 12.05 \pm 0.04$  mag, calibrated using several APASS stars in the field. From our Las Cumbres Observatory (LCO) optical monitoring (ATel #11418) we measure magnitudes (using PanStarrs calibration) of  $g' = 12.09 \pm 0.02$ ,  $r' = 12.14 \pm 0.03$ ,  $i' = 12.09 \pm 0.04$ ,  $y = 12.08 \pm 0.16$  on 2018 April 8 (~7 UTC).

We de-reddened the above flux densities using an extinction of  $E(B-V)=0.163$  (ATel #11418), and constructed the broadband, radio to optical spectral energy distribution (SED; figure linked below). For comparison, we also include a radio, near-IR and optical SED from 2018 March 18 (data from ATel #11439, #11458 and LCO). The mid-IR fluxes clearly represent an excess over the optical emission, and appear to lie close to the extrapolation of the near-IR excess previously reported in ATel #11458. The SED is similar to those of GX 339-4 (Gandhi et al. 2011, ApJ, 740, L13) and MAXI J1836-194 (Russell et al. 2014, MNRAS, 439, 1390) in which the IR was shown to be produced by synchrotron emission from the jet. The SED shape suggests that the jet break between optically thin and partially self-absorbed synchrotron may reside within the mid-IR to far-IR range. Assuming a simple power law, the radio to mid-IR spectral index is  $\alpha \sim +0.3$  (where  $F_\nu \propto \nu^\alpha$ ). By fitting the April 8 VISIR data and y-band point, we obtain  $\alpha \sim -0.7$ .

We encourage coordinated observations during the outburst decay to track the evolution of this spectrum. The LCO observations are part of an on-going **monitoring campaign** of ~ 40 low-mass X-ray binaries (Lewis et al. 2008) with LCO and the Faulkes Telescopes.

### *MAXI J1820+070 light curves and SEDs*

	18ey) during its recent state transition
11855	Polarimetric monitoring of the MAXI J1820+070 in optical and near-infrared wavelengths
11833	Declining near-infrared flux from the black-hole candidate MAXI J1820+070 (ASASSN-18ey) in transition
11831	(Sub)-millimetre Observations of MAXI J1820+070 (ASASSN-18ey) Suggest Jet Quenching on July 6
11827	AMI-LA 15.5 GHz observations of radio flaring from the black hole candidate MAXI J1820+070 in transition
11824	Other low-frequency optical QPO-like features in MAXI J1820+070 detected with IFI+IQUEYE@Galileo
11823	Continuing NICER observations of the state transition in ASASSN-18ey/MAXI J1820+070
11820	A rapid state transition in MAXI J1820+070
11756	17-Hour Period in V light from MAXI J1820+070 = ASASSN-18ey
11723	Low-frequency optical QPO in MAXI J1820+070 detected with IFI+IQUEYE@Galileo
11661	KEGS Discovery of 9 Supernova Candidates in the K2 Campaign 17 field with Pan-STARRS PS1
11609	Simultaneous LOFAR and AMI-LA observations of MAXI J1820+070
11596	MAXI J1820+070 has optical period of 3.4 hours
11591	Further detection of the optical low frequency QPO in the black hole transient MAXI J1820+070
11578	Exponential increase in X-ray QPO frequency with time in MAXI J1820+070
11576	NICER observations of MAXI J1820+070: Continuing evolution of X-ray variability properties
11574	Optical/X-ray Flux Decoupling in MAXI J1820+070
11540	VLITE meter-wavelength detection of MAXI J1820+070 at 339 MHz
11539	The 30-day monitoring of MAXI J1820+070 at 4.7 GHz
11533	A bright mid-infrared excess in MAXI J1820+070
11510	Detection of optical and X-ray QPOs at similar frequencies in MAXI J1820+070
11490	INTEGRAL observations of MAXI J1820+070: public data products
11488	Low-frequency QPOs in MAXI J1820+070 as seen by INTEGRAL/SPI
11482	Palomar 60-inch SEDM classification of optical transients
11481	MAXI J1820+070: VLT and GTC spectroscopic follow-up shows a significant spectral

	evolution from the early stages of the outburst
11480	ePESSTO spectroscopic classification of optical transients
11478	INTEGRAL observations of MAXI J1820+070
11462	Near Infrared JHKs observations of the transient MAXI J1820+070 / ASASSN-18ey: Erratum on 2MASS counterpart designation
11458	Near Infrared JHKs observations of the transient MAXI J1820+070 / ASASSN-18ey
11451	Fast infrared photometry of the black-hole candidate MAXI J1820+070
11445	First measurements of linear polarization of MAXI J1820+070
11440	NOEMA Sub-millimetre Detection of MAXI J1820+070
11439	A flat radio spectrum of MAXI J1820+070
11437	Red sub-second optical flaring in MAXI J1820+070 observed by ULTRACAM/NTT
11432	Correlated Optical/X-ray Timing Variations in MAXI J1820+070 found by Swift UVOT and XRT
11427	The hard X-ray spectrum of MAXI J1820+070 observed by Swift/BAT
11426	Detection of 10-msec scale optical flares in the black-hole binary candidate MAXI J1820+070 (ASASSN-18ey)
11425	Optical Spectra of MAXI J1820+070 with Keck
11424	SOAR/Goodman optical spectroscopy of MAXI J1820+070
11423	NICER observations of MAXI J1820+070 suggest a rapidly-brightening black hole X-ray binary in the hard state
11421	Fast optical flaring in the suspected black-hole binary MAXI J1820+070 (ASASSN-18ey)
11420	AMI radio observations of the black hole candidate MAXI J1820+070
11418	Optical observations of MAXI J1820+070 suggest it is a black hole X-ray binary
11406	MAXI J1820+070: Errata and updated XRT Position
11404	MAXI J1820+070: Swift/UVOT counterpart correction
11403	Swift detection of MAXI J1820+070
11400	Optical follow-up of MAXI J1820+070 and possible identity with ASASSN-18ey
11399	MAXI/GSC detection of a probable new X-ray transient MAXI J1820+070
10459	Ongoing radio monitoring of Cyg X-1 with the RATAN-600 radio telescope
10446	Change in radio behaviour of Cygnus X-1
10322	Unusual soft X-ray activity of Cygnus X-1 detected with MAXI/GSC
9089	New RATAN-600 data for Cygnus X-1

---

[ **Telegram Index** ]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org

Mansi M. Kasliwal, Co-Editor

mansi@astronomerstelegam.org