



UvA-DARE (Digital Academic Repository)

ATCA radio detection of the new X-ray transient MAXI J1813-095 as a candidate radio-quiet black hole X-ray binary

Russell, T.D.; Miller-Jones, J.C.A.; Sivakoff, G.R.; Tetarenko, A.J.; JACPOT XRB collaboration

Publication date

2018

Document Version

Final published version

Published in

The astronomer's telegram

License

Unspecified

[Link to publication](#)

Citation for published version (APA):

Russell, T. D., Miller-Jones, J. C. A., Sivakoff, G. R., Tetarenko, A. J., & JACPOT XRB collaboration (2018). ATCA radio detection of the new X-ray transient MAXI J1813-095 as a candidate radio-quiet black hole X-ray binary. *The astronomer's telegram*, 11356. <http://www.astronomerstelegam.org/?read=11356>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.


UvA-DARE is a service provided by the library of the University of Amsterdam (<https://dare.uva.nl>)

Outside

GCN
IAUCs

Other

ATel on [Twitter](#) and [Facebook](#)
[ATELstream](#)
[ATel Community Site](#)



[[Previous](#) | [Next](#) | [ADS](#)]

ATCA radio detection of the new X-ray transient MAXI J1813-095 as a candidate radio-quiet black hole X-ray binary

ATel #11356; *T. D. Russell (UvA), J. C. A. Miller-Jones (ICRAR-Curtin), G. R. Sivakoff, A. J. Tetarenko (UAlberta) and the JACPOT XRB collaboration*

on 26 Feb 2018; 10:08 UT

Credential Certification: Thomas Russell (t.d.russell@uva.nl)

Subjects: Radio, X-ray, Black Hole, Neutron Star, Transient

Referred to by ATel #: [11357](#)

We observed the new X-ray transient MAXI J1813-095 (ATels #[11323](#), #[11326](#), #[11332](#)) with the Australia Telescope Compact Array (ATCA) between 2018-02-22 20:52 UT and 2018-02-23 02:59 UT. Our observations were taken simultaneously at 5.5 and 9 GHz, with a bandwidth of 2 GHz at each frequency. We used 1934-658 for flux calibration and 1819-096 for phase calibration. Data were reduced in CASA version 4.7.0 (McMullin et al 2007). Imaging was carried out using a robustness of 0 and flux densities were measured by fitting for a point source in the image plane. Due to its compact configuration (750B), we did not include the isolated antenna 6 when imaging, leading to angular resolutions of 108" x 7.31" and 68.4" x 4.63" at 5.5 and 9 GHz, respectively, at a position angle of -177.8 degrees east of north.

We significantly detect MAXI J1813-095 at the position (measured at 9 GHz):

RA (J2000): 18:13:34.073 +/- 0.014

Dec (J2000): -9:32:07.3 +/- 3.7

where the 1-sigma errors are the larger of the statistical error from the fitted position (Declination error) or the theoretical statistical error of centroiding ($\text{Beam}/2 \cdot \text{SNR}$; Right Ascension error). There may be larger systematic errors that we do not include. Using Galactic disk and bulge stellar populations (Juric et al 2008; McMillan 2011) and excluding the potential effect of any natal supernova kick, a source along this line of sight has a likely distance of 8^{+6}_{-2} kpc (1 σ confidence interval). We measure flux densities of 0.69 +/- 0.06 mJy and 0.47 +/- 0.04 mJy at 5.5 and 9 GHz, respectively, implying a radio spectral index of $\alpha = -0.7 \pm 0.2$, where $S_\nu \propto \nu^\alpha$.

Given this spectral index, the radio emission converts to a 5-GHz luminosity of $4 \cdot \pi \cdot \nu \cdot f_\nu \cdot d^2 = (3.3 \pm 0.4) E29 \cdot (d/8\text{kpc})^2 \text{ erg/s}$.

Swift/XRT observed MAXI J1813-095 on 2018-02-21 (between 11:40:06 and 11:58:12 UT) and 2018-02-23 (22:25:00 to 22:44:00 UT), bracketing our radio observation. We extracted the data using the online XRT pipeline (Evans et al. 2009) and fit with an absorbed power-law model. On

Related	
11357	INTEGRAL observations of MAXI J1813-095
11356	ATCA radio detection of the new X-ray transient MAXI J1813-095 as a candidate radio-quiet black hole X-ray binary
11332	MAXI J1813-095: GROND discovery of a candidate optical/NIR counterpart
11326	MAXI J1813-095: Swift detection and localization
11323	MAXI/GSC detection of a new X-ray transient MAXI J1813-095

2018-02-21 we find a best fit with an N_H of $(9.6 \pm 0.2)E21$ and a photon index of 1.45 ± 0.07 , providing an unabsorbed 1-10 keV X-ray flux of $(7.8 \pm 0.2)E-10$ erg/s/cm², corresponding to a 1-10 keV X-ray luminosity of $(5.95 \pm 0.15)E36*(d/8kpc)^2$ erg/s. On 2018-02-23, we find an N_H of $(1.1 \pm 0.3)E21^2$ and a photon index of 1.5 ± 0.1 , providing an unabsorbed 1-10 keV X-ray flux of $(7.9 \pm 0.3)E-10$ erg/s/cm², corresponding to a 1-10 keV X-ray luminosity of $(6.05 \pm 0.15)E36*(d/8kpc)^2$ erg/s.

The X-ray luminosities and spectrum are consistent with the source being in the low-Eddington part of the hard state, typical of the early part of an outburst. This state is associated with a compact steady jet, as opposed to jet ejecta (which is associated with the transition between the hard and soft state at higher luminosities). Espinasse and Fender (2018) suggested that radio-quiet hard state black hole X-ray binaries (BHXBs) have steeper indices ($\alpha=-0.2\pm 0.3$) than radio-loud hard state BHXBs ($\alpha=0.2\pm 0.2$). We measure a spectral index that is consistent with the optically thin spectra expected from jet ejecta ($\alpha\sim 0.6$), potentially consistent with compact jets in radio-quiet hard-state BHXBs, and inconsistent with compact jets in radio-loud hard-state BHXBs.

Comparing the luminosities to those of hard-state XBs, MAXI J1813-095 is consistent with radio-quiet BHXBs and radio-loud neutron star XBs; however, we caution that this correlation would not apply to jet ejecta. Given the consistency of the X-ray properties, the radio spectral index and the radio/X-ray correlation with the properties of radio-quiet BHXBs, MAXI J1813-095 may be a radio-quiet BHXB.

We thank Jamie Stevens and ATNF staff for making these observations possible.

[**Telegram Index**]

R. E. Rutledge, Editor-in-Chief

rrutledge@astronomerstelegam.org

Derek Fox, Editor

dfox@astronomerstelegam.org

Mansi M. Kasliwal, Co-Editor

mansi@astronomerstelegam.org