



Animal Performance in Pastures of *Panicum maximum* cv. IPR 86 Fertilized with Nitrogen

S. M. B. Lugão

Instituto Agronômico do Paraná, Brazil

L. R. de A. Rodrigues

Universidade Estadual Paulista, Brazil

E. B. Malheiros

Universidade Estadual Paulista, Brazil

J. J. dos S. Abrahão

Instituto Agronômico do Paraná, Brazil

A. de Moraes

Universidade Federal do Paraná, Brazil

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/19/22/27>

This collection is currently under construction.

The XIX International Grassland Congress took place in São Pedro, São Paulo, Brazil from February 11 through February 21, 2001.

Proceedings published by Fundacao de Estudos Agrarios Luiz de Queiroz

**ANIMAL PERFORMANCE IN PASTURES OF *Panicum maximum* cv. IPR 86
FERTILIZED WITH NITROGEN**

S.M.B. Lugão¹, L.R. de A. Rodrigues², E. B. Malheiros³, J.J. dos S. Abrahão⁴, and
A. de Moraes⁵

¹Depto de Zootecnia, IAPAR, Paranavaí, PR, Brasil, 87701-970, simonyl@pr.gov.br

²Depto de Zootecnia, FCAV-UNESP, Jaboticabal, SP, Brasil, 14884-900

³Depto de Ciências Exatas, FCAV-UNESP, Jaboticabal, SP, Brasil, 14884-900

⁴Depto de Zootecnia, IAPAR, Paranavaí, PR, Brasil, 87701-970

⁵Depto de Fitotecnia, UFPR, Curitiba, PR, Brasil, 80035-050

Abstract

The experiment was carried out at the Experimental Station of the IAPAR in Paranavaí–PR, Brazil, from October 1998 to May 1999 to evaluate the response of crossbred steers grazing pastures of *Panicum maximum* cv. IPR 86 fertilized with four doses of N (0; 150; 300 and 450 kg.ha⁻¹.year). The grass was grazed according to a rotational grazing system with grazing cycles (GCs) of 40 days (grazing period of 5 days and rest period of 35 days). It was used a randomized complete block design, with split-plots, and two replications per treatment, being the N-doses studied in the main plots and the GCs in the sub-plots. The availability of green leaf-blades increased with the increase of N-doses, resulting in 4.09; 8.18; 11.18; and 13.92 steers.ha⁻¹.day⁻¹ and daily LWG of 2.73; 5.25; 7.40; and 9.24 kg.ha⁻¹ when the pastures were fertilized with 0; 150; 300 and 450 kg N.ha⁻¹, respectively. The efficiency of N-utilization was 3.35; 3.11; and 2.89 kg LWG kg.N⁻¹ when the pastures were fertilized with 150; 300; and 450 kg N.ha⁻¹, respectively.

Keywords: carrying capacity, liveweight gain, nitrogen, *Panicum maximum*, stocking rate

Introduction

Several cultivars of *Panicum maximum* Jacq. recently released in Brazil were well accepted by the farmers due to their adaptation to different edaphic and climatic conditions as well as their high potential for forage production. Other accesses, such as BRA-006998, registered as cv. IPR 86, are still being studied by the research Institutions and may be released in a near future. This species has a high requirement of N, especially in soils with low levels of organic matter. There are many papers reporting the responses of tropical grasses to the application of a wide range of N-doses. Unfortunately there are not too many papers reporting animal production under these conditions. The objective of this work was to determine the effects of four N-doses on forage production and animal performance in pastures established with the new cultivar IPR 86 in the northwest region of the State of Paraná, Brazil.

Material e Methods

The experiment is being conducted at the Experimental Station of the IAPAR in Paranavaí-PR, Brazil, (23⁰⁰'05'' S, 42⁰⁰'02'06'' W, and 480m above sea level). It was initiated in 11/1997 and will be concluded in 11/2001. The data presented in this paper were obtained from October 15, 1998 to May 03,1999 (200 days).

The region has a subtropical climate (Cfa – humid and mesothermic) with an annual average temperature of 22⁰ C and 1500 mm of rainfall (Muzilli et al., 1990). The soil of the experimental area was classified as red-yellow Podzolic, sand-texture, flat and well drained.

The treatments consisted of four N-doses (0; 150; 300; and 450 kg.ha⁻¹) applied as calcium nitrate and distributed throughout the growing season (Nov., Jan., and Mar.) according to a randomized complete block design, with split-plots, and two replications per treatment, being the N-doses studied in the main plots and the grazing cycles (GCs) in the sub-plots. The area was divided in 8 paddocks to allow a rotational grazing system (variable stocking) with a grazing period of 5 days and a rest period of 35 days. Treatments 0 and 150 kg N.ha⁻¹ were allocated to paddocks with 1.35 ha, and treatments 300 and 450 kg N.ha⁻¹ to paddocks with 0.68 ha. Crossbred steers (Nelore x Red Angus, Nelore x Marchigiana, and Nelore x Simental) with 12 months of age and an average LW of 248 kg were equally distributed by the treatments. Three animal-testers were maintained in each treatment throughout the experimental period. The stocking rate (SR) adjustment was done considering that 70 % of green leaf-blades (GLB) DM produced would be consumed by the animals, allowing an intake of 11 kg of GLB-DM.AU⁻¹. The SR was adjusted every 10 days, after sampling to determine forage availability. The animals received water and mineral mixture “ad-libitum” and preventive applications of vermifuge against endo and ectoparasites.

The effects of N-doses were evaluated by regression analysis using the procedures GLM and REG (SAS, 1990).

Results and Discussion

The DM availability of GLB before grazing increased with the increase in the N-doses presenting the following overall means for GC: 1136; 2857; 3663; and 4384 kg.ha⁻¹, respectively for the treatments 0; 150; 300; and 450 kg N.ha⁻¹. Increases in DM availability with the increase in N-doses up to 500 kg N.ha⁻¹ were also observed in a rotational grazing system using *P. maximum* (Crespo, 1986).

The average daily gain (ADG) was not affected ($P>0,01$) by N-doses with values of 0.668; 0.641; 0.644; and 0.664 $\text{kg}\cdot\text{animal}^{-1}\cdot\text{day}^{-1}$, when the pastures were fertilized with 0; 150; 300; and 450 $\text{kg N}\cdot\text{ha}^{-1}$, respectively. It has been shown that increases in beef cattle production can be obtained due to the increase in the SR associated with the N-fertilization (Setelich et al., 1998). The ADGs obtained in this work are similar to those reported by Euclides (1995) for several cultivars of *P. maximum* but lower than that registered by Tosi (1999) working with crossbred steers in pastures of Tanzania-grass (*P. maximum* cv. Tanzania).

The SR was affected ($P<0.01$) by N-doses (N), by GC and by the interaction NxGC. The effects of the N-doses in each GC is shown in Figure 1. In all GCs the N-fertilization increased the number of steers. $\text{ha}^{-1}\cdot\text{day}^{-1}$. Stocking rates responses to N-fertilization were better in GCs 2 and 1 (Figure 1) with the maximum doses of N resulting in an increase of 4.43 to 3.69 times in the SR when compared to the control. The $\text{LWG}\cdot\text{ha}^{-1}\cdot\text{day}^{-1}$ was affected ($P<0.01$) by N-fertilization (N), by GC and by the interaction N x GC. The effects of N-doses on the LWG were linear (Figure 2). In all GCs occurred an increase in the $\text{gain}\cdot\text{ha}^{-1}\cdot\text{day}^{-1}$ due to the N-fertilization. The maximum N-doses allowed an increase of 4.32; 3.66; 3.32; 3.19; and 2.77 times in the $\text{LWG}\cdot\text{ha}^{-1}$ in the GCs 2; 1; 4; 3; and 5, respectively. The LWGs of 2.73; 5.25; 7.40; and 9.24 $\text{kg}\cdot\text{ha}^{-1}\cdot\text{day}^{-1}$ in the treatments 0; 150; 300; and 450 $\text{kg}\cdot\text{ha}^{-1}$, respectively, indicate the high potential of IPR 86 for DM production of a good quality forage as well as the good performance of the crossbred steers used in the experiment.

The efficiency of N-utilization was 3.35; 3.11; and 2.89 $\text{kg LWG}\cdot\text{kg N}^{-1}$ when the pastures were fertilized with 150; 300; and 450 $\text{kg N}\cdot\text{ha}^{-1}$ respectively. These values are in agreement with those reported for tropical pastures by Gomide (1989).

It is concluded nitrogen fertilization stimulated forage growth resulting in higher pasture carrying capacity and weight gain per hectare.

References

- Crespo G.** (1986) Variation in the response of tropical pastures to nitrogenous fertilizers through the year. 3. Guinea grass (*Panicum maximum* Jacq.) with irrigation. Cuban Journal of Agricultural Science, **20**:78-81.
- Euclides V.P.B.** (1995) Valor alimentício de espécies forrageiras do gênero *Panicum*. In: Peixoto A.M., Moura J.C. de, Faria V.P. de (eds) Anais do Simpósio sobre Manejo da Pastagem, **12**:245-273.
- Gomide J.A.** (1989) Aspectos biológicos e econômicos da adubação de pastagens. In: Favoretto V. e Rodrigues L.R.deA. (eds) Anais do Simpósio sobre Ecossistema de Pastagens, **1**:237-270.
- Muzilli O., Laurente A.C., Llamillo R.F., Fagundes A.C., Fidalski J.A., Ribeiro M.F.S. e Lugão S.M.B.** (1990) Conservação do solo em sistema de produção nas microbacias hidrográficas do Arenito Caiuá do Paraná. (Boletim Técnico 33).
- SAS INSTITUTE INC. SAS / STAT User' s Guide. (1990) Version 6.12 ed., Cary, NC: SAS Institute Inc.. 943p.
- Setelich E.A., Almeida E.X. e Maraschin G.E.** (1998) Resposta à adubação nitrogenada de capim-elefante anão cv. Mott, sob pastejo. Anais da Reunião Anual da Sociedade Brasileira de Zootecnia, **35**:494-496.
- Tosi P.** (1999) Estabelecimento de parâmetros agrônômicos para o manejo e eficiência de utilização de *Panicum maximum* Jacq. cv Tanzânia 1 sob pastejo rotacionado. Tese (Mestrado), Escola Superior de Agricultura "Luiz de Queiroz", Universidade de São Paulo, Brazil.

GC1	$Y = -0.140 + 2.4870 N$	$P < 0.01$	$R^2 = 0.98$
GC2	$Y = -0.415 + 3.5700 N$	$P < 0.01$	$R^2 = 0.99$
GC3	$Y = 1.585 + 3.8020 N$	$P < 0.01$	$R^2 = 0.99$
GC4	$Y = 2.145 + 3.9560 N$	$P < 0.01$	$R^2 = 0.93$
GC5	$Y = 5.496 + 0.0172 N$	$P < 0.01$	$R^2 = 0.91$




Figure 1 - Effects of four nitrogen doses on the number of steers.ha⁻¹.day⁻¹ in five grazing cycles (GC1: □ ----- ; GC2: ○ ---- ; GC3: Δ - - - - - ; GC4: × ——— ; GC5: + ———) in *Panicum maximum* Jacq. cv. IPR 86 pastures from 10/15/1998 to 05/03/1999 (200 days).

GC1	$Y = 1.521 + 0.0109 N$	$P < 0.01$	$R^2 = 0.98$
GC2	$Y = 2.029 + 0.0157 N$	$P < 0.01$	$R^2 = 0.99$
GC3	$Y = 3.486 + 0.0167 N$	$P < 0.01$	$R^2 = 0.99$
GC4	$Y = 3.945 + 0.0174 N$	$P < 0.01$	$R^2 = 0.96$
GC5	$Y = 3.549 + 0.0114 N$	$P < 0.01$	$R^2 = 0.93$




Figure 2- Effects of four nitrogen doses on the liveweight gain of steers in five grazing cycles (GC1: \square ----- ; GC2: \circ ----; GC3: Δ -..... ; GC4: \times ——— ; GC5: +——) in *Panicum maximum* Jacq. cv. IPR 86 pastures from 10/15/1998 to 05/03/1999 (200 days).