

Reciprocal or independent hemispheric specializations: Evidence from cerebral dominance for fluency, faces, and bodies in right- and left-handers

Karlsson, Emma; Johnstone, Leah; Carey, David

Psychology & Neuroscience

DOI:
[10.1037/pne0000269](https://doi.org/10.1037/pne0000269)

E-pub ahead of print: 01/01/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Karlsson, E., Johnstone, L., & Carey, D. (2021). Reciprocal or independent hemispheric specializations: Evidence from cerebral dominance for fluency, faces, and bodies in right- and left-handers. *Psychology & Neuroscience*. <https://doi.org/10.1037/pne0000269>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Reciprocal or independent hemispheric specializations: Evidence
from cerebral dominance for fluency, faces, and bodies in right-
and left-handers**

Emma M. Karlsson, Leah T. Johnstone, and David P. Carey

Perception, Action and Memory Research Group, School of Psychology, Bangor University

Author Note

Emma M. Karlsson, <https://orcid.org/0000-0002-9605-1923>

David P. Carey, <https://orcid.org/0000-0001-8860-9731>

Leah T. Johnstone is now at Sports Psychology Group, UCFB Wembley

We have no known conflict of interest to disclose.

This work was supported by the Leverhulme Trust [grant number RPG-2019-102].

*Correspondence concerning this article should be addressed to Emma M. Karlsson, School of Psychology, Brigantia Building, Penrallt Road, Bangor, LL57 2AS, UK. Email:

e.karlsson@bangor.ac.uk

Page count: 26.

Number of figures: 1; number of tables: 4

Abstract

Objective: There are distinct cortical regions that respond preferentially to human faces and bodies. It is generally accepted that these face- and body-selective regions are lateralized with a preference for the right hemisphere, but unknown how frequently these biases occur or if they are lateralized in a complementary fashion to language processing.

Methods: fMRI was used to examine face and body lateralization in two samples of right-handers (n's = 31 and 18) and left-handers (n's = 43 and 24) with 'typical', left hemisphere, language dominance to examine the frequency of these biases. Crucially, we also recruited individuals with 'atypical', right hemisphere, language dominance (n's = 17 and 10) to examine complementarity with language.

Results: Language typical right-handers had consistent population-level and average right-sided biases for face- and body perception. Language typical left-handers had population-level biases for faces in sample 2, but not sample 1; and for bodies in sample 1 but not sample 2. Language typical left-handers were, on average, right-lateralized for faces in both samples, but right-lateralized for bodies in sample 1 only. Language atypicals did not have a population-level bias for body- or face perception, and were, on average, left-lateralized for faces in sample 1, but not in sample 2. Atypicals were not lateralized for body perception.

Conclusions: These results add to the growing literature which suggests that many right hemisphere processes are not lateralized in a fully complementary fashion to language. Left-handers seem to have more varied lateralization patterns even when language dominance is controlled for.

Keywords: Hemispheric lateralization; functional brain asymmetry; atypical language dominance; handedness

Public Significance statement

This research demonstrates differences in patterns of complementary brain processes across the hemispheres in left-handers and right-handers. We also show that the rare kind of brain dominance for language is not associated with specialization of body and face processing to the non-language hemisphere as a matter of routine. These data are important for models which claim unusual hemispheric dominance in individuals with neurodevelopmental conditions, as well as for our understanding of the nature of brain asymmetries in left-handed people.

Reciprocal or independent hemispheric specializations: Evidence from cerebral dominance for fluency, faces, and bodies in right- and left-handers

Speculation about hemispheric specialization in the human brain predates contemporary neuroimaging by some distance. After the discoveries of language localization and left-hemispheric dominance by Broca, Dax and Wernicke in the late 19th century, the idea of functional specialization in the cortex, within and between-hemispheres, became neurological canon (Hillis, 2007). As the majority of the 19th century cases described were right-handed individuals, the association of language dominance and contralateral control of the writing hand was assumed (Benton, 1972; Eling, 1984; Goodglass & Quadfasel, 1954). Nevertheless, by the 1970s, a general consensus emerged from Wada testing of epileptics (e.g., Rasmussen & Milner, 1977), as well as group studies of aphasia in left- and right-handed patients with unilateral brain damage (e.g., Zangwill, 1967), confirming that roughly 70% of left-handers are lateralized in this “typical” left hemisphere fashion for speech and language, as are 90-95% of right-handed individuals (Carey & Johnstone, 2014).

In sharp contrast, theories about functions that depend more on right hemisphere structures, such as face processing (Della Sala & Young, 2003; Kanwisher et al., 1997; Sergent et al., 1992) or visuospatial attention (Holmes, 1918; Riddoch, 1935; Buxbaum, 2006; Patterson & Zangwill, 1944) were much slower to develop. This literature became quite removed from the research on apraxia and aphasia, leaving a tendency for authors in either domain to refer to little more than right hemisphere dominance for attention (e.g., Blankenberg et al., 2010; Petersen & Posner, 2012), or left hemisphere dominance for speech/language (e.g., Friederici, 2017; or aspects of motor function; Kimura & Archibald, 1974; Geschwind, 1975). In neurology and neuropsychology, it is frequently assumed these two major classes of asymmetries “anti-localize” (i.e., if one function is over-represented in the left hemisphere, the other will be over-represented in the right hemisphere), but there is limited evidence for or against this idea. For example, unlike several papers on unilateral brain damage and presence/absence of aphasia (reviewed in Carey & Johnstone, 2014), very few clinics or

hospitals have published incidence data on neglect or prosopagnosia using standardized tests in the same individuals, recording both handedness (however specified) and side of lesion. A few early studies (e.g., Bryden et al., 1983) used multiple indices for visuospatial dysfunction in large patient groups, but these included measures of different dysfunctions (e.g., constructional apraxia), which can follow lesions of either cerebral hemisphere.

A limited number of neuroimaging studies have tackled questions of complementary specialization, and tend to be focused on language and visuospatial abilities (e.g., Ng et al., 2000; Powell et al., 2012) rather than face perception, the latter of which is implicated in recent models (see below). Nevertheless, attempts at comparing one or several functions in right- and left-handed groups are gradually becoming more common (Badzakova-Trajkov et al., 2010; Flöel et al., 2005; Mazoyer et al., 2016; Van der Haegen et al., 2012; Rosch et al., 2012; Whitehouse & Bishop, 2009; Woodhead et al., 2019, 2021). For instance, one of these experiments was designed to examine complementarity of hemispheric lateralization of attention and language (Cai et al., 2013). Individuals who were identified as right hemisphere dominant for verbal fluency in a previous experiment (Van der Haegen et al., 2011) were re-scanned using a landmark task requiring judgments of pre-bisected lines, known to depend more on right hemispheric mechanisms. Cai et al. (2013) found a near perfect complementarity of these two functions, in left-handed individuals with typical (left; 15 out of 16 individuals) and atypical (right; all 13 individuals) language dominance.

One caveat is that the authors selected individuals with very strong language dominance (discussed in Badzakova-Trajkov et al., 2016; Karlsson et al., 2019), where complementarity of function may be more likely. However, a strength of the Cai et al. (2013) report is that it acknowledges the importance of the individuals with the rarer form of cerebral asymmetry for tests of complementary hemispheric dominance(s). This group of individuals is hugely important, as data for language asymmetry, at the very least, is so skewed towards left hemispheric specialization in right-handed samples that many interesting questions regarding complementarity cannot be addressed by right-handers alone. Any model of complementarity

should apply in the same way to the rare individuals who are right-hemisphere dominant for speech and language.

Large sample neuroimaging studies that measure multiple functions in the same individuals allow for comparison of pairs of functions that depend on the right or the left hemisphere (Badzakova-Trajkov et al., 2010). Complementary specialization is the usual prediction; the lateralization of one function to one hemisphere causes, in some fashion, another function to lateralize to the opposite hemisphere (Bryden, 1990; Bryden et al., 1983). These functions may have been symmetrically located at an initial point, but growth or development of one system (often deemed to be language-related), has a causal role for the other function to lateralize in the opposite direction (Bradshaw & Nettleton, 1981). Many of these models are, in effect, “crowding” hypotheses, that suggest that functional specialization of some neural tissues in one hemisphere leaves “less brain” for the other function, which comes to depend on the other hemisphere.

Current variants of crowding hypotheses of left- and right-hemispheric function are almost exclusively focused on face processing, and their relegation to the right hemisphere as reading develops in foveal regions of occipitotemporal cortex in the left hemisphere (Behrmann & Plaut, 2015; Centanni et al., 2018; Dehaene, et al., 2010; Plaut & Behrmann, 2011). As most people are (largely) innately left hemispheric (Carey & Johnstone, 2014; Holowka & Petitto, 2002; Rasmussen & Milner, 1977) for some of the auditory, somatosensory and motor aspects of speech perception/production, when reading training begins, it makes sense for linguistic systems to co-opt occipitotemporal regions in the same hemisphere. As a result, largely bilateral face processing gradually becomes more right hemispheric (although see Lochy et al., 2017, for a recent critique of the electrophysiological evidence in particular).

In contrast, the statistical hypothesis (Bryden, 1990; Bryden & Allard, 1981) assumes that whatever underlies the lateralization of one function is independent of that of another function and may reflect independent probabilistic biases. Each function has an independent statistical probability of being lateralized to the right or the left hemisphere, and the fact a

population level bias for language lateralized to the left hemisphere and, for example, face processing in the right hemisphere may arise, but reflect probabilities relating to independent causal sources.

In the current context, a first step in the evaluation of complementary hemispheric specialization requires estimating the frequency with which face and language hemispheric bias in groups that are lateralized in the more common pattern (i.e., left hemispheric for language, right hemispheric for faces) and quantifying the precision of that estimate. A second crucial step in such an evaluation is to collect data from individuals with the rarer types of asymmetry, in this instance right hemispheric dominance for language and left hemispheric dominance for faces. An important recent example of this approach is by Gerrits et al. (2019). These colleagues capitalized on a pool of left-handers with previously identified atypical (right hemispheric) language dominance, to examine asymmetry related to face- and visual word recognition. Their sample included 12 left-handers with atypical dominance for language and 12 left-handers with typical (left hemisphere) dominance. In this sample, 75 percent of the typically-lateralized individuals processed faces in the non-language hemisphere, but only 42 percent of individuals with atypical language dominance did so. In a second study, Gerrits and colleagues (2020) examined hemispheric dominance for five different functions (language, praxis, emotional prosody, spatial attention, and face recognition) in a larger sample of individuals. In contrast to the earlier study, they found strong evidence for complementarity (ranging from 75 - 87%) for language typical and atypical participants for all functions, except for face perception in the language typical left-handers, who surprisingly did not have a population level bias on the task.

The studies from Gerrits et al. (2019; 2020) are important for efforts of quantifying how asymmetries relate to one another. However, both studies lack right-handed control groups. Right-handed participants are crucial for teasing apart potential handedness effects from potential language dominance effects (and are also important for relating handedness differences to genetic models, see below). There is some evidence that left-handers may be

more variable in their brain dominance for right-hemisphere functions in general (Karlsson et al., 2019; Liu et al., 2009). The ideal experiment would, of course, also include a group of right-handed language atypical participants, but these individuals are rare (about 5 percent). Although our and other labs have started to search for these individuals, these efforts will take time.

Here, we use fMRI to explore co- and anti-lateralization of three different functional asymmetries (language, face perception, body perception) in right- and left-handers with typical and atypical language dominance in two samples of participants. The data in sample 2 was collected as part of a larger study effort to examine multiple asymmetries in large numbers of language typical and atypical participants, but gave us the opportunity to replicate in part, the findings of sample 1.

Body perception was included as it has been linked to the right hemisphere from the first studies identifying specialized brain regions. Downing and colleagues (2001) found that human bodies and body parts preferentially activated an area in the lateral occipitotemporal cortex in the right hemisphere in 19 participants that were scanned, with a majority of these showing a significant, but weaker, activation in the left hemisphere. The lateralization of body selective activity is generally accepted in the body perception literature (it is common practice to only define and report data obtained from regions of interest in the right hemisphere, e.g., Peelen et al., 2009; Schwarzlose et al., 2005; Taylor et al., 2007), but not well characterized from an asymmetry perspective (although see Willems et al., 2010).

A verbal fluency task was also included to examine hemispheric dominance for language processing, to examine its' complementarity with these right hemisphere functions. The main goal of this project was to examine both the *breadth* (i.e., the frequency) of the asymmetry for face and body perception in right- and left-handers with known hemispheric dominance for speech/language, and its *depth* (i.e., magnitude). In other words, how many of the participants demonstrate right hemisphere dominance for face and body perception, and the size of the asymmetry. Most right-handed and left-handed participants that have confirmed

RECIPROCAL OR INDEPENDENT HEMISPHERIC SPECIALIZATIONS

9

leftward asymmetry for language should be right hemispheric for these functions, for strong evidence of complete complementarity. Likewise, individuals with right hemisphere (atypical) language processing should show the reversed relationship and the breadth of asymmetry favoring their non-language dominant hemisphere should be similar in magnitude to that seen in language typical individuals. Furthermore, left-handed language typicals should show similar breadth and depth as the right-handed typicals if these functions are related to language, but not handedness per se.

Methods

Participants

Sample 1

Ninety-three participants from the Bangor University community took part; 60 were left-handed (22 female), and 33 right-handed (21 female). Two participants (left-handed, one female and one male) were excluded from the final analysis due to excessive head movement ($> 4\text{mm}$). The left-handed participants had a mean age of 24.83 ($SD = 7.55$), and a mean Waterloo handedness questionnaire (WHQ; Steenhuis, & Bryden, 1989: ranging from -30 always left for each item, to +30 always right) score of -20.38 ($SD = 13.31$). The right-handed participants had a mean age of 26.09 ($SD = 5.92$) and a mean WHQ score of +28.13 ($SD = 2.58$). Data from these scans were subjected to a split-half reliability analysis in a previous paper (Johnstone et al., 2020). This study was approved by the Bangor Psychology ethics committee.

Sample 2

Data from 21 participants (8 right-handed and 13 left-handed) who had already taken part in sample 1 were excluded from this analysis. A total of 52 participants from the Bangor University community took part; 33 were left-handed (19 female), and 19 right-handed (15 female). The left-handed participants had a mean age of 23.91 ($SD = 6.93$), and a mean WHQ score of -20.75 ($SD = 13.69$). The right-handed participants had a mean age of 23.05 ($SD = 4.64$) and a mean WHQ score of +28.05 ($SD = 2.01$). This study was approved by the Bangor Psychology ethics committee.

Tasks***Verbal fluency***

A single letter verbal fluency paradigm was used in a blocked design. For sample 1, the paradigm consisted of 14 experimental, 14 control blocks, and 30 rest blocks: each with a duration of 15 seconds. In experimental blocks, participants were presented with a single letter in the middle of the screen which stayed on for the duration of the block. The letters used were those that begin most words in English: T, A, S, H, W, I, O, B, M, F, C, L, D, P (as reported in the Natural Language Toolkit 3.0 - <http://www.nltk.org/>). Participants were instructed to think of as many words as they could which began with that letter during the time the letter was shown on screen. A practice was run outside the scanner using the letter "D". In control blocks, participants were shown one of two letter strings ("RARA" or "LALA") and were instructed to covertly repeat these for as long as they were presented on the screen. A fixation cross was presented in the 30 rest blocks, and participants were instructed to relax and clear their minds. This experiment was presented across two runs, comprising seven experimental/control blocks respectively and 15 rest blocks per run. The letters were randomly presented in any order across these two runs.

The verbal fluency task for sample 2 was identical to the one used for sample 1, but with only one run of data collected. The seven different letters 'T' 'A' 'S' 'H' 'W' 'I' 'O' were used.

Face/body localizers

Brain activation for face- and body-related activation was obtained with a four-condition visual localizer in sample 1. This involved viewing blocks of images from the stimulus categories: faces, bodies, chairs, and scenes. Participants completed two runs of this task, with two different fixed stimulus orders, which were counterbalanced across participants. Each localizer run consisted of 16 active blocks (4 for each stimulus category) and 5 rest blocks where a central fixation point was shown on the screen. Each block lasted 16 seconds during which 16 images were displayed for 300 ms followed by a blank screen for 700 ms.

Participants completed a one-back task, pressing a button if they saw a consecutive, repeated image. Which hand participants held the button box in was counterbalanced within the right-handed and left-handed groups.

Two four-condition localizers were used to identify body- and face-selective brain areas for sample 2. These were identical in implementation to the localizer used in sample 1, but with different stimulus categories. The body localizer consisted of images of hands, hand-held tools, human bodies without heads, and chairs; the face localizer of images of faces with neutral expressions, faces with emotional expressions, butterflies, and flowers. Bodies were of 10 males and 10 females, and chairs were 20 various chairs in different styles. The neutral faces consisted of 10 male faces and 10 female faces from The Karolinska Directed Emotional Faces (KDEF) database (Lundqvist, Flykt, & Öhman, 1998) with neutral facial expressions. The butterflies and flowers were both of images depicting 20 unique variants of butterflies and flowers.

fMRI data acquisition

The scans were acquired in a Philips 3 Tesla Achieva magnetic resonance (MR) scanner at the Bangor Imaging Unit at Bangor University, using a 32-channel head coil. Functional images in sample 1 were acquired with a T2-weighted gradient-echo EPI sequence, field of view (FOV) = 220 x 220, acquisition matrix = 96 x 96, 36 slices; acquired voxel size (mm) = 2.3 x 2.3 x 2.5 (reconstructed voxel size (mm) = 2.3 x 2.3 x 2.5). Verbal fluency (repetition time (TR) = 2500 ms, echo time (TE) = 30 ms, flip angle (FA) = 90°) consisted of two runs of 435 seconds each, and the face/body localizer (TR = 2000 ms, TE = 30 ms, FA = 90°) consisted of two runs of 332 seconds. Functional images for verbal fluency in sample 2 were acquired with slightly different parameters: a T2-weighted gradient-echo single-shot EPI pulse sequence (SENSE, acceleration factor = 2); TR = 2500 ms, TE = 30 ms, acquisition time = 435 seconds, FA = 83°, FOV = 240 x 240 x 105, acquisition matrix = 80 x 79 x 35; 35 slices (width = 3 mm, no gap); acquired voxel size (mm) = 3 x 3 x 3 (reconstructed voxel size (mm)

RECIPROCAL OR INDEPENDENT HEMISPHERIC SPECIALIZATIONS

13

= 3 x 3 x 3). Functional images for the body- and face localizers in sample 2 were acquired with the following parameters: a T2-weighted gradient-echo single-shot EPI pulse sequence (SENSE, acceleration factor = 2); TR = 2000 ms, TE = 30 ms, acquisition time = 336 seconds, FA = 77°, FOV = 240 x 240 x 105, acquisition matrix = 80 x 79 x 35; 35 slices (width = 3 mm, no gap); acquired voxel size (mm) = 3 x 3.04 x 3 (reconstructed voxel size (mm) = 3 x 3 x 3). Fat suppression was implemented with SPIR. The first 5 scans of each functional run were discarded before image acquisition to establish steady-state magnetization.

T1-weighted structural images were obtained with the following parameters: T1-weighted image acquisition using a multi echo, multi-shot turbo field echo pulse sequence, with a five echo average, TR = 12 ms, TE = 3.5 ms, acquisition time = 329 seconds, FA = 8°, FOV (mm) = 240 x 240 x 175, acquisition matrix = 80 x 79; 175 contiguous slices were acquired, voxel size (mm) = 1 x 1 x 2 (reconstructed voxel size = 1mm³).

fMRI design and analysis

All MRI data was pre-processed and analyzed using SPM12 (Wellcome Department of Cognitive Neurology, University College London, <http://www.fil.ion.ucl.ac.uk/spm/>) implemented in MATLAB R2015b 8.6 (Mathworks Inc., Sherborn, MA, USA). Anatomical images were manually aligned to the anterior and posterior commissure (AC-PC). The functional images were corrected for head motion (spatial realignment; trilinear interpolation) and realigned to the first (sample 1) or last (sample 2) functional volume of the session (the one closest to the anatomical scan). The functional scans were then coregistered to their anatomical scans and then normalized to standard MNI space (3mm isotropic voxels). Normalized data were then spatially smoothed using a Gaussian kernel of 6 mm full-width at half-maximum.

The general linear model was used to map the hemodynamic response curve onto each experimental condition using boxcar regressors. This boxcar function was then fitted to the time series at each voxel resulting in a weighted beta-image. The fitted model was

converted to a t-statistic image, comprising the statistical parametric map. The LI-toolbox plugin for SPM (Wilke & Lidzba, 2007; Wilke & Schmithorst, 2006) was used to assess hemispheric contribution for processing each of the three tasks. This toolbox allows for comparison of right and left hemispheres without commonly cited problems such as complications that arise from statistical outliers, threshold-dependent comparisons, or data sparsity (Wilke & Lidzba, 2007).

The toolbox employs a bootstrapping method whereby 20 equally-sized thresholds are calculated between 0 and the maximum t-value in the dataset. At each threshold, 100 bootstrapped samples (with a resampling ratio of $k = 0.25$) are taken at each threshold in each hemisphere. The 10,000 LI combinations are calculated from these samples for all surviving voxels on the left and right, with the standard LI formula ($LI = (R-L)/(R+L)$), where a resulting positive score indicates more left hemisphere activity, and negative indicates more right. To avoid effects due to statistical outliers only the central 50% of data are kept. A final LI is calculated from all the LIs weighted to their corresponding threshold and provides an estimate of how lateralized a participant is for a given contrast.

Whole-brain LIs were calculated from the following contrasts: faces > scenes (sample 1), neutral faces > flowers and butterflies (sample 2), and bodies > chairs (Experiments 1 and 2). The cerebellum was excluded from the contrast fluency > letter string (sample 1 and 2), as it is activated in some language tasks, including verbal fluency (e.g., Häberling, Steinemann, & Corballis, 2016; Schlösser et al., 1998). Cerebellar activation in language processing is contralateral to that of the cerebral cortex (e.g., Gelinias et al., 2014; Jansen et al., 2005).

Participant classification and statistical analysis

Participants were first divided into three groups based on LI values for verbal fluency and based on hand preference. Zero was used as a cut-off for all tasks, thus each participant with an LI value of > 0 were categorized as left hemisphere lateralized, and participants with

RECIPROCAL OR INDEPENDENT HEMISPHERIC SPECIALIZATIONS

15

LIs of < 0 as right hemisphere lateralized. Language typical right-handers and language typical left-handers were separated into groups to examine for handedness differences independent of language dominance. The third group were the language atypically lateralized individuals. As language atypical right-handers are rare, this group included both right-handed and left-handed individuals.

Mean LI values were first compared against 0, using a one-sample t-test, to examine if the group, on average, was significantly lateralized for the specified contrast. Differences in mean LI values for each of the non-language functions were assessed using Kruskal-Wallis tests comparing the three language/handedness groups. If there is no complementarity, then differences defined by language processing and handedness should have no bearing on how asymmetrical anyone is for any of these functions.

To examine if there was a statistical majority of individuals with dominance in one hemisphere over the other (i.e., the breadth of asymmetry), the proportion of individuals with right and left hemisphere processing was compared against 50% using a binomial test. Furthermore, z-tests were used to examine proportional differences in 'typical' processing between the three language/handedness groups for all tasks. In other words, proportions of right hemisphere processing in the two language typical groups were compared with proportions of left hemisphere processing in the language atypical group. The z-test between the language typically lateralized groups and the language atypically lateralized group were two-tailed, as the proportions were assumed to be similar if complementarity of functions exists. The examination of proportional differences between language typical right-handers and language typical left-handers was two-tailed for the same reason. There was no reason to suspect that one group would be more lateralized than the other in terms of proportions. Lastly, we examine the different patterns of lateralization that are found across the two samples as a whole, and in how many individuals they are found.

Results

Table 1

Sample sizes, mean LI values, standard deviations, and t-values from one sample t-tests for each function (fluency, faces, bodies) and group in sample 1 and 2.

		Fluency LI			Face LI			Body LI		
		Mean	SD	t	Mean	SD	t	Mean	SD	t
Sample 1	n									
Typical RH	31	+0.67	0.18	21.10***	-0.42	0.37	-6.19***	-0.52	0.23	-12.46***
Typical LH	43	+0.58	0.22	17.28***	-0.13	0.39	-2.25*	-0.20	0.36	-3.65**
Atypicals	17	-0.63	0.24	-10.56***	+0.09	0.54	0.72	-0.00	0.47	-0.04
Sample 2	n									
Typical RH	18	+0.68	0.16	18.31***	-0.28	0.35	-3.39*	-0.39	0.31	-5.45***
Typical LH	24	+0.58	0.20	14.41***	-0.31	0.37	-4.08***	-0.11	0.34	-1.64
Atypicals	10	-0.50	0.23	-6.88***	+0.23	0.31	2.34*	+0.05	0.34	0.47

Note. RH = right-handers, LH = left-handers. * $p < .05$, ** $p < .01$, *** $p < .001$. Degrees of freedom = $n - 1$.

Sample 1


All tasks were found to be significantly lateralized in the sample as a whole; verbal fluency to the left hemisphere ($M = +.39$, $SD = .53$), $t(90) = 6.91$, $p < .001$, and face perception ($M = -.19$, $SD = .45$), $t(90) = -3.95$, $p < .001$) and body perception ($M = -.27$, $SD = .39$), $t(90) = -6.57$, $p < .001$) to the right hemisphere.

Participants were grouped according to handedness and language dominance. Three groups were created: language typical (left hemisphere lateralized) right-handers ($n = 31$), language typical left-handers ($n = 43$) and language atypical (right hemisphere lateralized) individuals ($n = 17$, 2 right-handers and 15 left-handers). Average LI values are illustrated in Figure 1 and Table 1. All three groups were, on average, significantly lateralized for verbal

fluency as assessed with a one sample t-test against 0 (see Table 1). There was no difference in the depth of fluency LIs between the groups, compared using absolute values ($p = .193$).

Figure 1

Pirate plots showing distributions of individual LI values for each language-defined group in sample 1 (left panel; typical right-handers in blue, $n = 31$; typical left-handers in red, $n = 43$; atypicals in green, $n = 17$) for each of the three fMRI measures (verbal fluency, faces and bodies) and sample 2 (right panel; typical right-handers in blue, $n = 18$; typical left-handers in red, $n = 24$; atypicals in green, $n = 10$) for each of the three fMRI measures (verbal fluency, faces, and bodies) in sample 2.



Note. The bold line indicates the mean and the lighter highlighted area the 95% CIs. All participants in the groups fall above and below the zero line in the verbal fluency graphs as this is how the groups were defined.

Face perception

A Kruskal-Wallis test revealed a significant difference in LI values between the groups, $H(2) = 14.84, p = .001$. Pairwise comparisons revealed statistically significant between-group differences in the median face LI scores for typical right-handers ($Mdn = -.57$) and typical left-handers ($Mdn = -.11; p = .009, r = .34$), typical right-handers and atypicals ($Mdn = .40; p = .001, r = .51$), but not for typical left-handers and atypicals ($p = .581$).

The proportion of individuals with typical hemispheric processing in each group can be seen in Table 2 and were compared against .50 using binomial tests. The proportion of right hemisphere processing in language typical right-handers (.77, 95% CI's .60, .89) was significantly higher, $p = .003$. The proportion of right hemisphere processing in language typical left-handers (.65, 95% CI's .50, .78; $p = .066$) or language atypicals (.59, 95% CI's .36, .78; $p = .629$) was not significantly higher. A z-test compared if there were differences in the number of participants with right hemisphere dominance in language typical right-handers (24/31) and language typical left-handers (28/43), and was not significant, $z = 1.14, p = .254$. To examine if there was a difference in the complementary patterns found for faces in the three groups, z-tests between proportion of right hemisphere processing in the typical groups and left hemisphere processing in the atypical group was compared. The proportion of complementarity in language typical right-handers was not significantly higher than the language atypicals (10/17), $z = 1.36, p = .174$, neither was the difference in language typical left-handers as compared to language atypicals $z = 0.46, p = .646$.

Body perception

LI values for each group can be seen in Figure 1 and Table 1. A Kruskal-Wallis test revealed a significant difference in LI values between the groups, $H(2) = 20.99, p < .001$. Pairwise comparisons revealed statistically significant between-group differences in the median face LI scores for typical right-handers ($Mdn = -.56$) and typical left-handers ($Mdn = -$

RECIPROCAL OR INDEPENDENT HEMISPHERIC SPECIALIZATIONS

19

.19), $p = .001$, $r = .43$, typical right-handers and atypicals ($Mdn = -.14$) was significant, $p < .001$, $r = .59$, but not typical left-handers and atypicals ($p = .612$).

The proportion of individuals with *typical* hemispheric processing in each group can be seen in Table 2 and was compared against .50. The proportion of right hemisphere processing in language typical right-handers (.97, 95% CI's .84, .99) was significantly higher, $p < .001$. The proportion of right hemisphere processing in language typical left-handers (.70, 95% CI's .55, .81) was significantly higher, $p = .014$. The proportion of left hemisphere processing in language atypicals of .47, 95% CI's [.26, .69] was not significantly different, $p = 1$. There was a significant difference in the proportion of individuals who processed bodies in the right hemisphere between language typical right-handers (30/31) and language typical left-handers (30/43), $z = 2.93$, $p = .003$, 95% CI of the difference (-0.27) did not overlap with zero [-0.42, -0.10]. There was a significant difference in complementarity of function between language typical right-handers and atypicals (8/17), $z = 4.06$, $p < .001$, 95% CI of the difference (-0.50) did not overlap with zero [-0.71, -0.24]. There was no significant difference between language typical left-handers and atypicals, $z = 1.64$, $p = .101$.

Table 2

Complementarity patterns for face- and body processing with language found in sample 1 and sample 2 and significance values for binomial tests against 50%.

	Face dominance			Body dominance		
	Right	Left	<i>p</i>	Right	Left	<i>p</i>
Sample 1						
Typical RH	24 (77%)	7 (23%)	.003	30 (97%)	1 (3%)	<.001
Typical LH	28 (65%)	15 (35%)	.066	30 (70%)	13 (30%)	.014
Atypicals	10 (59%)	7 (41%)	.629	8 (47%)	9 (53%)	1
Sample 2						
Typical RH	14 (78%)	4 (22%)	.031	16 (89%)	2 (11%)	.001
Typical LH	19 (79%)	5 (21%)	.007	17 (71%)	7 (29%)	.064
Atypical	3 (30%)	7 (70%)	.344	4 (40%)	6 (60%)	.754

Note. RH = right-handers, LH = left-handers.

Sample 2

Again, all tasks were significantly lateralized in the sample as a whole; verbal fluency to the left hemisphere ($M = +.41$, $SD = .49$, $t(51) = 6.04$, $p < .001$, and face perception ($M = -.19$, $SD = .40$, $t(51) = -3.47$, $p = .001$) and body perception ($M = -.18$, $SD = .36$, $t(51) = -3.55$, $p = .001$) to the right hemisphere. The sample consisted of 18 language typical right-handers, 24 language typical left-handers, and 10 language atypicals (9 left-handed and 1 right-handed). Average LI values, standard deviations, and one-sample t-tests against 0 can be seen in Table 1. There was a difference in median LI values between the group, $H(2) = 6.63$, $p = .036$, and this was found between the language typical right-handers ($Mdn = 0.73$) and atypicals ($Mdn = 0.56$, $p = .041$, $r = .47$), but not language typical right-handers and language and typical left-handers ($Mdn = 0.64$; $p = .230$) language typical left-handers and atypicals ($p = .788$).

Face perception

A Kruskal-Wallis test revealed a significant difference in LI values between the groups, $H(2) = 13.21$, $p = .001$. Pairwise comparisons adjusted for multiple comparisons revealed statistically significant between-group differences in the median face LI scores for typical right-handers ($Mdn = -.37$) and atypicals ($Mdn = .24$; $p = .006$, $r = .59$) typical left-handers ($Mdn = -.33$) and atypicals ($p = .001$, $r = .60$), but not typical right-handers and typical left-handers ($p = 1$).

The proportion of individuals with complementary hemispheric processing in each group can be seen in Table 2 and was compared against .50 using binomial tests. The proportion of right hemisphere processing in language typical right-handers (.78, 95% CI's .55, .91) was significantly higher, $p = .031$. The proportion of right hemisphere processing in language typical left-handers (.79, 95% CI's .60, .91) was significantly higher, $p = .007$. The proportion of left hemisphere processing in language atypicals of .7, 95% CI's [.40, .90] was not significantly different, $p = .344$.

Z-tests were used to compare if there were differences in complementarity processing in the different groups. There was no significant difference between language typical right-handers (14/18), and language typical left-handers (19/24, $p = .913$). The proportion of language typical right-handed participants with a right hemisphere dominance was not significantly different to the left hemisphere processing in the language atypicals (7/10), $p = .648$. The proportion of right hemisphere processing in language typical left-handers was also not significantly different compared to language atypicals ($p = .564$).

Body perception

LI values for each group can be seen in Figure 1, right panel and Table 1. A Kruskal-Wallis test revealed a significant difference in LI values between the groups, $H(2) = 10.47$, $p = .005$. Pairwise comparisons adjusted for multiple comparisons revealed statistically

significant between-group differences in the median face LI scores for typical right-handers ($Mdn = -.42$) and typical left-handers ($Mdn = -.13$; $p = .048$, $r = .37$) typical right-handers and atypicals ($Mdn = +.11$; $p = .008$, $r = .57$), but not for typical left-handers and atypicals ($p = .734$).

The proportion of individuals with typical hemispheric processing in each group can be seen in Table 2 and was compared against .50 using binomial tests. The proportion of right hemisphere processing in language typical right-handers (.89, 95% CI's .67, .97) was significantly higher, $p = .001$. The proportion of right hemisphere processing in language typical left-handers (.71, 95% CI's .51, .85) was not significantly higher, $p = .064$. The proportion of left hemisphere processing in language atypicals (.60, 95% CI's .31, .83) was not significantly different, $p = .754$. Z-tests were used to compare the number of participants with complementary patterns in the different groups. There was no significant difference in the proportion of individuals who processed bodies in the right hemisphere between language typical right-handers (16/18) and language typical left-handers (17/24), $p = .157$. There was no significant difference between language typical right-handers and atypicals (6/10 = .60), $p = .074$, or between language typical left-handers and atypicals, $p = .540$.

Hemispheric patterns of lateralization in individuals

The different combinations of hemispheric patterns found for fluency, faces, and bodies, were investigated for all individuals in sample 1 and sample 2 combined. The different patterns found in the two datasets can be seen in Table 3. The 'traditional' pattern of hemispheric dominance (with language in the left hemisphere and the other asymmetries in the right hemisphere) is also the most common pattern, with 74 of participants (64% of all language typical participants) exhibiting this pattern. The second most common pattern, found in 19 individuals is with language and faces processed in the left hemisphere, and bodies in the right hemisphere. Amongst language atypical individuals, the reversed 'anti-localizing' asymmetry was also seen most frequently (in 12 individuals, 44 % of all language atypicals).

RECIPROCAL OR INDEPENDENT HEMISPHERIC SPECIALIZATIONS

23

A z-test comparing the proportion of language typical and language atypical participants with these “traditional”, assumed, patterns (i.e., .64 versus .44) found no significant difference ($p = .056$). In total, all possible 8 different combinations of asymmetry patterns were seen in this dataset.

Table 3

Patterns of lateralization found for individuals in sample 1 and 2 combined.

Verbal fluency	Faces	Bodies	Percentage (n)
L	R	R	52 % (74)
L	R	L	8 % (11)
L	L	R	13 % (19)
L	L	L	8 % (12)
R	L	L	8 % (12)
R	L	R	4 % (5)
R	R	L	1 % (2)
R	R	R	6 % (8)

Note. L = left hemisphere; R = right hemisphere

Discussion

The aim of these two experiments was to examine the lateralization patterns of language, body perception and face perception in both left-handers and right-handers. To this aim we investigated lateralization of body- and face perception in three groups: language typical right-handed; language typical left-handed; and language atypical individuals. Of specific interest was the breadth (i.e., the proportion) of right hemisphere asymmetry in the language dominance groups, subdivided by handedness for language typicals. We specifically aimed to recruit individuals likely to have language dominance in the right hemisphere to examine rates of complementarity in right- and left-handers with typical and atypical language dominance. This inclusion meant that we focused on recruiting large numbers of strongly left-handed individuals, as they are more likely to exhibit atypical language dominance (Knecht et al., 2000).

Table 4

A summary of significant depth and breadth of LI values for the three different language and handedness defined groups (language typical right-handers, language typical left-handers, language atypical participants) in sample 1 and sample 2.

Face processing	Sample	Language typical RH	Language typical LH	Atypicals
Population-level breadth of bias?	1	✓	X	X
	2	✓	✓	X
Group-level depth of bias?	1	✓	✓	X
	2	✓	✓	✓
Body processing	Sample	Language typical RH	Language typical LH	Atypicals
Population-level breadth of bias?	1	✓	✓	X
	2	✓	X	X
Group-level depth of bias?	1	✓	✓	X
	2	✓	X	X

A summary of the depth and breadth of biases for each of the three groups in both samples can be seen in Table 4. For face processing, typical right-handers had population-level breadth to the right hemisphere in both samples. Typical left-handers and atypicals did not have population-level breadth of face processing in sample 1, and this was also the case for atypicals in sample 2. Language typical right-handers and left-handers were, on average, right lateralized in both samples. The language atypical group was, on average, not lateralized in sample 1 but left-lateralized in sample 2.

Although the atypical group in sample 1 was, on average, not lateralized for face processing, Figure 1 shows that there seems to be two clusters of participants, one left-lateralized and one right-lateralized, resulting in a no-asymmetry average. This idea suggests that face processing is still lateralized to the right or the left hemisphere in the majority of individuals. The LIs in the subgroups are similarly sized, even in individuals lateralized to the same hemisphere that is dominant for language. Altogether, these data suggest that face processing does not always lateralize to the non-language dominant hemisphere.

It can be deduced from the supplementary data in Badzakova-Trajkov et al. (2010) that a reduced number of language typical left-handers (77%) as compared to language typical right-handers (96%) were right hemisphere dominant for faces in their study. In their small sample of atypicals, only 44% (4 out of 9) showed a complementary relationship, which is similar to that seen in sample 1. Gerrits et al. (2019) also found that approximately 70% of their typical left-handers processed faces in the right hemisphere, but similarly reduced to 40% the language atypical sample. In contrast, Gerrits et al. (2020) found that their left-handed atypical sample showed higher rates of complementarity as compared to the typical left-handers (75% complementarity in atypicals versus 56% in typicals). Similarly, the atypical group in the current sample 2 showed high rates of complementarity (70%), although this was not as high as the rates seen for language typical left-handers (79%). The inconsistency seen for left-handers in the current samples, and across these studies, may suggest that they, regardless

of language dominance group, are more variable for other lateralized processes such as face perception.

One important detail to note about the data in the present study is that all three groups were found to be matched in terms of mean lateralization for verbal fluency in sample 1 (although there was a difference between typical right-handers and atypicals in sample 2). There was no difference in the depth of LI values for language typical left-handers and language typical right-handers. Language atypical participants did not show reduced depth of asymmetry when compared to language typical participants. That is, their laterality indices did not tend towards bilaterality as compared with the language typical group, and the depth of the asymmetry was at least equivalent to that of language typical left-handers. This 'symmetry' for language processing means that any differences between the groups on other measures cannot easily be explained by different degrees of language specialization.

These results do not fully support current crowding hypotheses that specifically theorize that face processing and language processing are linked (Behrmann & Plaut, 2015; Dehaene et al., 2010). Of course, it should be noted that these results may also depend on the tasks used. The current incarnations of the crowding hypotheses between faces and language are about reading and asymmetry in the vWFA. One caveat to the present analysis is that we used a verbal fluency task to estimate language dominance (as have several other laboratories, e.g., Badzakova-Trajkov et al., 2010; Gerrits et al., 2020). We describe the reliability and validity of this measure elsewhere (Johnstone et al., 2020). Arguably it might be more astringent to measure asymmetries in the so-called visual word form area (vWFA) for letters and words, for direct comparison with faces. Attempts of this sort have been made by our colleagues in Ghent. Van der Haegen et al. (2012) find a surprisingly poor match in concurrence of hemispheric dominance within the same hemisphere between lexical decision LIs from the temporal lobes and verbal fluency LIs from the frontal lobes (63% atypicals and 81% typicals, hemispheric dominance as defined by the verbal fluency task, all left-handed). Gerrits et al. (2019) found an even lower congruency of verbal fluency and vWFA; only in 58%

of their language typicals and 67% of language atypical left-handers (language typicality defined by verbal fluency). Surprisingly, these colleagues do not compare congruency of face asymmetry and vWFA asymmetry in their typical and atypical groups.

Language typical right-handers were, on average, right lateralized for body perception. Typical left-handers were right lateralized in sample 1 but not sample 2. Atypicals were not lateralized, on average. A large proportion (97 and 89 percent respectively) of right-handers had more activation in the non-language hemisphere for body perception, and this was statistically higher than 50 percent. Language atypicals (53% and 60%) did not have a significant breadth for body processing, but language typical left-handers had a population-level bias in sample 1 (70%) but not in sample 2 (71%). This pattern suggests the intriguing possibility of handedness moderating the effect of body asymmetry. This possibility is further substantiated by the fact that typical right-handers had consistently stronger right hemisphere LI values as compared to typical left-handers and atypicals.

This weakened asymmetry in body processing in left-handers is consistent with the results from Willems et al. (2010), who found that such participants were bilateral on average when measuring activation in the extrastriate body area (EBA), in comparison to right-handers who were right lateralized, as in the current study. Willems et al. (2010) did, however, also find that fusiform face area (FFA) was not lateralized, on average, in their sample of left-handed participants. Although FFA was not specifically targeted in the current study, left-handers were, on average, right-lateralized for the whole-brain activation pattern. Bukowski et al. (2013) also found that the FFA was activated bilaterally for their left-handed sample, but that the activation was overall right hemispheric when comparing the whole brain.

Of course, one limitation is that the sample sizes here are small for proportional analyses and is particularly true considering the skewed nature of the data. The small sample size is reflected in the large confidence intervals that accompanies each proportional difference. Nonetheless, by accumulating data from more participants, this proportional approach is an extremely useful analysis for determining constructs/processes that really

differentiate right-handers and left-handers (Carey & Johnstone, 2014; Karlsson et al., 2020).

When right- and left-handed participants are compared, reduced asymmetries in the left-handers are usually found (e.g., Karlsson et al., 2019; Willems et al., 2010). Taking breadth into account will help to disentangle the underlying cause of a mean difference (i.e., is it driven by a small subgroup of the left handers who show the atypical cerebral asymmetry or are they less lateralized as compared to their right-handed counterparts? Karlsson et al., 2019).

In this study a cut-off of 0 is used, rather than a band around zero labelled by many investigators as “bilateral”. The boundaries for classification of no dominance are difficult to justify and to date are not well agreed upon in the literature. In fact, test-retest with fMRI LI data is rare. Therefore, it is difficult to know how stable these LI values are from session to session. Jansen et al. (2006) found that if a bilateral category was used ($\pm .2$), participants who were classified as bilateral by one calculation were often not by a different calculation of LIs. Test-retest for participants with LIs in this range were not reproducible (in terms of dominant hemisphere) in a second session. It would perhaps be more appropriate to classify participants in more data-driven ways. The boundaries for categorical misclassification rates could be defined by using data to decide an uncertainty of lateralization interval (Carey & Karlsson, 2019). In other words, identify how often individuals with single session LI values of, for example, ± 0.1 would be misclassified in a second session or run.

Overall, these results add to the growing literature that does not fully support complementarity of functions when both language typical and language atypical participants are included. The strongest support for complementarity of face- and body processing with language processing in the current dataset came from the group of language typically lateralized right-handers. Interestingly, the highest rates of complementarity in language typical right-handers alone was seen for body perception, a lateralized function rarely discussed in the cerebral asymmetries literature. Rates of right hemisphere processing in typical left-handers were significantly reduced for bodies specifically. Again, these results

suggest that left-handers are more variable in their lateralization patterns in a way that is independent from language dominance.

These kinds of data might be important for evaluating models of individuals with fluctuating asymmetry. Ideas have been proposed in the handedness literature suggest that fluctuating asymmetry is much more common in left-handed individuals (e.g., McManus 1999; Annett, 2002). The extension of these accounts to underlying cerebral asymmetries could be fruitful. By cataloguing multiple brain asymmetries in large numbers of individuals, the most common phenotype(s) of cerebral patterns might be identifiable. According to this kind of model, some subset of individuals have handedness and different cerebral asymmetries randomly determined (subject to crowding constraints at some of the more extreme ends of the random distribution).

In conclusion, our results and other recent reports in the literature (Badzakova-Trajov et al., 2010; Gerrits et al., 2019), suggest that ideas of complementarity of function need revisiting in the unfortunate labor-intensive manner that these studies require. The effects of handedness versus lateralization typicality can be teased apart from one another, but such studies need to go even further than just enriching samples with left-handers. They need to include left-handers and right-handers with typical and atypical dominance for the functions in question. Finding atypical participants, in particular, is a challenge that to date is mainly met with sheer sample size. Our lab, and others (Gerrits et al., 2020), are working on ways to simplify the search for these elusive, but nevertheless, fascinating individuals.

Funding

This work was supported by the Leverhulme Trust [grant number RPG-2019-102].

References

- Annett, M. (2002). *Handedness and brain asymmetry: the Right Shift Theory*. Hove, UK: Psychology Press
- Badzakova-Trajkov, G., Corballis, M. C., & Häberling, I. S. (2016). Complementarity or independence of hemispheric specializations? A brief review. *Neuropsychologia*, *93B*, 386-393. <https://doi.org/10.1016/j.neuropsychologia.2015.12.018>
- Badzakova-Trajkov, G., Häberling, I. S., Roberts, R. P., & Corballis, M. C. (2010). Cerebral asymmetries: Complementary and independent processes. *PloS one*, *5*(3), e9682. <https://doi.org/10.1371/journal.pone.0009682>
- Behrmann, M., & Plaut, D. C. (2015). A vision of graded hemispheric specialization. *Annals of the New York Academy of Sciences*, *1359*(1), 30-46. <https://doi.org/10.1111/nyas.12833>
- Benton, A. L. (1972). The 'minor' hemisphere. *Journal of the History of Medicine and Allied Sciences*, *27*(1), 5-14. <https://doi.org/10.1093/jhmas/XXVII.1.5>
- Blankenburg, F., Ruff, C. C., Bestmann, S., Bjoertomt, O., Josephs, O., Deichmann, R., & Driver, J. (2010). Studying the role of human parietal cortex in visuospatial attention with concurrent TMS–fMRI. *Cerebral Cortex*, *20*(11), 2702-2711. <https://doi.org/10.1093/cercor/bhq015>
- Bradshaw, J. L., & Nettleton, N. C. (1981). The nature of hemispheric specialization in man. *Behavioral and Brain Sciences*, *4*(1), 51-63. <https://doi.org/10.1017/S0140525X00007548>
- Bryden, M. P. (1990). Choosing sides: The left and right of the normal brain. *Canadian Psychology*, *31*, 297-309. <https://doi.org/10.1037/h0078949>
- Bryden, M. P., & Allard, F. A. (1981). Shortcomings of the verbal/nonverbal dichotomy: Seems to us we've heard this song before.... *Behavioral and Brain Sciences*, *4*(1), 65-66. <https://doi.org/10.1017/S0140525X00007585>

- Bryden, M. P., Hécaen, H., & DeAgostini, M. (1983). Patterns of cerebral organization. *Brain and Language*, 20(2), 249-262 [https://doi.org/10.1016/0093-934X\(83\)90044-5](https://doi.org/10.1016/0093-934X(83)90044-5)
- Bukowski, H., Dricot, L., Hanseeuw, B., & Rossion, B. (2013). Cerebral lateralization of face-sensitive areas in left-handers: only the FFA does not get it right. *Cortex*, 49(9), 2583-2589. <https://doi.org/10.1016/j.cortex.2013.05.002>
- Buxbaum, L. J. (2006). On the right (and left) track: Twenty years of progress in studying hemispatial neglect. *Cognitive Neuropsychology*, 23(1), 184-201. <https://doi.org/10.1080/02643290500202698>
- Cai, Q., Van der Haegen, L., & Brysbaert, M. (2013). Complementary hemispheric specialization for language production and visuospatial attention. *Proceedings of the National Academy of Sciences*, 110(4), E322-E330. <https://doi.org/10.1073/pnas.1212956110>
- Carey, D. P., & Johnstone, L. T. (2014). Quantifying cerebral asymmetries for language in dextrals and adextrals with random-effects meta analysis. *Frontiers in Psychology*, 5, 1128. <https://doi.org/10.3389/fpsyg.2014.01128>
- Carey, D.P., & Karlsson, E.M. (2019). A bright future for the study of multiple cerebral asymmetries?: Comment on “Phenotypes in hemispheric functional segregation? Perspectives and challenges” by Guy Vingerhoets. *Physics of life reviews*, 30, 19-21. <https://doi.org/10.1016/j.plrev.2019.08.009>
- Centanni, T. M., Norton, E. S., Park, A., Beach, S. D., Halverson, K., Ozernov-Palchik, O., ... & Gabrieli, J. D. (2018). Early development of letter specialization in left fusiform is associated with better word reading and smaller fusiform face area. *Developmental Science*, e12658. <https://doi.org/10.1111/desc.12658>
- Dehaene, S., Pegado, F., Braga, L. W., Ventura, P., Nunes Filho, G., Jobert, A., ... & Cohen, L. (2010). How learning to read changes the cortical networks for vision and language. *Science*, 330(6009), 1359-1364. <https://doi.org/10.1126/science.1194140>

- Della Sala, S., & Young, A. W. (2003). Quaglino's 1867 case of prosopagnosia. *Cortex*, 39(3), 533-540. [https://doi.org/10.1016/S0010-9452\(08\)70263-6](https://doi.org/10.1016/S0010-9452(08)70263-6)
- Downing, P. E., Jiang, Y., Shuman, M., & Kanwisher, N. (2001). A cortical area selective for visual processing of the human body. *Science*, 293(5539), 2470-2473. <https://doi.org/10.1126/science.1063414>
- Eling, P. (1984). Broca on the relation between handedness and cerebral speech dominance. *Brain and Language*, 22(1), 158–159. [https://doi.org/10.1016/0093-934X\(84\)90085-3](https://doi.org/10.1016/0093-934X(84)90085-3)
- Flöel, A., Buyx, A., Breitenstein, C., Lohmann, H., & Knecht, S. (2005). Hemispheric lateralization of spatial attention in right-and left-hemispheric language dominance. *Behavioural Brain Research*, 158(2), 269-275. <https://doi.org/10.1016/j.bbr.2004.09.016>
- Friederici, A. D. (2017). *Language in our brain: The origins of a uniquely human capacity*. Cambridge, MA: MIT Press.
- Gelinas, J. N., Fitzpatrick, K. P., Kim, H. C., & Bjornson, B. H. (2014). Cerebellar language mapping and cerebral language dominance in pediatric epilepsy surgery patients. *NeuroImage: Clinical*, 6, 296-306. <https://doi.org/10.1016/j.nicl.2014.06.016>
- Gerrits, R., De Clercq, P., Verhelst, H., & Vingerhoets, G. (2020). Evaluating the performance of the visual half field paradigm as a screening tool to detect right hemispheric language dominance. *Laterality*, 25(6), 722-739. <https://doi.org/10.1080/1357650X.2020.1854279>
- Gerrits, R., Van der Haegen, L., Brysbaert, M., & Vingerhoets, G. (2019). Laterality for recognizing written words and faces in the fusiform gyrus covaries with language dominance. *Cortex*, 117, 196-204. <https://doi.org/10.1016/j.cortex.2019.03.010>
- Gerrits, R., Verhelst, H., & Vingerhoets, G. (2020). Mirrored brain organization: Statistical anomaly or reversal of hemispheric functional segregation bias?. *Proceedings of the*

National Academy of Sciences, 117(25), 14057-14065.

<https://doi.org/10.1073/pnas.2002981117>

Geschwind, N. (1975). The apraxias: Neural mechanisms of disorders of learned movement:

The anatomical organization of the language areas and motor systems of the human brain clarifies apraxic disorders and throws new light on cerebral dominance. *American Scientist*, 63(2), 188-195.

<https://www.jstor.org/stable/27845363>

Goodglass, H., & Quadfasel, F. A. (1954). Language laterality in left-handed

aphasics. *Brain*, 77(4), 521-548. <https://doi.org/10.1093/brain/77.4.521>

Häberling, I. S., Steinemann, A., & Corballis, M. C. (2016). Cerebral asymmetry for

language: comparing production with comprehension. *Neuropsychologia*, 80, 17-23.

<https://doi.org/10.1016/j.neuropsychologia.2015.11.002>

Hillis, A. E. (2007). Aphasia: progress in the last quarter of a century. *Neurology*, 69(2), 200-

213. <https://doi.org/10.1212/01.wnl.0000265600.69385.6f>

Holmes, G. (1918). Disturbances of visual orientation. *The British journal of*

ophthalmology, 2(9), 449. <https://doi.org/10.1136/bjo.2.9.449>

Holowka, S., & Petitto, L. A. (2002). Left hemisphere cerebral specialization for babies while

babbling. *Science*, 297(5586), 1515-1515. <https://doi.org/10.1126/science.1074941>

Jansen, A., Flöel, A., Van Randenborgh, J., Konrad, C., Rotte, M., Förster, A. F., ... &

Knecht, S. (2005). Crossed cerebro–cerebellar language dominance. *Human Brain Mapping*, 24(3), 165-172. <https://doi.org/10.1002/hbm.20077>

Jansen, A., Menke, R., Sommer, J., Förster, A. F., Bruchmann, S., Hempleman, J., ... &

Knecht, S. (2006). The assessment of hemispheric lateralization in functional MRI—robustness and reproducibility. *NeuroImage*, 33(1), 204-217.

<https://doi.org/10.1016/j.neuroimage.2006.06.019>

Johnstone, L. T., Karlsson, E. M., & Carey, D. P. (2020). The validity and reliability of

quantifying hemispheric specialisation using fMRI: Evidence from left and right

- handers on three different cerebral asymmetries. *Neuropsychologia*, 138, 107331.
<https://doi.org/10.1016/j.neuropsychologia.2020.107331>
- Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The fusiform face area: a module in human extrastriate cortex specialized for face perception. *Journal of Neuroscience*, 17(11), 4302-4311. <https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997>
- Karlsson, E. M., Johnstone, L. T., & Carey, D. P. (2019). The depth and breadth of multiple perceptual asymmetries in right handers and non-right handers. *Laterality: Asymmetries of Body, Brain and Cognition*, 24(6), 707-739.
<https://doi.org/10.1080/1357650X.2019.1652308>
- Kimura, D., & Archibald, Y. (1974). Motor functions of the left hemisphere. *Brain*, 97(2), 337-350. <https://doi.org/10.1093/brain/97.2.337>
- Knecht, S., Dräger, B., Deppe, M., Bobe, L., Lohmann, H., Flöel, A., ... & Henningsen, H. (2000). Handedness and hemispheric language dominance in healthy humans. *Brain*, 123(12), 2512-2518. <https://doi.org/10.1093/brain/123.12.2512>
- Liu, H., Stufflebeam, S. M., Sepulcre, J., Hedden, T., & Buckner, R. L. (2009). Evidence from intrinsic activity that asymmetry of the human brain is controlled by multiple factors. *Proceedings of the National Academy of Sciences*, 106(48), 20499-20503.
<https://doi.org/10.1073/pnas.0908073106>
- Lochy, A., de Heering, A., & Rossion, B. (2019). The non-linear development of the right hemispheric specialization for human face perception. *Neuropsychologia*, 126, 10-19.
<https://doi.org/10.1016/j.neuropsychologia.2017.06.029>
- Lundqvist, D., Flykt, A., & Öhman, A. (1998). *The Karolinska Directed Emotional Faces – KDEF*, CD ROM from Department of Clinical Neuroscience, Psychology section, Karolinska Institutet, ISBN 91-630-7164-9.
- Mazoyer, B., Mellet, E., Perchey, G., Zago, L., Crivello, F., Jobard, G., ... & Joliot, M. (2016). BIL&GIN: a neuroimaging, cognitive, behavioral, and genetic database for the study

of human brain lateralization. *NeuroImage*, 124, 1225-1231.

<https://doi.org/10.1016/j.neuroimage.2015.02.071>

McManus, I. C. (1999). Handedness, cerebral lateralization, and the evolution of language.

In M. C. Corballis & S. E. G. Lea (Eds.), *The descent of mind: Psychological perspectives on hominid evolution* (pp. 194-217). New York, NY, US: Oxford University Press.

Ng, V. W., Eslinger, P. J., Williams, S. C., Brammer, M. J., Bullmore, E. T., Andrew, C. M., ...

& Benton, A. L. (2000). Hemispheric preference in visuospatial processing: a complementary approach with fMRI and lesion studies. *Human Brain Mapping*, 10(2), 80-86. [https://doi.org/10.1002/\(SICI\)1097-0193\(200006\)10:2<80::AID-HBM40>3.0.CO;2-2](https://doi.org/10.1002/(SICI)1097-0193(200006)10:2<80::AID-HBM40>3.0.CO;2-2)

Paterson, A., & Zangwill, O. L. (1944). Disorders of visual space perception associated with

lesions of the right cerebral hemisphere. *Brain: A Journal of Neurology*, 67, 331–358. <https://doi.org/10.1093/brain/67.4.331>

Peelen, M. V., Glaser, B., Vuilleumier, P., & Eliez, S. (2009). Differential development of selectivity for faces and bodies in the fusiform gyrus. *Developmental Science*, 12(6),

F16-F25. <https://doi.org/10.1111/j.1467-7687.2009.00916.x>

Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years

after. *Annual review of Neuroscience*, 35, 73-89. <https://doi.org/10.1146/annurev-neuro-062111-150525>

Plaut, D. C., & Behrmann, M. (2011). Complementary neural representations for faces and

words: A computational exploration. *Cognitive Neuropsychology*, 28(3-4), 251-275. <https://doi.org/10.1080/02643294.2011.609812>

Powell, J. L., Kemp, G. J., & García-Finaña, M. (2012). Association between language and

spatial laterality and cognitive ability: an fMRI study. *NeuroImage*, 59(2), 1818-1829. <https://doi.org/10.1016/j.neuroimage.2011.08.040>

- Rasmussen, T., & Milner, B. (1977). The role of early left-brain injury in determining lateralization of cerebral speech functions. *Annals of the New York Academy of Sciences*, 299(1), 355-369. <https://doi.org/10.1111/j.1749-6632.1977.tb41921.x>
- Riddoch, G. (1935). Visual disorientation in homonymous half-fields. *Brain*, 58(3), 376-382. <https://doi.org/10.1093/brain/58.3.376>
- Rosch, R. E., Bishop, D. V., & Badcock, N. A. (2012). Lateralised visual attention is unrelated to language lateralisation, and not influenced by task difficulty—a functional transcranial Doppler study. *Neuropsychologia*, 50(5), 810-815. <https://doi.org/10.1016/j.neuropsychologia.2012.01.015>
- Schlösser, R., Hutchinson, M., Joseffer, S., Rusinek, H., Saarimaki, A., Stevenson, J., ... & Brodie, J. D. (1998). Functional magnetic resonance imaging of human brain activity in a verbal fluency task. *Journal of Neurology, Neurosurgery & Psychiatry*, 64(4), 492-498. <http://dx.doi.org/10.1136/jnnp.64.4.492>
- Schwarzlose, R. F., Baker, C. I., & Kanwisher, N. (2005). Separate face and body selectivity on the fusiform gyrus. *Journal of Neuroscience*, 25(47), 11055-11059. <https://doi.org/10.1523/JNEUROSCI.2621-05.2005>
- Sergent, J., Ohta, S., & Macdonald, B. (1992). Functional neuroanatomy of face and object processing: a positron emission tomography study. *Brain*, 115(1), 15-36. <https://doi.org/10.1093/brain/115.1.15>
- Steenhuis, R. E., & Bryden, M. P. (1989). Different dimensions of hand preference that relate to skilled and unskilled activities. *Cortex*, 25(2), 289-304. [https://doi.org/10.1016/S0010-9452\(89\)80044-9](https://doi.org/10.1016/S0010-9452(89)80044-9)
- Taylor, J. C., Wiggett, A. J., & Downing, P. E. (2007). Functional MRI analysis of body and body part representations in the extrastriate and fusiform body areas. *Journal of Neurophysiology*, 98(3), 1626-1633. <https://doi.org/10.1152/jn.00012.2007>

- Van der Haegen, L., Cai, Q., & Brysbaert, M. (2012). Colateralization of Broca's area and the visual word form area in left-handers: fMRI evidence. *Brain and Language*, *122*(3), 171-178. <https://doi.org/10.1016/j.bandl.2011.11.004>
- Van der Haegen, L., Cai, Q., Seurinck, R., & Brysbaert, M. (2011). Further fMRI validation of the visual half field technique as an indicator of language laterality: A large-group analysis. *Neuropsychologia*, *49*(10), 2879-2888. <https://doi.org/10.1016/j.neuropsychologia.2011.06.014>
- Whitehouse, A. J., & Bishop, D. V. (2009). Hemispheric division of function is the result of independent probabilistic biases. *Neuropsychologia*, *47*(8), 1938-1943. <https://doi.org/10.1016/j.neuropsychologia.2009.03.005>
- Wilke, M., & Lidzba, K. (2007). LI-tool: a new toolbox to assess lateralization in functional MR-data. *Journal of Neuroscience Methods*, *163*(1), 128-136. <https://doi.org/10.1016/j.jneumeth.2007.01.026>
- Wilke, M., & Schmithorst, V. J. (2006). A combined bootstrap/histogram analysis approach for computing a lateralization index from neuroimaging data. *NeuroImage*, *33*(2), 522-530. <https://doi.org/10.1016/j.neuroimage.2006.07.010>
- Willems, R. M., Peelen, M. V., & Hagoort, P. (2009). Cerebral lateralization of face-selective and body-selective visual areas depends on handedness. *Cerebral Cortex*, *20*(7), 1719-1725. <https://doi.org/10.1093/cercor/bhp234>
- Woodhead, Z. V. J., Bradshaw, A. R., Wilson, A. C., Thompson, P. A., & Bishop, D. V. (2019). Testing the unitary theory of language lateralization using functional transcranial Doppler sonography in adults. *Royal Society open science*, *6*(3), 181801. <https://doi.org/10.1098/rsos.181801>
- Woodhead, Z.V.J., Thompson, P.A., Karlsson, E.M., & Bishop, D.V.M. (2021). An updated investigation of the multidimensional structure of language lateralization in left- and right-handed adults: a test–retest functional transcranial Doppler sonography study

with six language tasks. *Royal Society Open Science*, 8(2), 200696.

<https://doi.org/10.1098/rsos.200696>

Zangwill, O. L. (1967). Speech and the minor hemisphere. *Acta Neurologica et Psychiatrica*

Belgica, 67(11), 1013–1020. <https://doi.org/10.1212/01.wnl.0000265600.69385.6f>