

Kent Academic Repository

Full text document (pdf)

Citation for published version

Bird, Emma E., Kivell, Tracy L. and Skinner, Matthew M. (2021) Internal bone architecture in the triquetrum of extant hominids and *Homo neanderthalensis*. In: European Society for the Study of Human Evolution.

DOI

Link to record in KAR

<https://kar.kent.ac.uk/92553/>

Document Version

Presentation

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

BONE ARCHITECTURE IN THE TRIQUETRUM OF EXTANT HOMINIDS AND *HOMO NEANDERTHALENSIS*

Emma E. Bird¹ (eeb8@kent.ac.uk), Tracy L. Kivell^{1,2}, Antonio Rosas³, Matthew M. Skinner^{1,2}

¹School of Anthropology and Conservation, University of Kent; ²Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology; ³Paleoanthropology Group, Museo Nacional de Ciencias Naturales - Consejo Superior de Investigaciones Científicas (CSIC), Madrid, Spain

Introduction

Soft and hard tissue structures at the ulna-side of the wrist, such as reduced ulna-triquetrum articulation and the Triangular Fibrocartilage Complex (TFCC), are conspicuous synapomorphies of hominoids [Fig.1]. These structures are historically linked to biomechanics characteristic of the hominoids such as high degrees of ulnar deviation and forearm supination [1,2]. Nevertheless, there is notable variation in the bony and soft tissue anatomy of the wrist among hominoids that affects how loads are transferred from the hand to the forearm. However, the functional consequences of these differences are not fully understood, in part because the functional morphology of the triquetrum is understudied. Although both committed tool users, it is unknown whether force is transferred through the triquetrum in a similar way between humans and Neanderthals, particularly as the Neanderthal triquetrum is closer to the morphology of the hypothesized human-Neanderthal last common ancestor [3]. Here we investigate for the first time the internal trabecular bone structure of the triquetrum and test the null hypothesis that great apes will exhibit similar distributions of relative trabecular bone volume to total volume (BV/TV) and degree of anisotropy (DA).





Fig. 1. Schematic of the radiocarpal joint adaptations in humans (left) relative to a baboon (right). The triquetrum is in pink and labelled with a t. p=pisiform; l=lunate; s=scaphoid. Adapted from [4]

Fig. 2. Morphology of the triquetrum (in pink) across the four study taxa. *Homo* is represented by *Homo sapiens*. Image is of the left hand, view from the palm.

Methods

- Internal bone structure related to strength (BV/TV and DA) were analysed in the triquetrum to test whether bone architecture correlates with soft and hard tissue structures in the ulna wrist
- The sample consisted of *Pongo* (n=12), *Gorilla* (n=10), *Pan* (n=10), Neanderthals (n=3; El Sidron SD-1227, Tabun C1, and Amud 1), and modern humans (n=21).
- Triquetra were μ CT scanned (30-50 microns) (Fig.3A) then segmented (Fig.3B,C). The triquetra were then analysed holistically in 3D using medtool (Fig 3D) [5].
- BV/TV and DA values were visualised across the entire bone using Paraview (3.89.0).
- Low DA was identified as values equal to or less than 20%. High BV/TV was identified as the top 20% of values for that individual's range.




Fig. 3. Steps of trabecular and cortical bone analysis using medtool. Dorsal is up, Proximal is left.

Predictions and Results

Joint	Predictions	Results	<i>Pongo</i>	<i>Gorilla</i>	<i>Pan</i>	Neanderthals	humans
Triquetrum-TFCC	Low DA		41% (5/12)	100% (10/10)	90% (9/10)	100% (3/3)	76% (16/21)
Triquetrum-Hamate	High BV/TV		75% (9/12)	90% (9/10)	70% (7/10)	100% (3/3)	61% (13/21)
Triquetrum-Lunate	High BV/TV		91% (11/12)	20% (2/10)	0% (0/10)	33% (1/3)	0.04% (1/21)

Table 1. Predictions are on the left of the table, results are on the right. The results follow the prediction in the same row. Eg: 41% of *Pongo* had low DA at the Triquetrum-TFCC




Fig. 4. Bone distribution of one representative individual from each extant group, and each Neanderthal (inside square brackets). Top row shows the top 20% of BV/TV in red. Dorsal is up, left is the lunate articulation and right is the hamate articulation. Bottom row shows DA \leq 20% in blue. View is of the proximal surface, dorsal is up.

Conclusions

Previous studies of trabecular architecture in great apes have often shown high degrees of variation both within and between species [6,7]. The results of this study exhibit relatively consistent inter- and intraspecific patterns of bone architecture.

In *Gorilla*, *Pan*, humans and Neanderthals, there was consistently greater BV/TV at the triquetro-hamate joint relative to the triquetro-lunate. This may indicate greater force transfer with the hamate relative to the lunate.

The architecture in Neanderthals appeared to have a similar distribution to humans, suggesting similar force transfer.

Pongo has distinct soft and hard tissue morphology at the ulna side of the wrist [1]. In this study they show a different pattern of bone architecture to the other great apes. *Pongo* exhibits greater bone volume at the triquetro-lunate joint which may reflect a reliance on the lunate for force transfer to the ulna. *Pongo* did not consistently exhibit low DA at the proximo-radial triquetrum, possibly due to a reduced capacity to transfer force due to the under-developed TFCC.

In all taxa except *Pongo*, the TFCC insertion site showed low DA. If force transfer with the synapomorphic TFCC results in characteristic isotropic trabeculae patterns, this would be a powerful framework in which to analyse fossil taxa whose radiocarpal joint configuration is unknown. Further research including non-hominoid primates would benefit this hypothesis.

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 819960), the Spanish government MICINN (PID2020-115062GB-I00), and the 50th Anniversary Research Scholarship, University of Kent. For providing access to the specimens we would like to acknowledge the following institutions: Bavarian State Collection of Zoology; Berlin Museum für Naturkunde; The Duckworth Collection, University of Cambridge; Frankfurt Senckenberg Museum; Georg-August-University Goettingen, Anthropology Collection; Max Planck Institute for Evolutionary Anthropology; Museo Nacional de Ciencias Naturales, Madrid Spain; Powell-Cotton Museum; Royal Museum for Central Africa; the Sackler School of Medicine at Tel Aviv University; University of Florence; Vienna Natural History Museum. For microCT scanning, we thank R. Freney, P. Schoenfeld, A. Silvester, K. Smithson, N. Stephens, H. Temming, Z. Tsegai, and A. Winzer.

References: [1] Lewis OJ (1989) *Functional morphology of the evolving hand and foot* [2] Sarmiento EE (1988) *Int. J. Primatol.* [3] Kivell TL (2011) *S. Afr. J. Sci.* [4] Kivell TL (2016) In Kivell TL, Lemelin P, Richmond BG, Schmitt D (Eds) *Evolution of the primate hand* [5] Pahr, D. <http://www.dr-pahr.at/medtool/> [6] Bird EE et al. (2021) *J. Anat.* [7] Bird EE et al. (2019) *ASHB*.