



## Pharmacological Activities of Three Kinds "Kayu kuning": *Arcangelisia flava*, *Fibraurea tinctoria* and *Coscinium fenestratum* – an Short Review

Riski Sulistiarini<sup>1,\*</sup>, Andreanus A Soemardji<sup>1</sup>, Elfahmi<sup>2</sup>, Maria Immaculata Iwo<sup>1</sup>

<sup>1</sup>Departement of Pharmacology - Clinical Pharmacy, School of Pharmacy, Bandung Institute of Technology, Jl. Ganesha 10 Bandung, 40132, Indonesia

<sup>2</sup>Department of Pharmaceutical Biology, School of Pharmacy, Bandung Institute of Technology, Jl. Ganesha 10 Bandung, 40132, Indonesia

\*E-mail: [riskisulistiarini@students.itb.ac.id](mailto:riskisulistiarini@students.itb.ac.id) and [kikyidris@gmail.com](mailto:kikyidris@gmail.com)

### Abstract

The literature-based review was constructed discussing three types of yellow woods plant from Indonesia, including *Arcangelisia flava*, *Fibraurea tinctoria*, and *Coscinium fenestratum*. Yellow wood plants are widely used as traditional medicine due to its activities that were pharmacologically studied. Those activities include antiplasmodial, cytotoxic, antioxidant, toxicity, antidiabetic, anticolostroleia, antihypertensive to liver activity, and health-behavior changes in experimental animals.

**Keywords:** review, pharmacologically studies, *Arcangelisia flava*, *Fibraurea tinctoria*, *Coscinium fenestratum*

**Submitted:** 25 April 2020

**Accepted:** 08 June 2020

**DOI:** <https://doi.org/10.25026/jtpc.v5i2.258>

### ■ Introduction

Indonesia widely uses plants as a source of natural-based medicines [1], one of which is a Kayu kuning or yellow wood plant or yellow root, namely *Arcangelisia flava*, *Fibraurea tinctoria*, and *Coscinium fenestratum* [2]. All three of these plants exhibit antimalarial effects with excellent IC<sub>50</sub> in inhibiting the growth of *P. falciparum* [3]. Researchers also found that the effects did not worked synergically toward the liver due to

inhibition of the cytochrome P3A4 by chloroform extract and chloroform insoluble fraction from *F. tinctoria* plants with IC<sub>50</sub> 3.4 µg/mL [4] and 5.1 µg/mL [5]. In contrast, the methanol extract of *C. fenestratum* (60 mg/kgBB) showed a hepatoprotective effect for 90 days on the induction of carbon tetrachloride [6] With a variety of the impact that corresponds, and some other effects are also the opposite of these three types of plants, the activities that were tested pharmacologically will be further discussed.

## ■ Discussion

### The taxonomy and morphology of Kayu kuning.

#### *Arcangelisia flava*

Kingdom	:	<i>Plantae</i>
Order	:	<i>Tracheophyta</i>
Subclass	:	<i>Magnoliopsida</i>
Suborder	:	<i>Ranunculales</i>
Family	:	<i>Menispermaceae</i>
Genus	:	<i>Arcangelisia</i>
Species	:	<i>Arcangelisia flava</i> (L.) Merr.

This plant exhibits a bright yellow wood color, and the leaves grow upwards with an almost round leaf shape [7]. These wild plants can be found on rocky beaches or on the edge of forests [8] (Figure 1).

#### *Fibraurea tinctoria*

Kingdom	:	<i>Plantae</i>
Order	:	<i>Tracheophyta</i>
Subclass	:	<i>Magnoliopsida</i>
Suborder	:	<i>Ranunculales</i>
Family	:	<i>Menispermaceae</i>
Genus	:	<i>fibraurea</i> Lour
Species	:	<i>Fibraurea tinctoria</i> Lour

This is a woody plant, with trees that can grow up to 40 m in diameter. This plant exhibits yellow stems and twigs. The oval-shaped leaves resemble betel leaves but do not exhibit a distinctive aroma [9] (Figure 2).

#### *Coscinium fenestratum*

Kingdom	:	<i>Plantae</i>
Order	:	<i>Tracheophyta</i>
Subclass	:	<i>Magnoliopsida</i>
Suborder	:	<i>Ranunculales</i>
Family	:	<i>Menispermaceae</i>
Genus	:	<i>Coscinium Colebra</i>
Species	:	<i>Coscinium fenesstratum</i> (Gaertn.)

This plant can grow very tall, with a stem diameter of up to 10 cm, in addition to being yellow on the inside and brown on the outside [10]. The shape of the leaves in this plant is like an elongated betel leaf (Figure 3).

### The traditional utilization of Kayu kuning

The recorded traditional utilization of kayu kuning plants is recorded below (Table 1).



Figure 1. *Arcangelisia flava* (Source : BKSDA Samboja, 2016)



Figure 2. *Fibraurea tinctoria* (Source : BKSDA Samboja, 2016)



Figure 3. *Coscinium fenestratum* (Source : BKSDA Samboja, 2016)

Table 1. The traditional utilization of Kayu kuning

Used	Species	Part(s)	Source
Fever, post-delivery recovery, hepatitis, digestive issue, and malaria	<i>A. flava</i>	shoot and root	[2]
Diarrhea, nyctalopia, and hepatitis	<i>A. flava</i>	(not defined)	[11]
Diabetes mellitus, kidney deseas, and eczema	<i>A. flava</i>	(not defined)	[12]
Sexual contagious disease	<i>F. tinctoria</i>	Root	[13]
Food poisoning and paralyze	<i>F. tinctoria</i>	Stem	[13]
Snakes' venom antidote	<i>F. tinctoria</i>	Root, stem	[13]
Eye diseases, diarrhea, dysentery, intestines inflammations, vaginomycosis, furunculosis, and burnt wound	<i>F. tinctoria</i>	All parts of plant	[13]
Malaria	<i>F. tinctoria</i>	(not defined)	[13]
Polyps and nose inflammation, headache	<i>F. tinctoria</i>	Stem	[13]
Insect and snakes' venom antidote	<i>F. tinctoria</i>	Bark	[13]
Eye disease, reumatism, hipertension, sinus, fever, cancer, headache, and snake venom antidote	<i>F. tinctoria</i>	Root, Bark	[13]
Food poisoning	<i>F. tinctoria</i>	Root, Stem	[13]
Diarrhea and hepatitis	<i>F. tinctoria</i>	Root, Stem	[14]
Hipocalemic, hipotensi, laxative, antidiabetic, digestive problems, hepatitis, fever, snakes' venom antidote, antiseptic, and inflammation.	<i>C. fenestratum</i>	Stem	[15], [16]

## Pharmacological activities of Kayu kuning

Kayu kuning or yellow wood plants demonstrate efficacy that has been tested pharmacologically. Some of the results of the study are as follows:

### Antimicrobial

Ethanol extract of *A. flava* root can inhibit *Bacillus cereus* ATCC 14579 and *Staphylococcus aureus* ATCC 25923, indicated by the clear zone formation of 9.6 mm at a dose of 10 mg/mL and 17.2 mm at a dose of 5 mg/mL, respectively [17]

Chloroform, n-hexane, and ethyl acetate extracts of *A. flava* can inhibit *Aeromonas hydrophilia* with clear zones of 17.25, 13.18, and 11.16 mm at a dose of 20 mg/mL, respectively [18]. The ethanol extract of *A. flava* vines of the branches resulted inactivity to *S. aureus* with a MIC value of 0.25 µg/mL [19]. The 96% methanol extract of the leaves provides antibacterial activity against *Pseudomonas fluorescens* ATCC 49642 with MIC values 14.17 µg/mL [20].

Dichloromethane extract: methanol (1: 1) leaves and stems of *F. tinctoria* can inhibit *Bacillus cereus* with 10 mm inhibition zone diameters at 400 µg/disc, MIC 25 µg/mL, and MBC > 100 µg/mL. This extract also demonstrates the ability to inhibit the growth of *S. aureus* with inhibition zone diameters of 10 mm at a dose of 400 µg/disc, MIC 50 µg/mL, and MBC > 100 µg/mL [21].

The methanol extract of *C. fenestratum* stems can inhibit *Nisseria gonorrhoeae* ATCC 49226 with MIC values of 47.39 µg/mL [22].

### Antifungal

The water extract of *A. flava* plant stem can inhibit the growth of *Candida albicans* with a MIC value of 10 mg/mL and a MFC of 4%. This extract also showed inhibitory effects on *Trichophyton mentographytes* with a MIC value of 10 mg/mL and MFC of 5% [8].

### Antiplasmodial

Plant extracts of *A. flava* and *F. tinctoria* demonstrate the ability to inhibit the growth of Plasmodium, which is shown in Table 2 [23].

### Cytotoxic

*A. flava* and *F. tinctoria* also contain cytotoxic properties, in addition to antiplasmodial properties (Table 3).

### Antiploriferation

The methanol extract and water fraction of *F. tinctoria* demonstrate the ability to inhibit the proliferation of Human Colon Cancer HT-29 cells with IC50 values of 17.12 and 9.29 µg/mL, respectively. The methanol extract and plant water fraction also demonstrate the ability to inhibit the proliferation of human skin fibroblasts with IC50 values of both > 100 µg/mL [25].

Table 2. The activity of antiplasmodial properties *A. flava* and *F. tinctoria*

Species	Used part	Extract	Activity	
			IC <sub>50</sub> (µg/mL)	% inhibition at 10 mg/mL
<i>A. flava</i>	Stem	Ethanol 80%	0.7	94 %
<i>A. flava</i>	Stem	Methylene chloride	0.4	98%
<i>A. flava</i>	Stem	Methanol fraction	0.9	98%
<i>F. tinctoria</i>	Root	Methanol fraction	1.0	94%
<i>F. tinctoria</i>	Root	Methylene chloride	0.7	96%
<i>F. tinctoria</i>	Root	Methanol	1.1	97%
<i>F. tinctoria</i>	Stem	Methanol	0.7	94%
<i>F. tinctoria</i>	Stem	Methylene chloride	0.5	97%
<i>F. tinctoria</i>	Stem	Methanol fraction	1.1	97%

Table 3. The activity of cytotoxic of *A. flava* and *F. tinctoria*

Species	Used part	Extract	Experimental cell	Dosage (µg/mL)	Source
<i>A. flava</i>	Stem	Ethanol 80%	HeLa cell	IC <sub>50</sub> 21.3	[3]
<i>A. flava</i>	Stem	Methylene chloride	HeLa cell	IC <sub>50</sub> 8.8	[3]
<i>A. flava</i>	Stem	Methanol	HeLa cell	IC <sub>50</sub> 40.7	[3]
<i>A. flava</i>	Stem	Ethanol 80%	<i>MRC5</i>	IC <sub>50</sub> 155.0	[3]
<i>A. flava</i>	Stem	Methylene chloride	<i>MRC6</i>	IC <sub>50</sub> 56.5	[3]
<i>A. flava</i>	Stem	Methanol	<i>MRC7</i>	IC <sub>50</sub> 147.7	[3]
<i>A. flava</i>	Stem	Ethanol	HeLa cell	IC <sub>50</sub> 467	[24]
<i>A. flava</i>	Leaf	Ethanol	HeLa cell	IC <sub>50</sub> 136	[24]
<i>A. flava</i>	Leaf	Ethanol	<i>WiDr</i> Cell line	IC <sub>50</sub> 213	[24]
<i>F. tinctoria</i>	Stem	Petroleum ether	<i>MRC7</i>	IC <sub>50</sub> >50	[14]
<i>F. tinctoria</i>	Stem	Chloroform	<i>MRC7</i>	IC <sub>50</sub> 11.2	[14]
<i>F. tinctoria</i>	Stem	Methanol	<i>MRC7</i>	IC <sub>50</sub> >50	[14]
<i>F. tinctoria</i>	Stem	Distilled water	<i>MRC7</i>	IC <sub>50</sub> >50	[14]
<i>F. tinctoria</i>	Stem	Ethanol	HeLa cell	IC <sub>50</sub> 70.6	[3]
<i>F. tinctoria</i>	Stem	Methylene chloride	HeLa cell	IC <sub>50</sub> 53.4	[3]
<i>F. tinctoria</i>	Stem	Methanol	HeLa cell	IC <sub>50</sub> 99.9	[3]
<i>F. tinctoria</i>	Stem	Ethanol	<i>MRC5</i>	IC <sub>50</sub> 70.6	[3]
<i>F. tinctoria</i>	Stem	Methylene chloride	<i>MRC5</i>	IC <sub>50</sub> 99.8	[3]
<i>F. tinctoria</i>	Stem	Methanol	<i>MRC5</i>	IC <sub>50</sub> 335.3	[3]

### Antioxidant

Petroleum ether, chloroform, methanol, and water extracts of *F. tinctoria* demonstrate the ability to reduce DPPH radicals with EC50 values > 100, 78.8, 83.6, and > 100 µg/mL [14]

### Anti-hypertension

A 50% ethanol stem extract of *C. fenestratum* shows antihypertensive activity in experimental dogs, which were anesthetized under normal blood pressure conditions. Extracts at doses of 5–40 mg/kg given intravenously reduce blood pressure by 20%–83% with effects lasting for 160 minutes [27].

### Anti-hypercholesterolemia

The methanol extract of *A. flava* stem demonstrates the ability to reduce total cholesterol by 25.49 mg/dL, total triglycerides by 5.5 mg/dL, and LDL by 9.14 mg/dL, in addition to increasing HDL values by 14.8 mg/dL in a dose of 500 mg/kg. This activity was obtained from testing on rat animals induced by high-fat and fructose foods for 45 days [28]

### Anti-diabetes

The ethanol extract of *F. tinctoria* stem demonstrates the ability to reduce plasma glucose concentration in the oral condition of maltose-loaded normal mice at a dose of 250 mg/kg BW [29]

### Analgesic

The methanol extract of *C. fenestratum* stem demonstrates the analgesic effect at a dose of 8 mg/kg BW. This effect is reported from the response of experimental Swiss albino mice to induce the tail flick method (response to snapping the tail) using heat as a pain inducer. The ability to withstand pain was demonstrated by experimental animals for 3.30 minutes after 60 minutes of extract administration [30]

### Hepatoprotector and Hepatotoxic

Chloroform extract and a non-polar portion of *F. tinctoria* chloroform showed inhibition of the cytochrome P3A4 enzyme with IC values of 3.4 µg/mL [5] and 5.1 µg/mL [4]. Conversely, the methanol extract of *C. fenestratum* at a dose of 60 mg/kg BW showed a protective effect on the liver of mice by reducing the free radicals produced by carbon tetrachloride for hepatotoxic conditions [6]

### Conclusion

*Arcangelisia flava*, *Fibraurea tinctoria* and *Cosciniium fenestratum* have various properties for the treatment and maintaining health, that have proven their biological activity in various research methods.

### Acknowledgement

Author thanks to Pharmacy School of Institut Teknologi Bandung, Pharmacy Faculty of Mulawarman University, and the funding from Beasiswa Unggulan Dosen Indonesia-LPDP program by Minister of Financial Indonesia.

### References

- [1] Katiyar C, Arun Gupta, Kanjilal S, et al. Drug Discovery from Plant Sources: An Intergrated Approach. *AYU*; 33.
- [2] Suzuki T, Kiyotani T, Maeda M, et al. Furanoditerpenes from *Arcangelisia flava* (L.) Merr. and their antifungal activity. *Phytochem Lett* 2011; **4**: 333–336.
- [3] Nguyen-Pouplin J, Tran H, Tran H. Antimalarial and Cytotoxic Activities of Ethnopharmacologically Selected Medicinal Plants from South Vietnam. *J Ethnopharmacol* 2007; **109**: 417–427.
- [4] Su CR, Ueng YF, Nguyen XD, et al. Cytochrome P3A4 Inhibitors and Other Constituents of *Fibraurea tinctoria*. *J Nat Prod* 2007; **70**: 1930–1933.
- [5] Su CR, Chen YF, Liou MJ, et al. Anti-inflammatory Activities of Furanoditerpenoids and Other Constituents from *Fibraurea tinctoria*. *Bioorg Med Chem* 2008; **16**: 9603–9609.
- [6] Venukumar MR LMS. Effect of *Cosciniium fenestratum* on hepatotoxicity in rats. *Indian J Exp Biol*; **42**, <https://www.ncbi.nlm.nih.gov/pubmed/15573529> (2004).
- [7] Soemardji AA. 2019. Beberapa Tumbuhan Obat Bermanfaat yang Tumbuh di Jawa. Bandung. ITB Press.
- [8] Setyowati R. The Effect Of Water-Soluble Stem Extract "Kayu Kuning" (*Arcangelisia flava* L.Merr) On The Growth Inhibition Of *Candida albicans* ATCC 10231 And *Trichophyton mentagrophytes* In Vitro. *Biol Med Nat Prod Chem* 2014; **3**: 2089–6514.
- [9] Rasadah Mat Ali, Zainon Abu Samah, Nik Musaadah Mustapha, Norhara Hussein. 2010. ASEAN Herbal and Medicinal Plants. Indonesia. ASEAN Secretariat Public Outreach and Civil Society Division.
- [10] K.V.Tushar, Balachandran I, Gorge S. *Cosciniium fenestratum* (Gaertn.) Colebr. - A Review on this Rare, Critically Endangered and Highly- Traded Medicinal Species. *J Plant Sci* 2008; **3**: 133–145.
- [11] Setyowati R. The Effect Of Water-Soluble Stem Extract " Kayu Kuning " (*Arcangelisia flava* L.Merr) On The Growth Inhibition Of *Candida albicans* ATCC 10231 And *Trichophyton mentagrophytes* In Vitro. *Biol Med Nat Prod Chem* 2014; **3**: 2089–6514.
- [12] Ulfa EU, Ema Rachmawati. Antihypercholesterolemic Effect of *Arcangelisia flava* Stem Extract In Hyperlipidemic Rats. In: *Prociding ICMHS 2016*. 2016, pp. 31–34.
- [13] Galappathie S, Palombo EA, Yeo TC, et al. Comparative Antimicrobial Activity of South East Asian plants used in Bornean folkloric medicine. *J Herb Med* 2014; **4**: 96–105.
- [14] Keawpradub N, Dej-adisai S. Antioxidant and Cytotoxic Activities Of Thai Medicinal Plants named *Khaminkhruea*: *Arcangelisia flava*, *Cosciniium blumeianum* and *Fibraurea tinctoria*. *Songklanakarin J Sci Technol* 2005; **2**: 455–467.
- [15] Wattanathorn J, Uabundit N, Itarat W, et al. Neurotoxicity of *Cosciniium fenestratum* stem, a Medicinal Plant Used in Traditional Medicine. *Food Chem Toxicol* 2006; **44**: 1327–1333.

- [16] Rai R V., Rajesh PS, Kim HM. Medicinal use of *Coscinium fenestratum* (Gaertn.) Colebr.: an short review. *Orient Pharm Exp Med* 2013; **13**: 1–9.
- [17] Soonthornchareonnon N, Wiwat C, Chuakul W. Biological Activities of Medicinal Plants from Mangrove and Beach Forests. *Mahidol Univ J Pharm Sci* 2012; **39**: 9–18.
- [18] Maryani, Marsoedi. The Phytochemistry and The Anti-Bacterial Activity of Yellow Root (*Arcangelisia flava* Merr.) against *Aeromonas hydrophila*. *J Biol Life Sci* 2013; **4**: 180–190.
- [19] Ongsakul M, Jindarat A. Antibacterial Effect of Crude Alcoholic And Aqueous Extracts of Six Medicinal Plants Against *Staphylococcus aureus* and *Escherichia coli*. *J Health Res* 2009; **23**: 153–156.
- [20] Maryani, Rosita, Monalisa SS, et al. In Vitro Test Of Natural Antibacterial Activity ff Yellow-Fruit Moonseed *Arcangelisia flava* Merr. Leaf on bacterium *Pseudomonas fluorescens* under different doses. *AACL Bioflux* 2018; **11**: 288–294.
- [21] Galappathie S, Palombo EA, Yeo TC, et al. Comparative Antimicrobial Activity of South East Asian Plants Used in Bornean Folkloric Medicine. *J Herb Med* 2014; **4**: 96–105.
- [22] Chomnawang MT, Trinapakul C, Gritsanapan W. In vitro antigonococcal activity of *Coscinium fenestratum* stem extract. *J Ethnopharmacol* 2009; **122**: 445–449.
- [23] Nguyen-Pouplin J, Tran H, Tran H. Antimalarial and Cytotoxic Activities of Ethnopharmacologically Selected Medicinal Plants From South Vietnam. *J Ethnopharmacol* 2007; **109**: 417–427.
- [24] Puspitasari E, Pangaribowo DA, Utami Y, et al. *Arcangelisia flava* Leaves Ethanolic Extract Suppresses. In: *Prociding ICMHS 2016*. 2016, pp. 83–86.
- [25] Manosroi A, Akazawa H, Pattamapun K, et al. Potent Anti-proliferative Effects Against Oral and Cervical Cancers of Thai Medicinal Plants Selected from the Thai/Lanna Medicinal Plant Recipe Database 'MANOSROI III'. *Pharm Biol* 2015; **53**: 1075–1081.
- [26] Singh B, Daneshvar C. Human Infections and Detection of Plasmodium knowlesi. *Clin Microbiol Rev* 2013; **26**: 165–184.
- [27] Ulfa EU, Ema Rachmawati. Antihypercholesterolemic Effect Of *Arcangelisia flava* Stem Extract In Hyperlipidemic Rats. In: *Prociding ICMHS 2016*. 2016, pp. 31–34.
- [28] Yibchok-anun S, Jittaprasatsin W, Sontir D, et al. Insulin Secreting And A -Glucosidase Inhibitory Activity ff *Coscinium Fenestratum* and Postprandial Hyperglycemia In Normal And Diabetic Rats. *J Med Plants Res* 2009; **3**: 646–651.
- [29] Kekuda P, Sudharshan, Chinmaya, et al. Central nervous system (CNS) depressant and Analgesic activity of methanolic extracts of *Nardostachys jatamansi* DC. and *Coscinium fenestratum* Colebr. in Experimental Animal Model. *J Pharm Res* 2009; **2**: 1716–1719.