



Article

Methane Mitigation Potential of Foliage of Fodder Trees Mixed at Two Levels with a Tropical Grass

Sara Stephanie Valencia-Salazar ^{1,2,*} , Guillermo Jiménez-Ferrer ¹ , Isabel Cristina Molina-Botero ³, Juan Carlos Ku-Vera ⁴, Ngonidzashe Chirinda ^{2,5} and Jacobo Arango ²

- ¹ The College of the Southern Border (ECOSUR), Livestock and Environment, Carrera Panamericana—Periferico sur, P.C., San Cristobal de las Casas 29290, Mexico; gjimenez@ecosur.mx
- ² International Center for Tropical Agriculture (CIAT), km 17, Recta Cali-Palmira, Palmira, Valle del Cauca, Cali, P.C., Cali 763537, Colombia; j.arango@cgiar.org
- ³ Department of Nutrition, Faculty of Animal Science, Universidad Agraria La Molina, La Molina, P.C., Lima 12456, Peru; hyzavell26@gmail.com
- ⁴ Faculty of Veterinary Medicine and Animal Science, University of Yucatan, Carretera Merida-Xmatkuil km 15.5, Apdo. 4-116 Itzimná, P.C., Merida 97100, Mexico; kvera@correo.uady.mx
- ⁵ AgroBioSciences (AgBS), Agricultural Innovations and Technology Transfer Centre (AITTC), Mohammed VI Polytechnic University, Benguerir, P.C., Ben Guerir 43150, Morocco; nchirinda@gmail.com
- * Correspondence: saraudea@gmail.com; Tel.: +57-323-2312470

Abstract: Enteric methane (CH₄) emitted by ruminant species is known as one of the main greenhouse gases produced by the agricultural sector. The objective of this study was to assess the potential the potential for CH₄ mitigation and additionally the chemical composition, in vitro gas production, dry matter degradation (DMD), digestibility and CO₂ production of five tropical tree species with novel forage potential including: *Spondias mombin*, *Acacia pennatula*, *Parmentiera aculeata*, *Brosimum alicastrum* and *Bursera simaruba* mixed at two levels of inclusion (15 and 30%) with a tropical grass (*Pennisetum purpureum*). The forage samples were incubated for 48 h, and a randomized complete block design was used. Crude protein content was similar across treatments (135 ± 42 g kg⁻¹ DM), while *P. purpureum* was characterized by a high content of acid detergent fiber (335.9 g kg⁻¹ DM) and *B. simaruba* by a high concentration of condensed tannins (20 g kg⁻¹ DM). Likewise, *A. pennatula* and *P. aculeata* were characterized by a high content of cyanogenic glycosides and alkaloids respectively. Treatments SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*) and BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*) resulted in superior degradability at 48h than *P. purpureum*, while in the AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*) was lower than the control treatment ($p \leq 0.05$). At 24 and 48 h, treatments that contained *P. aculeata* and *B. alicastrum* yield higher CH₄ mL g⁻¹ DOM than *P. purpureum* ($p \leq 0.05$). The inclusion of these forage species had no statistical effect on the reduction of CH₄ emissions per unit of DM incubated or degraded at 24 and 48 h with respect to *P. purpureum* although reductions were observed. The use of fodders locally available is an economic and viable strategy for the mitigation of the environmental impact generated from tropical livestock systems.

Keywords: in vitro; greenhouse gases; ruminal degradation; secondary metabolites; tropical livestock systems



Citation: Valencia-Salazar, S.S.; Jiménez-Ferrer, G.; Molina-Botero, I.C.; Ku-Vera, J.C.; Chirinda, N.; Arango, J. Methane Mitigation Potential of Foliage of Fodder Trees Mixed at Two Levels with a Tropical Grass. *Agronomy* **2022**, *12*, 100. <https://doi.org/10.3390/agronomy12010100>

Academic Editors: M. Jordana Rivero and Mary Bridget Lynch

Received: 22 November 2021

Accepted: 29 December 2021

Published: 31 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Methanogenesis in the gastrointestinal tract of ruminant species is the main sink for hydrogen, thus assuring the appropriate fermentation of fiber in the rumen [1]. Due to CH₄ synthesis, ruminants can make use of high-fiber diets (not edible for humans) growing abundantly on enormous land areas, marginal to crop agriculture and convert it into high quality food (e.g., milk, meat) for humans as well as other products derived from livestock [2]. Eructated CH₄ is part of natural cycles and can be transformed by methanotrophic bacteria and reactions with hydroxyl radicals (OH) in the air, to carbon

dioxide (CO₂) that will be used in photosynthesis [3]. However, high emission rates of CH₄ in cattle constitute an energy loss and production inefficiency, especially in tropical production systems where poorly managed grasses are used as feed [4,5]. Additionally, due to methane's global warming potential (GWP) and its contribution to greenhouse gas (GHG) emissions in the agricultural sector, it has gained attention in the last 20 years for research on mitigation alternatives in livestock production [1]. Some of these alternatives are focused on interfering with CH₄ synthesis in the rumen.

In the rumen there is a whole consortium of microorganisms that establish syntrophic relationships between them and of mutualism with their host [6]. The diversity and structure of microbial populations in the rumen will depend on the characteristics of the feed consumed by the ruminant [7]. These microbial populations use the compounds in ruminant diets for their growth and produce volatile fatty acids (VFA), metabolic hydrogen (H₂) and CO₂ as by-products [8]. Rumen bacteria are the most abundant and diverse group of microorganisms in the rumen ecosystem and carry out enzymatic activities for the fermentation of starch, cellulose, hemicellulose, proteins, and lipids [9]. Protozoa comprise a large proportion (approximately 20%) of the microbial mass of the rumen due to their size and can be up to 50% in some cases depending on the diet [9]. They can degrade fiber and have been directly related to the synthesis of CH₄ due to their hydrogenosomes that produce H₂ that serves as a substrate for methanogenic archaea [10]. Archaea are chemolithophilic organisms that use acetate, methyl groups, CO₂ and H₂ as a substrate for the synthesis of CH₄ [6]. Methanogenic archaea of the rumen produce CH₄ mainly through the hydrogenotrophic pathway and to a lesser extent through the methylotrophic and acetoclastic pathways [11]. These biochemical pathways have in common the presence of methyl-coenzyme M reductase (Mcr), responsible for the last step in the reduction of the methyl group to CH₄ [12]. Each of the steps of hydrogenotrophic methanogenesis are shown in Figure 1.




Figure 1. Hydrogenotrophic methanogenesis. Abbreviations: CH₄: methane, CO₂: carbon dioxide, H₂: dihydrogen.

Nutritional strategies for the reduction of enteric CH₄ production in ruminants have been widely studied [13]. Some of these strategies that have shown medium to high CH₄ mitigation potential include concentrates, lipids, algae, nitrates, and chemical inhibitors such as 3-nitrooxypropanol [1]. However, commercial CH₄ mitigation sources in tropical regions can represent a high cost for medium and small producers and be difficult to implement in extensive grazing systems [1,14]. However, biomass diversity in the tropics can offer a vast diversity of low-cost alternatives for CH₄ reduction and the improvement of cattle production to more sustainable systems. Many native tree and shrub species have a high forage potential due to their nutritional quality and secondary metabolite content that have shown to be beneficial to ruminant metabolism [15]. These metabolites from plants have the capacity to modulate the rumen microbiome and reduce CH₄ synthesis [16]. In addition, the use of forestry species due to their role in biogeochemical cycles can benefit the whole production system when used in combination with well managed forage-grass species [17]. Therefore, livestock production systems have a great GHG mitigation potential within the agricultural sector due to its close relation to natural biological cycles and its capacity of transformation to efficient and sustainable systems.

Livestock production in Latin America and the Caribbean is based on extensive systems that for years have promoted the deforestation of large forest areas for the sowing of pastures. Under these conditions, most of the dry matter consumed by cattle comes from medium to low quality pastures [18] and the dry seasons contribute to a scarce supply of forage and a decrease in consumption causing the animals to only cover their maintenance requirements. In this sense, five novel tropical tree species with forage potential evaluated in this study were collected at the Lacandon rainforest, one of the most important ecosystems in Mexico, however one of the main economic activities is extensive cattle production that has generated significant deforestation in the region [19]. By using the Global Forest Watch Pro platform (<https://www.globalforestwatch.org> (accessed on: 15 July 2020)), the loss of tree cover was monitored between 2011 to 2019. A loss of 145,000 ha was recorded, equivalent to a 12% decrease in tree cover since the year 2000, registering the greatest loss in 2019. Many Mayan indigenous and “mestizo” livestock producers use secondary vegetation in critical periods of the year to improve cattle diets due to the low nutritional quality of the pasture in those periods. However, most of the land used for cattle production is directed towards extensive grazing as most of the livestock systems in tropical regions.

Additionally, there is not much knowledge on the biodiversity that exists at the regional level that can be used for cattle production and information on the use of native resources for the design of pastoral systems and management of these species is needed. Introduced species and silvopastoral systems have been promoted and the use of native species has been minimized despite its potential for small and medium producers in tropical countries to improve the cattle’s diets and its capacity to regenerate soils and deforested regions. In this respect, the objective of this study was to quantify in vitro CH₄ mitigation potential, nutritional quality, and digestibility of ten experimental diets composed of *S. mombin*, *A. pennatula*, *P. aculeata*, *B. alicastrum* and *B. simaruba* replacing 15 and 30% (of DM) of grass-based rations designed to be fed to tropical cattle.

2. Materials and Methods

2.1. Description of the Study Area

Forage species were sampled in the Ocosingo Valley of the Lacandon Rainforest, in the East and Northeast of Chiapas, Mexico. The prevailing climate is warm-humid (23–27 °C) with an altitude that varies from 10 to 900 MASL [20]. The municipality of Ocosingo, covers the largest region of the rainforest and one of the major activities is extensive livestock production.

2.2. Sampling

Species were selected from a previous screening of fifteen species [19] for their nutritional quality and their potential to reduce in vitro CH₄ emissions when incubated alone.

The selected species were: *S. mombin*, *A. pennatula*, *P. aculeata*, *B. alicastrum* and *B. simaruba*. Species were harvested with the help of cattle producers from the area. Leaves were collected from 5 to 9 individuals per species. Botanical samples were taken to verify the species in the herbarium of the Southern Border College (ECOSUR). Samples of *P. purpureum* grass at 60 days regrowth were taken to use as a control treatment and basal ration. Samples were dried in a forced air oven at 55 °C or until constant weight to determine dry matter (DM) content [21]. Dried samples were ground in a Wiley Laboratory Mill (Thomas Scientific®, Swedesboro, NJ, USA) to a particle size of 1 mm and stored for transportation and chemical analysis.

2.3. Treatments

Experimental diets were the inclusion of foliage of fodder tree species at two levels (15 and 30%) mixed with *Pennisetum purpureum* sp. grass (85 and 70%, respectively) simulating feeding practices commonly used by some farmers in the region. Treatments were designated as following: SM15-PP85 (15% *S. mombin* + 85% *P. purpureum*), SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*), AP15-PP85 (15% *A. pennatula* + 85% *P. purpureum*), AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*), PA15-PP85 (15% *P. aculeata* + 85% *P. purpureum*), PA30-PP70 (30% *P. aculeata* + 70% *P. purpureum*), BA15-PP85 (15% *B. alicastrum* + 85% *P. purpureum*), BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*), BS15-PP85 (15% *B. simaruba* + 85% *P. purpureum*), BS30-PP70 (30% *B. simaruba* + 70% *P. purpureum*), and control treatment 100PP (100% *P. purpureum*).

2.4. Chemical Analysis

The in vitro gas production technique and chemical analysis were carried out at the Forage Quality and Animal Nutrition Laboratory at the International Center for Tropical Agriculture (CIAT), Palmira (Valle del Cauca, Colombia). This laboratory is certified by the FAO-IAG proficiency test of feed constituents 2017. Forage samples were incinerated in a muffle furnace at 500 °C for 4 h (method 942.05) [22] to determine ash and organic matter (OM = 100 – Ash (%) content). Crude protein was determined by Kjeldahl (AN 3001 FOSS; method 984.14) from the N content (CP = N × 6.25) [23]. Neutral and acid detergent fiber (NDF and ADF, respectively) content were analyzed according to Van Soest [24] and using the Ankom Fiber Analyzer AN 3805 (Ankom® Technology Corp., Macedon, NY, USA) and gross energy (GE) by ISO 9831 [25]. To determine digestibility, the two-stage in vitro technique was used [26]. Condensed tannins (TC) content of the species was determined by the vanillin extract assay [27] at the bromatology laboratory at ECOSUR, Chiapas, Mexico. Alkaloids, cyanogenic glycosides and saponins were qualitatively quantified by the methodologies proposed by Domínguez [28]. The content of ether extract (EE) was determined by the Soxhlet immersion method (NTC 668) [29]. Metabolizable energy (ME) was estimated according to the equation proposed by Menke et al. [30]:

$$\text{ME (MJ kg}^{-1} \text{ DM)} = 2.20 + 0.136 \times \text{GP} + 0.057 \times \text{CP} + 0.0029 \times \text{CP}^2$$

where GP is the net gas production in 24 h (mL/200 mg). Likewise, net energy (NE) was calculated from the equation proposed by Menke and Steingass [31]:

$$\text{NE (Mcal lb}^{-1}) = \frac{2.2 + (0.0272 \times \text{GP}) + (0.057 \times \text{CP}) + (0.149 \times \text{EE})}{14.64}$$

where: GP is the net gas production in 24 h (ml g⁻¹ DM), CP is crude protein (% DM), EE is the ether extract (% DM); results were transformed to MJ/kg⁻¹ DM. Short chain fatty acids (SCFA) were calculated according to Getachew et al. [32] and transformed to mmol per liter.

$$\text{SCFA} = \frac{0.0239 \text{ GP} - 0.0601 \text{ mL}}{200 \text{ mg DM}}$$

2.5. In Vitro Gas Production Technique

Three rumen cannulated *Brahman* bulls of 550 kg live weight fed *Cynodon plectostachyus* and minerals were used. Animals were treated in accordance with the Colombian normative num. 84 from 1989 following the protocol approved by the ethics committee of CIAT. In vitro gas production was quantified using the methodology proposed by Menke and Steingass [31] and modified by Theodorou et al. [33]. Rumen liquid and solid material were obtained from different locations in the rumen at 8 a.m. to assure a representative sample [34]. Rumen liquor was liquefied and filtered through 10 layers of gauze and mixed in a 1:9 ratio with a mineral solution [19,31]. Treatments were incubated in independent bottles of 160 mL by triplicate and kept under constant flow of CO₂. Bottles were placed in a water bath at 39 °C. Gas pressure and volume in the headspace of the bottles were measured with an 840,065 wide-range pressure gauge (Sper Scientific®, Scottsdale, AZ, USA) connected a PS100 2-bar pressure transducer (Lutron Electronic Enterprise Co. Ltd., Taipei, Taiwan) and a three-way valve connected to a hypodermic needle that was inserted into the bottles and a 60 mL syringe was used to measure the gas volume. Gas pressure and volume were measured at 0, 4, 8, 12, 24, 30, 36 and 48 h. The content of the bottles was withdrawn from fermentation at 24 and 48 h for the degradation of DM (DDM) and OM (DOM) as described by Valencia-Salazar et al. [19]. For the degradation and gas production, two sets of triplicate bottles were used (24 and 48 h), these were incubated from 0 to 24 h and from 0 to 48 h, respectively, blanks were also included. The pH was measured using a pH meter (AB15 Plus, Accumet®, Westford, MA, USA) at 24 and 48 h. Bottle content was filtered in crucibles with fiberglass filter and a vacuum pump, then dried in a forced air oven at 65 °C for 48 h and weighed with a precision balance. Data from the pressure and volume of the bottles was used to generate a polynomial equation for the correction of the volume of gas produced as described by Valencia-Salazar et al. [19]:

$$y = 0.0209 x^2 + 5.9023x - 2.984$$

$$R^2 = 0.9729$$

Gas production data was adjusted to the modified Gompertz model [35] to obtain time at the inflection point (TIP, h), gas at the inflection point (GIP, mL), maximum gas production rate (MGPR mL h⁻¹) and Lag phase (LP or the microbial establishment, h).

2.6. Methane Quantification

Gas volume was stored in amber bottles with a capacity of 125 mL from samples collected from the accumulated gas at 24 and 48 h of incubation. Methane and CO₂ concentrations were quantified in the Laboratory of GHG (CIAT) using a GC-2014 gas chromatograph (Shimadzu®, Tokyo, Japan) with 1/8" packed stainless-steel columns (1.0 m HayeSep T 80/100 mesh, 4 m HayeSep D 80/100, 1.5 P-N, 0.7m Shimalite Q 100/180), 80 °C temperature, column flow of 30.83 mL min⁻¹, injection volume handled by a loop with capacity of 2 mL and nitrogen was used as carrier gas.

2.7. Statistical Analysis

For the statistical analysis a randomized block design with 10 treatments (mixed feed substrates) and a control (100% *P. purpureum*) was used with three replicates per hour (24 and 48 h; two sets of triplicates as mentioned in 2.5) and three different inoculums from three different animals as blocking factor. The statistical model used was:

$$Y_{ij} = \mu + Ti + \beta j + \varepsilon_{ij}$$

where Y_{ij} are the observations of the response variables for treatment i and block j ; μ is the overall mean; i is the effect of the i -th treatment; βj is the effect of the j -th block; and ε_{ij} is the random error of treatment i in block j . To test treatment effects, the PROC GLM

procedure of SAS[®] software, version 9.4 was used [36]. The means of the treatments were compared by the Dunnett test with an alpha of 0.05 with respect to the control treatment.

3. Results

3.1. Chemical Composition, In Vitro Digestibility, and Phytochemical Screening

Chemical composition of forage samples and treatments is shown on Table 1. All evaluated species had higher crude protein (CP) contents than the control. The species with the lowest acid detergent fiber (ADF) content were *S. mombin* and *A. pennatula*, with 171.18 and 210.34 g kg⁻¹ DM, respectively. *P. purpureum* grass provides between 1.2 and 2 times more NDF than the other species. The above described is reflected in the mixed rations. Gross energy content of mixed rations ranged from 15.86 to 17.44 MJ kg⁻¹ DM. The highest metabolizable energy and net energy concentrations were obtained in the treatment BA30-PP70 with 7.82 and 4.82 MJ kg⁻¹ DM, respectively. In both inclusion levels of *P. aculeata* presented the highest in vitro digestibility and *A. pennatula* had the lowest digestibility (447.44 g kg⁻¹ DM). Regarding secondary metabolites, the highest content of CT was obtained with *B. simaruba* (20% CT) at inclusion of 15 and 30% in the ration with *P. purpureum*. In the phytochemical screening, presence of alkaloids was found in all species except for *B. simaruba*, and cyanogenic glycosides were found highly abundant only in *A. pennatula* as shown on Table 2. Also, saponin content was found in low abundance only in *A. pennatula*.

Table 1. Chemical composition, condensed tannin content and in vitro digestibility of forage species and treatments.

Species	g kg ⁻¹ DM								MJ kg ⁻¹ DM		
	DM	OM	NDF	ADF	CP	EE	IVDDM	CT	GE	ME	NE
<i>Spondias mombin</i> (SM)	259.75	852.29	307.77	171.18	126.95	44.78	638.59	9.90	16.25		
<i>Acacia pennatula</i> (AP)	505.42	924.85	492.56	210.34	192.69	39.25	447.44	31.1	20.92		
<i>Parmentiera aculeata</i> (PA)	308.95	874.47	614.35	268.81	183.17	13.85	548.37	0.00	18.04		
<i>Brosimum alicastrum</i> (BA)	489.18	821.28	298.17	269.22	116.21	29.92	686.38	0.00	15.65		
<i>Bursera simaruba</i> (BS)	356.71	900.82	354.37	249.23	99.07	25.05	471.37	200.1	18.92		
Treatments											
SM15-PP85	213.82	839.92	542.46	311.22	132.88	23.42	542.46	1.50	15.99	7.40	4.54
SM30-PP70	221.93	842.10	501.04	286.50	131.83	27.19	501.04	3.00	16.04	7.75	4.80
AP15-PP85	250.66	850.80	570.18	317.09	142.74	22.59	570.18	4.70	16.69	7.56	4.58
AP30-PP70	295.61	863.87	556.48	298.25	151.56	25.53	556.48	9.30	17.44	7.46	4.51
PA15-PP85	221.18	843.24	588.45	325.86	141.31	18.78	588.45	0.00	16.26	7.17	4.31
PA30-PP70	236.66	848.75	593.02	315.79	148.70	17.91	593.02	0.00	16.58	7.18	4.27
BA15-PP85	248.23	835.26	541.02	325.92	131.27	21.19	541.02	0.00	15.90	7.60	4.65
BA30-PP70	290.75	832.80	498.17	315.92	128.61	22.73	498.17	0.00	15.86	7.82	4.82
BS15-PP85	228.35	847.20	549.45	322.92	128.70	20.46	549.45	30.0	16.39	6.92	4.23
BS30-PP70	251.00	856.66	515.03	309.92	123.47	21.27	515.03	60.0	16.84	6.05	3.72
100PP	205.70	837.73	583.88	335.93	133.93	19.65	583.88	0.00	15.95	7.08	4.30

DM: Dry matter; OM: Organic matter; NDF: Neutral detergent fiber; ADF: Acid detergent fiber; CP: Crude protein; EE: Ether extract; IVDDM: In vitro digestibility of dry matter; CT: Condensed tannins; GE: Gross energy; ME: Metabolizable energy; NE: net energy; SM15-PP85 (15% *S. mombin* + 85% *P. purpureum*); SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*); AP15-PP85 (15% *A. pennatula* + 75% *P. purpureum*); AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*); PA15-PP85 (15% *P. aculeata* + 85% *P. purpureum*); PA30-PP70 (30% *P. aculeata* + 70% *P. purpureum*); BA15-PP85 (15% *B. alicastrum* + 85% *P. purpureum*); BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*); BS15-PP85 (15% *B. simaruba* + 85% *P. purpureum*); BS30-PP70 (30% *B. simaruba* + 70% *P. purpureum*); 100PP: 100% *P. purpureum*.

Table 2. Secondary metabolite content of tree species from southern Mexico.

Species	Alkaloids			Cyanogenic Glycosides	Saponins
	Mayer	Draggendorff	Wagner		
<i>Spondias mombin</i> (SM)	–	+	++	–	–
<i>Acacia pennatula</i> (AP)	–	++	++	++++	+
<i>Parmentiera aculeata</i> (PA)	++++	++++	++++	–	–
<i>Brosimum alicastrum</i> (BA)	+++	+	–	–	–
<i>Bursera simaruba</i> (BS)	–	–	–	–	–

– (No presence); + (low abundance); ++ (abundant); +++ (moderately abundant); ++++ (highly abundant).

3.2. In Vitro Gas Production Parameters

The Gompertz model parameters for in vitro gas production are shown on Table 3. Maximum gas production (*a*), time at the inflection point (TIP), gas inflection point (GIP), maximum gas production rate and Lag phase (LP) differed significantly ($p < 0.05$) between evaluated treatments and control ration (100PP). The highest maximum gas production (*a*) value was obtained in SM30-PP70; 10% above control ($p < 0.05$). The lowest values for MGPR were obtained when *B. simaruba* was included in the treatment with 13% (at 15%) and 37% (at 30%) below control treatment ($p < 0.05$). This treatment also presented the lowest values for maximum gas production compared to control (10% and 18% below) ($p < 0.05$). *S. mombin* showed a MGRP 13% above control ($p < 0.05$). Figure 2 shows the gas production per hour per gram of organic matter.

Table 3. Gompertz model parameters for in vitro gas production of forage species incorporated at two levels mixed with *P. purpureum*.

Rations	Parameters						
	<i>a</i>	<i>b</i>	<i>c</i>	TIP (h)	GIP (mL)	MGPR (mL h ⁻¹)	LP
100PP	261.304	1.009	0.079	12.786	96.110	7.580	0.120
SM15-PP85	266.533	1.101 *	0.084	13.170	98.033	8.200 *	1.206 *
SM30-PP70	287.369 *	1.048	0.083	12.690	105.697 *	8.730 *	0.583
AP15-PP85	263.756	1.017	0.079	12.866	97.010	7.670	0.216
AP30-PP70	240.909 *	0.953 *	0.078	12.160	88.607 *	6.947 *	–0.593 *
PA15-PP85	220.971 *	1.003	0.086 *	11.616 *	81.270 *	6.613 *	0.030
PA30-PP70	219.015 *	0.925 *	0.081	11.380 *	80.553 *	5.546 *	–0.926 *
BA15-PP85	277.702	0.998	0.081	12.396	102.140	8.220 *	–0.020
BA30-PP70	283.543 *	0.944 *	0.083	11.370 *	104.287 *	8.676 *	–0.676 *
BS15-PP85	235.527 *	0.979	0.076	12.826	86.620 *	6.610 *	–0.286
BS30-PP70	214.480 *	0.917 *	0.060 *	15.173 *	78.887 *	4.766 *	–1.386 *
MSE	8.089	0.0201	0.003	0.456	2.975	0.177	0.267
<i>p</i>-Value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.00001

Asterisks in the same column indicate statistical difference ($p < 0.05$) with respect to the control treatment. *a*: Maximum gas production (mL); *b*: difference between initial gas and final gas at an *x* time; *c*: specific gas accumulation rate; TIP: time at the inflection point (hours); GIP: Gas inflection point (mL); MGPR: Maximum gas production rate (mL h⁻¹); LP: Lag phase; SM15-PP85 (15% *S. mombin* + 85% *P. purpureum*); SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*); AP15-PP85 (15% *A. pennatula* + 75% *P. purpureum*); AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*); PA15-PP85 (15% *P. aculeata* + 85% *P. purpureum*); PA30-PP70 (30% *P. aculeata* + 70% *P. purpureum*); BA15-PP85 (15% *B. alicastrum* + 85% *P. purpureum*); BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*); BS15-PP85 (15% *B. simaruba* + 85% *P. purpureum*); BS30-PP70 (30% *B. simaruba* + 70% *P. purpureum*); 100PP: 100% *P. purpureum*.




Figure 2. Gas production in mL g⁻¹ organic matter per hour. Abbreviations: SM15-PP85 (15% *S. mombin* + 85% *P. purpureum*); SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*); AP15-PP85 (15% *A. pennatula* + 75% *P. purpureum*); AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*); PA15-PP85 (15% *P. aculeata* + 85% *P. purpureum*); PA30-PP70 (30% *P. aculeata* + 70% *P. purpureum*); BA15-PP85 (15% *B. alicastrum* + 85% *P. purpureum*); BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*); BS15-PP85 (15% *B. simaruba* + 85% *P. purpureum*); BS30-PP70 (30% *B. simaruba* + 70% *P. purpureum*); 100PP: 100% *P. purpureum*.

3.3. Dry Matter Degradability, pH, Short Chain Fatty Acids and CH₄ Production

The highest degradability at 48 h was observed when *S. mombin* and *B. alicastrum* species were included in the two levels. The lowest degradabilities at 48 h were obtained in PA30-PP70, AP30-PP70 and PA15-PP85 with 8.77, 7.91 and 7.34% below the control (PP100) ($p < 0.05$). The pH was not affected by the inclusion of the forage tree species at two levels and the concentrations of SCFA were 20% and 23% above control ($p < 0.05$) for treatments SM30-PP70 and BA30-PP70, respectively. Methane was expressed in mg g⁻¹ of incubated and DOM as suggested by Yáñez-Ruiz et al. [34] and is presented in Table 4. Methane production in mg g⁻¹ DOM and mg g⁻¹ IOM (incubated OM) was different between the treatments and *P. purpureum* ($p < 0.0001$). It was observed that PA30-PP70 had the major CH₄ mg g⁻¹ IOM produced at 24 h and at 48 h, 86% and 33% above control ($p < 0.05$). Methane emissions at 48 were also higher and significantly different ($p < 0.05$) in SM30-PP70, BA15-PP85 and BA30-PP70 when compared to control (100PP), the other treatments produced similar amounts of CH₄. On the other hand, BS30-PP70 had the lowest CH₄ mg g⁻¹ IOM production; 21% below control at 48 h, however, no statistical difference was observed.

Table 4. DM degradability, pH, SCFA and CH₄ production at 24 and 48 h of incubation of forage species mixed with *P. purpureum* at two levels.

	DM DEG %		pH		SCFA mMol l ⁻¹	CO ₂ mL g ⁻¹ IOM		CO ₂ mL g ⁻¹ DOM		CH ₄ mL g ⁻¹ IOM		CH ₄ mL g ⁻¹ DOM	
	24	48	24	48		24	48	24	48	24	48	24	48
100PP	47.01	55.84	6.68	6.69	77.45	168.07	155.89	102.03	112.27	14.08	26.74	8.55	19.27
SM15-PP85	47.13	60.69 *	6.78 *	6.79	84.86 *	176.37	202.58	153.78	158.15	13.58	26.20	8.22	20.47
SM30-PP70	56.65 *	63.21 *	6.67	6.63	92.80 *	221.38	268.74 *	185.08 *	219.06 *	16.75	37.51 *	12.44	30.97 *
AP15-PP85	46.29	55.02	6.72	6.68	85.29 *	148.72	166.34	87.30	116.35	12.56	25.81	7.38	18.04
AP30-PP70	46.04	51.43 *	6.69	6.74	80.50	156.73	225.55	109.89	139.02	12.32	29.47	9.74	18.91
PA15-PP85	43.77	51.73 *	6.72	6.83	77.25	142.47	244.72 *	79.88	149.69	11.37	29.87	6.39	19.86
PA30-PP70	41.92*	50.95 *	6.70	6.94	75.13	321.66 *	267.98 *	171.45	174.07 *	26.16 *	35.68 *	13.95 *	23.17 *
BA15-PP85	51.37	60.11 *	6.68	6.76	89.62 *	259.01	272.44 *	172.51	211.89 *	23.61 *	37.76 *	14.14 *	29.39 *
BA30-PP70	55.98 *	62.24 *	6.68	6.87	95.42 *	269.39	285.77 *	195.74 *	230.51 *	22.95 *	38.06 *	16.68 *	30.69 *
BS15-PP85	45.01	54.52	6.70	6.77	75.49	305.59 *	265.79 *	175.28 *	184.96 *	20.68	31.81	12.95	22.16
BS30-PP70	39.17 *	54.47	6.70	6.72	57.88 *	240.56 *	208.22	118.20	143.09	16.35	21.02	6.65	14.43
MSE	2.06	1.48	0.02	0.11	1.67	38.30	30.51	29.03	24.74	2.30	2.98	1.94	2.11
<i>p</i> -value	<0.0001	<0.0001	0.0028	0.1467	<0.0001	<0.0001	0.0003	0.0003	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

Asterisks in the same column indicate statistical difference ($p < 0.05$) with respect to the control treatment. DEG: degradation, IOM: incubated organic matter, DOM: Degraded organic matter; SCFA: Short chain fatty acids; SM15-PP85 (15% *S. mombin* + 85% *P. purpureum*); SM30-PP70 (30% *S. mombin* + 70% *P. purpureum*); AP15-PP85 (15% *A. pennatula* + 75% *P. purpureum*); AP30-PP70 (30% *A. pennatula* + 70% *P. purpureum*); PA15-PP85 (15% *P. aculeata* + 85% *P. purpureum*); PA30-PP70 (30% *P. aculeata* + 70% *P. purpureum*); BA15-PP85 (15% *B. alicastrum* + 85% *P. purpureum*); BA30-PP70 (30% *B. alicastrum* + 70% *P. purpureum*); BS15-PP85 (15% *B. simaruba* + 85% *P. purpureum*); BS30-PP70 (30% *B. simaruba* + 70% *P. purpureum*); 100PP: 100% *P. purpureum*; Means in the same column with same superscript are significantly different ($p < 0.05$) when compared to control (100PP).

4. Discussion

4.1. Chemical Composition and In Vitro Digestibility

In the present study the incorporation of the five forage tree species, independently at two different levels of inclusion (15 and 30%), into a diet based on forage grass *P. purpureum* improved nutritional composition, thus these tree forages are suitable for ruminant feeding. Crude protein content of the treatments was always above 7%, the minimum necessary for the correct function of rumen environment [24]. The mixture of grass with the foliage of native tree species at different levels can be a viable and a low-cost strategy to minimize the scarcity of nutrients in tropical cattle production systems. Gaviria-Urbe et al. [37] stated that including tree forages on ruminant's diet that are based on low quality grass, increases CP and decreases total carbohydrate intake. However, *P. purpureum* used in the present trial showed higher CP and lower values of structural carbohydrate content compared to other studies [18,38]. This is directly related to the age of regrowth of the pasture [39], as well as a higher digestibility and gas production as observed in this trial due to the high quality of the diets evaluated for tropical regions. The CP and EE values of *S. mombin* are similar to those obtained by Yusuf et al. [40] however, NDF and ADF contents were lower in the present study. The inclusion of *A. pennatula* at 30% had the highest CP content among the evaluated treatments, however the in vitro digestibility was reduced compared to the inclusion of 15% and this can be explained by its content of CT (31.1%) and cyanogenic (+++) glycosides that can alter the capacity of the microbiome to ferment nutrient components [15].

Metabolizable and net energy concentrations varied among evaluated treatments and was particularly high in BA30-PP70 (7.82 and 4.82 MJ kg⁻¹ DM, respectively), consistent with a higher gas production observed in this treatment. A higher net energy content promotes a higher digestibility of the total nutrients as a result of increased nutritive values. Metabolizable energy ranged between 6 and 7.8 MJ kg⁻¹ DM in the present investigation, this variations between species may be due to the differences in CP content and to a lesser extent to systematic errors made in the sampling of the gas produced as suggested by Tagliapietra et al. [41]. According to Krizsan et al. [42], microbial protein synthesis is favored by a higher ME input. Although, BA30-PP70 showed a higher net CH₄ production, its chemical composition and ME can improve animal performance and reduce CH₄ per kg of product.

4.2. Dry Matter Degradability, In Vitro Gas Production and pH

Dry matter degradation is inversely related to structural carbohydrate content [43,44], this is corroborated in treatments with species such as *S. mombin* or *B. alicastrum* which have higher digestibility and low NDF and/or FDA content or with mixtures constituted with PA, in which the opposite effect of carbohydrates was observed. Rumen degradation of DM at 48 h was improved in SM15-PP85, SM30-PP70, BA15-PP85, BA30-PP70 compared to control ($p \leq 0.0001$). The degradation of *A. pennatula* at 24 and 48 h was affected due to its content of secondary metabolites (CT: 31.1 g kg⁻¹ DM, alkaloids: ++, cyanogenic glycosides: +++++ and saponins: +) that can interfere in the degradation of carbohydrates and proteins. These findings are similar to those obtained by Albores-Moreno et al. [38] who supplemented *N. emargiata*, *L. leucocephala*, *P. piscipula* and *H. albicans* at 30% and increased CT content of a diet based on *P. purpureum*. Additionally, saponins can modify rumen fermentation by suppressing protozoa and selectively inhibiting some bacteria [45]. Secondary metabolites can make the diet less susceptible to the degradation of microorganism in the rumen due to their capacity to form complexes with the components of the diet, alter the mechanisms of degradation of the microorganism or reduce their population [46]. In addition, there is a direct relation between DMD and gas production, *i.e.*, the higher the digestibility, the higher the gas production rate expected. This is explained by the nutritional quality of the forages, especially by the structural carbohydrate content, *e.g.*, the diets that included *P. aculeata* presented a higher NDF content that will trigger a lower gas production. This observation agrees with authors such as Sánchez et al. [47] and Molina-Botero et al. [48].

The effect of plant secondary metabolites on maximum gas production can be observed when the fermentation parameters of *P. purpureum* are compared. Treatments with *A. pennatula*, *P. aculeata* and *B. simaruba* had the lowest gas productions and the highest content of secondary metabolites compared to *P. purpureum*. Regarding the CT content found in *A. pennatula* and *B. simaruba*, these have the capacity to inhibit enzymatic and microbial activity and consequently reduce fermentation [49]. Alkaloids present in *P. aculeata* can have negative effect on gas production as reported by Aguiar and Wink [50] due to a possible effect of these alkaloids on ruminal microorganisms. Aguiar and Wink [50] established that any outcome on the total gas production in the rumen can be interpreted as an effect on the microbial community. Fermentation kinetics of *A. pennatula*, *P. aculeata*, *S. mombin*, *B. alicastrum*, and *B. simaruba* incubated alone as single treatments were presented by Valencia-Salazar et al. [19]. The lowest maximum gas production observed by Valencia-Salazar et al. [19] was in *B. simaruba* and *A. pennatula* with 118.03 and 148.83 mL, respectively and *B. alicastrum* presented the highest maximum gas production (256.72 mL). These results were also observed in this study with *B. simaruba* and *B. alicastrum*, however, *A. pennatula* presented higher total gas production than *P. aculeata* in this study.

The observed pH values of the evaluated treatments are like those reported for tropical diets. Likewise, in the present investigation there is no relation between pH and SCFA variables, as described by Li et al. [51]. This observation is perhaps because pH values were not below 6.0, which is a critical value for the activity of the fibrolytic microbial population and, therefore, for fiber degradation. As the degradation of the fiber is the main precursor of SCFA in the rumen, the observation described above can also be related with methanogenesis [52]. On the other hand, and according to Meale et al. [53] SCFA had consistent concentrations on leguminous, non-leguminous and grass species (68.7–105.5 mMol). However, the methodology used for the quantification of SCFA must be considered since secondary metabolites can affect its production [52]. According to Li et al. [51], VFA production in the rumen is directly related to the ME consumed, and this study corroborated this postulate as the BS30-PP70 treatment had a low ME contribution (6.05 MJ kg⁻¹ DM) and resulted in a low content of SCFA (57.88 mMol L⁻¹), while with the BA30-PP70 treatment, the opposite occurred (7.2 MJ kg⁻¹ DM and 95.42 mMol L⁻¹).

4.3. Methane Production

Methane production of treatment with *P. aculeata* was low, perhaps due to the higher NDF content, and to the alkaloid content as described before. Cyanogenic glycosides present in *A. pennatula* are either toxic to methanogens or impair their growth potential by reducing the availability of sulphur [54]. Several studies have shown the reduction effect on CH₄ synthesis of cyanogenic glycosides from cassava [54] however, further work is needed to understand the mode of action of these components on the rumen microbiome and on animal performance. On the other hand, diet components that contain alkaloids have showed in other studies to reduce CH₄ formation [55,56]. Through the rumen simulation technique (RUSITEC), Khiaosa-ard et al. [56] showed that alkaloid supplementation at a low dose shifted the fermentation pathway to more propionate and less acetate and at a high dose an effect is observed on specific methanogenic archaea without affecting their abundance.

Results in CH₄ reduction were observed in the screening carried out by Valencia-Salazar et al. [19] with *S. mombin*, *P. aculeata* and *B. simaruba* and high CH₄ production with *B. alicastrum*). The high content of CT found *B. simaruba* can explain the lower CH₄ production at 24 and 48 h compared to control treatment, however, no statistical differences were observed in this study. Forages that contain CT have demonstrated to reduce CH₄ production both in vivo and in vitro trials [57]. Condensed tannins can reduce protozoal [58] and bacterial activity of archaea [59] by inhibiting enzyme activity, decreased degradation of substrates and direct action on the cell membrane [60]. The secondary effect of CT on CH₄ is the reduction in fiber digestion and fermentation, the decrease in the inter-specific transfer of hydrogens between protozoa and methanogenic archaea and

the consequent increase in the concentration of propionic acid in the rumen [61]. Piñeiro-Vázquez et al. [62] included 30% of *B. simaruba* in sheep diets and concluded that the effect of CT from this species reduced CH₄ emissions despite not observing a numerical difference, those results are similar to the ones obtained in the present investigation. However, different animal species may vary in their response to the same mitigation strategy [34] and the results obtained in the present study must be verified in a *in vivo* trial using cattle. Bhatta et al. [63] found a positive correlation ($R^2 = 0.98$) in CH₄ production measured using the SF₆ tracer technique and the *in vitro* gas production technique proposed by Menke and Steingass [31] as used in this trial. Among CH₄ quantification techniques, the *in vitro* fermentation technique involving incubation of substrates in rumen fluid has been extensively used for the evaluation of ruminant feeds, complement standard laboratory analysis, and constitute a cheaper alternative to evaluate large number of samples [34].

In developing countries, livestock is predominantly kept on high-roughage diets with no concentrate supplementation which increases ruminal methanogenesis, converting forage tree species that contain secondary metabolites in an alternative in many parts of the tropics for the improvement of animal performance and the reduction of enteric CH₄ emissions [18,45]. The extent of methane mitigation when feeding secondary metabolites, can be variable between *in vivo* and *in vitro* studies, so further research is necessary on the specific effect on rumen microorganisms and the chemical structure of these compounds.

5. Conclusions

The species evaluated in the present study presented nutritional quality that favored fermentation parameters such as total gas production, degradability, and short chain fatty acid production. Treatments that included *Bursera simaruba*, *Acacia pennatula* and *Parmentiera aculeata* contained considerable amounts of secondary metabolites such as condensed tannins, cyanogenic glycosides, and alkaloids. There was an inverse relation between NDF content and DM degradation, net gas and methane production and the content of secondary metabolites and methane production. However, the inclusion of these forage species had no statistical effect on the reduction of CH₄ emissions per unit of DM incubated or degraded at 24 and 48 h with respect to *P. purpureum*. However, their high nutritional quality can improve tropical diets based on low quality pastures.

Author Contributions: Conceptualization, S.S.V.-S., G.J.-F., J.C.K.-V.; methodology, S.S.V.-S., J.A. and I.C.M.-B.; software, I.C.M.-B. and S.S.V.-S.; validation, I.C.M.-B. and S.S.V.-S.; formal analysis, I.C.M.-B.; investigation, S.S.V.-S. and G.J.-F.; resources, J.A.; data curation, I.C.M.-B.; writing—original draft preparation, S.S.V.-S. and J.C.K.-V.; writing—review and editing, S.S.V.-S. and J.C.K.-V.; supervision, J.A., N.C. and G.J.-F.; project administration, J.A.; funding acquisition, J.A. and N.C. All authors have read and agreed to the published version of the manuscript.

Funding: This work was implemented as part of the CGIAR Research Program (CRP) on Climate Change, Agriculture and Food Security (CCAFS), and the Livestock CRP which are carried out with support from CGIAR Fund Donors and through bilateral funding agreements. For details, please visit <https://ccafs.cgiar.org/donors> (accessed on 1 December 2021). Likewise, the field work was financed by The College of the Southern Border (ECOSUR), Mexico www.ecosur.mx (accessed on 1 December 2021).

Institutional Review Board Statement: The work described here was conducted using rumen fluid obtained from fistulated cattle maintained in accordance with the requirements of Colombian law No 84/1989 and following protocols approved by the Ethics Committee of the International Center for Tropical Agriculture, assuring the welfare of animals used in the experiment. The manuscript does not contain clinical studies or patient data.

Data Availability Statement: All authors ensure that all data and materials support the findings and comply with field standards. The data presented in this study are available on request from the corresponding author.

Acknowledgments: This research was supported by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) and the Global Research Alliance on Agricultural

Greenhouse Gases (GRA) through their CLIFF-GRADS program. CCAFS capability building objectives are carried out with support from CGIAR Trust Fund and through bilateral funding agreements. For details, please visit <https://ccafs.cgiar.org/donors> (accessed on 1 December 2021). Thank you to CIAT for hosting the recipient and to the Government of New Zealand for providing financial support. We also acknowledge the financial assistance of OneCGIAR initiative on Livestock, Climate and System Resilience (LCSR). The doctoral scholarship granted to the first author by the National Council of Science and Technology of Mexico (CONACYT) is appreciated.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Beauchemin, K.; Ungerfeld, E.; Eckard, R.; Wang, M. Fifty years of research on rumen methanogenesis: Lessons learned and future challenges for mitigation. *Animal* **2020**, *14* (Suppl. S1), s2–s16. [[CrossRef](#)]
2. Glatzle, A. Questioning key conclusions of FAO publications ‘Livestock’s Long Shadow’ (2006) appearing again in ‘Tackling Climate Change Through Livestock’. *Pastoralism* **2013**, *4*, 1. [[CrossRef](#)]
3. Haque, N. Dietary manipulation: A sustainable way to mitigate methane emissions from ruminants. *Anim. Feed Sci. Technol.* **2018**, *60*, 15. [[CrossRef](#)]
4. Kurihara, M.; Magner, T.; Hunter, R.; McCrabb, G. Methane production and energy partition of cattle in the tropics. *Br. J. Nutr.* **1999**, *81*, 227–234. [[CrossRef](#)]
5. Hyland, J.J.; Styles, D.; Jones, D.J.; Williams, A.P. Improving livestock production efficiencies presents a major opportunity to reduce sectoral greenhouse gas emissions. *Agric. Syst.* **2016**, *147*, 123–131. [[CrossRef](#)]
6. Cammack, K.M.; Austin, K.J.; Lamberson, W.R.; Conant, G.C.; Cunningham, H.C. Ruminant nutrition symposium: Tiny but mighty: The role of the rumen microbes in livestock production. *Anim. Sci. J.* **2018**, *96*, 752–770. [[CrossRef](#)]
7. Henderson, G.; Cox, F.; Ganesh, S.; Jonker, A.; Young, W.; Janssen, P.H. Rumen microbial community composition varies with diet and host, but a core microbiome is found across a wide geographical range. *Sci. Rep.* **2015**, *5*, 1–15. [[CrossRef](#)] [[PubMed](#)]
8. Lan, W.; Yang, C. Ruminal methane production: Associated microorganisms and the potential of applying hydrogen-utilizing bacteria for mitigation. *Sci. Total Environ.* **2019**, *654*, 1270–1283. [[CrossRef](#)] [[PubMed](#)]
9. Huws, S.A.; Creevey, C.J.; Oyama, L.B.; Mizrahi, I.; Denman, S.E.; Popova, M.; Mayorga, O.L.; Morrisson, S.; Elliott, C.; Morgavi, D.P.; et al. Addressing global ruminant agricultural challenges through understanding the rumen microbiome: Past, present, and future. *Front. Microb.* **2018**, *9*, 2161. [[CrossRef](#)] [[PubMed](#)]
10. Belanche, A.; De La Fuente, G.; Newbold, C.J. Study of methanogen communities associated with different rumen protozoal populations. *FEMS Microbiol. Ecol.* **2014**, *90*, 663–677. [[CrossRef](#)]
11. Tapio, I.; Snelling, T.J.; Strozzi, F.; Wallace, R.J. The ruminal microbiome associated with methane emissions from ruminant livestock. *J. Anim. Sci. Biotechnol.* **2017**, *8*, 7. [[CrossRef](#)]
12. Berghuis, B.A.; Yu, F.B.; Schulz, F.; Blainey, P.C.; Woyke, T.; Quake, S.R. Hydrogenotrophic methanogenesis in archaeal phylum Verstraetearchaeota reveals the shared ancestry of all methanogens. *Proc. Natl. Acad. Sci. USA* **2019**, *116*, 5037–5044. [[CrossRef](#)]
13. Arndt, C.; Hristov, A.N.; Price, W.J.; McClelland, S.C.; Pelaez, A.M.; Cueva, S.F.; Martin, C.; Schwarm, A.; Veneman, J.B.; ZhongTang, Y.; et al. Strategies to mitigate enteric methane emissions by ruminants—a way to approach the 2.0 °C target. *Preprint* **2021**, *in press*.
14. Arango, J.; Ruden, A.; Martinez-Baron, D.; Loboguerrero, A.M.; Berndt, A.; Chacón, M.; Torres, C.F.; Oyhantcabal, W.; Gomez, C.A.; Ricci, P.; et al. Ambition Meets Reality: Achieving GHG Emission Reduction Targets in the Livestock Sector of Latin America. *Front. Sustain. Food Syst.* **2020**, *4*, 65. [[CrossRef](#)]
15. Ku-Vera, J.C.; Jiménez-Ocampo, R.; Valencia-Salazar, S.; Flores-Santiago, E.J.; Montoya-Flores, M.D.; Molina-Botero, I.C.; Arango, J.; Gomez-Bravo, C.A.; Aguilar-Perez, C.F.; Solorio-Sanchez, F.J. Role of Secondary Plant Metabolites on Enteric Methane Mitigation in Ruminants. *Front. Vet. Sci.* **2020**, *7*, 584. [[CrossRef](#)]
16. Patra, A.K.; Park, T.; Kim, M.; Yu, Z. Rumen methanogens and mitigation of methane emissions by anti-methanogenic compounds and substances. *J. Anim. Sci. Biotechnol.* **2017**, *8*, 13. [[CrossRef](#)]
17. Rao, I.; Peters, M.; Castro, A.; Schultze-Kraft, R.; White, D.; Fisher, M.; Jenet, A.; Schmidt, A.; Vivas, N.; Rudel, T. LivestockPlus—The sustainable intensification of forage-based agricultural systems to improve livelihoods and ecosystem services in the tropics. *Trop. Grassl.* **2015**, *3*, 59. [[CrossRef](#)]
18. Valencia-Salazar, S.; Piñeiro, A.T.; Molina, I.C.; Lazos, F.J.; Uuh, J.J.; Segura, M.; Ramírez, L.; Solorio, F.J.; Ku, J.C. Potential of *Samanea saman* pod meal for enteric methane mitigation in crossbred heifers fed low-quality tropical grass. *Agric. Forest. Meteorol.* **2018**, *258*, 108–116. [[CrossRef](#)]
19. Valencia-Salazar, S.S.; Jiménez-Ferrer, G.; Arango, J.; Molina-Botero, I.C.; Chirinda, N.; Piñeiro-Vázquez, A.; Jiménez-Ocampo, R.; Nahed-Toral, J.; Ku-Vera, J. Enteric methane mitigation and fermentation kinetics of forage species from Southern Mexico: In vitro screening. *Agroforest Syst.* **2021**, *95*, 293–305. [[CrossRef](#)]

20. García del Valle, Y.; Naranjo, E.; Caballero, J.; Martorell, C.; Ruan-Soto, F.; Enríquez, P. Cultural significance of wild mammals in mayan and mestizo communities of the Lacandon Rainforest, Chiapas, Mexico. *J. Ethnobiol. Ethnomed.* **2015**, *11*, 36. [[CrossRef](#)]
21. ISO. *ISO 6496. Animal Feeding Stuffs—Determination of Moisture and Other Volatile Matter Content*; International Organization for Standardization: Geneva, Switzerland, 1999.
22. AOAC. *Official Methods of Analysis*, 16th ed.; Association of Official Analytical Chemists (AOAC): Washington, DC, USA, 2005.
23. AOAC. *Official Methods of Analysis*, 15th ed.; Association of Official Analytical Chemists (AOAC): Washington, DC, USA, 1990.
24. Van Soest, P.J.; Robertson, J.B.; Lewis, B.A. Methods for dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* **1999**, *74*, 3583–3597. [[CrossRef](#)]
25. ISO. *ISO 9831. Animal Feeding Stuffs, Animal Products, and Faeces or Urine—Determination of Gross Calorific Value—Bomb Calorimeter method*; International Organization for Standardization: Geneva, Switzerland, 1998.
26. Tilley, J.M.A.; Terry, R.A. A two-stage technique for the in vitro digestion of forage crops. *J. Br. Grassl. Soc.* **1963**, *18*, 104–111. [[CrossRef](#)]
27. Hagerman, A.E.; Butler, L.G. Protein precipitation method for the quantitative determination of tannins. *J. Agric. Food Chem.* **1978**, *26*, 809–812. [[CrossRef](#)]
28. Domínguez, X.A. *Métodos de Investigación Fitoquímica*; Editorial Limusa: Mexico City, Mexico, 1979.
29. ICONTEC. NTC 668. *Alimentos y Materias Primas. Determinación de los Contenidos de Grasa y Fibra Cruda*; Instituto Colombiano de Normas Técnicas y Certificación (ICONTEC): Bogotá, Colombia, 1973; p. 8.
30. Menke, K.H.; Raab, L.; Salewski, A.; Steingass, H.; Fritz, D.; Schneider, W. The estimation of the digestibility and metabolisable energy content of ruminant feedstuffs from the gas production when they are incubated with rumen liquor. *J. Agric. Sci.* **1979**, *93*, 217. [[CrossRef](#)]
31. Menke, K.H.; Steingass, H. Estimation of the energetic feed value obtained from chemical analysis and in vitro gas production using rumen fluid. *Anim. Res. Dev.* **1988**, *28*, 7–55.
32. Getachew, G.; Makkar, H.P.S.; Becker, K. Tropical browses: Contents of phenolic compounds, in vitro gas production and stoichiometric relationship between short chain fatty acid and in vitro gas production. *J. Agric. Sci.* **2002**, *139*, 341–352. [[CrossRef](#)]
33. Theodorou, M.K.; Williams, B.A.; Dhanoa, M.S.; McAllan, A.B.; France, J.A. Simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Anim. Feed Sci. Technol.* **1994**, *48*, 185–197. [[CrossRef](#)]
34. Yáñez-Ruiz, D.R.; Bannink, A.; Dijkstra, J.; Kebreab, E.; Morgavi, D.P.; O’Kiely, P.; Reynolds, C.K.; Schwarm, A.; Shingfield, K.J.; Yu, Z.; et al. Design, implementation and interpretation of in vitro batch culture experiments to assess enteric methane mitigation in ruminants—A review. *Anim. Feed Sci. Technol.* **2016**, *216*, 1–18. [[CrossRef](#)]
35. Lavrenčič, A.; Mills, C.R.; Stefanon, B. Application of the Gompertz model to describe the fermentation characteristics of chemical components in forages. *Anim. Sci. J.* **1998**, *66*, 155–161. [[CrossRef](#)]
36. SAS Institute. *User’s Guide: Statistics Version 9.4*; SAS Inst. Inc.: Cary, NC, USA, 2012.
37. Gaviria-Urbe, X.; Bolívar, D.; Rosenstock, T.; Molina-Botero, I.C.; Chirinda, N.; Barahona, R.; Arango, J. Nutritional quality, voluntary intake and enteric methane emissions of diets based on novel Cayman grass and its associations with two *Leucaena* shrub legumes. *Vet. Sci. Front.* **2020**, *7*, 579189. [[CrossRef](#)] [[PubMed](#)]
38. Albores-Moreno, S.; Alayón-Gamboa, J.A.; Miranda-Romero, L.A.; Alarcón-Zúñiga, B.; Jiménez-Ferrer, G.; Ku-Vera, J.C.; Piñeiro-Vázquez, A.T. 2018 Effect of tree foliage supplementation of tropical grass diet on in vitro digestibility and fermentation, microbial biomass synthesis and enteric methane production in ruminants. *Trop. Anim. Health Prod.* **2018**, *58*, 893–904. [[CrossRef](#)]
39. Ortega-Gómez, R.; Castillo-Gallegos, E.; Jarillo-Rodríguez, J.; Escobar-Hernández, R.; Ocaña-Zavaleta, E.; Valles de la Mora, B. Nutritive Quality of Ten Grasses During the Rainy Season In A Hot-Humid Climate And Ultisol Soil. *Trop. Subtrop. Agroecosyst.* **2011**, *13*, 481–491.
40. Yusuf, A.O.; Egbinola, O.O.; Ekunseitan, D.A.; Salem, A.Z.M. Chemical characterization and in vitro methane production of selected agroforestry plants as dry season feeding of ruminants livestock. *Agroforest Syst.* **2020**, *94*, 1481–1489. [[CrossRef](#)]
41. Tagliapietra, F.; Cattani, M.; Hansen, H.H.; Hindrichsen, I.K.; Bailoni, L.; Schiavon, S. Metabolizable energy content of feeds based on 24 or 48 h in situ NDF digestibility and on in vitro 24 h gas production methods. *Anim. Feed Sci. Technol.* **2011**, *170*, 182–191. [[CrossRef](#)]
42. Krizsan, S.J.; Pang, D.; Fatehi, F.; Rinne, M.; Huhtanen, P. Metabolisable energy of grass and red clover silages fed to sheep at maintenance level. *Animal* **2020**, *14*, 753–762. [[CrossRef](#)] [[PubMed](#)]
43. Narváez, N.; Lascano, C.E. Caracterización química de especies arbóreas tropicales con potencial forrajero en Colombia. *Pasturas Trop.* **2004**, *26*, 1–8.
44. Lee, M.R.F.; Jordana Rivero, M.; Cone, J.W. The role of pasture in the diet of ruminant livestock. In *Improving Grassland and Pasture Management in Temperature Agriculture*; Marshall, A., Collins, R., Eds.; Burleigh Dodds Science Publishing Limited: Cambridge, UK, 2018. [[CrossRef](#)]
45. Goel, G.; Makkar, H.P. Methane mitigation from ruminants using tannins and saponins. *Trop. Anim. Health Prod.* **2012**, *44*, 729–739. [[CrossRef](#)] [[PubMed](#)]
46. Tedeschi, L.O.; Muir, J.P.; Naumann, H.D.; Norris, A.B.; Ramírez-Restrepo, C.A.; Mertens-Talcott, S.U. Nutritional Aspects of Ecologically Relevant Phytochemicals in Ruminant Production. *Front. Vet. Sci.* **2021**, *8*, 628445. [[CrossRef](#)] [[PubMed](#)]
47. Sánchez, D.E.; Arreaza, L.C.; Abadía, B. Estudio de la cinética de degradación in vitro de cuatro forrajes tropicales y una leguminosa de clima templado. *Cienc. Tecnol. Agropecu.* **2005**, *6*, 58–68. [[CrossRef](#)]

48. Molina-Botero, I.C.; Mazabel, J.; Arceo-Castillo, J.; Urrea-Benitez, J.L.; Olivera-Castillo, L.; Barahona-Rosales, R.; Chirinda, N.; Ku-Vera, J.; Arango, J. Effect of the addition of *Enterolobium cyclocarpum* pods and *Gliricidia sepium* forage on dry matter degradation, volatile fatty acid concentration, and in vitro methane production. *Trop. Anim. Health Prod.* **2020**, *52*, 2787–2798. [[CrossRef](#)]
49. Patra, A.K.; Yu, Z. Effect s of vanillin, quillaja saponin, and essential oils on in vitro fermentation and protein degrading microorganisms of the rumen. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 897–905. [[CrossRef](#)]
50. Aguiar, R.; Wink, M. Do naïve ruminants degrade alkaloids in the rumen? *J. Chem. Ecol.* **2005**, *31*, 761–787. [[CrossRef](#)] [[PubMed](#)]
51. Li, M.M.; White, R.R.; Guan, L.L.; Harthan, L.; Hanigan, M.D. Metatranscriptomic analyses reveal ruminal pH regulates fiber degradation and fermentation by shifting the microbial community and gene expression of carbohydrate-active enzymes. *Anim. Microbiome* **2021**, *3*, 32. [[CrossRef](#)]
52. Bodas, R.; Prieto, N.; García-González, R.; Andrés, S.; Giráldez, F.J.; López, S. Manipulation of rumen fermentation and methane production with plant secondary metabolites. *Anim. Feed Sci. Technol.* **2012**, *176*, 78–93. [[CrossRef](#)]
53. Meale, S.J.; Chaves, A.V.; Baah, J.; McAllister, T.A. Methane production of different forages in in vitro ruminal fermentation. *Asian-Aust. J. Anim. Sci.* **2012**, *25*, 86. [[CrossRef](#)]
54. Vongsamphanh, P.; Inthapanya, S.; Preston, T.; Van Dung, D.; Xuan Ba, N. Effect of leaves from sweet or bitter cassava and brewers' grains on methane production in an in vitro rumen incubation of cassava root pulp-urea. *Livest. Res. Rural Dev.* **2018**, *30*, 167.
55. Randami, D.; Chaudhry, A.S.; Seal, C.J. Alkaloid and polyphenol analysis by HPLC in green and black tea powders and their potential use as additives in ruminant diets. *AIP Conf. Proc.* **2018**, *1927*, 030008. [[CrossRef](#)]
56. Khiaosa-Ard, R.; Mahmood, M.; Lerch, F.; Traintinger, F.P.; Petri, R.M.; Münnich, M.; Zebeli, Q. Physicochemical stressors and mixed alkaloid supplementation modulate ruminal microbiota and fermentation in vitro. *Anaerobe* **2020**, *65*, 102263. [[CrossRef](#)]
57. Patra, A.K.; Saxena, J. A new perspective on the use of plant secondary metabolites to inhibit methanogenesis in the rumen. *Phytochemistry* **2010**, *71*, 1198–1222. [[CrossRef](#)] [[PubMed](#)]
58. Puchala, R.; Animut, G.; Patra, A.K.; Detweiler, G.D.; Wells, J.E.; Varel, V.H.; Sahlu, T. Methane emissions by goats consuming *Sericea lespedeza* at different feeding frequencies. *Anim. Feed Sci. Technol.* **2012**, *175*, 76–84. [[CrossRef](#)]
59. Min, B.R.; Solaiman, S.; Shange, R.; Eun, J.S. Gastrointestinal bacterial and methanogenic archaea diversity dynamics associated with condensed tannin-containing pine bark diet in goats using 16S rDNA amplicon pyrosequencing. *Int. J. Food Microbiol.* **2014**, *11*, 1–11. [[CrossRef](#)] [[PubMed](#)]
60. McSweeney, C.S.; Palmer, B.; Bunch, R.; Krause, D.O. Microbial interactions with tannins: Nutritional consequences for ruminants. *Anim. Feed. Sci. Technol.* **2001**, *91*, 83–93. [[CrossRef](#)]
61. Vargas, J.; Cárdenas, E.; Pabón, M.; Carulla, J. Emisión de metano entérico en rumiantes en pastoreo. *Arch. De Zootec.* **2012**, *61*, 51–66. [[CrossRef](#)]
62. Piñeiro-Vázquez, A.T.; Canul-Solis, J.R.; Casanova-Lugo, F.; Chay-Canul, A.J.; Ayala-Burgos, A.; Solorio-Sánchez, J.; Aguilar-Pérez, C.F.; Ku-Vera, J.C. Emisión de metano en ovinos alimentados con *Pennisetum purpureum* y árboles que contienen taninos condensados. *Rev. Mex. Cienc. Pecu.* **2017**, *2*, 111–119. [[CrossRef](#)]
63. Bhatta, R.; Tajima, K.; Takusari, N.; Higuchi, K.; Enishi, O.; Kurihara, M. Comparison of in vivo and in vitro techniques for methane production from Ruminant diets. *Asian-Aust. J. Anim. Sci.* **2007**, *20*, 1049–1056. [[CrossRef](#)]