

Impact of Comorbid Personality Disorders on Depression Treatment in Routine Outpatient Care

Citation for published version (APA):

van Bronswijk, S. C., van Dijk, D. A., van den Boogaard, T. M., Deen, M. L., Ruhe, H. G., Spijker, J., & Peeters, F. P. M. L. (2021). Impact of Comorbid Personality Disorders on Depression Treatment in Routine Outpatient Care. *American Journal of Psychotherapy*, 74(4), 150-156. <https://doi.org/10.1176/appi.psychotherapy.202120200046>

Document status and date:

Published: 01/12/2021

DOI:

[10.1176/appi.psychotherapy.202120200046](https://doi.org/10.1176/appi.psychotherapy.202120200046)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Impact of Comorbid Personality Disorders on Depression Treatment in Routine Outpatient Care

Suzanne C. van Bronswijk, M.D., Ph.D., Dyllis A. van Dijk, M.D., Th. Michael van den Boogaard, M.D., Ph.D., Mathijs L. Deen, M.Sc., Henricus G. Ruhé, M.D., Ph.D., Jan Spijker, M.D., Ph.D., Frenk P. M. L. Peeters, M.D., Ph.D.

Objective: The impact of personality disorder on treatment effectiveness for depression has been debated, and study results have been inconsistent. However, studies that report a negative impact of personality disorders on depression treatment outcomes are often characterized by uncontrolled treatment designs. Within such contexts, individuals with depression and personality disorders are at risk to receive suboptimal treatment. The aim of this retrospective observational study was to investigate whether and to what extent comorbid personality disorders were associated with the type and amount of depression treatment received in routine outpatient care.

Methods: Retrospectively extracted data from electronic records of 1,455 outpatients treated for depression at several sites of a nationwide mental health provider in the Netherlands were included. The type and number of treatment sessions and visits were analyzed by using regression models.

Results: Individuals with depression and comorbid personality disorders received more psychotherapy sessions than individuals without personality disorders, irrespective of depression severity. The number of pharmacotherapy sessions and supportive and crisis visits did not differ between individuals with and without comorbid personality disorders.

Conclusions: Individuals with depression and personality disorders received more intensive treatment than individuals without comorbid personality disorders. These results conflict with treatment guidelines and recommendations from high-quality studies and may be indicative of overtreatment among this large group of patients.

Am J Psychother 2021; 74:150–156;
doi: 10.1176/appi.psychotherapy.202120200046

Despite the many available and effective treatment options for depression (1, 2), a substantial number of individuals do not achieve remission (3, 4). Because individual treatment responses vary and are highly unpredictable, identifying risk factors associated with treatment outcome may help to personalize treatment selection and subsequently improve remission rates (5, 6). In addition to known risk factors for treatment resistance, including high depression severity and chronicity of the index episode (7, 8), the presence of comorbid personality disorder is often considered (9, 10). Its relevance is potentially large because personality disorders are a common comorbid condition of outpatients with depression (11, 12).

Two frequently cited meta-analyses (13, 14) have reported that personality disorders negatively affect acute-phase treatment outcomes in depression, concluding that the presence of these disorders is important in the prognosis of depression. In contrast, a review by Mulder (15), which focused solely on controlled studies, found no negative effects of personality disorders. According to Mulder, this difference in findings could be explained by patients with

these comorbid conditions receiving less optimal treatment in uncontrolled studies, because such studies are likely biased by clinicians who regard a personality disorder as a relevant factor in treatment selection. Three meta-analyses have contributed to Mulder's hypothesis, indicating no significant differences in outcomes between individuals with and without personality disorders in controlled trials of pharmacotherapy (16, 17) and cognitive-behavioral therapy

HIGHLIGHTS

- Individuals with personality disorders received more sessions of psychotherapy for depression than individuals without these disorders.
- Personality disorders did not affect the number of pharmacotherapy visits nor crisis visits for depression.
- The challenging treatment of patients with depression and a comorbid personality disorder requires focused depression therapy to reduce the risk of both under- and overtreatment.

(18) for depression. Their findings have been replicated and extended by a recent meta-analysis (19) that focused on treatment outcomes in only well-designed controlled trials of pharmacotherapy and psychotherapy for depression. Results showed no significant differences between individuals with and without comorbid personality disorders in terms of average depression severity change and rates of response and remission (19). Because individuals with personality disorders appear to have better treatment outcomes in studies with controlled treatment selection, the less satisfactory outcomes of naturalistic designs may be incorrectly attributed to the comorbid condition, instead of to insufficiently provided depression treatment. On the basis of these meta-analyses (16–19), one could hypothesize that with care as usual, treatment provision may be less optimal for individuals with personality disorders. However, only a few older studies (20, 21) have reported on this issue, indicating that individuals with comorbid personality disorders receive less pharmacotherapy and electroconvulsive therapy (ECT) for depression, compared with counterparts without comorbid personality disorders. Recent studies addressing this research question are lacking.

The aim of the current study was to examine whether, and to what extent, the presence of comorbid personality disorders was associated with the type and number of treatment sessions received in the context of naturalistic outpatient depression care. First, we examined whether individuals with comorbid personality disorders received different types and intensities of depression treatments than those recommended by international and national evidence-based guidelines (i.e., psychotherapy, pharmacotherapy, a combination of both, or ECT [22–24]). We hypothesized, on the basis of the limited evidence available on the receipt of less pharmacotherapy and ECT among individuals with depression and comorbid personality disorders, that these individuals would receive less optimal treatment (i.e., fewer therapy sessions, particularly for pharmacotherapy) than individuals with depression and no comorbid personality disorders, irrespective of the level of depression severity. Second, we investigated whether comorbid personality disorders affected the number of other types of care received. We hypothesized that individuals with depression and comorbid personality disorders would be more likely to receive a greater number of supportive and crisis visits than individuals without comorbid personality disorders.

METHODS

Design and Participants


This naturalistic cohort study was conducted at PsyQ, a nationwide organization in the Netherlands that provides outpatient secondary mental health care. The patients whose data were studied were referred to several PsyQ outpatient departments for specialized treatment for depression. Anonymized data were retrospectively extracted from the electronic patient record (EPR) of individuals

who completed a diagnostic workup between June 2014 and June 2016. Data from adults (ages ≥ 18 years) who met the *DSM-IV-TR* criteria for major depressive disorder or dysthymic disorder according to their EPR were selected for inclusion in the study. Information on treatments received was extracted for a maximum of 2 years after the diagnostic workup. The Medical Research Ethics Committee of the Leiden University Medical Centre waived formal informed consent requirements because only routinely collected information was used for the study (25). Data from individuals who had objected to the use of their information for scientific purposes were not included. Exclusion criteria were a primary diagnosis of bipolar disorder, psychotic disorder, or substance dependence (except for nicotine). All patients whose data were analyzed had received treatments by trained professionals (psychiatrists, psychologists, and psychiatric nurses) according to the Dutch national evidence-based guideline for depression treatment (24).

Measures

Outcomes. The primary outcome of this study was the type and number of face-to-face treatment sessions received in accordance with international and national evidence-based guidelines for depression treatment (e.g., pharmacotherapy, psychotherapy, a combination of both, and ECT). To further examine the other types of care received by individuals with depression, with or without a comorbid personality disorder, the number of supportive visits and crisis visits were examined as secondary outcomes. Supportive visits were defined as planned face-to-face appointments with social workers or nurses focusing on psychosocial problems. Crisis visits were unplanned face-to-face appointments regarding an acute situation (e.g., suicidality) and were provided by various health care professionals, including psychiatrists and psychologists. Primary and secondary outcomes were extracted from the EPR, which used a coding system to represent the type and number of treatment sessions received.

Comorbid personality disorder. The presence of a comorbid personality disorder was based on a patient's score on the Standardized Assessment of Personality–Abbreviated Scale (SAPAS), which was an element of the routine diagnostic workup and was available to the therapists. The SAPAS is a clinician-rated eight-item questionnaire that is used to screen for the presence of personality disorders in routine clinical care (26–28). It includes descriptive statements about the person that can be scored as 0 (not present) or 1 (present). A score of ≥ 3 has been found to correctly identify 73.1% of comorbid personality disorders among an outpatient sample of individuals with depression (26) and is used as a cutoff score to indicate the presence of a personality disorder. For readability purposes, we will refer to having this score as having a “comorbid personality disorder.” Individuals missing a SAPAS score according to their EPR were excluded from the analyses.

FIGURE 1. Flowchart of the sample studied^a

^a There were no significant differences in clinical and demographic parameters between analyzed electronic patient records (N=1,455) and records missing a completed baseline Quick Inventory of Depressive Symptomatology–Self Report (QIDS-SR) or Standardized Assessment of Personality–Abbreviated Scale (SAPAS) score (N=780), except for previously undergone inpatient or day treatment. However, portions were small in both groups (2.5% [N=37] versus 4.2% [N=33], $\chi^2=4.77$, $p=0.03$). Blue boxes indicate the number of individuals missing a complete assessment.

Additional pretreatment variables. Demographic variables (age and gender) were extracted from the EPRs. Baseline depression severity was measured with the Quick Inventory of Depressive Symptomatology–Self Report (QIDS-SR) (29), a 16-item self-report depression symptom severity scale. The QIDS-SR score (not included in the diagnostic workup but filled out separately online by the patients) was considered a baseline measure when it was obtained 45 days before or after the diagnostic workup. Individuals with missing baseline QIDS-SR scores according to their EPR were excluded from the analyses. Information on episode duration and severity, comorbid anxiety, previous treatment failures, functional impairment, and psychosocial stressors was obtained by using the Dutch Measure for Quantification of Treatment Resistance in Depression (25) during the diagnostic workup. Additionally, childhood adversity was assessed by the clinician during the diagnostic workup by asking the question, “Have you had to deal with severe affective neglect, (repeated) sexual abuse, or psychological or physical threats during your youth (before the age of 16 years)?” Information on the DSM diagnosis (major depressive disorder or dysthymic disorder) and the duration of treatment in days was also extracted from the EPRs.

Statistical Analyses

First, pretreatment characteristics of individuals who were missing baseline QIDS-SR and/or SAPAS scores (N=780) and of individuals included in the analyses (N=1,455) were compared by using t tests (with Cohen’s d effect sizes) and chi-square tests where appropriate. Second, pretreatment variables of individuals with and without

comorbid personality disorders were examined using t tests (with Cohen’s d effect sizes) and chi-square tests. Third, the association between the amount of depression treatment received in accordance with international and national evidence-based guidelines and the presence of a comorbid personality disorder was tested. For this analysis, two separate negative binomial regression models were applied. The numbers of psychotherapy and pharmacotherapy sessions were the dependent variables in the first and second models, respectively. For both models, the presence of a comorbid personality disorder was included as an independent variable. To examine the moderating impact of baseline depression severity on the association between number of sessions and the presence of comorbid personality disorders, QIDS-SR baseline score and a QIDS-SR baseline score \times comorbid personality disorder status interaction were added to the models as independent variables. After that, two negative binomial regression models were applied with the secondary outcomes (the numbers of supportive and crisis visits) as dependent variables and comorbid personality disorder status, QIDS-SR baseline score, and QIDS-SR baseline score \times comorbid personality disorder status interaction as independent variables. For all analyses, the independent variables were centered to prevent errors in statistical inference (30).

RESULTS

Pretreatment Comparisons Between Individuals With and Without Missing Baseline QIDS-SR and SAPAS Scores

As illustrated in Figure 1, 1,455 of 2,235 individuals completed baseline QIDS-SR and SAPAS assessments, according to their EPRs, and were therefore included in our analyses. Individuals with and individuals without a completed baseline QIDS-SR and/or SAPAS assessment did not significantly differ in terms of age; gender; baseline depression severity; previous treatment failures, including pharmacotherapy, psychotherapy, or ECT; functional impairment; comorbid anxiety; psychosocial stressors; or childhood adversity. Individuals with missing baseline QIDS-SR and/or SAPAS scores had significantly more failures of intensified treatment (i.e., inpatient and day treatment) than those with complete QIDS-SR and/or SAPAS scores; however, the proportion of individuals who underwent these intensified treatments was low in both groups (4.2% [N=33] versus 2.5% [N=37], $\chi^2=4.77$, $p=0.03$).

Pretreatment Comparisons Between Individuals With and Without Comorbid Personality Disorders

Table 1 shows the differences in pretreatment characteristics between individuals with and without comorbid personality disorders. Individuals without comorbid personality disorders were significantly older compared with individuals with personality disorders ($t=2.99$, $df=1,453$, $p=0.003$), although the effect size of this difference was small (Cohen’s $d=0.16$). Comorbid personality disorders were associated

with higher average QIDS-SR baseline scores ($t=3.80$, $df=1,453$, $p<0.001$), although the effect size of the difference between individuals with and without personality disorders was small (Cohen's $d=0.21$). The index episode of depression was significantly longer for individuals with comorbid personality disorders ($\chi^2=24.9$, $p<0.001$). In addition, higher levels of comorbid anxiety, psychosocial stressors, and childhood adversity were found among individuals with comorbid personality disorders compared to those without personality disorders ($\chi^2=27.1$, $p<0.001$; $\chi^2=4.6$, $p=0.03$; $\chi^2=46.7$, $p<0.001$), although the effect sizes of these differences were small ($\Phi=0.14$, $\Phi=0.06$, and $\Phi=0.18$, respectively). Finally, comorbid personality disorders were associated with a higher proportion of dysthymia diagnoses ($\chi^2=7.5$, $p=0.006$) and a higher average treatment length in days ($t=-2.18$, $df=1,453$, $p=0.03$).

Received Depression Treatment in Accordance With Evidence-Based Guidelines

Table 2 provides the mean number of pharmacotherapy and psychotherapy sessions for individuals with and without comorbid personality disorders. Table 3 summarizes the results of the negative binomial regression models on the association between comorbid personality disorders and the number of pharmacotherapy and psychotherapy sessions. There was no significant association between the presence of comorbid personality disorders and the number of pharmacotherapy sessions received. Higher baseline depression severity was associated with a higher number of pharmacotherapy sessions, but there was no significant "depression severity \times personality disorder status" interaction. Individuals with comorbid personality disorders received significantly more psychotherapy sessions than individuals without personality disorders. No significant effect of baseline severity or baseline depression severity \times personality disorder status interaction on the number of received psychotherapy sessions was found. The number of ECT sessions was not analyzed as a separate outcome because only four individuals received ECT.

TABLE 1. Pretreatment characteristics of individuals with depression with and without comorbid personality disorder (PD)^a

Characteristic	Individuals without PD (N=521)		Individuals with PD (N=934)	
	N	%	N	%
Age (M \pm SD years)**	41.3 \pm 13.2		39.2 \pm 12.4	
Female	353	67.8	611	65.4
QIDS-SR score (M \pm SD)**	16.3 \pm 4.9		17.2 \pm 4.4	
Episode duration**				
\leq 12 months	286	54.9	410	43.9
13–24 months	117	22.5	199	21.3
$>$ 24 months	118	22.6	325	34.8
Treatment failures				
Antidepressants	215	41.3	380	40.7
Psychotherapy	213	40.9	397	42.5
Electroconvulsive therapy	4	.8	4	.4
Intensified treatment ^b	13	2.5	24	2.6
Functional impairment				
No impairment (GAF 90–100)	2	.4	4	0.4
Mild (GAF 60–90)	84	16.1	104	11.1
Moderate (GAF 30–60)	427	82.0	808	86.5
Severe (GAF 0–30)	8	1.5	18	1.9
Comorbid anxiety*				
Not present	254	48.8	330	35.3
Present, but not fulfilling <i>DSM-IV</i> criteria	188	36.1	394	42.2
Fulfilling <i>DSM-IV</i> criteria	79	15.2	210	22.5
Psychosocial stressors	394	75.6	751	80.4
Childhood adversity**	125	24.0	391	41.9
Dysthymia diagnosis** ^c	31	6.0	95	10.2
Treatment length (M \pm SD days)* ^d	332.5 \pm 238.9		361.4 \pm 245.5	

^a GAF, Global Assessment of Functioning; QIDS-SR, Quick Inventory of Depressive Symptomatology–Self Report.

^b Intensified treatment was defined as day-patient treatment or inpatient treatment.

^c Proportion of individuals with a *DSM* diagnosis of dysthymia (and not major depressive disorder).

^d Treatment length in days within a 2-year follow-up period, including all treatments (pharmacotherapy, psychotherapy, crisis contacts, and supportive visits).

* $p<.05$, ** $p<.01$; significant difference between groups with and without PD.

Total Care Received, Including Non-Evidence-Based Treatments

Table 2 shows the mean number of supportive and crisis visits for individuals with and without comorbid personality disorders. Table 3 summarizes the estimates of two negative binomial regression models for the secondary outcomes (the number of supportive and crisis visits). For both models, higher baseline depression severity was (borderline) significantly associated with more visits. The presence of comorbid

TABLE 2. Amount and type of depression treatment received by individuals with and without comorbid personality disorder (PD)

Type of visit	Individuals without PD (N=521)		Individuals with PD (N=934)	
	M	SD	M	SD
Pharmacotherapy	3.90	5.64	3.91	5.09
Psychotherapy	15.90	19.76	22.08	28.37
Supportive	7.96	16.38	10.40	22.34
Crisis	.80	6.72	.60	4.05

TABLE 3. Effect of comorbid personality disorder (PD) on depression treatment received (negative binominal regression)

Type of visit	B	SE	Wald	p	OR	95% CI
Pharmacotherapy sessions						
Intercept	1.36	.06	505.89	<.001	3.91	3.47–4.40
Depression severity	.50	.01	16.64	<.001	1.05	1.03–1.08
Comorbid PD ^a	-.06	.08	.59	.441	.94	.81–1.09
Depression severity × comorbid PD	.02	.02	.84	.361	1.01	.98–1.05
Psychotherapy sessions						
Intercept	2.77	.06	2,083.60	<.001	15.99	14.20–18.01
Depression severity	.01	.01	.87	.350	1.01	.99–1.04
Comorbid PD ^a	.31	.08	16.30	<.001	1.36	1.17–1.57
Depression severity × comorbid PD	.02	.02	1.14	.285	1.02	.99–1.05
Supportive visits						
Intercept	2.07	.09	557.67	<.001	7.95	6.70–9.45
Depression severity	.06	.02	10.08	.002	1.06	1.02–1.10
Comorbid PD ^a	.19	.11	2.86	.091	1.20	.97–1.49
Depression severity × comorbid PD	.18	.02	.60	.439	1.02	.97–1.07
Crisis visits						
Intercept	-.39	.42	.85	.356	.68	.30–1.54
Depression severity	.15	.08	3.59	.058	1.16	1.00–1.35
Comorbid PD ^a	-.17	.52	.11	.746	.84	.30–2.35
Depression severity × comorbid PD	-.08	.12	.43	.511	.92	.73–1.17

^a Reference groups: comorbid personality disorder, 0.5; no comorbid personality disorder, -0.5.

personality disorders was not associated with more (or fewer) supportive and crisis visits, and there were no significant baseline depression severity × personality disorder status interactions.

DISCUSSION

The present retrospective observational study investigated whether, and to what extent, comorbid personality disorders were associated with the amount and type of treatment for depression received by 1,455 outpatients. Our main finding was that individuals with depression and comorbid personality disorders received more psychotherapy sessions and therefore more intensive treatment than individuals without comorbid personality disorders, irrespective of initial depression severity. In addition, there was no difference in the number of pharmacotherapy sessions, supportive visits, and crisis visits for depression between individuals with and without comorbid personality disorders.

The finding that individuals with depression and comorbid personality disorders received more psychotherapy sessions for depression was not in line with our hypothesis or with previous findings. Earlier studies (20, 21) have indicated that individuals with depression and comorbid personality disorders are less likely to receive evidence-based treatments, including ECT and antidepressant medication, than individuals without comorbid personality disorders. However, recent evidence addressing this issue is lacking. The results of our study also conflict with international and national evidence-based guidelines. These guidelines (22–24) demonstrate a consensus that treatment strategies for depression should not vary based on depression subtypes or individual characteristics, such as comorbid personality

disorders. One exception is depression severity: antidepressant medication (with or without psychotherapy) is considered to be the treatment of choice for individuals with severe depression, although the guidelines are not entirely consistent (31). In the present study, the total number of pharmacotherapy sessions was indeed related to baseline depression severity.

The finding that individuals with comorbid personality disorders did not receive more supportive and crisis visits did not align with our secondary hypothesis. Because individuals with comorbid personality disorders can have more interpersonal problems (32), lower social functioning (33), and sui-

cidality (34), we expected them to receive a greater number of supportive and crisis visits. However, one could speculate that these visits were prevented by or handled within the greater number of psychotherapy sessions.

A key question underlying our findings is whether the greater number of psychotherapy sessions is either a required solution for a complex clinical problem or an unnecessary intervention that adds to overtreatment of depression (35). The complexity of individuals with comorbid personality disorders in this sample was reflected by their initial clinical presentation with more depressive and anxiety symptoms, longer episode duration, and higher incidence of childhood trauma, which has been found in previous research (36–38) as well. One could speculate that therapists feel that extension of treatment increases the odds of a good outcome for patients with these complex conditions (39). This opinion is supported by a recent meta-analysis (18) that describes an association between adequate treatment duration (16–20 sessions) and treatment effects among individuals with comorbid personality disorders receiving cognitive-behavioral therapy for depression. Another possible explanation could be that the greater number of psychotherapy sessions was an indicator for the responsiveness of therapists to the comorbid personality disorder; possibly therapists were unconsciously focusing on personality disorder symptoms. In addition, the finding could also indicate that patients with comorbid personality disorders needed to receive more sessions for the process of ending the therapeutic relationship. However, despite the complex clinical problems, a significantly greater number of psychotherapy sessions could be a sign of overtreatment. First of all, the association between the number of psychotherapy sessions and a favorable outcome in depression is not significant and very small (2). In addition, although sufficient

treatment duration has been shown to be associated with better depression outcomes for individuals with comorbid personality disorders, a sufficient duration has been defined as 16–20 sessions (18). This range was exceeded in our study, which found the average length of psychotherapy to be 22 sessions. Moreover, findings from well-designed studies (16–19) with controlled treatment have not indicated that comorbid personality disorders negatively affect acute-phase depression outcomes; therefore the need for additional therapy remains questionable. Given the current problems in the timely availability of treatment in Dutch mental health care (40, 41), it is important not to unnecessarily extend therapy.

Although this study provided an interesting look at daily mental health care practice, there were some limitations. First, a substantial number of participants did not have a baseline QIDS-SR or SAPAS assessment and had to be excluded from the analyses. However, pretreatment comparisons indicated that these individuals did not significantly differ from those included in the analyses. Second, the presence of personality disorders was determined by a brief standardized clinical interview (i.e., SAPAS) (26, 27) and did not provide information on the different types of personality disorders. Validated methods, such as the Structured Clinical Interview for DSM-IV Axis II (SCID-II) (42), may be more appropriate for studying personality disorders. However, time and resources for the use of such instruments are typically not available in routine depression care. Considering this limitation, we think that the SAPAS is an appropriate compromise between the need for a validated instrument and the need for representativeness of diagnostic assessments in daily clinical practice. The third limitation concerns the lack of information on how treatment decisions were motivated; it is possible that the clinicians decided explicitly that more psychotherapy sessions were indicated because of the comorbid personality disorder status, but we cannot exclude other reasons for the positive association between the comorbidity and the number of psychotherapy sessions. The fourth limitation involves the information on the received depression treatment. Although reliable data on the type and number of depression treatment sessions were available, details were lacking, including information on the quality of the psychotherapy and pharmacotherapy and switches between different types of psychotherapy and pharmacotherapy. The fifth limitation concerns generalizability. Although our study sample came from several treatment sites within a nationwide organization in the Netherlands, these results may not be generalizable to all mental health care settings.

CONCLUSIONS

To our knowledge, this was the first retrospective observational study investigating the impact of comorbid personality disorders on the type and the amount of depression treatment received in naturalistic outpatient care. Our findings could be indicative of overtreatment of individuals with

depression and comorbid personality disorders. Further research is needed to study subgroups of patients diagnosed as having depression who are possibly undertreated or overtreated (35) and to identify indicators related to those subgroups. To achieve this goal, research should focus on the types and amounts of treatments received, as well as on the specific contents and quality of these treatments. If the goal is to effectively improve depression treatment strategies, we need to understand and evaluate our current clinical decision making more thoroughly. With this understanding, we could study and formulate suggestions on how to modify treatment decisions to allow for a more evidence-based practice.

AUTHOR AND ARTICLE INFORMATION

Department of Clinical Psychological Science, Maastricht University, Maastricht, the Netherlands (van Bronswijk, van Dijk, Peeters); Department of Psychiatry and Psychology, Maastricht University Medical Center+, Maastricht, the Netherlands (van Bronswijk); Department of Mood Disorders, PsyQ Haaglanden, the Hague, the Netherlands (van Dijk, van den Boogaard); Parnassia Psychiatric Institute, the Hague (Deen); Institute of Psychology, Leiden University, Leiden, the Netherlands (Deen); Department of Psychiatry, Radboud University Medical Center, Nijmegen, the Netherlands (Ruhé); Donders Institute for Brain, Cognition and Behavior, Radboud University, Nijmegen (Ruhé); Pro Persona Mental Healthcare, and Behavioural Science Institute, Radboud University, Nijmegen (Spijker). Send correspondence to Dr. van Bronswijk (suzanne.vanbronswijk@maastrichtuniversity.nl)

Dr. van Bronswijk and Dr. van Dijk share first authorship.

Dr. Ruhé has received speaking fees from Janssen and Lundbeck and grants from ZonMW, Horizon 2020, and Hersenstichting. The other authors report no financial relationships with commercial interests.

Received September 17, 2020; revision received January 27, 2021; accepted March 24, 2021; published online June 17, 2021.

REFERENCES

1. Cipriani A, Furukawa TA, Salanti G, et al: Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Focus Am Psychiatr Publ* 2018; 16:420–429
2. Cuijpers P, Sijbrandij M, Koole SL, et al: The efficacy of psychotherapy and pharmacotherapy in treating depressive and anxiety disorders: a meta-analysis of direct comparisons. *World Psychiatry* 2013; 12:137–148
3. Kolovos S, van Tulder MW, Cuijpers P, et al: The effect of treatment as usual on major depressive disorder: a meta-analysis. *J Affect Disord* 2017; 210:72–81
4. Rush AJ, Trivedi MH, Wisniewski SR, et al: Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am J Psychiatry* 2006; 163:1905–1917
5. Cohen ZD, DeRubeis RJ: Treatment selection in depression. *Annu Rev Clin Psychol* 2018; 14:209–236
6. Simon GE, Perlis RH: Personalized medicine for depression: can we match patients with treatments? *Am J Psychiatry* 2010; 167:1445–1455
7. Fekadu A, Wooderson SC, Markopoulou K, et al: The Maudsley Staging Method for treatment-resistant depression: prediction of longer-term outcome and persistence of symptoms. *J Clin Psychiatry* 2009; 70:952–957
8. Friedman ES, Davis LL, Zisook S, et al: Baseline depression severity as a predictor of single and combination antidepressant treatment outcome: results from the CO-MED trial. *Eur Neuropharmacol* 2012; 22:183–199

9. Berk M, Boyce P, Hamilton A, et al: Personality: distraction or driver in the diagnosis of depression. *Pers Ment Health* 2018; 12:126–130
10. Van HL, Kool M: What we do, do not, and need to know about comorbid depression and personality disorders. *Lancet Psychiatry* 2018; 5:776–778
11. Friberg O, Martinsen EW, Martinussen M, et al: Comorbidity of personality disorders in mood disorders: a meta-analytic review of 122 studies from 1988 to 2010. *J Affect Disord* 2014; 152–154:1–11
12. Zimmerman M, Rothschild L, Chelminski I: The prevalence of DSM-IV personality disorders in psychiatric outpatients. *Am J Psychiatry* 2005; 162:1911–1918
13. Newton-Howes G, Tyrer P, Johnson T: Personality disorder and the outcome of depression: meta-analysis of published studies. *Br J Psychiatry* 2006; 188:13–20
14. Newton-Howes G, Tyrer P, Johnson T, et al: Influence of personality on the outcome of treatment in depression: systematic review and meta-analysis. *J Pers Disord* 2014; 28:577–593
15. Mulder RT: Personality pathology and treatment outcome in major depression: a review. *Am J Psychiatry* 2002; 159:359–371
16. Kool S, Schoevers R, de Maat S, et al: Efficacy of pharmacotherapy in depressed patients with and without personality disorders: a systematic review and meta-analysis. *J Affect Disord* 2005; 88:269–278
17. Kavanagh BE, Ashton MM, Cowdery SP, et al: Systematic review and meta-analysis of the role of personality disorder in randomised controlled trials of pharmacological interventions for adults with mood disorders. *J Affect Disord* 2021; 279:711–721
18. Banyard H, Behn AJ, Delgadillo J: Personality disorders and their relation to treatment outcomes in cognitive behavioural therapy for depression: a systematic review and meta-analysis. *Cognit Ther Res* (Epub Jan 20, 2021)
19. van Bronswijk SC, Köster EM, Peeters FP: Effectiveness of acute-phase treatment of depression is not influenced by comorbid personality disorders: results from a meta-analysis and meta-regression. *Psychother Psychosom* 2020; 89:109–110
20. Black DW, Bell S, Hulbert J, et al: The importance of axis II in patients with major depression. A controlled study. *J Affect Disord* 1988; 14:115–122
21. Charney DS, Nelson JC, Quinlan DM: Personality traits and disorder in depression. *Am J Psychiatry* 1981; 138:1601–1604
22. Practice Guideline for the Treatment of Patients With Major Depressive Disorder, 3rd ed. Washington, DC, American Psychiatric Association, 2010. https://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf
23. Health NCCfM: Depression: the Treatment and Management of Depression in Adults (Updated Edition) Leicester, UK, British Psychological Society, 2010
24. Spijker J, Bockting C, Meeuwissen J, et al: Multidisciplinary Guideline on Depression (Third Revision): Guideline on the Diagnosis, Treatment and Guidance of Adult Patients With a Depressive Disorder [in Dutch]. Utrecht, Trimbos Instituut, 2013
25. van Dijk DA, van den Boogaard TM, Deen ML, et al: Predicting clinical course in major depressive disorder: the association between DM-TRD score and symptom severity over time in 1115 outpatients. *Depress Anxiety* 2019; 36:345–352
26. Bukh JD, Bock C, Vinberg M, et al: Clinical utility of Standardised Assessment of Personality–Abbreviated Scale (SAPAS) among patients with first episode depression. *J Affect Disord* 2010; 127:199–202
27. Moran P, Leese M, Lee T, et al: Standardised Assessment of Personality–Abbreviated Scale (SAPAS): preliminary validation of a brief screen for personality disorder. *Br J Psychiatry* 2003; 183:228–232
28. Gorwood P, Rouillon F, Even C, et al: Treatment response in major depression: effects of personality dysfunction and prior depression. *Br J Psychiatry* 2010; 196:139–142
29. Rush AJ, Trivedi MH, Ibrahim HM, et al: The 16-Item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a psychometric evaluation in patients with chronic major depression. *Biol Psychiatry* 2003; 54:573–583
30. Kraemer HC, Blasey CM: Centring in regression analyses: a strategy to prevent errors in statistical inference. *Int J Methods Psychiatr Res* 2004; 13:141–151
31. Zimmerman M, Morgan TA, Stanton K: The severity of psychiatric disorders. *World Psychiatry* 2018; 17:258–275
32. Markowitz JC, Skodol AE, Petkova E, et al: Longitudinal effects of personality disorders on psychosocial functioning of patients with major depressive disorder. *J Clin Psychiatry* 2007; 68:186–193
33. Newton-Howes G, Mulder R, Ellis PM, et al: Predictive utility of personality disorder in depression: comparison of outcomes and taxonomic approach. *J Pers Disord* 2018; 32:513–526
34. Oquendo MA, Galfalvy H, Russo S, et al: Prospective study of clinical predictors of suicidal acts after a major depressive episode in patients with major depressive disorder or bipolar disorder. *Am J Psychiatry* 2004; 161:1433–1441
35. Lorenzo-Luaces L: Heterogeneity in the prognosis of major depression: from the common cold to a highly debilitating and recurrent illness. *Epidemiol Psychiatr Sci* 2015; 24:466–472
36. Kelly BD, Nur UA, Tyrer P, et al: Impact of severity of personality disorder on the outcome of depression. *Eur Psychiatry* 2009; 24:322–326
37. van Bronswijk SC, Lemmens LHJM, Viechtbauer W, et al: The impact of personality disorder pathology on the effectiveness of cognitive therapy and interpersonal psychotherapy for major depressive disorder. *J Affect Disord* 2018; 225:530–538
38. van den Hout M, Brouwers C, Oomen J: Clinically diagnosed axis II co-morbidity and the short term outcome of CBT for axis I disorders. *Clin Psychol Psychother* 2006; 13:56–63
39. Delgadillo J, Gellatly J, Stephenson-Bellwood S: Decision making in stepped care: how do therapists decide whether to prolong treatment or not? *Behav Cogn Psychother* 2015; 43:328–341
40. Questions and Answers: Waiting Times in Mental Health Care [in Dutch]. Amersfoort, the Netherlands, GGZNederland, 2018. https://www.denederlandseggz.nl/getmedia/ac42b1d4-30d4-444e-a007-492b4d8f3967/denederlandseggz_q-awachttijden_nov2018.pdf
41. Factsheet: Waiting Time Information on Mental Health Care [in Dutch]. Zeist, the Netherlands, Vektis, 2021. <https://www.vektis.nl/intelligence/publicaties/factsheet-wachttijdinformatie-ggz>. Accessed April 28, 2021
42. First MB, Gibbon M, Spitzer RL, et al: Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II). Washington, DC, American Psychiatric Association, 1997