

COVID-19 Post-Vaccination Depression in Older Israeli Adults: The Role of Negative World Assumptions

Lee Greenblatt-Kimron¹<https://orcid.org/0000-0003-1039-3736>, Yaakov Hoffman²

<https://orcid.org/0000-0001-5158-012X>, Menachem Ben-Ezra¹ [https://orcid.org/0000-0002-](https://orcid.org/0000-0002-7890-2069)

[7890-2069](https://orcid.org/0000-0002-2407-7016), Robin Goodwin³ <https://orcid.org/0000-0002-2407-7016>, & Yuval Palgi⁴

<https://orcid.org/0000-0002-8675-5513>

¹ School of Social Work, Ariel University, Ariel 40700, Israel

² Interdisciplinary Department, Bar-Ilan University, Ramat-Gan, Israel

³ Department of Psychology, Warwick University, Coventry, UK

⁴ Department of Gerontology, University of Haifa, Haifa, Israel

Corresponding Author:

Lee Greenblatt-Kimron, School of Social Work, Ariel University, Ariel, Israel

Mail: leeegr@ariel.ac.il

Phone: 972508534354

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

COVID-19 Post-Vaccination Depression in Older Israeli Adults: The Role of Negative

World Assumptions

Abstract

Background: With the outbreak of SARS-CoV-2 pandemic, the aging population has been shown to be highly vulnerable. As a result, policy makers and the media urged older adults to restrict social interactions, placing them at greater risk of mental health problems, such as depression. However, there has been little previous attempt to examine COVID-19 vaccine related risk factors and depressive symptoms amongst older adults.

Methods: Participants (938 older adults, $Mage=68.99$, $SD=3.41$, range 65-85) answered an online questionnaire at the start of the COVID-19 vaccination program in Israel. Participants completed measures of background characteristics, world assumptions, COVID-19 vaccine related variables, and symptoms of depression.

Results: Univariate logistic regression revealed that more negative world assumptions were linked with clinical depression levels.

Conclusions: Older adults in our sample were susceptible to unique factors associated with clinical depression influenced by their world assumptions during their COVID-19 vaccination. The high level of depression following vaccination indicates that it may take time to recover from depression associated with pandemic distress. Cognitive interventions that focus on world assumptions are recommended.

Keywords

Depression; COVID-19 Vaccine; Word Assumptions; Older Adults

COVID-19 Post-Vaccination Depression in Older Israeli Adults: The Role of Negative World Assumptions

1. Introduction

With the outbreak of the SARS-CoV-2 pandemic, it was postulated that the pandemic be considered a (traumatic) stressor (Shevlin et al., 2020). Old age was confirmed as a risk for COVID-19 complications (Palgi et al., 2020); therefore, policy makers and media urged older adults to restrict social interactions (Ayalon, 2020). Consequently, evidence from France, Italy, and Spain, the first countries outside Asia affected by the virus, revealed increased depressive feelings during the lockdown in the older population in April 2020, approximately one month after the beginning of the first lockdown (Arpino et al., 2021). Likewise, older adults in Israel reported peritraumatic distress (Greenblatt-Kimron et al., 2021) and depressive symptoms during COVID-19 (Palgi et al., 2020) in the initial stage of the lockdown in Israel between March 15th and April 1st 2020. Despite vaccination programs (potentially) "signaling the light at the end of the pandemic tunnel" (Hoffman et al., 2021a), research reported strong links tying vaccination side effects with depression and high prevalence of vaccination hesitancy with anxiety, depression, peritraumatic distress (Palgi, et al., 2021a) and post-traumatic stress disorder even in those who were vaccinated (Palgi, et al., 2021b), suggesting that many older adults continue to be psychologically impacted. However, the association between psychological health and postvaccination depression in older adults has been little explored.

Depression is a prevalent mental disorder among older adults, which commonly causes substantial distress and reduced quality of life (Kok & Reynolds, 2017; Overvliet et al., 2021). The association found between the pandemic and depression in older adults (Arpino et al., 2021;

Palgi et al., 2020) can result from appraisals of traumatic events crucial to emotional responses and consequent psychological reaction (Lazarus, 1993). Cognitive processes have been found to maintain post-traumatic stress symptoms and depression in old age (Greenblatt-Kimron & Cohen, 2020). Beck's (2008) cognitive Developmental Model of Depression suggests that depressive symptoms manifest from negative interpretations of life events, which are rooted in maladjusted beliefs concerning the self, others and the world developed from early life experiences. In line with Beck's (2008) model, a study among older care givers found negative thought patterns to be associated with postvaccination depression (Segerstrom et al., 2008). One possible direction is that one may have adopted negative world assumptions (Janoff- Bulman, 1992), which are cognitive schemas that create a person's assumptive world regarding the self, others and the world. According to Janoff- Bulman (1992) these assumptions are inherently optimistic; nevertheless, they may be shattered by stressful life events, in this case, the COVID-19 global pandemic. People may then attempt to rebuild their assumptive worlds by incorporating the stressful experience (Janoff- Bulman, 1992), as demonstrated in a recent study among older adults (Greenblatt-Kimron, 2021). Research in Poland on the COVID-19 pandemic found an association between a positive world view and fewer panic thoughts and emotions evoked by the ongoing pandemic (Trzebiński et al, 2020). Moreover, older adults with internalized negative attitudes such as an older subjective age identity showed more adverse effects of loneliness during the COVID-19 pandemic (Shrira et al., 2021).

The rapid production of COVID vaccines may have likely caused mistrust about the safety, credibility and effectiveness of the vaccines that was enhanced by anti-vaccination movements (Shacham et al., 2021). Indeed, preliminary results in Israel indicate that vaccine hesitancy was widespread in Israel with negative attitudes towards the COVID-19 more prevalent than attitudes

towards general vaccines (Shacham et al., 2021). Therefore, it is assumed that alongside the hope that the vaccination project brought with it, many negative psychological reactions may have been elicited by the vaccination given to participants in this study (i.e., Pfizer-BioNTech), as such vaccinations adopted a new technology that was not used in the past.

Based on the above, and in conjunction with a preliminary study reporting high depression levels among Israelis vaccinated for COVID-19 (Palgi et al., 2021a), the current study aimed at understanding the cognitive processes associated with clinical depression among vaccinated older adults, with a particular focus on negative schemas, specifically negative world views. We predict that negative world assumptions will be associated with depressive symptomology in vaccinated older adults.

2. Methods

2.1. Participants and Procedure

Data was collected among older Jewish adults across Israel after Israel initiated its COVID-19 vaccine program. We employed a survey company using a web-based platform with representative proportional sampling, with data collected between January 25 to February 4, 2021. On the final day of data collection Israel had completed more than 5.1 million vaccine doses, out of which about 1.9 million Israeli's had received the second dose, and 84% of those aged 60+ had received at least one of the two Pfizer doses (Israeli Ministry of Health). After omitting 42 participants who were not vaccinated, and 38 participants who did not report depressive symptom levels, the remaining 938 older adults had all received at least one dose of the COVID-19 vaccine (mean age =68.99, $SD=3.41$, range 65-85). Most participants were women ($n=562$, 59.9%), married/with a partner ($n=709$, 75.6%), and had tertiary education and

above ($n=725$, 77.3%). All participants provided informed consent. Ethical approval was granted by the Ariel University Institutional Review Board (ref no. AU-SOC-MBE-20191029-2).

2.2. Measures

Background characteristics included age, sex, marital-status (1=not married, 2=married or living in cohabitant), level of religiosity, and education (two levels 1= up to secondary education, 2=tertiary education and above).

World assumptions were measured by four items developed for this study, based on the items suggested by (Blevins et al, 2016). These items evaluated effects of shattered assumptions, which according to Janoff- Bulman (1992) may be due to stressful life events (i.e., the global pandemic); these world assumptions items state: (*I feel that my life has no meaning or purpose, I fear a negative event that may occur in the future, I feel like I have less control over my life, the world is a dangerous place to live in*). Participants rated their basic assumptions regarding the world, other people and self since the initiation of the COVID-19 vaccine program in Israel. Responses were rated on a five- point Likert scale ranging from 1 (*not at all*) to 5 (*to a very great degree*). A higher score indicates more negative world assumption. Cronbach's α in the current study was 0.810.

Vaccination measures included the number of days since the first vaccine and vaccine hesitancy (8 adapted items taken from Giambi et al., 2018). Cronbach's α in the current study was 0.832.

Depressive symptoms were measured with a 9-items scale (PHQ-9, Kroenke & Spitzer, 2002), which indicates clinically significant depression with good reliability and construct validity (Kroenke & Spitzer, 2002). Participants rated how often they had been bothered by any of the given problems since the initiation of the COVID-19 vaccine program on a four-point

Likert scale from 1 (*not at all*) to 4 (*nearly every day*). Scores range from 0 to 28 and are categorized from no depression to mild (0-9) or moderate-to-severe depression (10 and above).

Cronbach's α in the current study was 0.846

2.3. Data analysis

A univariate logistic regression analysis was conducted using the IBM statistic package (SPSS-25) software to predict depression (See Table 1). We controlled for age, sex, marital status, level of religiosity, and level of education, subjective health, COVID-19 related exposure, and added vaccine hesitancy, days since vaccination and world assumptions to the equation. Odds Ratio (OR) with 95% CI indicated the independent association of each correlate with each outcome.

3. Results

As shown in Table 1, 103 (11.0%) of the participants reached the criteria for clinical depression after being vaccinated with at least one dose of a COVID-19 vaccine. Logistic regression analysis was employed to predict the probability that older adults would reach clinical depression levels. The full model containing all predictors was statistically significant $\chi^2(10, N=935) = 197.20, p < .001$. The model as a whole explained between 19% (Cox and Snell R square) and 38.0% (Nagelkerke R squared) of the variance in clinical depression.

A univariate logistic regression analysis indicated that females had a 3.6 times higher chance of suffering from clinical depression than males after receiving one dose of the COVID-19 vaccine. Better subjective health was associated with 50% lower chance of experiencing depressive symptoms. Findings for all other covariates were non-significant (all $p > 0.05$). Each score on the independent variable of more negative world assumptions rendered one 4 times

more likely to have clinical depression ($OR=3.98$, 95%, CI: 2.99-5.28). For further information see Table 1.

4. Discussion

The current study showed an association between negative world assumptions and clinically relevant depressive symptoms amongst older vaccinated adults. To the best of our knowledge, this study is the first study to find such an association amongst older vaccinated adults during the coronavirus vaccine rollout during the pandemic.

Those who reported negative world assumptions reported 4.4 times higher odds for clinical symptoms of depression than those with more positive assumptions. Beck's Developmental Model of Depression (Beck, 2008) claims that depressive symptoms emerge from negative interpretations of life events (i.e., the COVID-19 pandemic). These are rooted in maladjusted beliefs concerning the self, others and the world, developed from early life experiences and may have persisted latently until being triggered by the coronavirus pandemic. These are then manifested in negative thoughts that augment depressive symptoms even after receiving the COVID-19 vaccination. Older adults may have also internalized the widespread negative attitudes towards older people during the COVID-19 pandemic (see Ayalon, 2020), which may have affected their world assumptions, or triggered latent negative cognitions. The COVID-19 pandemic has been considered a robust stressor (Shevlin et al., 2020). Even after receiving a vaccination that is supposed to restore the world to being the "safe" place, older adults who hold negative world assumptions may be unable to perceive the world as secure and feel protected from the virus; and therefore, experienced depression (11.0%). Such older adults may have adopted a negative world view of "what-was-will-be" (see Hoffman et al., 2021b)

common after widespread stressors. In line with the suggestion of Hoffman et al. (2021b), this may express a view that equilibrium may take time to reconstruct, despite the initiation of vaccine programs.

We also note that health awareness increased during the COVID-19 pandemic (Shacham et al., 2021). Attitudes towards the COVID-19 vaccination were more negative compared to attitudes regarding other vaccines (Shacham et al., 2021). It may thus be that more negative world assumptions may have resulted in a lack of faith in the vaccine. The rollout of the Pfizer COVID-19 vaccine was in its nascent stage during our data collection, with conspiracies and mistrust ripe about the short amount of time for vaccination development and long-term safety, credibility and effectiveness of COVID-19 vaccines (Shacham et al., 2021). While the vaccine should have augured hope, negative cognitions may have elicited feelings of helplessness and more wariness of vaccinations and; therefore, these negative cognitions were linked with depression.

Despite the above, our current study had several limitations. First, we used a cross-sectional design making it difficult to determine causality. Therefore, the receipt of the vaccine may only be significant at this particular timepoint (vs. influencing negative world views). Longitudinal studies that examine the associations between pre-existing world assumptions and depression before and after vaccination are recommended. We used a self-report design and lacked a psychiatric evaluation of participants. Finally, we used a web-based sample, which may be biased towards older adults with digital literacy (Nimrod, 2018), and may also underrepresent minor, but sizable, groups of Israeli older adults (e.g., Israeli Arabs, immigrants from the former Soviet Union).

Nevertheless, the associations found in the current study between negative world assumptions with depressive symptoms among older adults have important implications for practitioners during the global implementation of a COVID-19 vaccine. On a practical level, the knowledge gained from this study may be utilized to build appropriate treatment interventions based on a cognitive approach directed at processing clients' world assumptions, and in this way increase a sense of belief in self, in others and in the world. More positive world assumptions could decrease depressive symptomology in older adults during global pandemic vaccination programs, with these are critical for global success in combating the virus.

Acknowledgments: None.

Financial Support: This work was supported by the Ariel University internal grant

RA2000000522

Conflict of Interest: None.

Ethical Standards: The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Arpino B, Pasqualini M, Bordone V, Solé-Auró A** (2021). Older people's nonphysical contacts and depression during the COVID-19 lockdown. *The Gerontologist*, 61(2), 176-186.
- Ayalon L** (2020). There is nothing new under the sun: Ageism and intergenerational tension in the age of the COVID-19 outbreak. *International Psychogeriatrics*, 32(10), 1221-1224.
- Beck AT** (2008). The evolution of the cognitive model of depression and its neurobiological correlates. *American Journal of Psychiatry*, 165(8), 969-977.
- Blevins CE, Wusik MF, Sullivan CP, Jones RT, Hughes M** (2016). Do negative changes in worldview mediate links between mass trauma and reckless behavior? A longitudinal exploratory study. *Community Mental Health Journal*, 52(1), 10-17.
- Giambi C, Fabiani M, D'Ancona F, Ferrara L, Fiacchini D, Gallo T, Martinelli D, Pascucci MA, Prato R, Filia A, Bella A, Del Manso M, Rizzo C, Rota MC** (2018). Parental vaccine hesitancy in Italy—results from a national survey. *Vaccine*, 36, 779-87.
- Greenblatt- Kimron L** (2021). World assumptions and post- traumatic growth among older adults: The case of Holocaust survivors. *Stress and Health*, 37(2), 353-363.
- Greenblatt-Kimron L, Cohen M** (2020). The role of cognitive processing in the relationship of posttraumatic stress symptoms and depression among older Holocaust survivors: A moderated-mediation model. *Anxiety, Stress, & Coping*, 33, 59-74.
- Greenblatt-Kimron L, Ring L, Hoffman Y, Shrira A, Bodner E, Palgi Y** (2021). Subjective accelerated aging moderates the association between COVID-19 health worries and peritraumatic distress among older adults. *Global Mental Health*, 1-30.

- Hoffman YS, Hamama-Raz Y, Ben-Ezra M, Lavenda, O** (2021b). A Dissociation Between Two Cognitive Biases in Acute Stress Disorder: Preliminary Evidence for the Reverse Gambling Fallacy. *Cognitive Therapy and Research*, 1-12.
- Hoffman Y, Palgi Y, Goodwin R, Ben-Ezra M, & Greenblatt-Kimron L.** (2021a). A storm in a teacup: older adults' low prevalence of COVID-19 vaccine side-effects and their link with vaccination anxiety. *International Psychogeriatrics*, 1-3.
- Janoff-Bulman R** (1992). *Shattered assumptions*. The Free Press.
- Kok RM, Reynolds CF** (2017). Management of depression in older adults: a review. *JAMA*, 317(20), 2114-2122.
- Kroenke K, Spitzer RL** (2002). The PHQ-9: a new depression and diagnostic severity measure. *Psychiatric Annals*, 32, 509–521.
- Lazarus RS** (1993). Coping theory and research: Past, present, and future. *Psychosomatic Medicine*, 55, 234–247.
- Ministry of Health** (2021). COVID-19 in Israel. Retrieved from https://www.gov.il/BlobFolder/reports/daily-report-20210204/he/daily-report_daily-report-20210204.pdf
- Nimrod G** (2018). Technophobia among older Internet users. *Educational Gerontology*, 44(2-3), 148-162.
- Overvliet GM, Jansen RA, van Balkom AJ, van Campen DC, Oudega ML, Van Der Werf YD, van Exel E, van den Heuvel OA, Dols A** (2021). Adverse events of repetitive transcranial magnetic stimulation in older adults with depression, a systematic review of the literature. *International Journal of Geriatric Psychiatry*, 36(3), 383-392.

- Palgi Y, Bergman YS, Ben-David B, Bodner E** (2021a). No psychological vaccination: Vaccine hesitancy is associated with negative psychiatric outcomes among Israelis who received COVID-19 vaccination. *Journal of Affective Disorders*, 287, 352-353.
- Palgi Y, Greenblatt-Kimron L, Hoffman Y, Goodwin R, & Ben-Ezra M** (2021b). Factors associated with current posttraumatic stress disorder among COVID-19 vaccinated older adults in Israel. *Journal of Psychiatric Research*, 142, 272-274.
- Palgi Y, Shrira A, Ring L, Bodner E, Avidor S, Bergman Y, Cohen-Fridel S, Keisari S, Hoffman Y** (2020). The loneliness pandemic: loneliness and other concomitants of depression, anxiety and their comorbidity during the COVID-19 outbreak. *Journal of Affective Disorders*, 275, 109–111.
- Segerstrom SC, Schipper LJ, Greenberg RN** (2008). Caregiving, repetitive thought, and immune response to vaccination in older adults. *Brain, Behavior, and Immunity*, 22(5), 744–752.
- Shacham M, Greenblatt-Kimron L, Hamama-Raz Y, Martin LR, Peleg O, Ben-Ezra M, Mijiritsky E** (2021). Increased COVID-19 Vaccination Hesitancy and Health Awareness amid COVID-19 Vaccinations Programs in Israel. *International Journal of Environmental Research and Public Health*, 18(7), 3804.
- Shevlin M, McBride O, Murphy J, Miller JG, Hartman TK, Levita L, Mason L, Martinez AP, McKay R, Stocks TVA, Bennett KM, Hyland P, Karatzias T, Bentall RP** (2020). Anxiety, depression, traumatic stress and COVID-19-related anxiety in the UK general population during the COVID-19 pandemic. *BJPsych Open*, 6(6), e125, 1-9.

Shrira A, Hoffman Y, Bodner, E, & Palgi Y (2020). COVID-19-related loneliness and psychiatric symptoms among older adults: the buffering role of subjective age. *The American Journal of Geriatric Psychiatry*, 28(11), 1200-1204.

Trzebiński J, Cabański M, Czarnecka JZ (2020). Reaction to the COVID-19 pandemic: the influence of meaning in life, life satisfaction, and assumptions on world orderliness and positivity. *Journal of Loss and Trauma*, 25(6-7), 544-557.

Table 1. Logistic Regression Analyses Predicting Likelihood of Clinical Depression

	B	S.E.	Odds Ratio	95.0% C.I. for Odds Ratio	
				Lower	Upper
Age	.01	.04	1.01	.93	1.09
Sex ^a	1.30	.32	3.67***	1.97	6.82
Marital status ^b	-.19	.19	.83	.58	1.20
Education ^c	-.62	.30	.94	.52	1.70
Religiosity	.06	.29	1.06	.60	1.87
Subjective health	-.55	.14	.58***	.44	.76
COVID-19 related exposure	.19	.25	1.21	.74	1.98
Vaccine hesitancy	.17	.16	1.18	.86	1.63
Days since vaccine	-.00	.01	1.00	.97	1.03
World Assumptions	1.38	.15	3.98***	3.00	5.29
Constant	-6.20	2.99	.00*		

Note: Total N=938; Nagelkerke R^2 =.35, clinical depression=103, negative depression= 831.

^aSex 1= male, 2=female. ^bMarital status 1=not married, 2= currently married, or living with a partner. ^c= two education levels 1=up to secondary education, 2=tertiary education and above.

* $p < .05$ *** $p < .001$.