



Influence of Material Deprivation on Clinical Outcomes Among People Living with HIV in High-Income Countries: A Systematic Review and Meta-analysis

Vasiliki Papageorgiou¹ · Bethan Davies² · Emily Cooper¹ · Ariana Singer³ · Helen Ward¹

Accepted: 27 November 2021
© The Author(s) 2021

Abstract

Despite developments in HIV treatment and care, disparities persist with some not fully benefiting from improvements in the HIV care continuum. We conducted a systematic review to explore associations between social determinants and HIV treatment outcomes (viral suppression and treatment adherence) in high-income countries. A random effects meta-analysis was performed where there were consistent measurements of exposures. We identified 83 observational studies eligible for inclusion. Social determinants linked to material deprivation were identified as education, employment, food security, housing, income, poverty/deprivation, socioeconomic status/position, and social class; however, their measurement and definition varied across studies. Our review suggests a social gradient of health persists in the HIV care continuum; people living with HIV who reported material deprivation were less likely to be virologically suppressed or adherent to antiretrovirals. Future research should use an ecosocial approach to explore these interactions across the lifecourse to help propose a causal pathway.

Keywords Antiretroviral therapy · HIV · Meta-analysis · Social determinants of health · Socioeconomic factors · Systematic review · Viral suppression

Resumen

A pesar de los avances en el tratamiento y la atención al VIH, ciertas desigualdades persisten en estas áreas. Como resultado, algunas personas que viven con el VIH no se benefician plenamente de las mejoras en la atención del VIH a todos los niveles. Realizamos una revisión sistemática para explorar las asociaciones entre los determinantes sociales de la salud y los resultados del tratamiento del VIH (supresión viral y tratamiento de adherencia) en países de ingresos altos. Completamos un metaanálisis siguiendo un modelo de efectos aleatorios e incluyendo medidas consistentes de exposiciones. Identificamos 83 estudios observacionales elegibles para inclusión. Los determinantes sociales vinculados a las circunstancias materiales (privación) se identificaron como educación, empleo, seguridad alimentaria, vivienda, ingresos, pobreza / privación, situación / posición socioeconómica y clase social; sin embargo, su medición y definición variaron entre los estudios. Nuestro estudio sugiere que un gradiente de salud social persiste en la atención sanitaria del VIH a todos los niveles; las personas que viven con el VIH que declaran privación material tenían menos probabilidades de alcanzar la supresión viral o de ser adherente a los antirretrovirales. Futuras investigaciones deberían utilizar un enfoque ecosocial para explorar estas interacciones a lo largo del curso de la vida para ayudar a proponer una vía causal.

✉ Vasiliki Papageorgiou
vasiliki.papageorgiou17@imperial.ac.uk

¹ Patient Experience Research Centre, School of Public Health, Imperial College London, London, UK

² Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

³ School of Public Health, Imperial College London, London, UK

Abbreviations

aHR	Adjusted Hazard Ratio
aOR	Adjusted Odds Ratio
aPR	Adjusted Prevalence Ratio
aRR	Adjusted Relative Risk
ART	Antiretroviral therapy/treatment
cART	Combination antiretroviral therapy
CI	Confidence interval
CSDH	WHO Commission on Social Determinants of Health

EPICES	Evaluation of Deprivation and Inequalities in Health Examination Centres
FPL	Federal Poverty Level
HAART	Highly active antiretroviral therapy
HFSSM	Household Food Security Survey Module
MEMS	Medication Event Monitoring Systems
NICE	National Institute for Health and Care Excellence (UK)
OECD	Organisation for Economic Co-operation and Development
SEP/SES	Socioeconomic position/status
VL	Viral load

Introduction

The social determinants of health describe the conditions in which a person is “born, grows, lives, works and ages” and operate alongside social hierarchy, socioeconomic position, power differentials and the wider socioeconomic and political context [1]. Less privileged and marginalised individuals experience poorer health outcomes than the general population which can, in part, be attributed to current and historical struggles with structural racism, discrimination, recurring inequalities and social exclusion [1–4]. Recent evidence suggests that a decade of government austerity has led to widening health, social and economic inequalities in countries, such as England, that have resulted in stalling life expectancy, declining social mobility and increased food insecurity [5, 6].

Inequalities persist among people living with HIV globally despite advancements in life expectancy, treatment and care. Some groups are disproportionately affected including migrants, the homeless and sex workers [7–11]. These disparities are underpinned by social and structural factors including gender, sexual orientation, ethnicity, racism and socioeconomic position which influence an individual’s agency and power within a specific context [12, 13]. A literature review by Burch et al. [14] reported that people living with HIV who had poorer socioeconomic status (SES) were more likely to have poorer virological and immunological responses to antiretroviral therapy (ART). The authors defined SES according to material factors (e.g., education, neighbourhood socioeconomic position) and the health system (e.g., health insurance) itself [1, 14]. In countries without universal healthcare systems, such as the US, publicly funded systems of care exist to support individuals who are uninsured access healthcare; for instance, the Ryan White HIV/AIDS Program [15, 16]. However, the funding, accessibility and eligibility requirements of these services are influenced by structural determinants including the political systems, structures, policies and leadership in which they exist and operate.

For people living with HIV, adhering to prescribed antiretrovirals is essential to maintain virological suppression and to reduce the risk of drug resistance [17, 18]. Early research [19] estimated that HIV-1 viral load (VL) can be reduced by approximately 99% within two weeks of treatment initiation (using protease inhibitors and reverse transcriptase inhibitors). The level of ART adherence required to reach viral suppression is now considered to be regimen-dependent and could be as low as 75% for some [20]. Successful treatment and viral suppression are the second and third UNAIDS 90:90:90 targets; several high-income countries including the UK, Denmark and the Netherlands met these targets before the deadline of 2020 [21, 22]. Referral to, and retention in, HIV care services is therefore critical. However, this is dependent on early diagnosis which is not always achieved; in the UK in 2019, it was estimated that approximately 42% of people living with HIV were diagnosed late, defined by a CD4 count at diagnosis of < 350 cells/mm³ [23, 24]. The percentage of late diagnoses varied according to age, ethnicity and mode of transmission, with the highest proportion (52%) among heterosexual men [23, 24].

Social determinants exist within complex, intersectoral systems, can be highly correlated and are driven by the context in which they are created and manifest [1, 25]. We build on the definition of social determinants by the WHO Commission on Social Determinants of Health (CSDH) framework to include the “intermediary determinants” of living circumstances, working conditions and food availability [1]. We frame our findings in relation to Krieger’s ecosocial theory which aims to understand how “health inequities constitute biological expressions of injustice” across societal and ecosystem levels, pathways and power differentials [13, 26–28]. Ecosocial theory attempts to unpick the complexity of interactions across ecologies; for instance, the scale of phenomena (including measured and unmeasured factors), how these are organised (hierarchies) and spatiotemporal dynamics which means they are restricted by the extent to which these have been previously theorised, conceptualised, inferred and explored [27]. Much like ecosocial theory, the social production of disease and/or political economy of health as well as psychosocial theory can help elucidate how and why diseases are unevenly distributed across societies as well as implications for action [27]. We focus on factors of material deprivation, which could be targeted by social and public health policies.

Our aim is to synthesise the evidence and identify the social determinants that have an impact on HIV treatment outcomes (specifically viral suppression, ART adherence) among people living with HIV in high-income countries.

Methods

We conducted a systematic review using the PRISMA 2020 checklist (Additional file 1); a full protocol is published on PROSPERO (identification number: CRD42020171850) [29, 30]. We adapted the approach of Burch et al. [14] to provide a more recent examination of the association between social determinants and HIV treatment outcomes; however, we focus on observational studies (cohorts and cross-sectional studies), rather than randomised controlled trials, as these replicate real-world settings.

Search Strategy and Selection Criteria

We searched MEDLINE, EMBASE, Global Health, HMIC, Cochrane Library, CINAHL, Web of Science, ProQuest and Scopus databases from date of creation (or first stored record) to 13 January 2020 using a search strategy developed with a University librarian. We also hand searched conference databases until March 2020 and searched the reference lists of relevant review articles and editorials. Further detail, including search strategies, are provided in Additional file 2.

Studies were assessed for eligibility using the criteria detailed in Additional file 3. To be included, study populations had to comprise adults (aged 18 or older) living with HIV in high-income countries. We defined high-income countries using the 2019 World Bank classification and Organisation for Economic Co-operation and Development (OECD) country membership [31, 32]. Social determinants focussed on measures of material deprivation and were broadly defined as education, employment, food security, housing, income, poverty (or deprivation), socioeconomic status (or position) and social class. They had to be compared across levels and recorded at either the individual, household, or neighbourhood level. The primary outcomes of interest were HIV treatment-related, specifically medication adherence and viral suppression, measured by VL or CD4 cell counts. We also extracted data of other social factors which may act as confounders, specifically age, gender, sexual orientation, ethnicity and migration status.

Screening, Data Collection and Analysis

Two authors (VP, AS) screened title and abstracts, followed by full-text, using Covidence [33] and original study authors were contacted by VP to provide any unavailable full-text articles. If the search identified a non-peer-reviewed and peer-reviewed publication for the same study, the most recently published was included. Selected studies were then exported and managed using Excel with a data extraction table initially piloted among 10 studies and subsequently

refined. VP extracted data items (Additional file 4) which were cross-checked by EC.

Forest plots were created using Revman 5 [34] for each social determinant and used to present relative effect sizes of comparable associations [adjusted odds ratios (aOR) with 95% confidence intervals (CIs)]. Data are mainly presented using a narrative synthesis as there were large amounts of heterogeneity between included studies. A random-effects meta-analysis was performed for studies where definitions and measurements across studies were consistent; we present the associated I^2 value for heterogeneity (proportion of variation in effect estimates due to heterogeneity rather than chance) [35]. Data presented in forest plots compare poorer social determinants (e.g., unstable housing) to improved determinants (e.g., stable housing).

Quality Assessment

VP and BD assess the risk of bias of included studies using the Newcastle–Ottawa scale and an adapted version for cross-sectional studies (Additional file 5) [36, 37]. Studies categorised as ‘other’ study design, such as secondary data analysis, are assessed using the most appropriate quality assessment tool. We define the most important confounders for studies to adjust for as gender, sexual orientation, age, race/ethnicity and social class, based on the literature [38]. We define adequate follow-up for outcomes based on WHO guidelines of routine VL monitoring of 6 months following ART initiation and measuring ART adherence at 30 days [39, 40]. No data from ‘low quality’ studies are included in the meta-analysis; therefore, a sensitivity analysis was not conducted.

Results

We screened 4031 records, following the removal of duplicates, of which 83 observational studies were eligible (Fig. 1).

Most included studies explored the social determinants of education ($n = 52$, 62.7%); followed by housing ($n = 39$, 47.0%), employment ($n = 33$, 39.8%) and income ($n = 33$, 39.8%) (Table 1). The studies included a total of 1,445,150 people living with HIV. Almost three-quarters ($n = 61$, 73.5%) of the included studies were based in North America. Some studies focussed on specific sub-populations of people living with HIV such as transgender women ($n = 5$, 6.0%), people who formerly/currently use drugs ($n = 3$, 3.6%), people in prison ($n = 1$, 1.2%), migrants ($n = 1$, 1.2%), people living with HIV and hepatitis C ($n = 1$, 1.2%), homeless/marginally housed individuals ($n = 1$, 1.2%) as well as individuals who hold multiple identities; for instance, being in




Fig. 1 PRISMA 2020 flow diagram. Adapted from Page et al. [29]

prison and homeless (n = 1, 1.2%) or being socioeconomically disadvantaged and using drugs (n = 1, 1.2%).

We identified 38 studies that explored the association between social determinants and virological suppression (or non-suppression); 35 that investigated adherence (or non-adherence) to ART including combination ART (cART) and highly active ART (HAART); and 10 that explored both primary outcomes. Study characteristics are detailed in Table 2.

Most studies measuring viral suppression predominantly recorded VL, although some also reported the immunological response of CD4 cell count. The VL thresholds to meet viral suppression varied across studies from 20 to 400 VL copies/mL or defined as ‘detectable’ or ‘undetectable’. Adherence had more varied measurements; some studies asked how often participants had missed medication in a specific time frame or used medication event monitoring systems (MEMS) to record the number of times a pill cap was opened which was sometimes also verified by an unannounced pill count by phone. One study [41] used an index of the “number of days out of medication” (MED-OUT) using pharmacy-refill based measures.

Social determinants primarily focussed on measures of material deprivation. This included education (e.g., level, attainment, student status); employment (e.g., status, grade, type); housing (e.g., status, stability, homelessness, living situation, condition, ownership); measures of food security;

income (e.g., annual household, financial stability, receiving benefits, financial hardship/concerns, economic situation); socioeconomic status/position (individual/neighbourhood); and measures of deprivation and social class.

We present data under the following headings: education, employment, housing, and material deprivation (which includes measures of deprivation, food security, income, and socioeconomic status/position). For some studies, we present the inverse of the data reported in the original manuscript as we were interested in the outcomes of virological suppression and ART adherence. A full summary of findings, including the confounders adjusted for and reference categories, is provided in Additional file 6.

Education

Education was typically defined by the highest educational level of attainment; from primary school in the UK (or elementary school in the US) to university qualifications.

Of 83 studies, 52 focussed on education with nine studies reporting a significant adjusted association with virological suppression [43, 84, 88, 97, 103, 106, 107, 116, 119]; four reported negative associations (lower education) and five positive (higher education). Overall, people who had lower educational levels were less likely to be virologically suppressed (Fig. 2a; Additional file 6) than individuals who had

Table 1 Summary of key characteristics of included studies

Characteristic	Total studies (%)
Publication type	
Peer-reviewed (e.g., journal article, short/brief report, short/concise communication)	77 (92.8)
Not peer-reviewed (e.g., conference abstract, editorial letter, thesis)	6 (7.2)
Setting	
North America	61 (73.5)
Europe	20 (24.1)
Asia	1 (1.2)
Australia	1 (1.2)
Study design	
Cohort/longitudinal	31 (37.3)
Cross-sectional ^b	30 (36.1)
Other ^c	22 (26.5)
Social determinants ^a	
Education	52 (62.7)
Employment	33 (39.8)
Food security	7 (8.4)
Housing	39 (47.0)
Income	33 (39.8)
Poverty/deprivation	9 (10.8)
Socioeconomic status/position	4 (4.8)
Social class	1 (1.2)
Primary outcome(s)	
Viral (non-)suppression ^d	38 (45.8)
Medication (non-)adherence ^e	35 (42.2)
Both	10 (12.0)
Secondary outcome(s) ^a	
Diagnosis-related	1 (1.2)
Medication-related (e.g., initiation, use, coverage, response)	8 (9.6)
HIV care-related (e.g., engagement, retention, missed visits)	13 (15.7)

^aMore than 1 option possible

^bIncludes cross-sectional surveys of cohort studies

^cMixed methods (observational data extracted), programme evaluations, needs assessment, cross-sectional surveys/analysis of cohort studies and chart/record/baseline intervention reviews

^dCD4 cell count, HIV viral load

^eART, cART, HAART

higher educational attainment after adjusting for confounders [53, 88, 106]. For instance, in a cohort of 1246 people living with HIV attending healthcare in France, D'Almeida et al. [106] found individuals whose highest educational attainment was elementary school were 60% less likely (aOR 0.40; 95% CI 0.18, 0.90) to have a VL < 50 copies/mL, compared to those with more than 2 years of a university degree. Shacham et al. [88] found that individuals presenting at an urban HIV clinic in the US, with a high school diploma or less were over 2-times more likely (aOR 2.32; 95% CI 1.08, 5.00) to be virologically non-suppressed; this study used a higher threshold (VL < 400 copies/mL).

The direction of the association between education and medication adherence was less clear. Four studies reported

a significant adjusted negative association with medication adherence and lower educational attainment [57, 93, 103, 118]; one [93] also reported positive associations among some sub-groups (Fig. 2b; Additional file 6). Sunil and McGehee [93] found different patterns across educational levels when stratifying by race/ethnicity. White Americans (n = 992) who had completed high school, some College or had an undergraduate qualification were 30% (aOR 1.30; 95% CI 1.24, 1.37), 41% (aOR 1.41; 95% CI 1.34, 1.48) and 58% (aOR 1.58; 95% CI 1.51, 1.66) more likely, respectively, to be adherent to ART compared to individuals who completed some high school only [93]. The same general trend was seen for African American participants (n = 581): individuals who completed high school and some College

Table 2 Characteristics of included studies

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Virological (non-) suppression measure ^b	Medication ^c (non-) adherence measure
North America (n = 61)					
Almeida-Brasil et al. 2018 [42] (article)	Canada	566 adults with hepatitis C, receiving cART; part of Food Security & HIV-HCV Study (FS) of the Canadian Co-infection Cohort (CCC) (prospective cohort)	Education; employment; food security; income	VL > 50 copies/mL	Missed ≥ 1 cART dose, past 4 days
Anderson et al. 2018 [43] (article)	USA	239 women at an urban HIV specialty clinic (cross-sectional)	Education; employment	CD4 < 200 cells/mm ³ VL > 20 copies/mL	–
Baguso et al. 2019 [44] (article)	San Francisco (USA)	123 self-identified transgender women; part of Transwomen Empowered to Advance Community Health 3 (TEACH 3) study (cross-sectional)	Education; housing	Detectable and unknown viral load	–
Berg et al. 2004 [45] (article)	Bronx, New York (USA)	113 people who currently/formerly use opioids; part of HIV Epidemiologic Research on Outcomes (HERO) study (prospective cohort)	Housing	–	MEMS caps openings divided by number of prescribed doses
Blank et al. 2015 [46] (article)	Brooklyn; Chicago; Los Angeles; Miami; San Antonio; Longview; Anniston; and Chapel Hill (USA)	944 women of colour; part of Special Projects of National Significance (SPNS) Enhancing Access to and Retention in Quality HIV Care for Women of Color initiative (prospective cohort)	Education; employment; housing	VL < 200 copies/mL	–
Chitsaz et al. 2013 [47] (article)	Connecticut; Georgia; Illinois; Massachusetts; New York; Ohio; Pennsylvania; Rhode Island; South Carolina (USA)	1166 adults with a diagnosis of HIV before being jailed in 10 jails in US across 9 states; part of Enhance Link initiative (cross-sectional)	Education; employment; food security; housing	–	$\geq 95\%$ ART adherence
Clemenzi-Allen et al. 2018 [48] (brief report)	San Francisco, USA	1222 adults at HIV clinic ("Ward 86") (cross-sectional)	Housing	VL < 200 copies/mL	–

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Creasy et al. 2019 [49] (article)	Atlanta; Detroit; Houston; Memphis; Philadelphia; and Washington, DC (USA)	5143 male and Black/African American adults recruited from Black Pride events; part of Promoting Our Worth, Equality, and Resilience (POWER) study (cross-sectional)	Housing	–	Missed doses (4–7/week, 2–3/week, 1/week, <1/week, never) Last missed dose (never, > 3 months, 1–3 months, 3–4 weeks, 1–2 weeks, or within the week)
Doshi et al. 2017 [50] (article)	USA	1,296,248 adults accessing >800 medical care providers, 2010–2014; part of Health Resources and Services Administration (HRSA) and served by Ryan White program (surveillance records review)	Housing	VL < 200 copies/mL	–
Dowshen et al. 2016 [51] (short report)	USA (including Puerto Rico), 15 cities	1584 “behaviourally-infected youth” (including 66 young transgender women) from a multisite study conducted at 20 Adolescent Medicine Trials Units; part of Adolescent Trials Network (ATN) (secondary data analysis—cross-sectional)	Housing	Detectable viral load	–
Fadul et al. 2017 [52] (conference abstract)	North Carolina, USA	184 newly diagnosed people living with HIV treated in East Carolina University clinic; part of Ryan White program (retrospective chart review)	Education; housing; poverty	VL < 200 copies/mL	–
Feldman et al. 2015 [53] (article)	Greater New York metropolitan area (USA)	2896 MSM; part of Ryan White Part A programme evaluation)	Education; housing	VL > 200 copies/mL CD4 < 350 cells/mm ³	–
Feller and Agins 2016 [54] (article)	New York State (USA)	11,252 adults receiving treatment across 186 HIV clinics (classification and regression tree analysis algorithm—cohort)	Housing	VL < 200 copies/mL % virally suppressed	–

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Gardner et al. 2015 [55] (conference abstract)	Ontario (Canada)	3322 people living with HIV receiving clinical care; part of Ontario HIV Treatment Network Cohort Study (OCS) (cohort)	Income	VL < 200 copies/mL	–
Gebro et al. 2003 [56] (article)	USA	196 people living with HIV taking at least 1 ART in an urban hospital clinic (John Hopkins University HIV Clinic) (cross-sectional)	Deprivation	–	< 90% ART adherence (stratified by injecting drug use and gender)
Golin et al. 2002 [57] (article)	Southern California (USA)	117 people living with HIV in a county hospital HIV clinic with newly initiated HAART; part of Adherence and Efficacy to Protease Inhibitor Therapy (ADEPT) study (prospective cohort)	Education; employment; income	–	Number of PI/NNRTI/HAART doses taken divided by prescribed over 4-weeks
Haider et al. 2019 [58] (article)	South Carolina (USA)	342 people living with HIV receiving HIV care in immunology centre; part of Ryan White Program (cross-sectional)	Education; employment; income	VL < 100 copies/mL	–
Hussen et al. 2018 [59] (article)	Southeastern city, USA	81 young Black gay, bisexual and other MSM aged 18–24 years old in a paediatric/adolescent clinic (cross-sectional)	Education; employment; housing	VL ≤ 40 copies/mL	–
Iralu et al. 2010 [60] (article)	Navajo Nation (USA)	36 American Indians under (NAN) case management; part of Four Corners American Indian Circle of Services Collaborative (4CC) (cross-sectional)	Education; employment; housing; income	CD4 < 200 cells/mm ³ Viral load (log)	–
Johnson et al. 2003 [61] (article)	San Francisco, Los Angeles; New York City; Milwaukee, (USA)	2765 adults taking ART; part of Healthy Living Project (baseline pre-intervention—cross-sectional)	Education; employment; housing	–	90% ART adherence, past 3 days

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Kacanek et al. 2019 [62] (article)	USA (including Puerto Rico)	122 18–22-year olds based across 15 clinical sites; part of Pediatric HIV/AIDS Cohort Study (longitudinal)	Income	VL > 400 copies/mL	Self-reported missed ≥ 1 ART dose, past week
Kalichman and Grebler 2010 [63] (article)	Atlanta, Georgia (USA)	188 people living with HIV/AIDS who demonstrated poor health literacy (cross-sectional)	Education; employment; income; poverty	–	85% ART adherence 75% ART adherence
Kalichman et al. 2010 [64] (article)	Atlanta, Georgia (USA)	344 people living with HIV/AIDS (cross-sectional)	Education; employment; food security; housing; income	Undetectable viral load Change in most recent viral load Change in most recent T cells Most recent CD4 cell count	80% ART adherence 90% ART adherence Self-report adherence rating Unannounced pill count adherence
Kalichman et al. 2014 [65] (article)	Atlanta, Georgia (USA)	364 men and 157 women living with HIV (cross-sectional)	Food security	CD4 < 500 Detectable viral load	85% ART adherence
Keith McInnes et al. 2013 [41] (article)	Atlanta; Baltimore; Bronx; Brooklyn/Manhattan; Houston; Los Angeles; Pittsburgh; and Washington, DC (USA)	1871 veterans living with HIV; part of Veterans Aging Cohort Study (VACS) and completed 5th follow-up survey in 2010–2011 (cross-sectional analysis of longitudinal cohort)	Education; employment; housing; income	–	ART adherence (1-MED-OUT ≥ 0.90)
Kleeberger et al. 2004 [66] (concise communication)	USA	597 men living with HIV reporting use of HAART; part of Multicenter AIDS Cohort Study (prospective cohort)	Education; income	–	Negative change (%): visit-pairs with 100% adherence Positive change (%): visit-pairs with < 100% adherence
Koehn et al. 2020 [67] (short communication)	Vancouver (Canada)	99 Dr Peter Centre clients who completed a 12-month follow-up interview (longitudinal cohort)	Education; employment; food security; housing; income	–	$\geq 95\%$ cART adherence

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Kyser et al. 2011 [68] (article)	Denver; Minneapolis; Providence; and St. Louis (USA)	528 people living with HIV taking cART; part of study to understand the natural history of HIV/AIDS in the era of effective therapy (SUN) (cross-sectional analysis of longitudinal, prospective cohort)	Education; employment	–	cART adherence, past 3 days (missed ≥ 1 ART doses)
Lacombe-Duncan et al. 2019 [69] (article)	British Columbia, Ontario, and Quebec (Canada)	50 transgender women; part of Canadian HIV Women's Sexual and Reproductive Health Cohort Study (CHIWOS) (cross-sectional / mixed methods)	Education; housing; income	VL < 50 copies/mL	$\geq 95\%$ ART adherence
Lim et al. 2015 [70] (article)	New York City (USA)	1698 people living with HIV /AIDS with both jail incarceration and homelessness (retrospective cohort)	Housing	VL < 400 copies/mL	–
Ludema et al. 2016 [71] (article)	Bronx; Brooklyn; Washington DC; Los Angeles; San Francisco; Chicago (USA)	1481 women; part of Women's Interagency HIV Study (WIHS) (prospective cohort)	Income	VL > 200 copies/mL	–
Marshall et al. 2016 [72] (article)	Vancouver (Canada)	706 people who use drugs; part of AIDS Care Cohort to Evaluate Exposure to Survival Services (ACCESS) (prospective cohort)	Education; employment; housing	Undetectable VL or VL < 50 copies/mL	–
McCoy et al. 2016 [73] (article)	Arizona, California, Illinois, Massachusetts, Michigan, Pennsylvania, Texas, Washington, and Wisconsin (USA)	426 people living with HIV aged 50 years and older; part of PRIME baseline study (baseline pre-intervention—cross-sectional)	Education; employment; income	–	$\geq 95\%$ ART adherence
Miller et al. 2006 [74] (article)	British Columbia (Canada)	892 people living with HIV; part of HIV/AIDS Drug Treatment Program of the British Columbia Centre for Excellence in HIV/AIDS (programme evaluation)	Housing; income	VL < 500 copies/mL	–

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Mimiaga et al. 2019 [75] (article)	Rhode Island (USA)	296 adults; part of Ryan White Part B HRSA (needs assessment)	Education; employment; housing; income	Virological non-suppression	Sometimes/never take ART as prescribed, past 12 months
Mohammed et al. 2004 [76] (article)	Louisiana (USA)	273 adults from 8 areas in non-urban Louisiana (cross-sectional)	Education; employment	–	Self-report missed any HAART doses, previous week
Moore et al. 2016 [77] (article)	Vancouver (Canada)	719 MSM; part of Momentum Health study (cross-sectional)	Education; income	VL ≥ 200 copies/mL	–
Nyaku, Beer and Shu 2019 [78] (article)	USA (including Puerto Rico)	18,423 adults who self-reported taking ART; part of Medical Monitoring Project (MMP) (secondary data analysis—surveillance records review from cross-sectional sample)	Education; housing; poverty	–	ART non-persistence
O’Neil et al. 2012 [79] (article)	British Columbia (Canada)	566 people living with HIV who have accessed ART; part of Longitudinal Investigations into Supportive and Ancillary health services (LISA) cohort study (cross-sectional analysis of cohort)	Education; housing; income	–	≥ 95% HAART adherence
Oliver et al. 2019 [80] (article)	Nashville, Tennessee (USA)	248 women living with HIV with ≥ 1 prenatal visit; part of Vanderbilt Obstetrics Comprehensive Care Clinic (VCCC) (observational cohort)	Education	VL ≥ 200 copies/mL	–
Phillips 2011 [81] (article)	USA	160 Black men living with HIV/AIDS who use illicit drugs (cross-sectional)	Housing	–	Self-reported mean number of ART doses missed, past 4 days
Phillips et al. 2013 [82] (article)	Canada; USA (including Puerto Rico)	1873 adults; part of International Nursing Network for HIV Research Study V (cross-sectional)	Education	–	≥ 99% ART adherence (self-reported visual analogue scale for 30 days)
Rebeiro et al. 2018 [83] (article)	Southern USA	2541 adults in viral suppression analysis; part of VCCC (observational cohort)	Socioeconomic status/position	VL < 200 copies/mL among those with ≥ 1 clinic visit	–

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Robinson and Knowlton 2016 [84] (article)	Baltimore (USA)	383 people who inject drugs formerly/currently (secondary data analysis—cross-sectional)	Education	VL ≤ 40 copies/mL	–
Santos et al. 2014 [85] (short report)	San Francisco (USA)	314 transgender women; part of San Francisco Transfemale Respondent Driven Sampling Study (secondary data analysis—cross-sectional)	Housing	VL ≤ 200 copies/mL	–
Sayles et al. 2012 [86] (article)	Los Angeles county (USA)	11,397 people living with HIV who were uninsured; part of Ryan White program (cohort)	Housing; income	VL > 200 copies/mL	–
Schafer et al. 2012 [87] (article)	Virginia (USA)	251 men and women; part of UVA Ryan White clinic (RWC) (cross-sectional cohort)	Education; socioeconomic status	CD4 < 200 Detectable HIV viral load	–
Shacham et al. 2010 [88] (article)	St Louis, Missouri (USA)	514 people living with HIV presenting at Washington University HIV Clinic in 2007 (cross-sectional)	Education; employment; housing	VL ≥ 400 copies/mL	–
Shacham et al. 2013 [89] (article)	St Louis, Missouri (USA)	762 people living with HIV presenting at Washington University HIV Clinic in 2008 (cross-sectional)	Employment; poverty	CD4 < 200 cells/μl VL < 400 copies/mL	–
Singh et al. 1999 [90] (article)	USA	123 people who completed refill-methodology assessment of adherence (prospective cohort)	Education; employment; income	–	< 90% ART adherence
Stone et al. 2001 [91] (article)	Rhode Island; Maryland; New York State; Michigan (USA)	289 women living with HIV/AIDS; part of HIV Epidemiologic Research (HER) longitudinal cohort study (cross-sectional survey of cohort)	Education	–	Missed ≥ 1 HAART doses, past 3 days

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Storholm et al. 2019 [92] (article)	Los Angeles, California (USA)	239 African American adults living with HIV recruited in community settings (246: Project Mednet and 108 from Project Rise) (longitudinal)	Education; employment; housing; income	–	ART adherence trajectory group: high-stable (mean > 90%), moderately low-stable (> 60%), low-decreasing (< 27%) at 2, 4, 6 months or 1.5, 4.5, 6 months
Sunil and McGehee 2007 [93] (article)	USA	1910 people living with HIV taking ART; part of HIV Cost and Services Utilization Study (HCSUS) (cross-sectional survey of cohort)	Education	–	ART adherence, past 7 days (non-adherent: ≥ 1 medication missed)
Surratt et al. 2015 [94] (article)	Miami; Fort Lauderdale (USA)	503 socioeconomically disadvantaged substance users (cross-sectional—structured interviews)	Housing; income; poverty	–	Missed ARTs because of diversion, past 90 days
Tymejczyk et al. 2018 [95] (article)	New York City (USA)	1045 people living with HIV (sexual health clinic electronic medical records matched with longitudinal data from NYC HIV Surveillance Registry in 12 months before and after clinic visits) (secondary data analysis—linked surveillance and medical records review)	Poverty	VL ≤ 200 copies/mL	–
Vyas et al. 2014 [96] (article)	San Diego, California (USA)	350 adults attending a HIV clinic (prospective longitudinal)	Income	–	≥ 90% ART adherence
Wagoner et al. 2016 [97] (article)	Birmingham, Alabama (USA)	382 people living with HIV at University of Alabama clinic (1917 Clinic) with a HIV viral load available; part of Project Client-Oriented New Patient Navigation to Encourage Connection to Treatment (CONNECT) (cohort)	Education	VL ≥ 200 copies/mL	–

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Weiser et al. 2013 [98] (conference communication)	San Francisco (USA)	284 homeless and marginally housed individuals; part of San Francisco Research on Access to Care in the Homeless cohort (observational cohort)	Education; food security; housing; income	VL > 100 copies/mL CD4 < 200 cells/mm ³	< 90% ART adherence
Whelan et al. 2019 [99] (article)	King County, Washington (USA)	549 newly diagnosed people living with HIV who received a partner services interview (retrospective cohort)	Housing	VL ≥ 200 copies/mL	–
Wilson et al. 2018 [100] (article)	San Francisco Bay Area (USA)	159 transgender women of colour (55 participated at TransAccess, 46 at Brandy Martell Project, and 58 at Butterfly Project) as part of the Special Projects of National Significance (SPNS) Program, “Enhancing Engagement in Retention and Quality of Care for Transgender Women of Color” initiative (programme evaluation)	Food security; housing; income	CD4 < 500, past 6 months Undetectable VL, past 6 months	–
Yehia et al. 2014 [101] (article)	Philadelphia, Pennsylvania (USA)	12,759 adults using multiple clinics for primary HIV care; part of Ryan White program (retrospective cohort)	Income	VL ≤ 200 copies/mL	–
Europe (n = 20) Abgrall et al. 2014 [102] (letter)	France	200 individuals from Sub-Saharan Africa adherent to cART at enrollment; part of Agence Nationale de Recherche sur le SIDA et les hépatites virales (ANRS)-VIHVO study (cross-sectional)	Housing	–	< 80% cART adherence

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Burch 2018 [103] (thesis)	UK	2704 adults recruited from eight HIV outpatient clinics; part of Antiretrovirals Sexual Transmission Risk and Attitudes (ASTRA) study (cross-sectional)	Education; employment; housing; income	VL > 50 copies/mL VL > 200 copies/mL	≥ 2 ART consecutive missed days, past 3 months or ≥ 1 missed, past 2 weeks
Carrieri et al. 2003 [104] (article)	Marseilles; Avignon; Nice; Paris suburbs (France)	96 people who inject drugs (30 women, 66 men) living with HIV initially adherent to HAART in hospital clinics; part of French MANIF 2000 cohort (cohort)	Education; employment; housing	–	< 80% HAART adherence, past 18-months
Collazos et al. 2009 [105] (article)	Spain	1352 people living with HIV from 69 hospitals; part of Grupo Español para el Estudio Multifactorial de la Adherencia (GEEMA) cohort (prospective cohort)	Education	CD4 + cell count (cells/ μ l) at baseline Undetectable VL (baseline and 12 months) VL (log copies/mL) at baseline	HAART adherence rates
D'Almeida et al. 2016 [106] (article)	France	1246 people living with HIV who were HIV treatment-naïve at cART initiation and on cART for at least 12 months across 73 hospital departments; part of ANRS VESPA-2 study (cross-sectional)	Education; employment; deprivation	Sustained viral suppression (or undetectable VL < 50 copies/mL) for at least 6 months	–
Del Amo et al. 2017 [107] (article)	Austria, France, Germany, Greece, Italy, Spain, Switzerland, and the Netherlands	24,069 people living with HIV with data on education from 15 cohorts of patients initiating cART; part of Collaborator of Observational HIV Epidemiological Europe (COHERE) within EuroCoord Network of Excellence (multi-site cohort)	Education	VL < 400 copies/mL (1 year after cART initiation) Immunological response (change in CD4 + count) at baseline, first 6 months and after 6 months	–
Dorz et al. 2003 [108] (article)	Italy	88 male and 21 female adults living with HIV undergoing protease inhibitor treatment (cross-sectional)	Education; employment; income	–	< 80% adherence, past 7 days

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Gordillo et al. 1999 [109] (article)	Madrid (Spain)	366 people living with HIV on ART treatment (cross-sectional)	Education; employment	–	> 90% ART adherence ('good')
Gueler et al. 2015 [110] (article)	Switzerland	2694 ART-naïve for virologic response at six months; part of Swiss HIV Cohort Study (prospective cohort)	Socioeconomic position	Virological response to cART or viral suppression (VL < 50 copies/mL) at 6 months	–
Jansen et al. 2009 [111] (conference abstract)	Germany	2045 people living with HIV; part of Competence Network for HIV/AIDS (KompNet) cohort (cohort)	Income	VL < detection limit CD4 + cell count	–
Papadopoulou 2000 [112] (thesis)	Enfield/Haringey, London (England)	56 people living with HIV taking cART (cross-sectional)	Education	–	cART adherence (4 categories—"adhering to correct dose", "taking drugs at right time", "following dietary instructions" and "overall adherence")
Parruti et al. 2006 [113] (article)	Abruzzo Region (Italy)	171 people living with HIV clinically monitored for at least 24 weeks (cohort)	Employment; housing; SES	–	90% HAART adherence
Persson et al. 1994 [114] (article)	Malmö (Sweden)	47 MSM living with HIV (cross-sectional)	Social class	Low CD4 cell count (< median)	–
Raho-Moussa et al. 2019 [115] (article)	Paris (France)	475 people living with HIV attending two hospital outpatient clinics on ART for at least 6 months (cross-sectional)	Education; employment; deprivation	VL > 50 copies/mL	–
Saracino et al. 2018 [116] (article)	Italy	8023 ART naïve people living with HIV with Italian nationality; part of Italian Cohort Naïve Antiretrovirals (ICONA) cohort (observational cohort)	Education; employment	VL < 50 copies/mL	ART discontinuation
Sellier et al. 2006 [117] (brief report)	Paris (France)	61 adults born in Sub-Saharan Africa and living in France attending 3 infectious diseases departments (cross-sectional)	Education; employment	–	Never missed ART doses compared to frequently or rarely

Table 2 (continued)

Study (publication type)	Population		Exposure	Outcomes	
	Setting (city/state, country)	Population of people living with HIV; project name (study design)		Social determinant(s) ^a	Virological (non-) suppression measure ^b
Sherr et al. 2012 [118] (article)	London (UK); South East (UK)	259 adults living with HIV attending 5 HIV clinics (cross-sectional)	Education	–	All HAART doses taken at correct time and under correct conditions
Sobriño-Vegas et al. 2012 [119] (article)	Spain	4549 people living with HIV from 27 public health centres; part of CoRIS (AIDS Research Network Cohort) (prospective cohort)	Education	Immunological response to treatment (CD4 + T-cell count) at 6 months and 1 year VL < 50 copies/mL (6 months and 1 year)	–
Spire et al. 2002 [120] (article)	France	445 people living with HIV in 47 hospital departments; part of APROCO cohort (prospective cohort)	Housing; income	–	HAART non-adherence at 4 months follow up
Uusküla et al. 2012 [121] (article)	Kohtla-Järve (Estonia)	144 people living with HIV receiving outpatient HIV care (cross-sectional)	Education; income	–	< 100% ART adherence
Asia and Australia (n = 2)					
Sieffried et al. 2017 [122] (article)	Australia	522 adults on ART at 7 clinics; part of Parameters Associated with Adherence to Antiretroviral Therapy (PAART) study (cross-sectional analysis of cohort)	Employment; housing; income	–	Self-reported missed ≥ 3 ART, past 3 months
Yang and Bang 2017 [123] (article)	South Korea	300 people living with HIV from 6 hospitals; part of Nationwide Specialized Counseling Program (cross-sectional)	Education; employment; income	–	$\geq 95\%$ ART adherence

ART antiretroviral therapy, cART combination antiretroviral therapy, HAART highly active antiretroviral therapy, MEMS medication event monitoring system, MSM men who have sex with men, PI protease inhibitor, VL viral load

^aMeasures of social determinants vary across studies and may be compared at individual, household, or neighbourhood level

^bMeasures of viral suppression include CD4 cell count and viral load. ^cMedication includes ART, cART, and HAART



(a) Educational attainment and virological suppression**(b) Educational attainment and medication adherence**

Fig. 2 Forest plots of likelihood (aOR) of **a** virological suppression and **b** medication adherence among people living with HIV with low, compared to high, educational attainment after adjusting for sociode-

mographic factors. *ART* antiretroviral therapy, *GED* general educational diploma, *VL* viral load

were 96% (aOR 1.96; 95% CI 1.87, 2.05) and 45% (aOR 1.45; 95% CI 1.38, 1.52) more likely to be adherent to ART than those who only completed some high school, however those with an undergraduate qualification were 12% less likely (aOR 0.88; 95% CI 0.82, 0.95) to be adherent [93]. For Hispanic American participants ($n = 272$), the direction of association was less sequential; individuals who had completed high school or some College were 22% (aOR 0.78; 95% CI 0.72, 0.84) and 29% (aOR 0.71; 95% CI 0.66, 0.77) less likely, respectively, to be adherent whereas university graduates were 40% more likely (aOR 1.40; 95% CI 1.27,

1.55) to be adherent compared to those who only completed some high school [93].

Employment

Employment was defined either by status, occupation type or whether work was paid.

Of 83 studies, 33 looked at employment with only three studies reporting a significant association with virological suppression after adjustment for confounders [58, 103, 116]; two associations were negative (those unemployed), one was

positive (those employed) (Fig. 3a; Additional file 6). Overall, there was an inconsistent association between employment and virological suppression. For instance, D'Almeida et al.[106] reported no difference in viral suppression between people living with HIV of a lower employment grade or unemployed working status compared with executive occupational grades or employed. However, Saracino et al. 2018 [116] found that unemployed people living with HIV had lower rates (adjusted Hazard Ratio [aHR] 0.87; 95% CI 0.79, 0.96) of virological suppression, compared to full-time workers. Similarly, Burch [103] reported that the prevalence of virological non-suppression (VL > 50 copies/mL) was almost 2-times greater among the unemployed, compared to employed (adjusted Prevalence Ratio [aPR] 1.98; 95% CI 1.51, 2.61).


Similarly, the association between medication adherence and employment varied. Four studies reported a significant adjusted association with medication adherence

[47, 68, 103, 116]; three reported a negative association (those unemployed) and one a positive association (those employed) (Fig. 3b; Additional file 6). Saracino et al.[116] found unemployed people living with HIV were more at risk of discontinuing ART (aHR 1.18; 95% CI 1.04, 1.34) and when they looked at specific job types, “housewives” were found to be less at risk of ART discontinuation (aHR 0.73; 95% CI 0.59, 0.90), after adjusting for CD4 count, VL, pregnancy status and smoking. Kyser et al.[68] reported that people living with HIV who were unemployed or looking for work were over 2-times more likely (aOR 2.03; 95% CI 1.14, 3.61) to report having missed cART doses in the past 3 days.


Housing

Some studies described housing as the physical environment; some grouped individuals into categories of ‘unstable’

(a) Employment grade/status and virological suppression



(b) Employment grade/status and medication adherence



*Adjusted for different confounders


Fig. 3 Forest plots of likelihood (aOR) of **a** virological suppression and **b** medication adherence among people living with HIV with lower employment grade or unemployed status, compared to high, after adjusting for sociodemographic factors. VL viral load

and ‘stable’ housing; some used time of residence and others used home ownership. We grouped studies which explored the association between people living with HIV in unstable compared to stable housing and differing VL thresholds. Housing ‘stability’ was either explicitly described by the authors or was defined in relation to the type of housing (e.g., living in subsidised housing or in a shelter).

Of 83 studies, 39 looked at housing with eight studies reporting a significant adjusted association with virological suppression, all of which were a negative association, i.e., more unstable housing was associated with lower viral

suppression [44, 48, 50, 53, 72, 85, 98, 103] (Fig. 4a; Additional file 6). As measurements and definitions of housing status and virological suppression were consistent, we conducted a random-effects meta-analysis of this determinant and outcome. Most included studies in the meta-analysis used a VL threshold < 200 copies/mL [46, 48, 53, 85, 86] however, one used < 100 copies/mL [98]. The pooled aOR of studies that used a VL threshold of < 200 copies/mL was 0.48 (95% CI 0.33, 0.70) with high heterogeneity between the studies ($I^2=90\%$). All but 5 reported subgroups found unstable housing to be significantly associated with lower

(a) Housing status and virological suppression



(b) Housing status and medication adherence




Fig. 4 Forest plots of likelihood (aOR) of **a** virological suppression and **b** medication adherence among people living with HIV living in unstable, compared to stable, housing after adjusting for sociodemo-

graphic factors. A meta-analysis is presented for virological suppression. *M0* month 0, *M4* month 4, *SRO* single room occupancy, *VL* viral load

viral suppression with the strongest relative effect reported by Santos et al. [85]. Santos et al. [85] found that homeless or marginally, compared to stably, housed transgender women living with HIV were 95% less likely (aOR 0.05; 95% CI 0.01, 0.25) to be virally suppressed after adjusting for social factors including age and race/ethnicity. Overall, the pooled aOR was 0.49 (95% CI 0.34, 0.69) with the random-effects model for meta-analysis displaying considerable heterogeneity across studies ($I^2 = 89%$) [35].

Nine of 39 studies looking at housing found a significant adjusted association with medication adherence [45, 81, 94, 98, 102, 103, 113, 120, 122]; eight reported a negative association (unstable housing or homelessness) and one positive (long-term housing) (Fig. 4b; Additional file 6). Individuals living in unstable housing across a range of settings, including subsidised housing [122], living with others [102], being recently homeless [81, 94, 98, 113] or renting [103], were at greater risk of experiencing medication adherence failure, compared to those with more stable living situations. Spire et al. [120] looked at housing quality and found individuals living in stable, but poor housing or unstable housing at baseline have a greater likelihood of being non-adherent to HAART following 4-months of follow-up after adjusting for age, marital status and other intermediary determinants [120].

Other Measures of Material Deprivation

Some studies grouped social determinants to report one overall deprivation measure; for instance, Kalichman and Grebler [63] grouped social determinants of housing, food and financial security as “poverty-related experiences” or “stressors.” We present comparable aORs exploring the association between measures of material deprivation, including food security, with virological suppression and medication adherence (Fig. 5; Additional file 6) and a narrative synthesis provided for each factor. Of 83 studies, 53 explored a measure of deprivation: specifically, food security ($n = 7$), poverty/deprivation explicitly ($n = 9$), income ($n = 33$) and socioeconomic status/position ($n = 4$).

Food Security

Food security is measured at the individual level using specific measurement tools such as the Household Food Security Survey Module (HFSSM). Food security was sometimes reported as a combined measure with income. One of seven food security studies reported a significant negative adjusted association with viral suppression [98]. Weiser et al. [98] found that individuals who reported any food insecurity were 29% more likely to be virologically non-suppressed (aOR 1.29; 95% CI 1.04, 1.61); however, the same study [98] found no difference in virological suppression among

food insecure individuals after adjusting for adherence. Six studies investigating food security and adherence found a significant negative adjusted association [42, 56, 64, 65, 67, 98]. For instance, Almeida-Brasil et al. [42] examined a cohort of adults in Canada living with HIV and hepatitis C and established that there was no difference in virological suppression (VL > 50 copies/mL) among individuals reporting income-related food insecurity (aOR 1.44; 95% CI 0.95, 2.19) but were more likely to be non-adherent to medication (aOR 1.77; 95% CI 1.26, 2.48).


Poverty and Deprivation

Poverty and deprivation are reported as fundamentally neighbourhood or area level factors and based on context-specific measurement tools, including the Evaluation of Deprivation and Inequalities in Health Examination Centres (EPICES). Two of nine studies reported a significant adjusted negative association with virological suppression and deprivation [95, 115]. For instance, Raho-Moussa et al. [115] found that individuals who reported either individual determinants of deprivation (specifically financial difficulties in past month) or had an EPICES score indicating residence in a deprived state were 66% (aOR 0.34; 95% CI 0.16, 0.72) and 65% (aOR 0.35; 95% CI 0.14, 0.85) less likely, respectively, to achieve a VL < 50 copies/mL, compared to people living with HIV who did not meet these measures of deprivation, after adjusting for age and medication-, and clinically-related factors. Only one study reported a significant adjusted association with medication adherence which was in a negative direction [63]. Kalichman and Grebler [63] report that individuals reporting poverty-related stress are 28% (aOR 0.72; 95% CI 0.59, 0.89) and 26% (aOR 0.74; 95% CI 0.62, 0.88) less likely to report 85% and 75% ART adherence, respectively, after adjusting for social stressors, depression, internalised AIDS stigma and drug use.

Income

Income included components of temporality (e.g., annual), ecological level (e.g., household), sources (e.g., social benefits) and quantities (e.g., < \$15,000). Four of 33 studies looking at income found a significant adjusted association with virological suppression [58, 77, 86, 101, 103]; three reported a negative association (lower income) and the other two a positive association (higher/mid-range income). For instance, the likelihood of virological non-suppression among Canadian men who have sex with men earning < \$15,000 annually were 6-times greater (aOR 6.43; 95% CI 2.08, 19.89), than those earning more, after adjusting for age, ethnicity, sexual orientation, country of birth and other characteristics [77]. However, one study [58] found people living with HIV with an annual household

(a) Deprivation, food insecurity, income and virological suppression



(b) Deprivation, food insecurity, income and medication adherence




Fig. 5 Forest plots of likelihood (aOR) of **a** virological suppression and **b** medication adherence among people living with HIV who were disadvantaged, compared to more advantaged after adjusting for sociodemographic factors. *EPICES* Evaluation of Deprivation and Inequalities in Health Examination Centres, *FPL* federal poverty level, *VL* viral load

income of < \$10,000 were significantly more likely to report virological suppression, compared to those earning \$10,000–\$24,999 (aOR 0.21; 95% CI 0.06, 0.73) and \$25,000–\$49,999 (aOR 0.11; 95% CI 0.03, 0.52) [58]. The authors suggest that this may be due to those with lower annual household incomes being more likely to be receiving support linked to the Ryan White Program [58]. Sayles et al. [86] used federal poverty level (FPL) as a measure of income and found that uninsured people living with HIV in Los Angeles, who were receiving publicly funded healthcare through the Ryan White program, and had an income \leq 100% FPL were 27% more likely (aOR 1.27; 95% CI 1.15, 1.41) to be virologically non-suppressed than those with an income greater than the FPL. Five studies reported a significant adjusted association between income and medication adherence [41, 57, 79, 96, 103]; negative associations were found among low income individuals [57, 79, 103] whilst, comparatively, mid-level/higher income were found to be positively associated [41, 96]. Vyas et al. [96] found individuals with a higher annual household income (\geq \$10,000) were significantly more likely to be \geq 90% ART adherent. The situation is less clear within the Veterans Aging Cohort Study whereby only individuals earning a mid-range annual household income (\$25,000–\$49,999) were significantly more likely to be adherent to medication, compared to those earning < \$6000 [41]. A significant association was not found among the other 3 income categories [41]. Finally, Burch [103] reported whether individuals in the UK had financial stability through a proxy of having “enough money for basic needs”; they found that those who reported mostly, sometimes or not having enough were more likely to be non-adherent, compared to always having enough money.

Socioeconomic Status/Position (SES/SEP)

SES/SEP were reported across ecological levels. Two of four studies reported a significant positive adjusted association with higher/mid-range SES/SEP and viral suppression [87, 110]. Interestingly, a US study [87] found that individuals who had a “mid-range” SES which was defined by their payscale and whether they received support with healthcare costs (5–70% co-pay) were less likely (adjusted Relative Risk [aRR] 0.39; 95% CI 0.16, 0.94) to have a detectable HIV viral load, although this did not remain significant when considering CD4 counts < 200. Only one study looked at

adjusted associations between SES and medication adherence; Parruti et al. [113] report no significant difference (OR 0.76; 95% CI 0.30, 2.00) in HAART adherence between people living with HIV in Italy with very low or low SES, compared to those with medium or high SES.

Social Class

One study [114] explored the association between social class and treatment outcomes. Persson et al. [114] found no difference in the likelihood of having CD4 counts lower than the median value among skilled and unskilled workers (defined as social class III) compared to middle range civil servants (social class II) (OR 1.5; 95% CI 0.5, 4.9).

Marginalised Sub-groups

Some studies followed specific subgroups of individuals who are often disproportionately affected by HIV; including, the homeless, people in prisons, people who use drugs and transgender women. Oftentimes, people will identify with multiple identities (intersectionality). For instance, Marshall et al. [72] reported that homeless people living with HIV who use drugs were almost half as likely (aPR 0.55; 95% CI 0.42, 0.71) to have an undetectable VL compared to those with housing, even after adjusting for sociodemographic factors and factors related to their substance use including addiction treatment. Berg et al. [45] found ART adherence rate was greater among individuals who were current or former opioid users who had lived in long-term housing even after adjusting for gender and intermediary determinants including alcohol and substance use. Finally, a small cross-section of transgender women living with HIV ($n = 123$) living in unstable housing had one of poorest virological outcomes, specifically their risk of having a detectable VL was over 7-times (aRR 7.37; 95% CI 1.07, 50.88) that of transgender women living in stable housing [44].

Risk of Bias

Overall, 8 (9.6%) studies [51, 60, 69, 81, 102, 111, 117, 123] included in the review were scored ‘low quality’ or had a high risk of bias. Notably, several of the cross-sectional studies were possibly affected by selection bias; for instance, investigating a small sample size or not stating the frequency of non-respondents (Additional file 7). There was heterogeneous reporting within and between studies with no standardised approach for measuring and classifying social factors (exposures), outcomes or associations which limited possible study comparisons.

Discussion

A small proportion (10–20%) of studies observed significant associations between material deprivation and poorer clinical outcomes. Overall, they suggest that people living with HIV who are the most materially deprived (housing, employment, deprivation, or income) display poorer viral suppression and medication adherence compared to those more advantaged. The strongest evidence is present for housing whereby 1 in 5 studies found unstable housing status was associated with poorer viral suppression; however, we found that 89% of the variability in the ORs could be explained by heterogeneity between the studies. An inconsistent association was observed for studies measuring education and adherence outcomes. A higher proportion of included studies observed significant, consistent associations between SES/SEP and virological suppression (50%) and food security and adherence (86%). We also found evidence that intersectionality worsens outcomes. The magnitude of associations were compounded within specific subgroups; for example, US transgender women who were homeless or marginally housed were 20-times more likely to be virologically non-suppressed than transgender women living in stable housing, after adjusting for age, race/ethnicity, history of injection drug use and health insurance status [85].

Our findings are in line with other studies including reviews which have found worsening outcomes associated with material deprivation, including unstable housing status, food insecurity and lower socioeconomic status [14, 124–126]. We find that these associations persist in high-income countries even as advanced generations of antiretrovirals become widely available, HIV prevention programmes continue to be scaled-up and HIV care evolves significantly through the digital age. For instance, in line with Krieger's ecosocial theory [27, 28], issues of agency for HIV care exacerbate inequities among US women living with HIV [127]. This may be explained by the context-specific and interrelated nature of social determinants. For instance, school leavers ages vary across countries (from 15 to 18 years old) as does minimum wage, entitlements to welfare and state benefits and other support services available. In other words, social and cultural capital go beyond the measures of "relative deprivation" but can be more difficult to measure [128, 129]. Unlike Burch et al. [14], we did not explore health insurance as this is not a considerable barrier in countries with a universal healthcare system, such as the UK. Rather, factors such as social class drive health inequalities seen in the UK which were first described in the 1980 Black Report and later by Marmot's Reports [3, 5, 130]. Associations between structural factors (HIV-related laws), interpersonal factors (perceived social capital) and individual outcomes (ART adherence) have also been

identified across ecosocial context levels in North America which highlights the potential mechanisms of relationships between social and structural determinants [82]. Additionally, some determinants may be more directly associated with outcomes than others; for instance, some medication should be taken with food which may not be possible for somebody who is food insecure/insufficient, in turn influencing ART adherence [126].

The review had several strengths including a comprehensive search strategy and the inclusion of independent reviewers at screening, data extraction and quality assessment stages. We also grouped duplicate reports at the final stage of screening as one study and included the most recent report in the review to avoid possible publication bias. The majority of included studies were scored as low or uncertain risk of bias; however, the modified Newcastle–Ottawa scale used for cross-sectional studies has not been validated [37, 131]. Additionally, no grey literature was included, nor publications not published in English, which may have inadvertently excluded relevant studies, particularly from HIV community groups and charities. We only included observational studies, limiting the ability to make any causal inferences and potentially introducing social desirability bias of studies reliant on self-reported data only. Our review contributes to this field of knowledge but is unable to determine how determinants may or may not be causally linked but begins to suggest which determinants may interact with treatment outcomes of people living with HIV. We also recognise that social determinants are heavily interlinked, with collinearity between variables; however, we do not attempt to develop a causal pathway in this paper.

These findings re-emphasise the need for well-designed measures of social determinants in studies with evidence-based, context-specific definitions; for instance, higher-income countries (e.g., in Europe) use a 'class structural' approach to define occupation compared to low- and middle-income countries whereby occupation is highly dependent on working conditions (e.g., formal/informal sectors or environmental factors) [132]. As previously suggested by Krieger et al. [38] studies should be collecting socioeconomic data across all ecological levels and the lifecourse, treat income and poverty as dynamic, consider types of assets and wealth but also not conflate between SES and social class. We found that these recommendations were not met in most included studies and rather studies measured determinants at single timepoints or only at one ecological level. There is a clear gap for well-designed research looking at the impact of social class on treatment outcomes of people living with HIV. For instance, research into using subjective social class has found that individuals often use measures of socioeconomic position to assign status however, also take into account their current and future material and economic

prospects which may be a more suitable, composite socioeconomic indicator [133].

Additionally, to develop a more conclusive understanding, a need remains to standardise thresholds of viral suppression across studies to allow for more detailed analyses of the direction of reported effect sizes. This would enable more robust random effects meta-analyses, and subsequent meta-synthesis, to be conducted despite issues relating to the generalisability of results and heterogeneous nature of studies. As a result, this would support the design of more appropriate, system-wide interventions and clinics to identify and support the most socioeconomically disadvantaged, and marginalised, people living with HIV. For instance, interventions focussing on the provision of stable and secure housing, rather than limiting this to adherence support for people continuing to live in precarious housing conditions which could have significant implications for HIV outcomes as well as wider mental health and wellbeing outcomes [134]. The Housing and Health Study for rental assistance found that HIV-related housing services were cost-effective as a HIV prevention intervention but also in relation to quality of life; a cost-per-QALY-saved estimated to be \$62,493 for homeless and unstably housed people living with HIV across three US cities [135, 136]. At today's exchange rate, this exceeds the UK National Institute for Health and Care Excellence (NICE) evidence-based guidelines for public health and social care services in for which as a cost-effectiveness threshold between £20,000–£30,000 per QALY gained [137].

Further research is required to determine to what extent person-centred and holistic HIV care improves the health and wellbeing outcomes among people living with HIV. For instance, the provision of social support services including housing, welfare and benefits advice, food and transport vouchers, alongside routine clinical monitoring to help address stigma, reduce health inequalities and ensure equitable access to HIV treatment and care.

Conclusions

Our study has shown that the contribution of measures of material deprivation on HIV treatment outcomes remains complex. There is a need to propose a causal pathway of the relationship between these factors. An ecosocial analysis would help to establish the impact of social determinants between and across ecological levels (e.g., individual, household, neighbourhood) and how these are 'embodied' by individuals across the lifecourse. Further research might then be able to disentangle how social determinants are driven by inequalities in gender, sexual orientation, social class, and race/ethnicity allowing us to build a deeper understanding of how, when, and to what extent, these determinants interact

with one another. Subsequently, this would ensure the support systems and services in places for the most marginalised groups living with HIV are acceptable and have the greatest impact on service users.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10461-021-03551-y>.

Acknowledgements We thank the staff in Library Services (Imperial College London) for their support with designing our search strategy. We also thank Dr Valerie Delpech (Public Health England) for her feedback on our initial approach to this review. Finally, thank you to Dr Lidia Luna Puerta for her translation of our abstract into Spanish.

Author Contributions VP, BD and HW, designed the research study. VP led the systematic review and drafted the first version of this manuscript. AS and VP conducted title, abstract and full text screening. VP extracted data which was cross-checked by EC. VP conducted the quality assessment which was cross-checked by BD. VP, BD and HW critically interpreted the results and revised the manuscript. All authors have read and approved the final manuscript.

Funding VP is supported by an ESRC LISS-DTP CASE Studentship. BD acknowledges funding from the NIHR HPRU in Chemical and Radiation Threats and Hazards (NIHR-200922); HDR UK Hub DISCOVER-NOW and infrastructure support for the Department of Epidemiology and Biostatistics provided by the NIHR Imperial Biomedical Research Centre. HW is a NIHR Senior Investigator and acknowledges support from NIHR Imperial Biomedical Research Centre, NIHR School of Public Health, NIHR Applied Research Collaborative North West London, and Wellcome Trust (UNS32973). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Data Availability Supporting data are available from the corresponding author upon request.

Code Availability Not applicable.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Ethical Approval Not applicable.

Informed Consent Not applicable.

Consent for Publication Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Solar O, Irwin A. A conceptual framework for action on the social determinants of health. Geneva: World Health Organization; 2010. (Social Determinants of Health Discussion Paper 2 (Policy and Practice)).
2. Graham H. Social determinants and their unequal distribution: clarifying policy understandings. *Milbank Q*. 2004;82(1):101–24.
3. Marmot M, Allen J, Goldblatt P, Boyce T, McNeish D, Grady M, et al. Fair society, healthy lives. Amsterdam: Elsevier; 2010.
4. Marmot M, Allen J, Goldblatt P, Boyce T, McNeish D, Grady M, et al. Chapter 3: Lessons to be learned from the current Health Inequality Strategy, targets and indicators. In: Fair society, healthy lives. Amsterdam: Elsevier; 2010. p. 85–91.
5. Marmot M, Allen J, Boyce T, Goldblatt P, Morrison J. Health equity in England: the marmot review 10 years on. London: Institute of Health Equity; 2020.
6. Stuckler D, Reeves A, Loopstra R, Karanikolos M, McKee M. Austerity and health: the impact in the UK and Europe. *Eur J Public Health*. 2017;27(suppl_4):18–21.
7. Caiola C, Docherty SL, Relf M, Barroso J. Using an intersectional approach to study the impact of social determinants of health for African American mothers living with HIV. *ANS Adv Nurs Sci*. 2014;37(4):287–98.
8. Geter A, Sutton MY, McCree DH. Social and structural determinants of HIV treatment and care among black women living with HIV infection: a systematic review: 2005–2016. *AIDS Care*. 2018;30(4):409–16.
9. Robinson RS. From population to HIV: the organizational and structural determinants of HIV outcomes in sub-Saharan Africa. *J Int AIDS Soc*. 2011;14(2):S6.
10. Alvarez-del Arco D, Monge S, Azcoaga A, Rio I, Hernando V, Gonzalez C, et al. HIV testing and counselling for migrant populations living in high-income countries: a systematic review. *Eur J Public Health*. 2012;23(6):1039–45.
11. Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet*. 2014;384(9953):1529–40.
12. Zierler S, Krieger N. HIV infection in women: social inequalities as determinants of risk. *Crit Public Health*. 1998;8(1):13–32.
13. Krieger N. Measures of racism, sexism, heterosexism, and gender binarism for health equity research: from structural injustice to embodied harm: an ecosocial analysis. *Annu Rev Public Health*. 2020;41(1):37–62.
14. Burch LS, Smith CJ, Phillips AN, Johnson MA, Lampe FC. Socioeconomic status and response to antiretroviral therapy in high-income countries: a literature review. *AIDS*. 2016;30(8):1147–62.
15. Cahill SR, Mayer KH, Boswell SL. The Ryan white HIV/AIDS program in the age of health care reform. *Am J Public Health*. 2015;105(6):1078–85.
16. Bradley H, Viall AH, Wortley PM, Dempsey A, Hauck H, Skarbinski J. Ryan white HIV/AIDS program assistance and HIV treatment outcomes. *Clin Infect Dis*. 2016;62(1):90–8.
17. Benson C, Wang X, Dunn K, Li N, Mesana L, Lai J, et al. Antiretroviral adherence, drug resistance, and the impact of social determinants of health in HIV-1 patients in the US. *AIDS Behav*. 2020;24:3562.
18. Nachega JB, Uthman OA, Peltzer K, Richardson LA, Mills EJ, Amekudzi K, et al. Association between antiretroviral therapy adherence and employment status: systematic review and meta-analysis. *Bull World Health Organ*. 2014;93:29–41.
19. Perelson AS, Essunger P, Cao Y, Vesanen M, Hurley A, Saksela K, et al. Decay characteristics of HIV-1-infected compartments during combination therapy. *Nature*. 1997;387(6629):188–91.
20. Byrd KK, Hou JG, Hazen R, Kirkham H, Suzuki S, Clay PG, et al. Antiretroviral adherence level necessary for HIV viral suppression using real-world data. *J Acquir Immune Defic Syndr*. 2019;82(3):245–51.
21. Joint United Nations Programme on HIV/AIDS (UNAIDS). 90-90-90: An ambitious treatment target to help end the AIDS epidemic. p. 40. https://www.unaids.org/sites/default/files/media_asset/90-90-90_en.pdf (2014).
22. Kirby T. The UK reaches UNAIDS 90–90–90 targets. *Lancet*. 2018;392(10163):2427.
23. Public Health England. Trends in HIV testing, new diagnoses and people receiving HIV-related care in the United Kingdom: data to the end of December 2019. 2020;14 20:16.
24. National AIDS Trust. HIV in the UK statistics. UK HIV Statistics. <https://www.nat.org.uk/about-hiv/hiv-statistics>. Accessed 20 Mar 2021.
25. Eckersley R. Beyond inequality: acknowledging the complexity of social determinants of health. *Soc Sci Med*. 2015;147:121–5.
26. Krieger N. Epidemiology and the web of causation: has anyone seen the spider? *Soc Sci Med* (1982). 1994;39(7):887–903.
27. Krieger N. Theories for social epidemiology in the 21st century: an ecosocial perspective. *Int J Epidemiol*. 2001;30(4):668–77.
28. Krieger N. Epidemiology and the people's health: theory and context. Oxford: Oxford University Press; 2011.
29. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
30. Papageorgiou V, Singer A, Cooper E, Davies B, Ward H. A systematic review of social determinants and HIV treatment outcomes among people living with HIV in high-income countries. PROSPERO. https://www.crd.york.ac.uk/prospéro/display_record.php?ID=CRD42020171850 (2020). Accessed 9 June 2020.
31. World Bank. World Bank Country and Lending Groups. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups> (2019). Accessed 19 Mar 2020.
32. Organisation for Economic, Co-operation and Development (OECD), OECD. Our global reach. <https://www.oecd.org/about/members-and-partners/>. Accessed 19 Mar 2020.
33. Covidence systematic review software. Melbourne, Australia: Veritas Health Innovation. www.covidence.org
34. Review Manager (RevMan) [Computer program]. Version 5.4. The Cochrane Collaboration; 2020.
35. Deeks JJ, Higgins J, Altman DG, editors. Chapter 10: Analysing data and undertaking meta-analyses. In: Cochrane handbook for systematic reviews of interventions version 62 (updated February 2021). Cochrane. www.training.cochrane.org/handbook (2021).
36. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 2013. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp (2016).
37. Modesti PA, Reboldi G, Cappuccio FP, Agyemang C, Remuzzi G, Rapi S, et al. Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis. *PLoS ONE*. 2016;11(1):e0147601.
38. Krieger N, Williams DR, Moss NE. Measuring social class in US public health research: concepts, methodologies, and guidelines. *Annu Rev Public Health*. 1997;18(1):341–78.
39. World Health Organization. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: recommendations for a public health approach. http://apps.who.int/iris/bitstream/10665/208825/1/9789241549684_eng.pdf (2016). Accessed 10 July 2020.

40. World Health Organization. Patient monitoring guidelines for HIV care and antiretroviral therapy (ART); 2006.
41. Keith McInnes D, Shimada SL, Rao SR, Quill A, Duggal M, Gifford AL, et al. Personal health record use and its association with antiretroviral adherence: survey and medical record data from 1871 US veterans infected with HIV. *AIDS Behav.* 2013;17(9):3091–100.
42. Almeida-Brasil CC, Moodie EEM, McLinden T, Hamelin A-M, Walmsley SL, Rourke SB, et al. Medication nonadherence, multitablet regimens, and food insecurity are key experiences in the pathway to incomplete HIV suppression. *AIDS.* 2018;32(10):1323.
43. Anderson JC, Campbell JC, Glass NE, Decker MR, Perrin N, Farley J. Impact of intimate partner violence on clinic attendance, viral suppression and CD4 cell count of women living with HIV in an urban clinic setting. *AIDS Care.* 2018;30(4):399–408.
44. Baguso GN, Turner CM, Santos G-M, Raymond HF, Dawson-Rose C, Lin J, et al. Successes and final challenges along the HIV care continuum with transwomen in San Francisco. *J Int AIDS Soc.* 2019;22(4):e25270.
45. Berg KM, Demas PA, Howard AA, Schoenbaum EE, Gourevitch MN, Arnsten JH. Gender differences in factors associated with adherence to antiretroviral therapy. *J Gen Intern Med.* 2004;19(11):1111–7.
46. Blank AE, Fletcher J, Verdecias N, Garcia I, Blackstock O, Cunningham C. Factors associated with retention and viral suppression among a cohort of HIV+ women of color. *AIDS Patient Care STDs.* 2015;29(Suppl 1):S27–35.
47. Chitsaz E, Meyer JP, Krishnan A, Springer SA, Marcus R, Zaller N, et al. Contribution of substance use disorders on HIV treatment outcomes and antiretroviral medication adherence among HIV-infected persons entering jail. *AIDS Behav.* 2013;17(Supplement 2):S118–27.
48. Clemenzi-Allen A, Geng E, Christopoulos K, Hammer H, Buchbinder S, Havlir D, et al. Degree of housing instability shows independent ‘dose-response’ with virologic suppression rates among people living with human immunodeficiency virus. *Open Forum Infect Dis.* 2018;5(3):ofy035.
49. Creasy SL, Henderson ER, Bukowski LA, Matthews DD, Stall RD, Hawk ME. HIV Testing and ART adherence among unstably housed black men who have sex with men in the United States. *AIDS Behav.* 2019;23(11):3044–51.
50. Doshi RK, Milberg J, Jumento T, Matthews T, Dempsey A, Cheever LW. For many served by the Ryan white HIV/AIDS program, disparities in viral suppression decreased, 2010–14. *Health Aff.* 2017;36(1):116–23.
51. Dowshen N, Matone M, Luan X, Lee S, Belzer M, Fernandez MI, et al. Behavioral and health outcomes for HIV+ young transgender women (YTW) linked to and engaged in medical care. *LGBT Health.* 2016;3(2):162–7.
52. Fadul N, Taylor P, Kearney G. Predictors of viral load suppression in HIV-infected patient in rural Eastern North Carolina. *Open Forum Infect Dis.* 2017;4(Supplement 1):S421.
53. Feldman M, Thomas J, Alexy E, Irvine M. Crystal methamphetamine use and HIV medical outcomes among HIV-infected men who have sex with men accessing support services in New York. *Drug Alcohol Depend.* 2015;147:266–71.
54. Feller DJ, Agins BD. Understanding determinants of racial and ethnic disparities in viral load suppression: a data mining approach. *J Int Assoc Provid AIDS Care.* 2016;16(1):23–9.
55. Gardner S, Rachlis B, Light L, Raboud J, Cooper C, Kendall C, et al. Impact of social determinants of health and substance use on HIV viral suppression (SVL). *Can J Infect Dis Med Microbiol.* 2015;26(Suppl. SB):38B.
56. Gebo KA, Keruly J, Moore RD. Association of social stress, illicit drug use, and health beliefs with nonadherence to antiretroviral therapy. *J Gen Intern Med.* 2003;18(2):104–11.
57. Golin CE, Liu H, Hays RD, Miller LG, Beck CK, Ickovics J, et al. A prospective study of predictors of adherence to combination antiretroviral medication. *J Gen Intern Med.* 2002;17(10):756–65.
58. Haider MR, Brown MJ, Harrison S, Yang X, Ingram L, Bhochohibhoya A, et al. Sociodemographic factors affecting viral load suppression among people living with HIV in South Carolina. *AIDS Care.* 2019;33:1–9.
59. Hussen SA, Easley KA, Smith JC, Shenvi N, Harper GW, Camacho-Gonzalez AF, et al. Social capital, depressive symptoms, and HIV viral suppression among young black, gay, bisexual and other men who have sex with men living with HIV. *AIDS Behav.* 2018;22(9):3024–32.
60. Iralu J, Duran B, Pearson CR, Jiang YZ, Foley K, Harrison M. Risk factors for HIV disease progression in a rural southwest American Indian population. *Public Health Rep.* 2010;125(Supplement 4):43–50.
61. Johnson M, Catz S, Remien RH, Rotheram-Borus M, Morin CE, et al. Theory-guided, empirically supported avenues for intervention on HIV medication nonadherence: findings from the Healthy Living Project. *AIDS Patient Care STDs.* 2003;17(12):645–56.
62. Kacanek D, YanLing H, Malee K, Mellins CA, Smith R, Garvie PA, et al. Nonadherence and unsuppressed viral load across adolescence among US youth with perinatally acquired HIV. *AIDS.* 2019;33(12):1923–34.
63. Kalichman SC, Grebler T. Stress and poverty predictors of treatment adherence among people with low-literacy living with HIV/AIDS. *Psychosom Med.* 2010;72(8):810–6.
64. Kalichman SC, Cherry C, Amaral C, White D, Kalichman MO, Pope H, et al. Health and treatment implications of food insufficiency among people living with HIV/AIDS, Atlanta, Georgia. *J Urban Health.* 2010;87(4):631–41.
65. Kalichman S, Hernandez D, Cherry C, Kalichman M, Washington C, Grebler T. Food insecurity and other poverty indicators among people living with HIV/AIDS: effects on treatment and health outcomes. *J Community Health.* 2014;39(6):1133–9.
66. Kleeberger CA, Buechner J, Palella F, Detels R, Riddler S, Godfrey R, et al. Changes in adherence to highly active antiretroviral therapy medications in the Multicenter AIDS Cohort Study*. *AIDS.* 2004;18(4):683.
67. Koehn K, McLinden T, Collins AB, McDougall P, Baltzer-Turje R, Miewald C, et al. Assessing the impact of food insecurity on HIV medication adherence in the context of an integrated care programme for people living with HIV in Vancouver, Canada. *Public Health Nutr.* 2020;23(4):683–90.
68. Kyser M, Buchacz K, Bush TJ, Conley LJ, Hammer J, Henry K, et al. Factors associated with non-adherence to antiretroviral therapy in the SUN study. *AIDS Care.* 2011;23(5):601–11.
69. Lacombe-Duncan A, Bauer GR, Logie CH, Newman PA, Shokoohi M, Kay ES, et al. The HIV care cascade among transgender women with HIV in Canada: a mixed-methods study. *AIDS Patient Care STDs.* 2019;33(7):308–22.
70. Lim S, Nash D, Hollod L, Harris TG, Lennon MC, Thorpe LE. Influence of jail incarceration and homelessness patterns on engagement in HIV care and HIV viral suppression among New York City adults living with HIV/AIDS. *PLoS ONE.* 2015;10(11):e0141912.
71. Ludema C, Cole SR, Eron JJJ, Edmonds A, Holmes GM, Anastos K, et al. Impact of health insurance, ADAP, and income on HIV viral suppression among US women in the women’s interagency HIV study, 2006–2009. *JAIDS.* 2016;73(3):307–12.
72. Marshall BDL, Elston B, Dobrer S, Parashar S, Hogg RS, Montaner JSG, et al. The population impact of eliminating homelessness on HIV viral suppression among people who use drugs. *AIDS.* 2016;30(6):933–41.

73. McCoy K, Waldrop-Valverde D, Balderson BH, Mahoney C, Catz S. Correlates of antiretroviral therapy adherence among HIV-infected older adults. *J Int Assoc Provid AIDS Care*. 2016;15(3):248–55.
74. Miller CL, Spittal PM, Wood E, Chan K, Schechter MT, Montaner JSG, et al. Inadequacies in antiretroviral therapy use among Aboriginal and other Canadian populations. *AIDS Care*. 2006;18(8):968–76.
75. Mimiaga MJ, August Oddleifson D, Meersman SC, Silvia A, Hughto JMW, Landers S, et al. Multilevel barriers to engagement in the HIV care continuum among residents of the state of Rhode Island living with HIV. *AIDS Behav*. 2019;24:1133.
76. Mohammed H, Kieleyka L, Richardson-Alston G, Magnus M, Fawal H, Vermund SH, et al. Adherence to HAART among HIV-infected persons in Rural Louisiana. *AIDS Patient Care STDs*. 2004;18(5):289–96.
77. Moore DM, Cui Z, Lachowsky N, Raymond HF, Roth E, Rich A, et al. HIV community viral load and factors associated with elevated viremia among a community-based sample of men who have sex with men in Vancouver, Canada. *JAIDS*. 2016;72(1):87–95.
78. Nyaku M, Beer L, Shu F. Non-persistence to antiretroviral therapy among adults receiving HIV medical care in the United States. *AIDS Care*. 2019;31(5):599–608.
79. O'Neil CR, Palmer AK, Coulter S, O'Brien N, Shen A, Zhang W, et al. Factors associated with antiretroviral medication adherence among HIV-positive adults accessing highly active antiretroviral therapy (HAART) in British Columbia, Canada. *J Int Assoc Phys AIDS Care*. 2012;11(2):134–41.
80. Oliver C, Rebeiro PF, Hopkins MJ, Byram B, Carpenter L, Clouse K, et al. Substance use, demographic and socioeconomic factors are independently associated with postpartum HIV care engagement in the Southern United States, 1999–2016. *Open forum Infect Dis*. 2019;6(2):ofz023.
81. Phillips JC. Antiretroviral therapy adherence: testing a social context model among black men who use illicit drugs. *J Assoc Nurses AIDS Care*. 2011;22(2):100–27.
82. Phillips JC, Webel A, Rose CD, Corless IB, Sullivan KM, Voss J, et al. Associations between the legal context of HIV, perceived social capital, and HIV antiretroviral adherence in North America. *BMC Public Health*. 2013;13(1):1–16.
83. Rebeiro PF, Howe CJ, Rogers WB, Bebawy SS, Turner M, Kheshti A, et al. The relationship between adverse neighborhood socioeconomic context and HIV continuum of care outcomes in a diverse HIV clinic cohort in the Southern United States. *AIDS Care*. 2018;30(11):1426–34.
84. Robinson AC, Knowlton AR. Gender differences in psychosocial factors associated with HIV viral suppression among African-American injection drug users. *AIDS Behav*. 2016;20(2):385–94.
85. Santos GM, Wilson EC, Rapues J, Macias O, Packer T, Raymond HF. HIV treatment cascade among transgender women in a San Francisco respondent driven sampling study. *Sex Transm Infect*. 2014;90(5):430–3.
86. Sayles JN, Rurangirwa J, Kim M, Kinsler J, Oruga R, Janson M. Operationalizing treatment as prevention in Los Angeles county: antiretroviral therapy use and factors associated with unsuppressed viral load in the Ryan white system of care. *AIDS Patient Care STDs*. 2012;26(8):463–70.
87. Schafer KR, Brant J, Gupta S, Thorpe J, Winstead-Derlega C, Pinkerton R, et al. Intimate partner violence: a predictor of worse HIV outcomes and engagement in care. *AIDS Patient Care STDs*. 2012;26(6):356–65.
88. Shacham E, Nurutdinova D, Onen N, Stamm K, Overton ET. The interplay of sociodemographic factors on virologic suppression among a U.S. outpatient HIV clinic population. *AIDS Patient Care STDs*. 2010;24(4):229–35.
89. Shacham E, Lian M, Onen NF, Donovan M, Overton ET. Are neighborhood conditions associated with HIV management? *HIV Med*. 2013;14(10):624–32.
90. Singh N, Berman SM, Swindells S, Justis JC, Mohr JA, Squier C, et al. Adherence of human immunodeficiency virus: infected patients to antiretroviral therapy. *Clin Infect Dis*. 1999;29(4):824–30.
91. Stone VE, Hogan JW, Schuman P, Rompalo AM, Howard AA, Korkontzelou C, et al. Antiretroviral regimen complexity, self-reported adherence, and HIV patients' understanding of their regimens: survey of women in the her study. *JAIDS*. 2001;28(2):124–31.
92. Storholm ED, Bogart LM, Mutchler MG, Klein DJ, Ghosh-Dastidar B, McDavitt B, et al. Antiretroviral adherence trajectories among Black Americans living with HIV. *AIDS Behav*. 2019;23(8):1985–97.
93. Sunil TS, McGehee MA. Social and religious support on treatment adherence among HIV/AIDS patients by race/ethnicity. *J HIV/AIDS Soc Serv*. 2007;6(1/2):83–99.
94. Surratt HL, Kurtz SP, Levi-Minzi MA, Chen MX. Environmental influences on HIV medication adherence: the role of neighborhood disorder. *Am J Public Health*. 2015;105(8):1660–6.
95. Tymejczyk O, Jamison K, Pathela P, Braunstein S, Schillinger JA, Nash D. HIV care and viral load suppression after sexual health clinic visits by out-of-care HIV-positive persons. *AIDS Patient Care STDs*. 2018;32(10):390–8.
96. Vyas KJ, Limneos J, Qin HF, Mathews WC. Assessing baseline religious practices and beliefs to predict adherence to highly active antiretroviral therapy among HIV-infected persons. *AIDS Care*. 2014;26(8):983–7.
97. Wagoner N, Elope L, Westfall A, Mugavero M, Turan J, Hook E. Reported church attendance at the time of entry into HIV care is associated with viral load suppression at 12 months. *AIDS Behav*. 2016;20(8):1706–12.
98. Weiser SD, Yuan C, Guzman D, Frongillo EA, Riley ED, Bangsberg DR, et al. Food insecurity and HIV clinical outcomes in a longitudinal study of urban homeless and marginally housed HIV-infected individuals. *AIDS*. 2013;27(18):2953–8.
99. Whelan BM, Hebert PL, Ahrens KR, Katz DA, Buskin SE, Golden MR, et al. Predictors of failure to reach viral suppression within 1 year after human immunodeficiency virus diagnosis: a surveillance-based analysis. *Sex Transm Dis*. 2019;46(11):728–32.
100. Wilson EC, Turner C, Arayasirikul S, Woods T, Nguyen T, Lin R, et al. Housing and income effects on HIV-related health outcomes in the San Francisco Bay Area: findings from the SPNS transwomen of color initiative. *AIDS Care*. 2018;30(11):1356–9.
101. Yehia BR, Schranz AJ, Momplaisir F, Keller SC, Gross R, Frank I, et al. Outcomes of HIV-infected patients receiving care at multiple clinics. *AIDS Behav*. 2014;18(8):1511–22.
102. Abgrall S, Fugon L, Lele N, Carde E, Bentata PO, et al. Risk factors for adherence failure in HIV-infected sub-Saharan migrants living in France and travelling back to their native country. *Int J STD AIDS*. 2014;25(5):389–90.
103. Burch LS. Impact of gender, sexual orientation and socio-economic factors on HIV treatment outcomes in the UK [Internet] [Thesis (Doctoral)]. University of London, University College London (United Kingdom). <https://discovery.ucl.ac.uk/id/eprint/10041026> (2018).
104. Carrieri M, Chesney M, Spire B, Loundou A, Sobel A, Lepou G, et al. Failure to maintain adherence to HAART in a cohort of French HIV-positive injecting drug users. *Int J Behav Med*. 2003;10(1):1–14.
105. Collazos J, Asensi V, Carton JA, Ibarra S. The influence of the patients' educational levels on socioeconomic,

- clinical, immunological and virological endpoints. *AIDS Care*. 2009;21(4):511–9.
106. D'Almeida KW, Lert F, Spire B, Dray-Spira R. Determinants of virological response to antiretroviral therapy: socio-economic status still plays a role in the era of cART Results from the ANRS-VESPA 2 study, France. *Antivir Ther*. 2016;21(8):661–70.
 107. Del Amo on behalf of The Socio-economic Inequalities and HIV Working Group for Collaboration of Observational HIV Epidemiological Research in Europe (COHERE) in Euro-Coord. Inequalities by educational level in response to combination antiretroviral treatment and survival in HIV-positive men and women in Europe. *AIDS*. 2017;31(2):253–62.
 108. Dorz S, Lazzarini L, Cattelan A, Meneghetti F, Novara C, Concia E, et al. Evaluation of adherence to antiretroviral therapy in Italian HIV patients. *AIDS Patient Care STDs*. 2003;17(1):33–41.
 109. Gordillo V, del Amo J, Soriano V, Gonzalez-Lahoz J. Sociodemographic and psychological variables influencing adherence to antiretroviral therapy. *AIDS*. 1999;13(13):1763–9.
 110. Gueler A, Schoeni-Affolter F, Moser A, Bertisch B, Bucher HC, Calmy A, et al. Neighbourhood socio-economic position, late presentation and outcomes in people living with HIV in Switzerland. *AIDS (London, England)*. 2015;29(2):231–8.
 111. Jansen K, Brockmeyer NH, Dlugay V, Stoll M, Pauli R, Golz J, et al. Poorer clinical outcomes of HIV-positive patients (PLWHA) having lower income: results of an analysis on the basis of the KompNet cohort. *HIV Med*. 2009;10(SUPPL. 2):153.
 112. Papadopoulou A. Adherence to combination therapies in people with HIV/AIDS [Internet]. [Thesis (Doctoral)]. University of London, University College London (United Kingdom). <https://discovery.ucl.ac.uk/id/eprint/10105419/> (2000)
 113. Parruti G, Manzoli L, Marani TP, D'Amico G, Rotolo S, Graziani V, et al. Long-term adherence to first-line highly active antiretroviral therapy in a hospital-based cohort: predictors and impact on virologic response and relapse. *AIDS Patient Care STDs*. 2006;20(1):48–57.
 114. Persson L, Gullberg B, Hanson BS, Moestrup T, Ostergren PO. HIV infection: social network, social support, and CD4 lymphocyte values in infected homosexual men in Malmo, Sweden. *J Epidemiol Community Health*. 1994;48(6):580–5.
 115. Raho-Moussa M, Guiguet M, Michaud C, Honoré P, Palacios C, Boué F, et al. Respective roles of migration and social deprivation for virological non-suppression in HIV-infected adults on antiretroviral therapy in France. *PLoS ONE*. 2019;14(3):e0213019.
 116. Saracino A, Zaccarelli M, Lorenzini P, Bandera A, Marchetti G, Castelli F, et al. Impact of social determinants on antiretroviral therapy access and outcomes entering the era of universal treatment for people living with HIV in Italy. *BMC Public Health*. 2018;18(1):870.
 117. Sellier P, Clevenbergh P, Ljubicic L, Simoneau G, Evans J, Delcey V, et al. Comparative evaluation of adherence to antiretroviral therapy in sub-Saharan African native HIV-infected patients in France and Africa. *Clin Infect Dis*. 2006;43(5):654–7.
 118. Sherr L, Clucas C, Lampe F, Harding R, Johnson M, Fisher M, et al. Gender and mental health aspects of living with HIV disease and its longer-term outcomes for UK heterosexual patients. *Women Health*. 2012;52(3):214–33.
 119. Sobrino-Vegas P, Rodriguez-Urrego J, Berenguer J, Caro-Murillo AM, Blanco JR, Viciano P, et al. Educational gradient in HIV diagnosis delay, mortality, antiretroviral treatment initiation and response in a country with universal health care. *Antivir Ther*. 2012;17(1):1–8.
 120. Spire B, Duran S, Souville M, Leport C, Raffi F, Moatti J-P, et al. Adherence to highly active antiretroviral therapies (HAART) in HIV-infected patients: from a predictive to a dynamic approach. *Soc Sci Med*. 2002;54(10):1481–96.
 121. Uuskula A, Laisaar K-T, Raag M, Smidt J, Semjonova S, Kogan J, et al. Antiretroviral therapy (ART) adherence and correlates to nonadherence among people on ART in Estonia. *AIDS Care*. 2012;24(12):1470–9.
 122. Siefried KJ, Mao L, Kerr S, Cysique LA, Gates TM, McAllister J, et al. Socioeconomic factors explain suboptimal adherence to antiretroviral therapy among HIV-infected Australian adults with viral suppression. *PLoS ONE*. 2017;12(4):e0174613.
 123. Yang HJ, Bang JH. Factors associated with medication adherence in patients with human immunodeficiency virus in South Korea. *AIDS Care*. 2017;29(10):1315–9.
 124. Aidala AA, Wilson MG, Shubert V, Gogolishvili D, Globerman J, Rueda S, et al. Housing status, medical care, and health outcomes among people living with HIV/AIDS: a systematic review. *Am J Public Health*. 2015;106(1):e1–23.
 125. Leaver CA, Bargh G, Dunn JR, Hwang SW. The effects of housing status on health-related outcomes in people living with HIV: a systematic review of the literature. *AIDS Behav*. 2007;11(2):85–100.
 126. Singer AW, Weiser SD, McCoy SI. Does food insecurity undermine adherence to antiretroviral therapy? A systematic review. *AIDS Behav*. 2015;19(8):1510–26.
 127. Adimora AA, Ramirez C, Potat T, Archin NM, Averitt D, Auerbach JD, et al. HIV and women in the USA: what we know and where to go from here. *Lancet*. 2021;397(10279):1107–15.
 128. Scambler G. Sociology, social class, health inequalities, and the avoidance of “Classism.” *Front Sociol*. 2019;4:56.
 129. Bourdieu P. The forms of capital. In: Apple MW, Ball SJ, Gandin LA, editors. *Handbook of theory and research for the sociology of education*. Westport: Greenwood Press; 1986. p. 241–58.
 130. Black SD. Inequalities in health: report of a Research Working Group. Department of Health and Social Security; 1980.
 131. Moskalewicz A, Oremus M. No clear choice between Newcastle-Ottawa Scale and Appraisal Tool for Cross-Sectional Studies to assess methodological quality in cross-sectional studies of health-related quality of life and breast cancer. *J Clin Epidemiol*. 2020;120:94–103.
 132. Bonnefoy J, Morgan A, Kelly MP, Butt J, Bergman V. Constructing the evidence base on the social determinants of health: a guide. Measurement and Evidence Knowledge Network (MEKN); 2007.
 133. Singh-Manoux A, Adler NE, Marmot MG. Subjective social status: its determinants and its association with measures of ill-health in the Whitehall II study. *Soc Sci Med*. 2003;56(6):1321–33.
 134. Quinn KG, DiFranceis W, Spector A, Bendixen A, Peters A, Dickson-Gomez J. The effect of various supportive housing models on ART adherence among persons living with HIV in Supportive housing. *Med Care*. 2021;59:S124.
 135. Holtgrave DR, Briddell K, Little E, Bendixen AV, Hooper M, Kidder DP, et al. Cost and threshold analysis of housing as an HIV prevention intervention. *AIDS Behav*. 2007;11(2):162–6.
 136. Holtgrave DR, Wolitski RJ, Pals SL, Aidala A, Kidder DP, Vos D, et al. Cost-utility analysis of the housing and health intervention for homeless and unstably housed persons living with HIV. *AIDS Behav*. 2013;17(5):1626–31.
 137. National Institute for Health and Care Excellence. Guide to the methods of technology appraisal. p. 35. <http://nice.org.uk/process/pmg9> (2013)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.