

eCommons@AKU

Population Health, East Africa

Medical College, East Africa

11-2021

Invited Perspective: Most Affected by Climate Change; Least Studied

Kristie L. Ebi University of Washington, USA

Stanley Luchters Aga Khan University, stanley.luchters@aku.edu

Follow this and additional works at: https://ecommons.aku.edu/eastafrica_fhs_mc_popul_health



Part of the Environmental Public Health Commons

Recommended Citation

Ebi, K. L., Luchters, S. (2021). Invited Perspective: Most Affected by Climate Change; Least Studied. Environmental Health Perspectives, 129(11), 1-2.

Available at: https://ecommons.aku.edu/eastafrica_fhs_mc_popul_health/121

Invited Perspective

Invited Perspective: Most Affected by Climate Change; Least Studied

Kristie L. Ebi¹ and Stanley Luchters²

¹Center for Health and the Global Environment, University of Washington, Seattle, Washington, USA

²Department of Population Health, Aga Khan University, Nairobi, Kenya

https://doi.org/10.1289/EHP10384

Refers to https://doi.org/10.1289/EHP9391

Each country that is a signatory to the Paris Agreement under the United Nations Framework Convention on Climate Change (UNFCCC) produces a nationally determined contribution (NDC), outlining national plans for mitigating and adapting to climate change. The NDCs of low- and middle-income countries (LMICs) have greater coverage of the health risks of and responses to climate change, in terms of risks, adaptation, mitigation, co-benefits, and trade-offs, whereas higher-income countries tend to focus on energy, the economy, and related sectors (Dasandi et al. 2021). This difference reflects the expectation that the largest health risks of climate change will occur in LMICs, where rising ambient temperatures, changing precipitation patterns, and rising sea levels are increasingly affecting air, water, and food systems. Such changes will affect the geographic and seasonal patterns of major causes of morbidity and mortality—particularly undernutrition, diarrheal diseases, malaria, dengue, and injuries-and will disproportionately affect the most vulnerable population groups, such as the lowest-income families, older individuals, pregnant women, and neonates (Smith et al. 2014; Watts et al. 2015). The timeliness and effectiveness of adaptation and mitigation efforts will determine the extent of increase in risks with additional climate change (Ebi and Hess 2020). There is a critical need for more evidence on effective locally led adaptation and mitigation

Although substantial proportions of current and future climate-related adverse health outcomes are preventable, research and implementation funding are well below what is needed to minimize morbidity and mortality (Ebi et al. 2016; Green et al. 2017), with very little published research evidence from LMICs. Using machine learning, a comprehensive global mapping of Englishlanguage research published 1 January 2013-9 April 2020 predicted there were nearly 16,000 studies on climate and health (Berrang-Ford et al. 2021). Overall, 79% of studies that identified place names focused on high- and upper middle-income nations, particularly China. The numbers of publications from high-income countries were two times greater than from lower middle-income countries and close to 10 times those from low-income nations. For example, focusing just on the impact of high ambient temperatures on health, a review of epidemiological studies in LMICs from January 1980 to August 2018 identified 146 studies, of which 82 were conducted in China, 9 in other countries of East Asia and the Pacific, 12 in South Asia, 10 in sub-Saharan Africa, 9 in the

Address correspondence to Kristie L. Ebi, Center for Health and the Global Environment, University of Washington, Seattle, WA 98105 USA. Email: krisebi@uw.edu

Note to readers with disabilities: EHP strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in EHP articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Middle East and North Africa, and 7 each in Latin America and Europe (Green et al. 2019). More than 92% reported positive associations, noting that older people, women, and individuals within low socioeconomic brackets were the most vulnerable. Developing efficient and effective heat action plans is urgently needed yet requires more robust evidence.

Burrows et al. (2021) contributes to the meager evidence of the extent to which climate change is affecting the mental well-being of populations in LMICs. In this study, individuals displaced by landslides were more likely than those not displaced to self-report improvements in several aspects of mental well-being. Although not providing the final word, the research importantly adds to our understanding of the complex nature, duration, and extent to which landslides and other climate- and weather-related events affect health and well-being in these communities, directly and indirectly. More, and more rigorous, research is needed to create a deeper understanding of the challenges and opportunities to effectively plan for a world with more landslides with additional climate change.

The broad research agenda in climate change and health includes quantifying the burden of climate-sensitive health outcomes, and projecting risks at spatial and temporal scales useful for decision-making, under a range of climate and development scenarios. Research needs to be conducted in the context of other drivers of climate-sensitive health outcomes, such as population growth and aging, urbanization, and socioeconomic change. Climate change can also disrupt provision of and access to health care, with consequences for human health. Locally relevant adaptation and mitigation interventions, especially protecting the most vulnerable, need to be developed and evaluated. Research is needed to inform developing, implementing, and evaluating strategies, policies, and programs to proactively prepare for and manage health risks that will evolve with climate change and development. The health co-benefits of mitigation policies and technologies also need to be quantified.

The UNFCCC recognizes climate change as a problem with common but differentiated responsibilities; everyone is responsible for climate change (United Nations 1992). Because the impacts are inequitably distributed across communities and countries, high-income countries agreed to lead on providing resources to support adaptation and mitigation. Funding increases for climate change and health research from Wellcome Trust, Horizon Europe, the U.S. National Institutes of Health, and others should develop and strengthen research capacity in LMICs and build it through South–South and South–North partnerships, knowledge exchanges, and research implementation. The need is urgent because climate change is already damaging health and well-being, with greater impacts on vulnerable populations in LMICs who have contributed the least to the largest challenge of this century.

References

Berrang-Ford L, Sietsma AJ, Callaghan M, Minx JC, Scheelbeek PFD, Haddaway NR, et al. 2021. Systematic mapping of global research on climate and health: a machine learning review. Lancet Planet Health 5(8):e514–e525, PMID: 34270917, https://doi.org/10.1016/S2542-5196(21)00179-0.

The authors declare they have no actual or potential competing financial interests.

Received 24 September 2021; Revised 30 September 2021; Accepted 4 October 2021; Published 8 November 2021.

- Burrows K, Pelupessy DC, Khoshnood K, Bell ML. 2021. Environmental displacement and mental well-being in Banjarnegara, Indonesia. Environ Health Perspect 129(11):117002, https://doi.org/10.1289/EHP9391.
- Dasandi N, Graham H, Lampard P, Jankin Mikhaylov S. 2021. Engagement with health in national climate change commitments under the Paris Agreement: a global mixed-methods analysis of the nationally determined contributions. Lancet Planet Health 5(2):e93-e101, PMID: 33581071, https://doi.org/10.1016/ S2542-5196(20)30302-8.
- Ebi KL, Hess JJ. 2020. Health risks due to climate change: inequity in causes and consequences. Health Aff (Millwood) 39(12):2056–2062, PMID: 33284705, https://doi.org/10.1377/hlthaff.2020.01125.
- Ebi KL, Semenza JC, Rocklöv J. 2016. Current medical research funding and frameworks are insufficient to address the health risks of global environmental change. Environ Health 15(1):108, PMID: 27835959, https://doi.org/10.1186/ s12940-016-0183-3.
- Green H, Bailey J, Schwarz L, Vanos J, Ebi K, Benmarhnia T. 2019. Impact of heat on mortality and morbidity in low and middle income countries: a review of the epidemiological evidence and consideration for future research. Environ

- Res 171:80–91, PMID: 30660921, https://doi.org/10.1016/j.envres.2019.01.
- Green D, Pitman A, Barnett A, Kaldor J, Doherty P, Stanley F. 2017. Advancing Australia's role in climate change and health research. Nat Clim Chang 7(2):103–106, https://doi.org/10.1038/nclimate3182.
- Smith KR, Woodward A, Campbell-Lendrum D, Chadee DD, Honda Y, Liu Q, et al. 2014. Human health: impacts, adaptation, and co-benefits. In: Climate Change 2014: Impacts, Adaptation, and Vulnerability Part A: Global and Sectoral Aspects Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, et al., eds. New York, NY: Cambridge University Press, 709–754.
- United Nations. 1992. United Nations Framework Convention on Climate Change. FCCC/INFORMAL/84. GE.05-62220 (E) 200705. https://unfccc.int/resource/docs/convkp/conveng.pdf [accessed 30 September 2021].
- Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P, Cai W, et al. 2015. Health and climate change: policy responses to protect public health. Lancet 386(10006):P1861–P1914, https://doi.org/10.1016/S0140-6736(15)60854-6.