



LUND UNIVERSITY

Experimental assessment of aerosolized murine noroviruses

Alsved, Malin; Widell, Anders; Dahlin, Henrik; Medstrand, Patrik; Löndahl, Jakob

2019

[Link to publication](#)

Citation for published version (APA):

Alsved, M., Widell, A., Dahlin, H., Medstrand, P., & Löndahl, J. (2019). *Experimental assessment of aerosolized murine noroviruses*. Abstract from Swedish National Infection Biology Meeting, Sweden.

Total number of authors:

5

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Experimental assessment of infectivity of aerosolized murine noroviruses

Malin Alsved¹, Anders Widell², Henrik Dahlin¹, Patrik Medstrand², Jakob Löndahl¹

¹Ergonomics and Aerosol Technology, Lund University, Sweden

²Translational Medicine, Lund University, Sweden

Introduction

Noroviruses are the cause for the majority of viral gastroenteritis cases in the world. Due to their low infectious dose and long stability in the environment, they often gives rise to outbreaks, even in hospital wards where legitimate infection control measures are practiced by the staff. Resent research has shown that noroviruses are present in air during outbreaks (Bonifait et al. 2014); however more research is needed to investigate the ability of noroviruses to transmit disease via the airborne route.

Method

In this study, a method for evaluating viral infectivity after aerosolization was developed. As there is no well-established cell-cultivation technique for human noroviruses, a murine norovirus (MNV) strain was used as model virus in this study. MNV was aerosolized from a liquid suspension into a flow tube (Figure 1) and collected into phosphate buffer saline (PBS) after 10 seconds as airborne particles. MNV infectivity after aerosolization was assessed by detection of intracellular negative sense RNA in RAW 264.7 cells after 24 h incubation. The complementary negative sense RNA strand of the MNV genome is only present during active replication and can therefore be used as an indication of infection. The collected aerosol samples were added in 10-fold end point dilution series to 24-well cell culture plates. After 24 hours, the supernatant was removed, cells lysed, RNA was extracted, and negative sense RNA detected by a tagged primer according to the method in Vashist et al. (2012).

Result

MNV preserved its infectivity after aerosolization, shown by the detection of negative sense specific RT-qPCR in the cell infectivity assay. With a dilution factor in the aerosolisation setup of about 500 times, the viral titers in the collected samples were relatively low, and the infectivity incubation time was therefore optimized for those, resulting in 24 h. Our aerosolization methodology can be further applied to other virus types, or to evaluate the effect of specific parameters in the air environment, such as temperature or humidity.




Figure 1. Experimental setup for aerosolization of murine noroviruses (by atomizer or sparging liquid aerosol generator) into a flow tube, and collection into PBS using a BioSampler for further molecular analysis.

References

Bonifait L, Charlebois R, Vimont A, et al. (2015). Clin Infect Dis, 61(3), ss 299-304.
Vashist, S., Urena, L. & Goodfellow, I. (2012). J Virol Methods, 184(1-2), ss. 69-76.