



TITLE:

# COHOMOLOGY OF IGUSA CURVES -- A SURVEY (Automorphic forms, Automorphic representations, Galois representations, and its related topics)

AUTHOR(S):

SHIN, SUG WOO

---

CITATION:

SHIN, SUG WOO. COHOMOLOGY OF IGUSA CURVES -- A SURVEY (Automorphic forms, Automorphic representations, Galois representations, and its related topics). 数理解析研究所講義録 2021, 2204: 166-176

ISSUE DATE:

2021-12

URL:

<http://hdl.handle.net/2433/267818>

RIGHT:

## COHOMOLOGY OF IGUSA CURVES – A SURVEY

SUG WOO SHIN

ABSTRACT. We illustrate the strategy to compute the  $\ell$ -adic cohomology of Igusa varieties in the setup of ordinary modular curves, with updates on the literature towards a generalization.

## 1. INTRODUCTION

Igusa curves were introduced by Igusa [Igu68] to understand the mod  $p$  geometry of modular curves when the level is divisible by  $p$ , for each prime  $p$ . As a generalization, we have Igusa varieties in the setup of Shimura varieties of Hodge type thanks to [HT01, Man05, Ham17, Zha, HK19]. Igusa varieties shed light on the mod  $p$  and  $p$ -adic geometry of Shimura varieties via the so-called product structure. Moreover they play a vital role in the applications to the Langlands correspondence, vanishing results on the cohomology of Shimura varieties, and  $p$ -adic automorphic forms. We refer to [KS, §1] for a detailed introduction to Igusa varieties and further references. For an application to (an extension of) the Kottwitz conjecture on the cohomology of Rapoport–Zink spaces, see [Shi12, BM].

A fundamental problem on Igusa varieties is to compute their  $\ell$ -adic cohomology (with or without compact support) for primes  $\ell \neq p$ . To this end, the Langlands–Kottwitz (LK) method for Shimura varieties has been adapted to Igusa varieties in [HT01, Shi09, MC21], at least when the level structure at  $p$  is hyperspecial. (In [HT01], one can go a little further.) While there are excellent expositions<sup>1</sup> on the LK method for modular curves (with good reduction mod  $p$ ) by Clozel [Clo93, §3] and Scholze [Sch11, §5], in addition to Langlands’s original papers [Lan73, Lan76], there is no counterpart for Igusa curves. The goal of this article is to spell out the LK method for Igusa curves in a somewhat informal style, thereby to serve as a friendly entry point for the subject.

**Acknowledgments.** The author was partially supported by NSF grant DMS-1802039 and NSF RTG grant DMS-1646385. He is grateful to Takuya Yamauchi and Kazuki Morimoto for inviting him to speak on cohomology of Igusa varieties at the RIMS conference “Automorphic forms, Automorphic representations, Galois representations, and its related topics” in January 25–29, 2021. The present article works out the details of that talk in the simpler setup of elliptic modular curves. The author thanks Yoichi Mieda for giving him an opportunity to present the contents as part of the “Berkeley-Tokyo lectures on Number Theory”, January 12–15, 2021.

**Notation and Conventions.** When  $R$  is a commutative ring with unity, we often use  $R$  to mean  $\text{Spec}R$  when there is no danger of confusion. For example, a scheme  $X$  over  $R$  means a scheme over  $\text{Spec}R$ , and  $X \times_R S$  means  $X \times_{\text{Spec}R} \text{Spec}S$  when a ring homomorphism  $R \rightarrow S$  is given. By (Set) (resp. (Sch/ $R$ )) we denote the category of sets (resp. schemes over  $R$ ). We also write  $X_S$  for  $X \times_R S$  if  $R \rightarrow S$  is clear from the context. Similarly if  $X$  is a scheme, we write (Sch/ $X$ ) for the

---

*Date:* July 4, 2021.

<sup>1</sup>There are also several valuable surveys on the LK method for more general Shimura varieties with different emphases, such as [BR94, Clo93, GN09, Zhu20].

category of schemes over  $X$ . Write  $\widehat{\mathbb{Z}}^p := \varprojlim_{(N,p)=1} \mathbb{Z}/N\mathbb{Z}$  and  $\mathbb{A}^{\infty,p} := \widehat{\mathbb{Z}}^p \otimes_{\mathbb{Z}} \mathbb{Q}$  for the ring of adèles away from  $\infty, p$ . By  $C_c^\infty(X)$ , we mean the space of smooth compactly supported functions on a locally compact group  $X$  (with values in  $\mathbb{C}$  or  $\overline{\mathbb{Q}}_\ell$ ).

2. MODULAR CURVES

Let  $N \in \mathbb{Z}_{\geq 3}$ . Consider the moduli functor

$$Y_N : (\text{Sch}/\mathbb{Z}[1/N]) \longrightarrow (\text{Set})$$

sending  $S$  to the set of isomorphism classes of pairs  $(E, \alpha)$ , where  $E$  is an elliptic curve over  $S$ , and  $\alpha : (\mathbb{Z}/N\mathbb{Z})^2 \xrightarrow{\sim} E[N]$  is an isomorphism of group schemes over  $S$ .

**Theorem 2.1** (Igusa, Deligne–Rapoport). *The functor  $Y_N$  is represented by a smooth affine curve over  $\mathbb{Z}[1/N]$ .*

We keep writing  $Y_N$  for the curve it represents. Denote by  $\mathcal{E} \rightarrow Y_N$  the universal elliptic curve. Consider the following inverse limit

$$Y_{\mathbb{C}} := \varprojlim_{N \geq 3} Y_N \times_{\mathbb{Z}[1/N]} \mathbb{C},$$

which exists in the category of  $\mathbb{C}$ -schemes as the transition maps are finite étale. We have

$$\pi_0(Y_{\mathbb{C}}) := \varprojlim_{N \geq 3} \pi_0(Y_{N,\mathbb{C}}) = \varprojlim_{N \geq 3} (\mathbb{Z}/N\mathbb{Z})^\times = \widehat{\mathbb{Z}}^\times. \tag{2.1}$$

From now on, fix a prime  $p$  once and for all. We restrict the level  $N$  to integers coprime to  $p$ . Recall that an elliptic curve  $E$  over a field  $k$  of characteristic  $p$  is said to be **supersingular** if  $\#E[p](\overline{k}) = 1$ . Otherwise  $E$  is said to be **ordinary**, in which case  $\#E[p](\overline{k}) = p$ . Accordingly we have a partition of the topological space

$$Y_{N,\mathbb{F}_p} = Y_{N,\mathbb{F}_p}^{\text{ord}} \coprod Y_{N,\mathbb{F}_p}^{\text{ss}}, \tag{2.2}$$

where  $Y_{N,\mathbb{F}_p}^{\text{ord}}$  (resp.  $Y_{N,\mathbb{F}_p}^{\text{ss}}$ ) is the subset of  $x \in Y_{N,\mathbb{F}_p}$  such that the fiber  $\mathcal{E}_x$  is an ordinary (resp. supersingular) elliptic curve. Thus we can view  $Y_{N,\mathbb{F}_p}^{\text{ord}}$  as an open subscheme of  $Y_{N,\mathbb{F}_p}$  (and  $Y_{N,\mathbb{F}_p}^{\text{ss}}$  as a closed 0-dimensional subscheme). As  $N \in \mathbb{Z}_{\geq 3}$  varies over prime-to- $p$  integers, the transition maps are finite étale and compatible with the partition (2.2).

In this survey we will concentrate on the ordinary case though there is a parallel story in the supersingular case.

*Remark 2.2.* The stratification (2.2) admits a vast generalization to general Shimura varieties. The reader is referred to excellent articles such as [Man20, HR17].

3. IGUSA CURVES

We keep fixing a prime  $p$  and let  $N \geq 3$  be an integer coprime to  $p$ . We still write  $\mathcal{E}$  for the universal elliptic curve over  $Y_{N,\mathbb{F}_p}^{\text{ord}}$ . For each integer  $m \geq 1$ , we have the slope filtration  $0 \rightarrow \mathcal{E}[p^m]^\circ \rightarrow \mathcal{E}[p^m] \rightarrow \mathcal{E}[p^m]^{\text{ét}} \rightarrow 0$  such that  $\mathcal{E}[p^m]^{\text{ét}}$  is the maximal étale quotient. We introduce the Igusa functor of level  $Np^m$  as

$$\text{Ig}_{N,m}^{\text{ord}} : (\text{Sch}/Y_{N,\mathbb{F}_p}^{\text{ord}}) \longrightarrow (\text{Set}), \quad S \mapsto \{(j_m^{\text{ét}}, j_m^\circ)\}, \tag{3.1}$$

where  $j_m^{\text{ét}} : \mathbb{Z}/p^m\mathbb{Z} \xrightarrow{\sim} \mathcal{E}[p^m]_S^{\text{ét}}$  and  $j_m^\circ : \mu_{p^m} \xrightarrow{\sim} \mathcal{E}[p^m]_S^\circ$  are isomorphisms of groups schemes over  $S$ . A fundamental theorem by Igusa (reproduced by Katz–Mazur) is the following.

**Theorem 3.1.** *The functor  $\mathrm{Ig}_{N,m}^{\mathrm{ord}}$  is represented by a scheme, which is an étale  $\mathrm{GL}_1(\mathbb{Z}/p^m\mathbb{Z}) \times \mathrm{GL}_1(\mathbb{Z}/p^m\mathbb{Z})$ -torsor over  $Y_{N,\mathbb{F}_p}^{\mathrm{ord}}$ .*

As  $N \in \mathbb{Z}_{\geq 3}$  and  $m \in \mathbb{Z}_{\geq 0}$  vary,  $\{\mathrm{Ig}_{N,m}^{\mathrm{ord}}\}$  forms a projective system with finite étale transition maps, equipped with a prime-to- $p$  Hecke action of  $\mathrm{GL}_2(\mathbb{A}^{\infty,p})$  defined in the same way for modular curves. Define a  $\mathbb{Q}_p$ -group  $J := \mathrm{GL}_1 \times \mathrm{GL}_1$ , so that  $J(\mathbb{Q}_p) = \mathbb{Q}_p^\times \times \mathbb{Q}_p^\times$  is the automorphism group of  $\mathbb{Q}_p/\mathbb{Z}_p \times \mu_{p^\infty}$  in the isogeny category of  $p$ -divisible groups. The obvious action of  $\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times$  on  $\mathrm{Ig}_{N,m}^{\mathrm{ord}}$  by translating  $(j_m^{\mathrm{ét}}, j_m^\circ)$  induces an action on

$$H_c^i(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) := \varprojlim_{\substack{N \geq 3, \\ m \geq 1, \\ (N,p)=1}} H_c^i(\mathfrak{I}\mathfrak{g}_{N,m,\mathbb{F}_p}^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell), \quad i \geq 0, \tag{3.2}$$

which uniquely extends to an action of  $J(\mathbb{Q}_p)$ . (Mantovan [Man05] proved that the action extends. The same also follows from Caraiani–Scholze’s approach [CS17, CS].) The  $J(\mathbb{Q}_p)$ -action turns out to commute with the  $\mathrm{GL}_2(\mathbb{A}^{\infty,p})$ -action. Finite-dimensionality of cohomology for each  $N$  and  $m$  tells us that  $H_c^i(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell)$  is an admissible representation of  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ . Now we can state the goal of this article:

**Goal:** Compute (3.2) as a representation of  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ .

Caraiani and Scholze [CS17, CS] defined another version of Igusa varieties directly at level  $Np^\infty$ . In our case, their definition specializes to the following functor, where  $(\mathrm{Perf}/Y_{N,\mathbb{F}_p}^{\mathrm{ord}})$  means the category of perfect schemes over  $Y_{N,\mathbb{F}_p}^{\mathrm{ord}}$ :

$$\mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ord}} : (\mathrm{Perf}/Y_{N,\mathbb{F}_p}^{\mathrm{ord}}) \longrightarrow (\mathrm{Set}),$$

sending  $S$  to the set of isomorphisms  $\mathbb{Q}_p/\mathbb{Z}_p \times \mu_{p^\infty} \xrightarrow{\sim} \mathcal{E}[p^\infty]_S$  between  $p$ -divisible groups over  $S$ . This can be compared with the scheme

$$\mathrm{Ig}_{N,\infty}^{\mathrm{ord}} := \varprojlim_{m \geq 1} \mathrm{Ig}_{N,m}^{\mathrm{ord}},$$

where the “level-decreasing” transition maps are finite étale. As  $\mathrm{Ig}_{N,\infty}^{\mathrm{ord}}$  form a projective system with finite étale transition maps as  $N$  varies, we can take the limit scheme  $\mathrm{Ig}_\infty^{\mathrm{ord}}$ .

**Theorem 3.2** (Caraiani–Scholze). *The functor  $\mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ord}}$  is represented by a perfect scheme over  $Y_{N,\mathbb{F}_p}^{\mathrm{ord}}$  and canonically isomorphic to the perfection of  $\mathrm{Ig}_{N,\infty}^{\mathrm{ord}}$ .*

Since perfection does not affect topological information such as étale cohomology or the set of  $\overline{\mathbb{F}}_p$ -points, we can use either  $\mathrm{Ig}_{N,\infty}^{\mathrm{ord}}$  or  $\mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ord}}$ . Since the former is built out of finite-level Igusa curves, it is useful for applying a fixed point formula. On the other hand,  $\mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ord}}$  is a little more convenient for defining group actions and describing the  $\overline{\mathbb{F}}_p$ -points.

*Remark 3.3.* We can define  $\mathrm{Ig}_{N,\infty}^{\mathrm{ss}}$  and  $\mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ss}}$  in analogy with the ordinary case. Then  $\mathrm{Ig}_{N,\infty}^{\mathrm{ss}}$  is already perfect and  $\mathrm{Ig}_{N,\infty}^{\mathrm{ss}} = \mathfrak{I}\mathfrak{g}_{N,\infty}^{\mathrm{ss}}$ .



4.  $\overline{\mathbb{F}}_p$ -POINTS OF IGUSA CURVES

In order to achieve the aforementioned goal via a fixed-point formula, we need to describe the set of  $\overline{\mathbb{F}}_p$ -points of Igusa curves

$$\mathfrak{I}g_{\infty}^{\text{ord}}(\overline{\mathbb{F}}_p) = \varprojlim_{(N,p)=1, N \geq 3} \mathfrak{I}g_{\infty}^{\text{ord}}(\overline{\mathbb{F}}_p) = \varprojlim_{(N,p)=1, N \geq 3} \text{Ig}_{\infty}^{\text{ord}}(\overline{\mathbb{F}}_p)$$

with the  $\text{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -action.

Let us set up some more notation. Write  $\check{\mathbb{Z}}_p := W(\overline{\mathbb{F}}_p)$  and  $\check{\mathbb{Q}}_p := W(\overline{\mathbb{F}}_p)[1/p]$ . Denote by  $\text{Ell}^0$  the set of isogeny classes of elliptic curves over  $\overline{\mathbb{F}}_p$ . Those of ordinary elliptic curves define a subset  $\text{Ell}^{0,\text{ord}}$ . We identify  $\text{Ell}^{0,\text{ord}}$  with a set of representatives by fixing a representative in each isogeny class.

When  $E$  is an elliptic curve over  $\overline{\mathbb{F}}_p$ , define

- $I(E) := (\text{End}_{\overline{\mathbb{F}}_p}(E) \otimes_{\mathbb{Z}} \mathbb{Q})^{\times}$ ,
- $T^p(E) := \varprojlim_{(N,p)=1} E[N](\overline{\mathbb{F}}_p)$ ,
- $\check{T}_p(E)$  to be the covariant Dieudonné module of  $E[p^{\infty}]$ .

As we are concerned with the ordinary case,  $I(E) = F^{\times}$  for an imaginary quadratic field  $F$ . (As an algebraic group over  $\mathbb{Q}$ ,  $I(E) = \text{Res}_{F/\mathbb{Q}} \text{G}_m$ .) A standard fact is that  $T^p E$  is a free  $\widehat{\mathbb{Z}}^p$ -module of rank 2, and  $\check{T}_p E$  is free of rank 2 over  $\check{\mathbb{Z}}_p$  (which is the  $\check{\mathbb{Z}}_p$ -linear dual of  $H_{\text{cris}}^1(E/\check{\mathbb{Z}}_p)$ ). At  $p$ , we have the extra structure of semi-linear maps  $F^{-1}, V^{-1}$  on  $\check{T}_p E$  such that  $F^{-1}V^{-1} = V^{-1}F^{-1} = p$ . (The minus sign comes from the covariant convention.) It is useful to think of  $T^p E$  as a  $\widehat{\mathbb{Z}}^p$ -lattice in the free  $\mathbb{A}^{\infty,p}$ -module  $V^p E := T^p E \otimes_{\mathbb{Z}} \mathbb{Q}$  of rank 2. Similarly  $\check{T}_p E$  is an  $F^{-1}, V^{-1}$ -invariant lattice in  $\check{V}_p E := \check{T}_p E \otimes_{\mathbb{Z}} \mathbb{Q}$ . (We have linear extensions of  $F^{-1}, V^{-1}$  to self-bijections on  $\check{V}_p E$ .)

Now we start our analysis of  $\overline{\mathbb{F}}_p$ -points from (3.1).

$$\begin{aligned} \mathfrak{I}g_{\infty}^{\text{ord}}(\overline{\mathbb{F}}_p) &= \left\{ \begin{array}{l} E : \text{elliptic curve}/\overline{\mathbb{F}}_p, \\ \alpha : (\widehat{\mathbb{Z}}^p)^2 \xrightarrow{\sim} T^p E, \\ j : \mathbb{Q}_p/\mathbb{Z}_p \times \mu_{p^{\infty}} \xrightarrow{\sim} E[p^{\infty}] \end{array} \right\} / \simeq \\ &= \prod_{E_0 \in \text{Ell}^{0,\text{ord}}} \left\{ \begin{array}{l} (E, \alpha, j) \text{ as above,} \\ \text{s.t. } \exists \text{ an isogeny } f : E \rightarrow E_0 \end{array} \right\} / \simeq \\ &= \prod_{E_0 \in \text{Ell}^{0,\text{ord}}} I(E_0) \setminus \left\{ \begin{array}{l} (L^p, \phi^p, L_p, \phi_p) : \\ L^p \subset V^p E_0 \text{ is a } \widehat{\mathbb{Z}}^p\text{-lattice, } \phi^p : (\widehat{\mathbb{Z}}^p)^2 \xrightarrow{\sim} L^p, \\ L_p \subset \check{V}_p E_0 \text{ is an } F^{-1}, V^{-1}\text{-invariant } \check{\mathbb{Z}}_p\text{-lattice,} \\ \phi_p : \check{\mathbb{Z}}_p^2 \xrightarrow{\sim} L_p \text{ carries } (1, p^{-1})\sigma \text{ on } \check{\mathbb{Z}}_p^2 \text{ to } F \text{ on } L_p. \end{array} \right\}. \end{aligned}$$

In the last expression,  $\phi^p$  and  $\phi_p$  are respectively  $\widehat{\mathbb{Z}}^p$ -linear and  $\check{\mathbb{Z}}_p$ -linear. Each equality above is natural and equivariant with respect to the natural action of  $G(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ . To see the last equality, one starts from  $(E, \alpha, j)$  and chooses an isogeny  $f : E \rightarrow E_0$ . Then take  $L^p = f(T^p E)$  and  $L_p = f(\check{T}_p E)$ . We leave it as an exercise to give  $\phi^p$  and  $\phi_p$  from  $(E, \alpha, j)$  and to show that the left quotient by  $I(E_0)$  cancels out the choice of  $f$  (so that the quotient set is independent of the choice).

To proceed, we give more convenient parametrizations of  $(L^p, \phi^p)$  and  $(L_p, \phi_p)$ . We describe the right  $\mathrm{GL}_2(\mathbb{A}^{\infty,p})$ -set (which is a torsor for the action)

$$X^p(E_0) := \{(L^p, \phi^p) \text{ as above}\} = \{(\mathbb{A}^{\infty,p})^2 \xrightarrow{\sim} V^p E_0\},$$

where  $\mathrm{GL}_2(\mathbb{A}^{\infty,p})$  acts on the last set through its natural action on  $(\mathbb{A}^{\infty,p})^2$ . To obtain the inverse map, notice that an isomorphism  $(\mathbb{A}^{\infty,p})^2 \xrightarrow{\sim} V^p E_0$  determines  $(L^p, \phi^p)$  by restriction to  $(\widehat{\mathbb{Z}}^p)^2$ . The above identification is also equivariant for the left action of  $I(E_0)$ , which naturally acts on  $V^p E_0$ .

Similarly we have a bijection of right  $J(\mathbb{Q}_p)$ -sets (which are  $J(\mathbb{Q}_p)$ -torsors)

$$\check{X}_p(E_0) := \{(L_p, \phi_p) \text{ as above}\} = \{(\check{\mathbb{Q}}_p)^2 \xrightarrow{\sim} \check{V}_p E_0, \text{ s.t. } (1, p^{-1})\sigma \leftrightarrow F\},$$

where  $J(\mathbb{Q}_p)$  acts as automorphisms of the isocrystal  $((\check{\mathbb{Q}}_p)^2, (1, p^{-1})\sigma)$  associated with the  $p$ -divisible group  $\mathbb{Q}_p/\mathbb{Z}_p \times \mu_{p^\infty}$ . The above equality is equivariant for the left action of  $I(E_0)$ , which acts as automorphisms of the isocrystal  $(\check{V}_p E_0, F)$ .

The progress so far may be summarized as follows. As right  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -sets,

$$\mathfrak{I}\mathfrak{g}_{\infty}^{\mathrm{ord}}(\overline{\mathbb{F}}_p) = \coprod_{E_0 \in \mathrm{Ell}^{0,\mathrm{ord}}} I(E_0) \backslash (X^p(E_0) \times X_p(E_0)). \tag{4.1}$$

By choosing a base point, we can identify the right  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -torsor  $X^p(E_0) \times X_p(E_0)$  with  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$  equipped with an embedding of groups

$$I(E_0) \hookrightarrow \mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p),$$

well defined up to  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -conjugacy. On the other hand, a special case of Honda–Tate theory over  $\overline{\mathbb{F}}_p$  (cf. [HT01, V.2] or [Shi09, §8]) tells us that  $\mathrm{Ell}^{0,\mathrm{ord}}$  is in bijection with the set  $\mathrm{IQF}(p)(p)$  of imaginary quadratic fields (up to isomorphism) in which  $p$  splits, where we assign  $\mathrm{End}(E) \otimes_{\mathbb{Z}} \mathbb{Q}$  (which is an imaginary quadratic field since  $E$  is ordinary) to each  $E \in \mathrm{Ell}^{0,\mathrm{ord}}$ . Thus we can rewrite (4.1) as follows.

**Proposition 4.1.** *As right  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -sets,*

$$\mathfrak{I}\mathfrak{g}_{\infty}^{\mathrm{ord}}(\overline{\mathbb{F}}_p) = \coprod_{F \in \mathrm{IQF}(p)} F^{\times} \backslash (\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)),$$

where the quotient is taken with respect to the embedding of  $I(E_0) = F^{\times}$  above.

*Remark 4.2.* Mack-Crane [MC21] recently obtained the analogue for Igusa varieties in the setup of Hodge-type Shimura varieties with hyperspecial level at  $p$ , generalizing [Shi09] on the PEL case.

### 5. FROM $\overline{\mathbb{F}}_p$ -POINTS TO THE TRACE FORMULA

Before we go from Proposition 4.1 to compute the  $\ell$ -adic cohomology, we need some preparation. Let  $N \geq 3$  and  $m \geq 1$ . Define

$$\begin{aligned} K^p &= K^p(N) := \ker(\mathrm{GL}_2(\widehat{\mathbb{Z}}^p) \rightarrow \mathrm{GL}_2(\widehat{\mathbb{Z}}^p/N\widehat{\mathbb{Z}}^p)) \subset \mathrm{GL}_2(\mathbb{A}^{\infty,p}), \\ K_p &= K_{p,m} := (1 + p^m\mathbb{Z}_p) \times (1 + p^m\mathbb{Z}_p) \subset J(\mathbb{Q}_p). \end{aligned}$$

Then  $\mathrm{Ig}_{N,m}^{\mathrm{ord}} = \mathrm{Ig}_{\infty}^{\mathrm{ord}}/K^p \times K_p$ . Let

$$g^p \in \mathrm{GL}_2(\mathbb{A}^{\infty,p}), \quad g_p = (g_{p,1}, g_{p,2}) \in J(\mathbb{Q}_p) = \mathbb{Q}_p^{\times} \times \mathbb{Q}_p^{\times}.$$

We say that  $g_p$  is *acceptable* if the additive  $p$ -adic valuations satisfy the inequality  $v_p(g_{p,1}) > v_p(g_{p,2})$ .

Write  $\mathbf{1}_{K^p g^p K^p}$  and  $\mathbf{1}_{K_p g_p K_p}$  for the characteristic functions on the corresponding double cosets, viewed as elements of Hecke algebras for  $\mathrm{GL}_2(\mathbb{A}^{\infty,p})$  and  $J(\mathbb{Q}_p)$ , respectively. Let  $[K^p g^p K^p]$  and  $[K_p g_p K_p]$  denote the double coset operators on the set of  $\overline{\mathbb{F}}_p$ -points or cohomology of  $\mathrm{Ig}_{N,m}^{\mathrm{ord}}$ . Denote by  $H_c$  the alternating sum  $\sum_{i \geq 0} (-1)^i H_c^i$  in the Grothendieck group of representations. To achieve the goal stated in §3, we compute

$$\mathrm{tr} \left( \mathbf{1}_{K^p g^p K^p} \times \mathbf{1}_{K_p g_p K_p} \mid H_c(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) \right),$$

which is equal to (the volume of  $K^p \times K_p$  times)

$$\mathrm{tr} \left( [K^p g^p K^p] \times [K_p g_p K_p] \mid H_c(\mathrm{Ig}_{N,m,\overline{\mathbb{F}}_p}^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) \right). \quad (5.1)$$

Let  $\mathrm{Fix}(A|B)$  denote the set of fixed points of an operator  $A$  acting on a mathematical object  $B$ . We apply the fixed-point formula for non-proper varieties à la Fujiwara and Varshavsky<sup>2</sup> to obtain the following.

$$(5.1) = \# \mathrm{Fix} \left( [K^p g^p K^p] \times [K_p g_p K_p] \mid \mathrm{Ig}_{N,m}^{\mathrm{ord}}(\overline{\mathbb{F}}_p) \right) \\ \stackrel{\text{Prop. 4.1}}{=} \# \mathrm{Fix} \left( [K^p g^p K^p] \times [K_p g_p K_p] \mid \sum_{F \in \mathrm{IQF}(p)} F^\times \backslash (\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)) / K^p \times K_p \right).$$

The details are omitted, but this is turned into the following via the combinatorial lemma of [Mil92, §5]:

$$= \sum_{F \in \mathrm{IQF}(p)} \sum_{a \in F^\times} \# \left( F^\times \backslash (Y^p(a) \times \check{Y}_p(a)) \right), \quad (5.2)$$

where

$$Y^p(a) := \{y^p \in \mathrm{GL}_2(\mathbb{A}^{\infty,p})/K^p : y^p g^p = a y^p \text{ in } \mathrm{GL}_2(\mathbb{A}^{\infty,p})/K^p\} \\ = \{y^p \in \mathrm{GL}_2(\mathbb{A}^{\infty,p})/K^p : (y^p)^{-1} a y^p \in K^p g^p K^p\}, \\ \check{Y}_p(a) := \{y_p \in J(\mathbb{Q}_p)/K_p : y_p g_p = a y_p \text{ in } J(\mathbb{Q}_p)/K_p\} \\ = \{y_p \in J(\mathbb{Q}_p)/K_p : y_p^{-1} a y_p \in K_p g_p K_p\}.$$

Of course  $y_p^{-1} a y_p = a$  in our setup since  $J(\mathbb{Q}_p)$  is abelian, but we chose to write  $y_p^{-1} a y_p$  since this is the correct expression for general Igusa varieties where  $J$  is not a torus. Thus we can rewrite (5.2) as

$$= \sum_{F \in \mathrm{IQF}(p)} \sum_{a \in F^\times} \int_{F^\times \backslash \mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)} \mathbf{1}_{K^p g^p K^p}((y^p)^{-1} a y^p) \times \mathbf{1}_{K_p g_p K_p}(y_p^{-1} a y_p) d(y^p, y_p).$$

Since the integrand depends only on the  $F_{\mathbb{A}^\infty}^\times$ -coset of  $(y^p, y_p)$ , we can rewrite  $\int_{F^\times \backslash \mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)}$  as  $\mathrm{vol}(F^\times \backslash F_{\mathbb{A}^\infty}^\times) \int_{F^\times \backslash \mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)}$ , and then express the integral as an orbital integral at  $a$ :

$$= \sum_{F \in \mathrm{IQF}(p)} \sum_{a \in F^\times} \mathrm{vol}(F^\times \backslash F_{\mathbb{A}^\infty}^\times) O_a^{\mathrm{GL}_2(\mathbb{A}^{\infty,p})}(\mathbf{1}_{K^p g^p K^p}) O_a^{J(\mathbb{Q}_p)}(\mathbf{1}_{K_p g_p K_p}).$$

<sup>2</sup>To apply this formula, we need to twist the double coset operator by a sufficiently high power of Frobenius. In this article we will gloss over this point, but this turns out to be harmless for computing the cohomology. See [HT01, V.1] or [Shi09, §6] for details.

Now we reparametrize the pairs  $(F, a)$  in the sum. We will view  $J(\mathbb{Q}_p) = \mathrm{GL}_1(\mathbb{Q}_p) \times \mathrm{GL}_1(\mathbb{Q}_p)$  as the diagonal subgroup of  $\mathrm{GL}_2(\mathbb{Q}_p)$  below. Recall that an element  $\gamma_0 \in \mathrm{GL}_2(\mathbb{Q})$  is said to be  $\mathbb{R}$ -elliptic if  $\gamma_0$  is either central or has imaginary eigenvalues. We use the symbol  $\sim$  to designate the conjugacy relation. Then

**Lemma 5.1.** *There is a natural bijection between the following two sets:*

- (i)  $\{(F, a) : F \in \mathrm{IQF}(p), a \in F^\times, \text{ s.t. } a \text{ is conjugate to an acceptable element of } J(\mathbb{Q}_p)\}$ ,
- (ii)  $\mathcal{C} := \{(\gamma_0, \delta) : \gamma_0 \in \mathrm{GL}_2(\mathbb{Q})/\sim \text{ } \mathbb{R}\text{-elliptic}, \delta \in J(\mathbb{Q}_p)/\sim \text{ acceptable}\}$ ,

given as follows. For each pair  $(F, a)$ , write  $a = (a_1, a_2) \in F_{\mathbb{Q}_p}^\times \cong \mathbb{Q}_p^\times \times \mathbb{Q}_p^\times$ ; then  $v_p(a_1) \neq v_p(a_2)$  by the condition on  $a$ . The pair is sent to  $(a, \delta)$ , where  $\delta = (a_1, a_2)$  if  $v_p(a_1) > v_p(a_2)$ , and  $\delta = (a_2, a_1)$  otherwise.

*Proof.* Left as an exercise. □

We apply the lemma to the preceding formula to obtain

$$(5.1) = \sum_{(\gamma_0, \delta) \in \mathcal{C}} \mathrm{vol}(F^\times \backslash F_{\mathbb{A}^\infty}^\times) O_{\gamma_0}^{\mathrm{GL}_2(\mathbb{A}^\infty, p)}(\mathbf{1}_{K^p g^p K^p}) O_\delta^{J(\mathbb{Q}_p)}(\mathbf{1}_{K_p g_p K_p}).$$

As we can take finite linear combinations of test functions of the form  $\mathbf{1}_{K^p g^p K^p}$  (resp.  $\mathbf{1}_{K_p g_p K_p}$ ), we arrive at the following.

**Theorem 5.2.** *Let  $f^p \in C_c^\infty(\mathrm{GL}_2(\mathbb{A}^\infty, p))$ ,  $f'_p \in C_c^\infty(J(\mathbb{Q}_p))$ , and assume that  $f'_p$  is supported on acceptable elements. Then*

$$\mathrm{tr} \left( f^p \times f'_p \mid H_c(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) \right) = \sum_{(\gamma_0, \delta) \in \mathcal{C}} \mathrm{vol}(F^\times \backslash F_{\mathbb{A}^\infty}^\times) O_{\gamma_0}^{\mathrm{GL}_2(\mathbb{A}^\infty, p)}(f^p) O_\delta^{J(\mathbb{Q}_p)}(f'_p).$$

### 6. $\ell$ -ADIC COHOMOLOGY OF IGUSA CURVES

To achieve the goal stated in §3, the final step is to extract spectral information about the cohomology from the trace formula above. We need some input from the theory of automorphic forms. There are two key ingredients, local and global, which we state without proofs but include references. We fix an isomorphism  $\iota : \overline{\mathbb{Q}}_\ell \cong \mathbb{C}$  to identify  $\overline{\mathbb{Q}}_\ell$  and  $\mathbb{C}$  coefficients of representations.

**(Local)** There exists a “transfer” from  $f'_p \in C_c^\infty(J(\mathbb{Q}_p))$ , supported on acceptable elements, to  $f_p \in C_c^\infty(\mathrm{GL}_2(\mathbb{Q}_p))$  such that for every semisimple element  $\gamma \in \mathrm{GL}_2(\mathbb{Q}_p)$ ,

$$O_\gamma^{\mathrm{GL}_2(\mathbb{Q}_p)}(f_p) = \begin{cases} O_\delta^{J(\mathbb{Q}_p)}(f'_p), & \text{if } \exists \text{ acceptable } \delta \in J(\mathbb{Q}_p) \text{ s.t. } \delta \sim \gamma, \\ 0, & \text{otherwise.} \end{cases} \tag{6.1}$$

Moreover a character identity is satisfied by  $f'_p$  and  $f_p$ :

$$\mathrm{tr} \pi_p(f_p) = \mathrm{tr} (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2})(f'_p), \quad \forall \pi_p : \text{irred. adm. representation of } \mathrm{GL}_2(\mathbb{Q}_p), \tag{6.2}$$

where  $P$  is the upper triangular Borel subgroup of  $\mathrm{GL}_2$ ,  $N^{\mathrm{op}}$  is the unipotent radical of the opposite parabolic of  $P$ ,  $\delta_{P(\mathbb{Q}_p)}$  is the modulus character on  $P(\mathbb{Q}_p)$ , and  $J_{N^{\mathrm{op}}}$  is the normalized Jacquet module relative to  $N^{\mathrm{op}}$  from representations of  $\mathrm{GL}_2(\mathbb{Q}_p)$  to those of  $J(\mathbb{Q}_p)$ . We remark that [Shi10, Lem. 3.9] proves this local fact in a more general setup. (See the proof of [KS, Lem. 3.1.2] for a small correction to the statement and proof of [Shi10, Lem. 3.9].)

**(Global)** There exists an Euler–Poincaré function  $f_\infty \in C_c^\infty(\mathrm{GL}_2(\mathbb{R})/\mathbb{R}_{>0}^\times)$ , which encodes the “weight 2 condition” for classical modular forms in the following sense: for each irreducible unitary representation  $\pi_\infty$  of  $\mathrm{GL}_2(\mathbb{R})$  whose central character is trivial on  $\mathbb{R}_{>0}^\times$ ,

$$\mathrm{tr} \pi_\infty(f_\infty) = \begin{cases} -1, & \text{if } \pi_\infty \text{ is the weight 2 discrete series representation,} \\ 1, & \text{if } \pi_\infty = \chi \circ \det \text{ for } \chi = \mathbf{1} \text{ or } \chi = \mathrm{sgn}, \\ 0, & \text{otherwise.} \end{cases} \tag{6.3}$$

Here we have written  $\mathrm{sgn}$  for the sign character on  $\mathbb{R}^\times$ . Moreover, a simple trace formula of the following form holds:

$$\begin{aligned} & \mathrm{tr} \left( f^p f_p f_\infty | L_{\mathrm{disc}}^2(\mathrm{GL}_2(\mathbb{Q}) \backslash \mathrm{GL}_2(\mathbb{A}) / \mathbb{R}_{>0}^\times) \right) \\ &= \sum_{\substack{\gamma \in \mathrm{GL}_2(\mathbb{Q}) / \sim \\ \text{R-elliptic}}} (\text{volume}) \cdot O_\gamma^{\mathrm{GL}_2(\mathbb{A}^\infty)}(f^p f_p) + (\text{error terms}). \end{aligned} \tag{6.4}$$

When  $\gamma$  is noncentral so that it generates an imaginary quadratic field  $F$  over  $\mathbb{Q}$ , the volume term is precisely  $\mathrm{vol}(F^\times \backslash F_{\mathbb{A}^\infty}^\times)$  that we saw before. (The existence of  $f_\infty$  and the simple trace formula are respectively due to Clozel–Delorme [CD85] and Arthur [Art89] in quite a general setup.)

Write  $\mathcal{A}_1(\mathrm{GL}_2)$  for the set of 1-dimensional automorphic representations  $\pi$  of  $\mathrm{GL}_2(\mathbb{A})$  such that  $\pi_\infty \in \{\mathbf{1}, \mathrm{sgn} \circ \det\}$ . By  $\mathcal{A}_{\mathrm{wt} 2}(\mathrm{GL}_2)$  we denote the set of cuspidal automorphic representations  $\pi$  of  $\mathrm{GL}_2(\mathbb{A})$  arising from weight 2 modular forms. By  $(\ )^{\mathrm{ss}}$ , we mean the semisimplification of a representation with respect to the given action.

**Theorem 6.1.** *As  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -representations, we have:*

$$\begin{aligned} H_c^2(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) &= \bigoplus_{\pi \in \mathcal{A}_1(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}), \\ H_c^1(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell)^{\mathrm{ss}} &= \bigoplus_{\pi \in \mathcal{A}_{\mathrm{wt} 2}(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}) + (\text{error terms}), \\ H_c^0(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell) &= 0. \end{aligned}$$

*Remark 6.2.* One can think of  $\delta_{P(\mathbb{Q}_p)}^{1/2}$  as “raising weight by 1”. So when  $\pi_p$  is tempered (thus so is  $J_{N^{\mathrm{op}}}(\pi_p)$ ), we expect  $J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}$  to appear in  $H_c^1$  (except that  $H_c^i$  need not be pure of weight  $i$  due to non-properness). When  $\dim \pi_p = 1$  (nontempered),  $J_{N^{\mathrm{op}}}(\pi_p)$  is not unitary but has “weight 1”, so  $J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}$  is expected to contribute to  $H_c^2$ . To make the use of weight precise, the point is that  $\mathrm{Ig}_\infty^{\mathrm{ord}}$  is defined over  $\mathbb{F}_p$  (not just  $\overline{\mathbb{F}}_p$ ) and the geometric Frobenius action coincides with the action of  $(1, p) \in J(\mathbb{Q}_p)$ . Notice that indeed  $\delta_{P(\mathbb{Q}_p)}^{1/2}((1, p)) = p$ .

*Remark 6.3.* The error terms in  $H_c^1$  arise from spectral interpretation of the geometric error terms (on the proper Levi subgroup  $\mathrm{GL}_1 \times \mathrm{GL}_1$  of  $\mathrm{GL}_2$ ) in (6.4) as can be seen in the proof below. We invite the reader to explicitly describe the error terms in  $H_c^1$ .

*Remark 6.4.* The top-degree cohomology with compact support classifies the set of irreducible components; in our setup, this coincides with the set of connected components by (formal) smoothness. We leave it to the reader to notice the following: the description of  $H_c^2(\mathrm{Ig}_\infty^{\mathrm{ord}}, \overline{\mathbb{Q}}_\ell)$  implies that  $\pi_0(\mathrm{Ig}_\infty^{\mathrm{ord}})$  is in  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -equivariant bijection with  $\pi_0(Y_C) = \widehat{\mathbb{Z}}^\times$ , cf. (2.1).

*Proof.* Since  $\mathrm{Ig}_{N,m}^{\mathrm{ord}}$  is affine of dimension 1, we have  $H_c^i(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) = 0$  unless  $i \in \{1, 2\}$ . To understand

$$H_c^2(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) - H_c^1(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell})$$

in the Grothendieck group of admissible  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$ -representations, we rewrite the formula in Theorem 5.2 in terms of automorphic representations.

$$\begin{aligned} & \mathrm{tr} \left( f^p \times f'_p \mid H_c(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) \right) \\ = & \sum_{(\gamma_0, \delta) \in \mathcal{C}} \mathrm{vol}(F^{\times} \backslash F_{\mathbb{A}^{\infty}}^{\times}) O_{\gamma_0}^{\mathrm{GL}_2(\mathbb{A}^{\infty,p})}(f^p) O_{\delta}^{J(\mathbb{Q}_p)}(f'_p) \\ \stackrel{(6.1)}{=} & \sum_{\gamma_0 \in \mathrm{GL}_2(\mathbb{Q})/\sim} \mathrm{vol}(F^{\times} \backslash F_{\mathbb{A}^{\infty}}^{\times}) O_{\gamma_0}^{\mathrm{GL}_2(\mathbb{A}^{\infty,p})}(f^p) O_{\delta}^{\mathrm{GL}_2(\mathbb{Q}_p)}(f'_p) \\ \stackrel{(6.4)}{=} & \mathrm{tr} \left( f^p f_p f_{\infty} \mid L_{\mathrm{disc}}^2(\mathrm{GL}_2(\mathbb{Q}) \backslash \mathrm{GL}_2(\mathbb{A})/\mathbb{R}_{>0}^{\times}) \right) + (\text{error terms}) \\ \stackrel{(6.2)}{=} & \mathrm{tr} \left( f^p f'_p f_{\infty} \mid J_{N^{\mathrm{op}}} \left( L_{\mathrm{disc}}^2(\mathrm{GL}_2(\mathbb{Q}) \backslash \mathrm{GL}_2(\mathbb{A})/\mathbb{R}_{>0}^{\times}) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2} \right) \right) + (\text{error terms}) \\ \stackrel{(6.3)}{=} & \mathrm{tr} \left( f^p f'_p \mid \sum_{\pi \in \mathcal{A}_1(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}) \right) \\ & - \mathrm{tr} \left( f^p f'_p \mid \sum_{\pi \in \mathcal{A}_{\mathrm{wt} 2}(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}) \right) + (\text{error terms}) \end{aligned}$$

At this point, we can remove the assumption that  $f'_p$  is supported on acceptable elements. Indeed, [Shi09, Lem. 6.4] tells us that the trace identity between the first and last expressions in the displayed formula above holds for all  $f^p$  and all  $f'_p$ . Therefore, we have the identity in the Grothendieck group

$$\begin{aligned} & H_c^2(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) - H_c^1(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) \tag{6.5} \\ = & \sum_{\pi \in \mathcal{A}_1(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}) - \sum_{\pi \in \mathcal{A}_{\mathrm{wt} 2}(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2}) + (\text{error terms}). \end{aligned}$$

To separate  $H_c^2$  from  $H_c^1$ , the basic idea is that there is no cancellation between  $H_c^2$  and  $H_c^1$  since the geometric Frobenius action (encoded by  $(1, p) \in J(\mathbb{Q}_p)$ ; see Remark 6.2) has weight=2 in  $H_c^2$  and weight  $\leq 1$  in  $H_c^1$ . There are at least a couple of ways to proceed.

- (i) Show that the geometric Frobenius action has weight=2 in the first summation and  $< 2$  apart from it. (This method generalizes to study the top-degree  $H_c$ , or dually  $H^0$ , of Igusa varieties in the Hodge-type setting, as carried out in [KS].)
- (ii) Verify that everything in the error terms has the negative sign. (This is harder to generalize to higher dimensions when there are many cohomological degrees.)

Either way, we obtain

$$H_c^2(\mathrm{Ig}_{\infty}^{\mathrm{ord}}, \overline{\mathbb{Q}}_{\ell}) = \sum_{\pi \in \mathcal{A}_1(\mathrm{GL}_2)} \pi^{\infty,p} \otimes (J_{N^{\mathrm{op}}}(\pi_p) \otimes \delta_{P(\mathbb{Q}_p)}^{1/2})$$

in the Grothendieck group. Since distinct 1-dimensional representations of  $\mathrm{GL}_2(\mathbb{A}^{\infty,p}) \times J(\mathbb{Q}_p)$  have no extensions between each other, we obtain the formula for  $H_c^2$  in the theorem. From this, we can compute  $H_c^1$  up to semisimplification from (6.5).  $\square$

Finally we remark that endoscopy complicates the whole computation when considering general Igusa varieties, just like endoscopy intervenes in the computation of cohomology of Shimura varieties. See [Shi20] for an illustrative account of endoscopic calculation for Igusa varieties associated with certain unitary similitude groups. Endoscopy does not show up in the present article only because we are restricting to the  $GL_2$ -case.

## REFERENCES

- [Art89] James Arthur, *The  $L^2$ -Lefschetz numbers of Hecke operators*, Invent. Math. **97** (1989), no. 2, 257–290.
- [BM] Alexander Bertolini Meli, *An averaging formula for the cohomology of PEL-type Rapoport–Zink spaces*, <https://arxiv.org/abs/2102.10690>.
- [BR94] Don Blasius and Jonathan D. Rogawski, *Zeta functions of Shimura varieties*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Amer. Math. Soc., Providence, RI, 1994, pp. 525–571.
- [CD85] L. Clozel and P. Delorme, *Pseudo-coefficients et cohomologie des groupes de Lie réductifs réels*, C. R. Acad. Sci. Paris Sér. I Math. **300** (1985), no. 12, 385–387.
- [Clo93] Laurent Clozel, *Nombre de points des variétés de Shimura sur un corps fini (d’après R. Kottwitz)*, no. 216, 1993, Séminaire Bourbaki, Vol. 1992/93, pp. Exp. No. 766, 4, 121–149.
- [CS] Ana Caraiani and Peter Scholze, *On the generic part of the cohomology of non-compact unitary Shimura varieties*, <https://arxiv.org/abs/1909.01898>.
- [CS17] ———, *On the generic part of the cohomology of compact unitary Shimura varieties*, Ann. of Math. (2) **186** (2017), no. 3, 649–766.
- [GN09] Alain Genestier and Bao Châu Ngô, *Lectures on Shimura varieties*, *Autour des motifs—École d’été Franco-Asiatique de Géométrie Algébrique et de Théorie des Nombres/Asian-French Summer School on Algebraic Geometry and Number Theory. Volume I*, Panor. Synthèses, vol. 29, Soc. Math. France, Paris, 2009, pp. 187–236.
- [Ham17] Paul Hamacher, *The almost product structure of Newton strata in the deformation space of a Barsotti-Tate group with crystalline Tate tensors*, Math. Z. **287** (2017), no. 3-4, 1255–1277.
- [HK19] Paul Hamacher and Wansu Kim,  *$l$ -adic étale cohomology of Shimura varieties of Hodge type with non-trivial coefficients*, Math. Ann. **375** (2019), no. 3-4, 973–1044.
- [HR17] X. He and M. Rapoport, *Stratifications in the reduction of Shimura varieties*, Manuscripta Math. **152** (2017), no. 3-4, 317–343.
- [HT01] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich.
- [Igu68] Jun-ichi Igusa, *On the algebraic theory of elliptic modular functions*, J. Math. Soc. Japan **20** (1968), 96–106.
- [KS] Arno Kret and Sug Woo Shin,  *$H^0$  of Igusa varieties via automorphic forms*, <https://arxiv.org/abs/2102.10690>.
- [Lan73] R. P. Langlands, *Modular forms and  $\ell$ -adic representations*, Modular functions of one variable, II (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), 1973, pp. 361–500. Lecture Notes in Math., Vol. 349.
- [Lan76] ———, *Some contemporary problems with origins in the Jugendtraum*, Mathematical developments arising from Hilbert problems (Proc. Sympos. Pure Math., Vol. XXVIII, Northern Illinois Univ., De Kalb, Ill., 1974), 1976, pp. 401–418.
- [Man05] Elena Mantovan, *On the cohomology of certain PEL-type Shimura varieties*, Duke Math. J. **129** (2005), no. 3, 573–610.
- [Man20] ———, *The Newton stratification*, Shimura Varieties, London Mathematical Society Lecture Note Series, vol. 457, Cambridge University Press, 2020, pp. 166–191.
- [MC21] Sander Mack-Crane, *Counting points on Igusa varieties of Hodge type*, UC Berkeley PhD thesis, [https://math.berkeley.edu/~sander/writing/mack-crane\\_thesis.pdf](https://math.berkeley.edu/~sander/writing/mack-crane_thesis.pdf).
- [Mil92] J. S. Milne, *The points on a Shimura variety modulo a prime of good reduction*, The zeta functions of Picard modular surfaces, Univ. Montréal, Montréal, QC, 1992, pp. 151–253.
- [Sch11] Peter Scholze, *The Langlands-Kottwitz approach for the modular curve*, Int. Math. Res. Not. IMRN (2011), no. 15, 3368–3425.
- [Shi09] Sug Woo Shin, *Counting points on Igusa varieties*, Duke Math. J. **146** (2009), no. 3, 509–568.

- [Shi10] ———, *A stable trace formula for Igusa varieties*, J. Inst. Math. Jussieu **9** (2010), no. 4, 847–895.
- [Shi12] ———, *On the cohomology of Rapoport-Zink spaces of EL-type*, Amer. J. Math. **134** (2012), no. 2, 407–452.
- [Shi20] ———, *Construction of automorphic Galois representations: the self-dual case*, Shimura Varieties, London Mathematical Society Lecture Note Series, vol. 457, Cambridge University Press, 2020, pp. 209–250.
- [Zha] Chao Zhang, *Stratifications and foliations for good reductions of Shimura varieties of Hodge type*, preprint, arXiv:1512.08102.
- [Zhu20] Yihang Zhu, *Introduction to the Langlands–Kottwitz method*, Shimura Varieties, London Mathematical Society Lecture Note Series, vol. 457, Cambridge University Press, 2020, pp. 115–150.

*Email address:* [sug.woo.shin@berkeley.edu](mailto:sug.woo.shin@berkeley.edu)

DEPARTMENT OF MATHEMATICS, UC BERKELEY, BERKELEY, CA 94720, USA / KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 02455, REPUBLIC OF KOREA