



3-2011

Microbiota of Deciduous Endodontic Infections Analyzed by MDA and Checkerboard DNA-DNA Hybridization

W. L.F. Tavares

LC Neves de Brito

R. P. Teles

M.L. A. Massara

A.P. Ribeiro Sobrinho

See next page for additional authors

Follow this and additional works at: https://repository.upenn.edu/dental_papers

 Part of the [Dentistry Commons](#)

Recommended Citation

Tavares, W. L., de Brito, L. N., Teles, R. P., Massara, M. A., Sobrinho, A. R., Haffajee, A. D., Socransky, S. S., & Teles, F. R. (2011). Microbiota of Deciduous Endodontic Infections Analyzed by MDA and Checkerboard DNA-DNA Hybridization. *International Journal of Endontology*, 44 (3), 225-235. <http://dx.doi.org/10.1111/j.1365-2591.2010.01805.x>

At the time of publication, author Flavia Teles was affiliated with the Forsyth Institute. Currently, she is a faculty member at the School of Dental Medicine at the University of Pennsylvania.

This paper is posted at ScholarlyCommons. https://repository.upenn.edu/dental_papers/101
For more information, please contact repository@pobox.upenn.edu.

Microbiota of Deciduous Endodontic Infections Analyzed by MDA and Checkerboard DNA-DNA Hybridization

Abstract

Aims

To evaluate the microbiota of endodontic infections in deciduous teeth by checkerboard DNA-DNA hybridization after uniform amplification of DNA in samples by multiple displacement amplification (MDA).

Methodology

Forty samples from the root canal system of deciduous teeth exhibiting pulp necrosis with or without radiographically detectable periradicular/interadicular bone resorption were collected and 32 were analyzed, with 3 individuals contributing 2 samples; these were MDA- amplified and analyzed by Checkerboard DNA-DNA hybridization for levels of 83 bacterial taxa. Two outcome measures were used: the percentage of teeth colonized by each species; and the mean proportion of each bacterial taxon present across all samples were computed.

Results

The mean amount of DNA in the samples prior to amplification was 5.2 (\pm 4.7) ng and 6.1 (\pm 2.3) μ g after MDA. The mean number of species detected per sample was 19 (\pm 4) (range: 3–66) to the nearest whole number. The most prevalent taxa were *Prevotella intermedia* (96.9%), *Neisseria mucosa* (65.6%), *Prevotella nigrescens* (56.2%) and *Tannerella forsythia* (56.2%). *Aggregatibacter (Haemophilus) aphrophilus* and *Helicobacter pylori* were not detected. *P. intermedia* (10%), *Prevotella tanneriae* (7%) and *Prevotella nigrescens* (4.3%) presented the highest mean proportions of the target species averaged across the positive samples.

Conclusion

Root canals of infected deciduous teeth had a diverse bacterial population. *Prevotella* sp were commonly found with *P. intermedia*, *Prevotella tanneriae* and *Prevotella nigrescens* among the most prominent species detected.

Disciplines

Dentistry

Comments

At the time of publication, author Flavia Teles was affiliated with the Forsyth Institute. Currently, she is a faculty member at the School of Dental Medicine at the University of Pennsylvania.

Author(s)

W. L.F. Tavares, LC Neves de Brito, R. P. Teles, M.L. A. Massara, A.P. Ribeiro Sobrinho, A. D. Haffajee, S. S. Socransky, and F. R. Teles



Published in final edited form as:

Int Endod J. 2011 March ; 44(3): 225–235. doi:10.1111/j.1365-2591.2010.01805.x.

Microbiota of deciduous endodontic infections analyzed by MDA and Checkerboard DNA-DNA hybridization

WLF Tavares², LC Neves de Brito², RP Teles¹, MLA Massara², AP Ribeiro Sobrinho², AD Haffajee¹, SS Socransky¹, and FR Teles¹

¹Department of Periodontology, The Forsyth Institute, Boston, Massachusetts, USA

²Federal University of Minas Gerais School of Dentistry, Belo Horizonte, Minas Gerais, Brazil

Abstract

Aims—To evaluate the microbiota of endodontic infections in deciduous teeth by checkerboard DNA-DNA hybridization after uniform amplification of DNA in samples by multiple displacement amplification (MDA).

Methodology—Forty samples from the root canal system of deciduous teeth exhibiting pulp necrosis with or without radiographically detectable periradicular/interadicular bone resorption were collected and 32 were analyzed, with 3 individuals contributing 2 samples; these were MDA-amplified and analyzed by Checkerboard DNA-DNA hybridization for levels of 83 bacterial taxa. Two outcome measures were used: the percentage of teeth colonized by each species; and the mean proportion of each bacterial taxon present across all samples were computed.

Results—The mean amount of DNA in the samples prior to amplification was 5.2 (\pm 4.7) ng and 6.1 (\pm 2.3) μ g after MDA. The mean number of species detected per sample was 19 (\pm 4) (range: 3–66) to the nearest whole number. The most prevalent taxa were *Prevotella intermedia* (96.9%), *Neisseria mucosa* (65.6%), *Prevotella nigrescens* (56.2%) and *Tannerella forsythia* (56.2%). *Aggregatibacter (Haemophilus) aphrophilus* and *Helicobacter pylori* were not detected. *P. intermedia* (10%), *Prevotella tannerae* (7%) and *Prevotella nigrescens* (4.3%) presented the highest mean proportions of the target species averaged across the positive samples.

Conclusion—Root canals of infected deciduous teeth had a diverse bacterial population. *Prevotella* sp were commonly found with *P. intermedia*, *Prevotella tannerae* and *Prevotella nigrescens* among the most prominent species detected.

INTRODUCTION

Necrosis of pulp tissue is usually initiated by caries and/or trauma (Raslan & Wetzel 2006). Necrosis of the pulp in primary teeth might lead to periapical disease and could potentially affect the permanent tooth germ. Pulp therapy in primary teeth with necrotic pulps aims at eradicating the endodontic infection and preventing this early loss. In this way, the health of the succeeding tooth germ can be preserved (Pazelli *et al.* 2003, Bijoor & Kohli, 2005, da Silva *et al.* 2006). Since most pulp pathoses are caused by microbial infections, knowledge of their diversity in the infected root canals of primary teeth should underpin development of more efficacious endodontic therapies.

Several studies have investigated the microbiota associated with endodontic infections in adults (Sundqvist 1976, Baumgartner & Falkler 1991, Siqueira *et al.* 2000, Lana *et al.* 2001, Rolph *et al.* 2001, Munson *et al.* 2002, de Souza *et al.* 2005, Brito *et al.* 2007, Sassone *et al.* 2007, 2008). In contrast, few studies have assessed the microbial composition of deciduous endodontic infections (da Silva *et al.* 2006, Ruviere *et al.* 2007, Cogulu *et al.* 2008), utilizing different detection methods and focusing on a variable number of bacterial taxa. Knowledge of the endodontic infections of primary teeth is therefore far from complete (Marsh & Largent 1967, Toyoshima *et al.* 1988, Pazelli *et al.* 2003, da Silva *et al.* 2006).

Checkerboard DNA-DNA hybridization is a high-throughput molecular method that allows the identification and quantification of a wide range of bacterial species in multiple samples on a single nylon membrane. This technique has been employed in the study of the microbiota in saliva (Sachdeo *et al.* 2008), supragingival plaque (Haffajee *et al.* 2008), subgingival plaque (Haffajee *et al.* 2008, Teles *et al.* 2008), on oral soft tissue (Mager *et al.* 2003, Sachdeo *et al.* 2008), on dentures (Sachdeo *et al.* 2008), from dental implants (Gerber *et al.* 2006) and from root canals (Siqueira *et al.* 2000, Brito *et al.* 2007, Sassone *et al.* 2007, 2008).

The level of detection of the checkerboard DNA-DNA hybridization technique is between 10^4 and 10^7 bacterial cells of a given species in each sample. The bacterial content of samples from endodontic disease may be below this detection level, so a DNA amplification step called Multiple Displacement Amplification (MDA) can be used to enhance detection limits (Dean *et al.* 2002, Brito *et al.* 2007, Teles *et al.* 2007). MDA can amplify DNA present in oral biofilm samples with minimal bias (Dean *et al.* 2002, Hawkins *et al.* 2002, Yan *et al.* 2004). MDA enables the whole genomic amplification of DNA targets (Dean *et al.* 2002). The template is replicated again and again by a “hyperbranching” mechanism of strand displacement synthesis (Lizardi *et al.* 1998), with the polymerase laying down a new copy as it displaces previously made copies. Samples as small as 1 ng can be amplified 1000–10,000 fold (Mai *et al.* 2004). MDA uniformly amplifies the entire genomes (Hawkins *et al.* 2002) with minimal amplification bias (Hawkins *et al.* 2002, Nelson *et al.* 2002, Yan *et al.* 2004). This method allows the uniform amplification of the whole genomes present in a sample and has been effectively used as an aid in Checkerboard DNA-DNA Hybridization (Brito *et al.* 2007, Teles *et al.* 2007). Since bacterial genomes are uniformly amplified, the proportion that each taxon comprises of the test species can be calculated.

The aim of the present study was to combine MDA and Checkerboard DNA-DNA Hybridization to qualitatively and quantitatively evaluate the microbiota of endodontic infections in deciduous teeth.

MATERIAL AND METHODS

Subject population and sample collection

Thirty five subjects ranging in age from 4 to 10 years were recruited in the Department of Paediatric Dentistry, Federal University of Minas Gerais (UFMG), Belo Horizonte, Brazil. This study was approved by the Ethics Committee of the Federal University of Minas Gerais (ETIC132/07). Informed consent was obtained from the parents of the children enrolled in the study. The children had deciduous teeth exhibiting pulp necrosis, with or without radiographically detected periradicular/interadicular bone resorption. The selected teeth had intact roots or less than 2/3 of physiological root resorption. Selected teeth had clinical crowns that permitted effective rubber dam isolation. There was an absence of history of trauma associated with the selected teeth, periodontal involvement and previous root canal treatment. Thirty one primary teeth were molars and nine were single-rooted teeth. Samples

from multi-rooted teeth were taken from the largest root canal always associated with the periapical lesion.

Forty selected teeth were isolated using a rubber dam. Complete asepsis was employed, using the methodology proposed by Moller (1966); hydrogen peroxide (30%) was applied on the isolated crown, followed by 5% iodine, which was inactivated by 5% sodium thiosulfate solution. The samples were taken by scraping or filing the root canal walls with a size 15 K-type hand file (Dentsply Maillefer, Ballaigues, Switzerland). The file was introduced into the canal up to the radiographically determined working length, which was taken as approximately 1 mm short of the tooth apex in cases with intact roots, or to the limit of physiologic root resorption. Where there was radiographic image interposition of the primary root and the permanent germ, the file was inserted up to the level of the cusp of the permanent germ.

After removal from the canal, the final 2 mm of the file was removed using a sterile pair of surgical scissors and dropped into a microcentrifuge tube containing 20 μ L of alkaline lysis buffer (400 mM KOH, 100 mM DTT, 10 mM EDTA). After 10 min of incubation on ice, 20 μ L of neutralization solution (400 mM HCl, 600 mM Tris HCl, pH =0.6) were added, and the sample was kept at -20°C until MDA was performed. Among the teeth sampled, 3 were symptomatic cases, 15 exhibited exudation, 22 were associated with sinus tracts and 26 radiographically detectable periapical radiolucencies.

Multiple displacement amplification (MDA) of root canal samples

MDA was performed as described by Teles *et al.* (2007). GenomiphiTM (Amersham Biosciences, Arlington Heights, IL, USA) was used for whole genomic amplification as described by the manufacturer. In brief, 1 μ L of each of the DNA templates (i.e. endodontic samples) was added to 9 μ L of sample buffer (50 mM Tris-HCl pH 8.2, 0.5 mM EDTA containing random hexamer primers) in 200 μ L microcentrifuge tubes (Stratagene, La Jolla, CA, USA). Templates in sample buffer were heat denatured at 95°C for 3 minutes in a Perkin-Elmer Thermocycler and cooled to 4°C . One μ L of phi 29 DNA polymerase mix including additional random hexamers was mixed on ice with 9 μ L of reaction buffer containing dNTPs. The mixture was then added to the denatured sample to make a final volume of 20 μ L and incubated at 30°C for 16–18 hours. Ten ng of Lambda DNA (contained in 1 μ L) was used as a control. The amplification reaction was terminated by incubation of the samples at 65°C for 10 min. The amplified material was either immediately used, stored short-term at 4°C or at -20°C for longer storage.

The DNA content of the samples was measured prior to and after amplification using the PicogreenTM dsDNA quantification assay (Invitrogen, Carlsbad, CA, USA). PicogreenTM is a fluorescent nucleic acid stain that allows the quantification of as little as 25 pg/mL of double stranded DNA in samples.

Checkerboard DNA-DNA Hybridization

Preparation of probes and standards for quantification—Checkerboard DNA-DNA Hybridization was performed as previously described (Socransky *et al.* 1994, Socransky *et al.* 2004). For the preparation of probes and standards, each of the species listed in Table 1 were grown on the surface of blood agar plates (except the two spirochaetes, which were grown in broth) for 3–7 days. The cells were harvested and placed in 1.5 mL microcentrifuge tubes containing 1 mL of TE buffer (10 mM Tris-HCl, 0.1 mM EDTA, pH=7.6). Cells were washed twice by centrifugation in TE buffer at $1300\times g$ for 10 min. The cells were resuspended and lysed with either 10% SDS and Proteinase K (20 mg/mL) for Gram-negative strains or in 150 μ L of an enzymemixture containing 15 mg/mL

lysozyme (Sigma) and 5 mg/mL achromopeptidase (Sigma) in TE buffer (pH=8.0) for gram-positive strains. The pelleted cells were resuspended by 15 s of sonication and incubated at 37°C for 1 h. DNA was isolated and purified using the method of Smith *et al.* (1989). The concentration of the purified DNA was determined by spectrophotometric measurement of the absorbance at 260 nm. The purity of the preparations was assessed by the ratio of the absorbances at 260 and 280 nm. Whole genomic DNA probes were prepared from each of the 83 test strains by labeling 1–3µg DNA with digoxigenin (Boehringer Mannheim, Indianapolis, IN, USA) using a random primer technique (Feinberg & Vogelstein 1983).

Sample preparation and microbial analysis—In brief, following amplification and quantification, amplified samples were boiled for 10 min. Approximately 1500 ng of DNA (5 µL) of the amplified sample were placed in a microcentrifuge tube containing 1 mL of TE buffer prior to boiling. The samples were placed into the extended slots of a Minislot 30 apparatus (Immunelectrics, Cambridge, MA, USA), concentrated onto a nylon membrane (Boehringer Mannheim) by vacuum and fixed onto the membrane by cross-linking using ultraviolet light (Stratalinker 1800, La Jolla, CA, USA) followed by baking at 120°C for 20 min. The Minislot device permitted the deposition of 28 different samples in individual lanes on a single membrane, as well as two control lanes containing the standards for quantification: 1 and 10 ng of DNA of each bacterial species tested, equivalent to 10^5 and 10^6 cells, respectively.

The membrane with fixed DNA was placed in a Miniblotter 45 apparatus (Immunelectrics) with the lanes of DNA at 90° to the channels of the device. A 30 × 45 “checkerboard” pattern was produced. Each channel was used as an individual hybridization chamber for separate DNA probes. Bound probes were detected by anti-digoxigenin antibody conjugated with alkaline phosphatase and a chemifluorescent substrate. Signal intensities of the endodontic samples and the standards (containing 10^5 and 10^6 cells of each species) on the same membrane were measured using a Storm FluorImager (Molecular Dynamics, Sunnyvale, CA, USA). Signals were converted to absolute counts by comparison with standards on the membrane (Socransky *et al.* 2004). Failure to detect a signal was recorded as zero.

Two membranes were run for each sample: one containing the “standard” 40 DNA probes used to examine periodontal samples as well as a probe to detect *Streptococcus mutans*. A second membrane employed 42 probes to species thought to be implicated in endodontic infections. Sensitivity and specificity tests were performed for all probes before performing the Checkerboard DNA-DNA Hybridization analysis of the root canal samples. The protocol to validate the specificity of these 83 probes was similar to that described by Socransky *et al.* (2004). If cross-reactions were observed, the cross-reacting probes were discarded and new probes constructed and validated (Socransky *et al.* 2004).

Data analysis

The microbial data were expressed in two ways. The prevalence of each species, reflected by presence/absence data, indicated the proportion of samples in which the species were detected at $>10^4$ cells in MDA amplified samples. Since the sample DNA was amplified, absolute numbers could not be determined. Thus, proportions that each species comprised of the total DNA probe count were computed for each sample. Proportion data were expressed as percentage of total DNA probe count for each species and averaged across samples.

RESULTS

Quantification of DNA before and after MDA of endodontic samples

DNA from each root canal sample was amplified using MDA. Eight samples failed to provide good quality amplified DNA. The amount of DNA (\pm Standard Deviation - SD) present in the 32 remaining samples before the amplification averaged 5.2 (\pm 4.7, SD) ng and 6.1 (\pm 2.3) μ g after amplification, an approximately 1,000 - fold amplification. Among the 32 samples include in this study, 5 were from single rooted and 27 from multi-rooted teeth. Twelve of the samples were from females (n=12) and the mean age (\pm SD) was 6.8 years (\pm 1.9) (range: 4–12 years). Of the 8 samples excluded due to absence of amplification, 2 were from single rooted and 6 from multi-rooted teeth. Five of the samples were from females (n=5) and the mean age (\pm SD) was 6.4 years (\pm 1.8) (range: 4–9 years).

Microbial species in root canal samples

The mean number of species (\pm Standard Error of the mean - SEM) detected in the amplified samples at a threshold equivalent to $>10^4$ bacterial cells (in 5 μ l of amplified sample) was 19 (\pm 4) (range: 3–66). *Aggregatibacter (Haemophilus) aphrophilus* and *Helicobacter pylori* were not detected in any of the samples.

Figure 1 demonstrates the mean percentage of samples exhibiting counts of each of the 81 bacterial species detected at the level of $> 10^4$ bacterial cells. The most prevalent bacterial species was *Prevotella intermedia*, detected in 96.9% of the sampled teeth. It was followed by *Neisseria mucosa* (65.2%), *Prevotella nigrescens* (56.2%), *Tannerella forsythia* (56.2%), *Prevotella denticola* (53.1%) and *Fusobacterium nucleatum ss vincentii* (50.0%). The least prevalent species detected at the $> 10^4$ bacterial cells level were *Enterococcus faecalis* (3.2%) and *Eikenella corrodens* (3.1%).

Figure 2 presents the mean proportions (percentage of DNA probe counts) of the target species averaged across the positive samples. *P. intermedia* showed the highest mean proportions (10.0%) followed by *Prevotella tanneriae* (7.0%) and *P. nigrescens* (4.3%), while *Campylobacter ureolyticus* (0.05%), *Legionella pneumophila* (0.04%), and *E. faecalis* (0.03%) showed the lowest mean proportions.

DISCUSSION

The purpose of the present investigation was to evaluate the microbiota of endodontic infections in deciduous teeth. The bacterial diversity in any environment is underestimated when assessed by culture-based techniques (Papapanou 2002, Munson *et al.* 2002). Infected root canals yielded a maximum of 10–12 species when assessed by culture methods (Sundqvist *et al.* 1976, Gomes *et al.* 2004), while this range raises to 42–51 species in studies using culture-independent molecular methods (Siqueira *et al.* 2000, Brito *et al.* 2007). Conventional PCR and Quantitative PCR (qPCR) are often used for direct detection of taxa in root canal samples. However, the number of samples and taxa that can be analyzed is limited by the costs and complexity of the procedure, particularly when using qPCR.

Checkerboard DNA-DNA hybridization enables the quantitative analysis of multiple taxa in large numbers of samples (Socransky & Haffajee 2005). However, individual bacterial species may be present in the infected root canal system in numbers below the level of detection of the Checkerboard DNA-DNA hybridization technique. The lack of detection of such taxa would underestimate their possible role in the endodontic microbial ecosystem. MDA provides a simple and reliable method to amplify the sample DNA with minimal bias (Hawkins *et al.* 2002, Nelson *et al.* 2002, Yan *et al.* 2004, Brito *et al.* 2007, Teles *et al.*

2007). The characteristics of the enzyme used in MDA and the reaction *per se* contribute to minimizing the bias. The $\Phi 29$ DNA polymerase has a very low error rate of 1 in 10^6 – 10^7 nucleotides in its intrinsic enzymatic activity (Esteban *et al.* 1993) and during amplification (Nelson *et al.* 2002), in contrast to 3×10^4 for Taq DNA polymerase (Eckert & Kunkel 1991). Thus, the accumulation of mutations following a 10,000-fold amplification is just 3 per 10^6 nucleotides (Nelson *et al.* 2002). PCR-based amplification results in bias that varies from 10^2 to 10^6 , while MDA bias for human genomic DNA has been estimated to be less than 3-fold (Dean *et al.* 2002). Additionally, the MDA is an isothermal reaction, in contrast to PCR-based techniques, which require multiple cycles at different temperatures and thus can be biased by differences in % GC content of different genomes (Teles *et al.* 2007). In the present study, MDA amplification of DNA from the root canal samples led to an approximately 1000-fold amplification. Since the amplified sample was obtained from an aliquot of the original sample and the detection limit of the checkerboard DNA-DNA hybridization is 10^4 bacterial cells, it was possible that as few as 10 cells of a given species were present in the sample prior to amplification and which could be detected after amplification.

On average, 19 species were detected per amplified sample, substantially more than previously detected by culture (Lana *et al.* 2001, Gomes *et al.* 2004) and by clonal analysis (Jacinto *et al.* 2007) in adults. This figure is lower than the average of 51.2 species recently reported in endodontic infections in adults, using a similar methodology to the one employed in the present study (Brito *et al.* 2007). Ruvieri *et al.* (2007) used whole-genomic probes to detect 34 bacterial taxa in root canal samples from deciduous teeth. The authors found, on average 7.47 bacterial species (range: 0–10) in teeth exhibiting irreversible pulpitis and 11.45 species (range: 0–27) in teeth with pulp necrosis and chronic apical periodontitis. It suggests that the difference in numbers may be due to the primary teeth harboring a less complex microbiota or that infection in adults may have been present longer, allowing time for greater diversity to develop.

Studies focusing on root canal infections often employ “K” or “H” type files followed by two to four paper points to collect the samples (Siqueira *et al.* 2000, Pazelli *et al.* 2003, de Souza *et al.* 2005, da Silva *et al.* 2006, Ruvieri *et al.* 2007, Sassone *et al.* 2007, 2008, Cogulu *et al.* 2008). This method may not target the microbiota in the apical third of the root canal, since the entire content of the canal could be collected. To ensure that the apical portion was the main area sampled in this study, a K file was inserted using a reaming motion to the working length and only the apical two mm were sectioned. The difference in sampling may account for the prevalence of streptococci in the studies using paper points (da Silva *et al.* 2006, Pazelli *et al.* 2003, Ruvieri *et al.* 2007). The facultative anaerobic species, such as the *Streptococcus* species, may be present in higher counts in the coronal third of the canal, where conditions may favor their growth. In the present study, this genus was identified in higher counts only in teeth with pulp chamber exposure. Differences in sampling might also explain the lower bacterial cell numbers found in comparison to 10^5 – 10^7 cells per sample reported by other authors using H-files associated with paper points to collect samples analyzed by checkerboard DNA-DNA hybridization (Sassone *et al.* 2007, 2008), paper points and real-time PCR (Blome *et al.* 2008), and paper points and culture techniques (Vianna *et al.* 2008) to study endodontic infection in adults.

Root canal infections are polymicrobial and predominated by obligate and facultative anaerobes (Sundqvist 1992, Lana *et al.* 2001). In this study, the most prevalent bacterial species was *P. intermedia*, present in 96.9% of the samples. It was followed by other obligate anaerobes, such as *P. nigrescens*, *T. forsythia*, *P. denticola*, *F. nucleatum ss vincenti*, as well as *N. mucosa*, a facultative organism. These findings were in contrast with the results reported by Ruvieri *et al.* (2007), where *Campylobacter rectus*, *Treponema*

denticola, and *Gemella morbillorum* where the most prevalent taxa and Cogulu *et al.* (2008), who found that *T. denticola* and *Porphyromonas gingivalis* were the most prevalent species. The sampling procedures and methods for bacterial detection employed in those studies may account for those differences. Sassone *et al.* (2007, 2008) showed that more than 70% of root canal samples in permanent teeth were colonized by *N. mucosa* and *F. nucleatum* ss *vincentii*, which is in accord with the present study, even though Sassone *et al.* (2007, 2008) performed those studies on samples from adult teeth. However, *E. faecalis*, which was detected infrequently in this study, was found in high prevalence by those authors. *E. faecalis* has been frequently found in association with secondary endodontic infection in adults (Pirani *et al.* 2008, Vianna *et al.* 2008). Cogulu *et al.* (2008) reported the high prevalence of this species in primary infection of permanent teeth of children. Conceivably, the anatomy of permanent teeth favors colonization by this taxon. It is possible that it is a transient species introduced by food or human contact, since *E. faecalis* detection is a rare event at the advancing front of dentinal lesions (Zehnder & Guggenheim 2009).

The combination of MDA and checkerboard DNA-DNA hybridization permitted a better appreciation of the microbiota associated with root canal infection in deciduous teeth. The present study confirmed the polymicrobial nature of these infections and that they comprised a large number of facultative and anaerobic species. The data shows that deciduous endodontic infections are highly diverse, presenting, on average, 19 taxa per sample (range 3–66 taxa). The data also suggest that *Prevotella* sp. may play an important role in the microbiota of endodontic infections in primary teeth. *P. intermedia*, *P. nigrescens* and *P. denticola* were highly prevalent in the samples analyzed. *P. intermedia*, *P. tanneriae* and *P. nigrescens* represented the highest proportions of the total DNA probe count. However, more studies analyzing a larger number of samples using standardized sampling methods are needed to foster a better understanding of the pathogenesis of deciduous apical periodontitis, as well as aid the design of more efficient endodontic treatments.

Conclusions

Root canals of infected deciduous teeth present wide bacterial diversity. *Prevotella* sp. were the most prevalent amongst the samples and *P. intermedia*, *Prevotella tanneriae* and *Prevotella nigrescens* gave the highest proportion of hybridization of the total DNA probe counts.

References

- Baumgartner JC, Falkler WA. Bacteria in the apical 5 mm of infected root canals. *Journal of Endodontics*. 1991; 17:380–3. [PubMed: 1809801]
- Bijoor RR, Kohli K. Contemporary space maintenance for the pediatric patient. *The New York State Dental Journal*. 2005; 71:32–5. [PubMed: 15887467]
- Blome B, Braun A, Sobarzo V, Jepsen S. Molecular identification and quantification of bacteria from endodontic infections using real-time polymerase chain reaction. *Oral Microbiology and Immunology*. 2008; 23:384–90. [PubMed: 18793361]
- Brito LC, Teles FR, Teles RP, et al. Use of multiple-displacement amplification and checkerboard DNA-DNA hybridization to examine the microbiota of endodontic infections. *Journal of Clinical Microbiology*. 2007; 45:3039–49. [PubMed: 17634304]
- Cogulu D, Uzel A, Oncag O, Eronat C. PCR-based identification of selected pathogens associated with endodontic infections in deciduous and permanent teeth. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*. 2008; 106:443–9.
- da Silva LA, Nelson-Filho P, Faria G, de Souza-Gugelmin MC, Ito IY. Bacterial profile in primary teeth with necrotic pulp and periapical lesions. *Brazilian Dental Journal*. 2006; 17:144–8.

- de Souza CA, Teles RP, Souto R, Chaves MA, Colombo AP. Endodontic therapy associated with calcium hydroxide as an intracanal dressing: microbiologic evaluation by the checkerboard DNA-DNA hybridization technique. *Journal of Endodontics*. 2005; 31:79–83. [PubMed: 15671813]
- Dean FB, Hosono S, Fang L, et al. Comprehensive human genome amplification using multiple displacement amplification. *Proceedings of National Academy of Sciences of the United States of America*. 2002; 99:5261–6.
- Eckert KA, Kunkel TA. DNA polymerase fidelity and the polymerase chain reaction. *PCR Methods and Applications*. 1991; 1:17–24. [PubMed: 1842916]
- Esteban JA, Salas M, Blanco L. Fidelity of phi 29 polymerase. Comparison between protein-primed initiation and DNA polymerization. *The Journal of Biological Chemistry*. 1993; 268:2719–26. [PubMed: 8428945]
- Feinberg AP, Vogelstein B. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Analytical Biochemistry*. 1983; 1:6–13. [PubMed: 6312838]
- Gerber J, Wenaweser D, Heitz-Mayfield L, Lang NP, Persson GR. Comparison of bacterial plaque samples from titanium implant and tooth surfaces by different methods. *Clinical Oral Implants Research*. 2006; 17:1–7. [PubMed: 16441779]
- Gomes BP, Pinheiro ET, Gade-Neto CR, et al. Microbiological examination of infected dental root canals. *Oral Microbiology and Immunology*. 2004; 19:71–6. [PubMed: 14871344]
- Haffajee AD, Socransky SS, Patel MR, Song X. Microbial complexes in supragingival plaque. *Oral Microbiology and Immunology*. 2008; 23:196–205. [PubMed: 18402605]
- Hawkins TL, Detter JC, Richardson P. Whole genome amplification: applications and advances. *Current Opinion in Biotechnology*. 2002; 13:65–7. [PubMed: 11849960]
- Jacinto RC, Gomes BPFA, Desai M, Rajendram D, Shah HN. Bacterial examination infections by clonal analysis in concert with denaturing high-performance liquid chromatography. *Oral Microbiology and Immunology*. 2007; 22:403–10. [PubMed: 17949344]
- Lana MA, Ribeiro-Sobrinho AP, Stehling R, et al. Microorganisms isolated from root canals presenting necrotic pulp and their drug susceptibility in vitro. *Oral Microbiology and Immunology*. 2001; 16:100–5. [PubMed: 11240863]
- Lizardi PM, Huang X, Zhu Z, Bray-Ward P, Thomas DC, Ward DC. Mutation detection and single-molecule counting using isothermal rolling-circle amplification. *Nature Genetics*. 1998; 19:225–32. [PubMed: 9662393]
- Mager DL, Haffajee AD, Socransky SS. Effects of periodontitis and smoking on the microbiota of oral mucous membranes and saliva in systemically healthy subjects. *Journal of Clinical Periodontology*. 2003; 30:1031–7. [PubMed: 15002888]
- Mai M, Hoyer JD, McClure RF. Use of multiple displacement amplification to amplify genomic DNA before sequencing of the alpha and beta haemoglobin genes. *Journal of Clinical Pathology*. 2004; 57:637–40. [PubMed: 15166271]
- Marsh SJ, Largent MD. A bacteriological study of the pulp canals of infected primary molars. *Journal of Dentistry for Children*. 1967; 34:460–70. [PubMed: 4864179]
- Moller AJ. Microbiological examination of root canals and periapical tissues of human teeth. Methodological studies. *Odontologisk Tidskrift*. 1966; 74:Suppl, 380. [PubMed: 5335186]
- Munson MA, Pitt-Ford T, Chong B, Weightman A, Wade WG. Molecular and cultural analysis of the microflora associated with endodontic infections. *Journal of Dental Research*. 2002; 81:761–6. [PubMed: 12407091]
- Nelson JR, Cai YC, Giesler TL, et al. TempliPhi, phi29 DNA polymerase based rolling circle amplification of templates for DNA sequencing. *BioTechniques*. 2002; (suppl):44–7. [PubMed: 12083397]
- Papapanou PN. Population studies of microbial ecology in periodontal health and disease. *Annals of Periodontology*. 2002; 7:54–61. [PubMed: 16013217]
- Pazelli LC, Freitas AC, Ito IY, Souza-Gugelmin MC, Medeiros AS, Nelson-Filho P. Prevalence of microorganisms in root canals of human deciduous teeth with necrotic pulp and chronic periapical lesions. *Brazilian Oral Research*. 2003; 17:367–71. [PubMed: 15107921]

- Pirani C, Bertacci A, Cavrini F, Foschi F, Acquaviva GL, Prati C, Sambri V. Recovery of *Enterococcus faecalis* in root canal lumen of patients with primary and secondary endodontic lesions. *The New Microbiologica*. 2008; 31:235–40. [PubMed: 18623989]
- Raslan N, Wetzel WE. Exposed human pulp caused by trauma and/or caries in primary dentition: a histological evaluation. *Dental Traumatology*. 2006; 22:145–53. [PubMed: 16643290]
- Rolph HJ, Lennon A, Riggio MP, Saunders WP, MacKenzie D, Coldero L, Bagg J. Molecular identification of microorganisms from endodontic infections. *Journal of Clinical Microbiology*. 2001; 39:3282–9. [PubMed: 11526164]
- Ruviere DB, Leonardo MR, da Silva LA, Ito IY, Nelson-Filho P. Assessment of the microbiota in root canals of human primary teeth by checkerboard DNA-DNA hybridization. *Journal of Dentistry for Children*. 2007; 74:118–23. [PubMed: 18477431]
- Sachdeo A, Haffajee AD, Socransky SS. Biofilms in the edentulous oral cavity. *Journal of Prosthodontics*. 2008; 17:348–56. [PubMed: 18355168]
- Sassone L, Fidel R, Figueiredo L, Fidel S, Faveri M, Feres M. Evaluation of the microbiota of primary endodontic infections using checkerboard DNA-DNA hybridization. *Oral Microbiology and Immunology*. 2007; 22:390–7. [PubMed: 17949342]
- Sassone LM, Fidel R, Faveri M, Fidel S, Figueiredo L, Feres M. Microbiological evaluation of primary endodontic infections in teeth with and without sinus tract. *International Endodontic Journal*. 2008; 41:508–15. [PubMed: 18422583]
- Siqueira JF Jr, Roças IN, Souto R, Uzeda M, Colombo AP. Checkerboard DNADNA hybridization analysis of endodontic infections. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics*. 2000; 89:744–8.
- Smith GL, Socransky SS, Smith CM. Rapid method for the purification of DNA from subgingival microorganisms. *Oral Microbiology and Immunology*. 1989; 4:47–51. [PubMed: 2628868]
- Socransky SS, Smith C, Martin L, Paster BJ, Dewhirst FE, Levin AE. “Checkerboard” DNA-DNA hybridization. *BioTechniques*. 1994; 17:788–92. [PubMed: 7833043]
- Socransky SS, Haffajee AD, Smith C, et al. Use of checkerboard DNA-DNA hybridization to study complex microbial ecosystems. *Oral Microbiology and Immunology*. 2004; 19:352–62. [PubMed: 15491460]
- Socransky SS, Haffajee AD. Periodontal microbial ecology. *Periodontology 2000*. 2005; 38:135–87. [PubMed: 15853940]
- Sundqvist, G. PhD Dissertation. University of Umea; Umea, Sweden: 1976. Bacteriologic Studies of Necrotic Dental Pulp.
- Sundqvist G. Ecology of the root canal flora. *Journal of Endodontics*. 1992; 18:427–30. [PubMed: 9796509]
- Teles F, Haffajee AD, Socransky SS. Multiple displacement amplification as an aid in checkerboard DNA-DNA hybridization. *Oral Microbiology and Immunology*. 2007; 22:118–25. [PubMed: 17311635]
- Teles FR, Haffajee AD, Socransky SS. The reproducibility of curet sampling of subgingival biofilms. *Journal of Periodontology*. 2008; 79:705–13. [PubMed: 18380565]
- Toyoshima Y, Fukushima H, Inoue JI, et al. A bacteriological study of periapical pathosis on deciduous teeth. *Shoni Shikagaku Zasshi*. 1988; 26:449–58. [PubMed: 3272472]
- Vianna ME, Horz HP, Conrads G, Feres M, Gomes BP. Comparative analysis of endodontic pathogens using checkerboard hybridization in relation to culture. *Oral Microbiology and Immunology*. 2008; 23:282–90. [PubMed: 18582327]
- Yan J, Feng J, Hosono S, Sommer SS. Assessment of multiple displacement amplification in molecular epidemiology. *BioTechniques*. 2004; 37:136–8. [PubMed: 15283212]
- Zehnder M, Guggenheim B. The mysterious appearance of enterococci in filled root canals. *International Endodontic Journal*. 2009; 42:277–87. [PubMed: 19220511]





Figure 1. Mean prevalence (% of teeth colonized by counts of $>10^4$) of individual species in root canal samples from primary teeth. The data are ordered in descending order of prevalence in the amplified samples.

**Figure 2.**

Quantitative analysis of Microbiota of 32 deciduous endodontic infections analyzed by multiple displacement amplification (MDA) and Checkerboard DNA-DNA hybridization represented by the percentage of mean proportion DNA probe counts (\pm SEM) of test species. The percentage of the DNA probe count was computed for each species for each sample and averaged across samples. The data are ordered in descending order of mean percentages of DNA probe counts detected in amplified samples.

Table 1

Strains of bacterial species used to prepare DNA probes and standards

Strain *	Strain *
<i>Acinetobacter baumannii</i> (19606) [#]	<i>Lactobacillus casei</i> (393) [#]
<i>Actinomyces georgiae</i> (49285) [#]	<i>Legionella pneumophila</i> (33153) [#]
<i>Actinomyces gerencseriae</i> (23860) [#]	<i>Leptotrichia buccalis</i> (14201) [#]
<i>Actinomyces israelii</i> (12102) [#]	<i>Neisseria mucosa</i> (19696) [#]
<i>Actinomyces meyeri</i> (35568) [#]	<i>Peptostreptococcus anaerobius</i> (27337) [#]
<i>Actinomyces naeslundii I</i> (12104) [#]	<i>Parvimonas micra</i> (33270) [#]
<i>Actinomyces naeslundii II</i> (43146)	<i>Porphyromonas endodontalis</i> (35406) [#]
<i>Actinomyces odontolyticus</i> (17929) [#]	<i>Porphyromonas gingivalis</i> (33277) [#]
<i>Aggregatibacter (Actinobacillus) actinomycetemcomitans</i> **	<i>Prevotella denticola</i> (35308) [#]
<i>Aggregatibacter (Actinobacillus) actinomycetemcomitans</i> **	<i>Prevotella nigrescens</i> (33563) [#]
<i>Aggregatibacter (Haemophilus) aphrophilus</i> (33389) [#]	<i>Prevotella heparinolytica</i> (35895) [#]
<i>Aggregatibacter (Haemophilus) paraphrophilus</i> (29242) [#]	<i>Prevotella intermedia</i> (25611) [#]
<i>Aggregatibacter (Haemophilus) segnis</i> (33393) [#]	<i>Prevotella loescheii</i> (15930) [#]
<i>Atopobium parvulum</i> (33793) [#]	<i>Prevotella melaninogenica</i> (25845) [#]
<i>Bacteroides fragilis</i> (25285) [#]	<i>Prevotella oris</i> (33573) [#]
<i>Campylobacter concisus</i> (33237) [#]	<i>Prevotella tanneriae</i> (51259) [#]
<i>Campylobacter ureolyticus</i> (33387) [#]	<i>Propionibacterium propionicum</i> (14157) [#]
<i>Campylobacter gracilis</i> (33236) [#]	<i>Propionibacterium acnes I</i> ***
<i>Campylobacter rectus</i> (33238) [#]	<i>Propionibacterium acnes II</i> ***
<i>Campylobacter showae</i> (51146) [#]	<i>Rothia dentocariosa</i> (17931) [#]
<i>Capnocytophaga gingivalis</i> (33624) [#]	<i>Selenomonas noxia</i> (43541) [#]
<i>Capnocytophaga ochracea</i> (33596) [#]	<i>Selenomonas sputigena</i> (35185) [#]
<i>Capnocytophaga sputigena</i> (33612) [#]	<i>Slackia exigua</i> (700122) [#]
<i>Corynebacterium matruchotii</i> (14266) [#]	<i>Staphylococcus aureus</i> (33591)
<i>Dialister pneumosintes</i> (GBA27)	<i>Staphylococcus epidermidis</i> (14990) [#]
<i>Eikenella corrodens</i> (23834) [#]	<i>Staphylococcus warneri</i> (27836) [#]
<i>Enterococcus faecalis</i> (29212)	<i>Streptococcus anginosus</i> (33397) [#]
<i>Enterococcus aerogenes</i> (13048) [#]	<i>Streptococcus constellatus</i> (27823) [#]
<i>Escherichia coli</i> (10799)	<i>Streptococcus gordonii</i> (10558) [#]
<i>Eubacterium limosum</i> (8486) [#]	<i>Streptococcus intermedius</i> (27335) [#]
<i>Eubacterium nodatum</i> (33099) [#]	<i>Streptococcus mitis</i> (49456) [#]
<i>Eubacterium saburreum</i> (33271) [#]	<i>Streptococcus mutans</i> (25175) [#]
<i>Filifactor alocis</i> (35896) [#]	<i>Streptococcus oralis</i> (35037) [#]

Strain*	Strain*
<i>Fusobacterium necrophorum</i> (25286) [#]	<i>Streptococcus parasanguinis</i> (15912) [#]
<i>Fusobacterium nucleatum ss nucleatum</i> (25586) [#]	<i>Streptococcus salivarius</i> (27945)
<i>Fusobacterium nucleatum ss polymorphum</i> (10953) [#]	<i>Streptococcus sanguinis</i> (10556) [#]
<i>Fusobacterium nucleatum ss vincentii</i> (49256) [#]	<i>Streptococcus vestibularis</i> (49124) [#]
<i>Fusobacterium periodonticum</i> (33693) [#]	<i>Tannerella forsythia</i> (43037) [#]
<i>Gemella haemolysans</i> (10379) [#]	<i>Treponema denticola</i> (B1)
<i>Gemella morbillorum</i> (27824) [#]	<i>Treponema socranskii</i> (SI)
<i>Haemophilus influenza</i> (33533) [#]	<i>Veillonella dispar</i> (17748) [#]
<i>Helicobacter pylori</i> (43504) [#]	<i>Veillonella parvula</i> (10790) [#]
<i>Lactobacillus acidophilus</i> (4356) [#]	

* All strains were obtained from the American Type Culture Collection (ATCC number in parenthesis) except for *Treponema denticola* (B1) and *Treponema socranskii* (SI), which were obtained from The Forsyth Institute

** ATCC strains 43718 and 29523

*** ATCC strains 11827 and 11828

[#]Type strains