



Acute Imidacloprid Exposure Alters Mitochondrial Function in Bumblebee Flight Muscle and Brain

Chloe Sargent^{1*}, Brad Ebanks², Ian C. W. Hardy^{1,2,3}, T. G. Emyr Davies⁴,
Lisa Chakrabarti^{2,5} and Reinhard Stöger¹

¹ School of Biosciences, Sutton Bonington Campus, University of Nottingham, Loughborough, United Kingdom, ² School of Veterinary Medicine and Science, Sutton Bonington Campus, University of Nottingham, Loughborough, United Kingdom, ³ Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland, ⁴ Department of Biointeractions and Crop Protection, Rothamsted Research, Harpenden, United Kingdom, ⁵ Medical Research Council Versus Arthritis Centre for Musculoskeletal Ageing Research, Birmingham, United Kingdom

OPEN ACCESS

Edited by:

Léa Tison,
INRAE Nouvelle-Aquitaine
Bordeaux, France

Reviewed by:

Michael B. Powner,
City University of London,
United Kingdom
Harmen P. Hendriksma,
Julius Kühn-Institut-
Braunschweig, Germany

*Correspondence:

Chloe Sargent
Chloe.Sargent@nottingham.ac.uk

Specialty section:

This article was submitted to
Insect Neurobiology,
a section of the journal
Frontiers in Insect Science

Received: 26 August 2021

Accepted: 12 November 2021

Published: 01 December 2021

Citation:

Sargent C, Ebanks B, Hardy ICW,
Davies TGE, Chakrabarti L and
Stöger R (2021) Acute Imidacloprid
Exposure Alters Mitochondrial
Function in Bumblebee Flight Muscle
and Brain.
Front. Insect Sci. 1:765179.
doi: 10.3389/finsc.2021.765179

Mitochondria are intracellular organelles responsible for cellular respiration with one of their major roles in the production of energy in the form of ATP. Activities with increased energetic demand are especially dependent on efficient ATP production, hence sufficient mitochondrial function is fundamental. In bees, flight muscle and the brain have particularly high densities of mitochondria to facilitate the substantial ATP production required for flight activity and neuronal signalling. Neonicotinoids are systemic synthetic insecticides that are widely utilised against crop herbivores but have been reported to cause, by unknown mechanisms, mitochondrial dysfunction, decreasing cognitive function and flight activity among pollinating bees. Here we explore, using high-resolution respirometry, how the neonicotinoid imidacloprid may affect oxidative phosphorylation in the brain and flight muscle of the buff-tailed bumblebee, *Bombus terrestris*. We find that acute exposure increases routine oxygen consumption in the flight muscle of worker bees. This provides a candidate explanation for prior reports of early declines in flight activity following acute exposure. We further find that imidacloprid increases the maximum electron transport capacity in the brain, with a trend towards increased overall oxygen consumption. However, intra-individual variability is high, limiting the extent to which apparent effects of imidacloprid on brain mitochondria are shown conclusively. Overall, our results highlight the necessity to examine tissue-specific effects of imidacloprid on respiration and energy production.

Keywords: *Bombus terrestris*, mitochondria, imidacloprid, oxidative phosphorylation, high-resolution respirometry

INTRODUCTION

Intensification of agriculture has driven a global increase in pesticide use and fragmentation of pollinator habitats, often leading to a sparsity of resources within the foraging range of bees (1, 2). This requires bees to fly greater distances, thus increasing the energy demand of foraging. At the cellular level, energy is produced in the form of adenosine triphosphate (ATP) *via* oxidative phosphorylation (OXPHOS) within mitochondria (3). During OXPHOS, ATP is synthesised *via* the electron transport chain located in the inner mitochondrial membrane; electrons are transported *via* a series of carriers and protons are pumped into the intermembrane space producing a proton gradient which

drives ATP synthesis *via* ATP synthase (4). It has been reported that the rate of electron transfer between OXPHOS enzymes in bees are the highest measured in any animal (5).

Carbohydrate, predominately in the form of trehalose, is the main source of energy in the nervous system and flight muscles of bees, however tissue-specific differences have been shown (5, 6). In the brain, evidence of β -oxidation to metabolise fatty acids and contribute to the high energetic demands has been reported, however there is little evidence of fatty acids fuelling flight in the thoracic muscle (6). Furthermore, glycogen is stored in the flight muscles which may be utilised to extend flight duration (5). These differences reinforce the importance of looking at multiple tissue types when analysing the mitochondrial function in bees.

In bees and other hymenopterans, the brain and flight muscle have particularly high densities of mitochondria to facilitate the substantial energy demands of neuronal signalling and flight activity (3, 5, 7). There is increasing evidence of an association between brain mitochondrial OXPHOS and behaviour in bees, suggesting that alterations in brain mitochondrial function, such as those caused by neonicotinoids, could induce changes in behaviours, such as aggression (7). External factors that affect mitochondrial function may have the greatest impact within tissues with the highest energy demands.

Neonicotinoids are a widely used group of systemic insecticides which have been shown to affect pollinator learning, memory, homing, and flight capacity (8, 9). These compounds target nicotinic acetylcholine receptors (nAChRs); the most prevalent excitatory neurotransmitter in the central nervous system of insects (10, 11). Neonicotinoids impair mitochondrial function and structure in bumblebees and honeybees (12–14) and exposure to the neonicotinoid imidacloprid leads to premature flight exhaustion and altered cognitive performance (15, 16). This may be associated with a transient excitatory affect caused by the overstimulation of nAChRs by imidacloprid and potentially lead to a reduction in flight activity and foraging capacity (15). The decline in flight activity associated with pesticide exposure may be due to insufficient ATP production in the flight muscle leading to premature exhaustion and flight inactivity. As mitochondria are responsible for the majority of ATP production during insect flight, an imidacloprid-induced impairment of mitochondrial function may impact on flight performance and neuronal signalling (17).

Here we seek to determine if and how acute exposure to field-realistic doses of imidacloprid may affect mitochondrial function and OXPHOS in the bumblebee, *Bombus terrestris*. Using high-resolution respirometry we analyse mitochondrial respiration in the brain and flight muscle of female worker bees (18). We analyse the oxygen consumption of three different respiration states: (i) Routine—the rate at which OXPHOS occurs in cells in the physiological coupling state, where ATP synthesis is coupled with the electron transport chain (ii) LEAK—a non-phosphorylating resting state of uncoupled respiration by the inhibition of ATP synthase; hence, oxygen consumption is associated with proton leak through the inner mitochondrial membrane and not *via* ATP synthesis-linked respiration; and (iii) maximum electron transport capacity (ET capacity)—the maximum rate of the electron transport pathway when not

coupled to ATP synthase and at an optimum concentration of uncoupler (18). By analysing these three states we aim to identify if and how imidacloprid may affect the OXPHOS system.

MATERIALS AND METHODS

Bee Husbandry

Commercial queen-right *B. terrestris audax* colony were obtained from Biobest® (Westerlo, Belgium) between November 2020 and June 2021, and maintained at 26°C and 33% relative humidity. We fed the colony and age-matched laboratory reared bees on 2.0M sucrose solution and pollen (purchased from Agralan, UK) provided *ad libitum*. To create cohorts of bees of a similar age, pupae were harvested and stored in a separate plastic container adjacent to the colony until eclosion. Once eclosed, bees were placed into cohorts of 1–3 bees and contained in plastic deli pots (115 × 75 mm) until used for high-resolution respirometry (HRR).

Imidacloprid Treatment

Individual bees were taken from their cohort and placed in a queen marker tube where they were starved for 1 h to increase the likelihood of feeding on the diet subsequently provided. For the insecticide exposed bees, a 2.92 M sucrose solution containing 10 ppb imidacloprid was then used to feed the bees. This field-realistic concentration was selected as it has been previously reported in pollen and nectar, foraging bees, and inside colonies (15). For the control bees, feeding was with 2.92 M sucrose solution only. Bees were allowed to feed for 10 mins and the quantify consumed was recorded. Any bees which did not feed during this period were excluded from the experiment. Thirty minutes after the end of feeding, bees were cold anaesthetised (flight muscle $n = 4$, and brain $n = 7$; for both control and IMD groups).

Flight Muscle and Brain Tissue Dissection

Cold-anaesthetised bees were killed by decapitation. The thorax was photographed to later measure the inter-tegular distance (representing body size) for each bee using ImageJ software (32). The brain or a small section of flight muscle were removed, using a scalpel and forceps, and then individually weighed in 1.5 ml tubes. Individual bees representing biological replicates were used for each tissue type sample. Then 100 μ l of MiR05 buffer (0.5 mM EGTA, 3 mM MgCl₂, 60 mM Lactobionic acid, 20 mM Taurine, 10 mM KH₂PO₄, 20 mM HEPES, 110 mM D-Sucrose, 1g/L BSA, pH 7.1) per 1 mg of muscle, or one whole brain, was added and the tissue mechanically homogenised. Then 100 μ l (1 mg) of muscle tissue homogenate and all the brain homogenate were transferred into a high-resolution respirometry chamber for analysis.

High-Resolution Respirometry

High-resolution respirometry (HRR) was carried out using an Oroboros Oxygraph-2k (Oroboros® Instruments, Innsbruck, Austria). The respirometer electrodes were calibrated daily to ensure oxygen concentration remained consistent for the duration of the experiment. All respiratory analyses were

performed at 35°C as previous studies have reported this close to the average thoracic and head temperature of *B. terrestris* during flight (19, 20). Oxygen consumption of one brain and 1 mg flight muscle were analysed using the SUT-003 protocol (18, 21). Routine respiration was measured followed by the addition of 1.0 μl of 5 μM oligomycin to determine LEAK state (inhibition of ATP synthase). Uncoupled maximal electron transport state was then measured by subsequent 10.0 μl titrations of 0.5 μM carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP). Finally, 1 μl of 2.5 μM Antimycin A to determine the residual oxygen consumption (ROX) (Figure 1).

Data Analysis

Raw data outputs were acquired using the O2k-Software (DatLab v7.4.0.4, Oroboros), for data acquisition and analysis to determine instrumental background corrected oxygen flux values. Data were subsequently analysed using R Team (22). Unpaired Student's *t*-tests were performed to determine whether there were differences between the mean oxygen flux at routine, LEAK and maximum ET capacity states, as well as the mean FCRs for routine and LEAK. The FCR can be defined as the ratio of oxygen flux in the different respiration states (routine and LEAK) that are normalised to obtain lower [residual oxygen

consumption (ROX)] and upper (maximum ET capacity) limits of 0–1 (23). Each of these tests used adopted a Type I error rate of $\alpha = 0.5$, with interpretations controlled for multiple comparisons as detailed in the supplementary materials. Oxygen consumption was also analysed in relation to the amount of sucrose consumed, bee age, and bee size (inter-tegular distance, ITD). There was no significant difference found in mean bee age or size between treatment groups in either tissue (Supplementary Table 2). However, as a range of ages check were used in both groups, one-way ANOVAs were used to check for potential age-related effects on mitochondrial function: age did not significantly affect any of the respiration states in the flight muscle or brain. To determine whether the amount of diet (sucrose solution with or without imidacloprid) consumed correlated with oxygen flux, Pearson's correlation coefficients were calculated; there was no effect in either the flight muscle or brain (Supplementary Table 3).

RESULTS

Flight Muscle

Acute oral exposure of imidacloprid in worker bees significantly increased routine respiration in flight muscle (Oxygen flux: $t_{df=6} = -3.58$, $p = 0.012$; FCR: $t_{df=6} = -2.50$, $p = 0.047$,




FIGURE 1 | Annotated representation of the O₂k oxygraph output. The red line is the oxygen flux. Routine is shown before the addition of Omy. LEAK state can be determined after the inhibition of ATP synthase by Omy. The maximum electron capacity is determined as the highest peak after CCCP titrations. The spare respiratory capacity and its relative value can be calculated *via* the calculation, [(ET capacity specific flux)/(Routine specific flux × 100)]. AmA, a complex III inhibitor, is added to determine the residual oxygen consumption (ROX) as the baseline state and allow for background correction. Abbreviations: Omy: oligomycin; CCCP: carbonyl cyanide *m*-chlorophenyl hydrazone; AmA: antimycin A.

Figure 2; Table 1, note that the difference in FCR did not remain significant after correction for multiple comparisons, **Supplementary Table 1**). There were no significant differences found between the LEAK and maximum ET capacity states when comparing exposed and control bees (LEAK: oxygen flux, $t_{df=6} = -0.96$, $p = 0.376$; FCR, $t_{df=6} = -1.45$, $p = 0.196$. Maximum ET capacity: oxygen flux, $t_{df=6} = 1.72$, $p = 0.136$, **Figure 2; Table 1**).

The spare respiratory capacity (SRC) was also calculated for the flight muscle. The SRC describes the mitochondrial capacity to reach additional ATP demands that are greater than routine respiration levels and can be expressed as a quantitative value using the formula, $SRC = [(ET \text{ capacity specific flux}) / (\text{Routine specific flux} \times 100)]$ (24) (**Figure 1**). Hence, the mean SRCs were 0.069 and 0.035 for the control and imidacloprid groups respectively. This may be explained by the increase in routine respiration but not the maximum ET capacity by imidacloprid which would result in a lower SRC.

Brain

In brain tissue we found that the oxygen flux for imidacloprid fed bees had a significantly higher maximum ET capacity ($t_{df=12} = -2.76$, $p = 0.017$), but not in Routine and LEAK ($t_{df=12} = -2.08$, $p = 0.060$ and $t_{df=12} = -1.80$, $p = 0.097$, respectively) (**Figure 3; Table 1**). However, there were no significant differences between

the treatment groups in terms of FCRs (Routine: $t_{df=12} = 0.48$, $p = 0.639$; LEAK: $t_{df=12} = 0.43$, $p = 0.674$). The SRC values were similar for both groups; 0.011 and 0.009 for control and imidacloprid treated bees, respectively.

DISCUSSION

Acute imidacloprid exposure at the established field-realistic concentration of 10 ppb has been previously shown to affect flight activity in *B. terrestris* by increasing short-term velocity and decreasing the overall duration and distance flown (15, 25, 26). This study has further shown that exposure to imidacloprid, at a concentration of 10 ppb, increases routine oxygen consumption in the flight muscles of workers. Imidacloprid targets acetylcholine receptors and can result in over-stimulation of the nervous system, initiating an excitatory response (10). We suggest that an increase in oxygen consumption and respiration initiated by this response enables the short-term increase in flight velocity that has been previously reported (15, 27). The enhanced flight velocity in turn may lead to premature exhaustion resulting in a shorter flight duration (15, 28, 29). Hyperactivity caused by neonicotinoids, such as imidacloprid, could also result in long-term muscle exhaustion and impaired energy metabolism as shown with thiamethoxam by Tosi et al. (27).





FIGURE 2 | Effects of imidacloprid on flight muscle mitochondria at Routine, LEAK, and maximum ET capacity. States **(A)** Mean oxygen flux; and **(B)** Mean FCR, in flight muscle of imidacloprid and control worker bees. An unpaired students *t*-test was performed to determine significance between treatment groups. * $p < 0.05$. *p*-values marked with † were also significant after controlling for Type I error rates. Error bars show 95% confidence interval. $n = 4$ for each of the control and imidacloprid groups. Routine: routine respiration rate; LEAK: electron/H⁺ leak; ET capacity: maximum electron transport capacity.

TABLE 1 | Mean oxygen flux and flux control ratio values in the brain and flight muscle of imidacloprid and non-imidacloprid fed worker bees.

	Flight Muscle			Brain		
	Control <i>n</i> = 4	IMD <i>n</i> = 4	<i>p</i>	Control <i>n</i> = 7	IMD <i>n</i> = 7	<i>p</i>
Routine						
Oxygen flux						
Mean (s.e.)	6.48 (0.32)	9.20 (0.69)	0.012*[†]	2.75 (0.60)	7.23 (2.07)	0.060
Variance	0.40	1.91		2.54	30.02	
FCR						
Mean (s.e.)	0.16(0.03)	0.30(0.05)	0.047*	1.18 (0.38)	0.99 (0.13)	0.639
LEAK						
Oxygen flux						
Mean (s.e.)	3.33 (0.56)	4.24 (0.76)	0.376	1.62 (0.37)	4.82 (1.74)	0.097
Variance	1.24	2.32		0.97	21.24	
FCR						
Mean (s.e.)	0.08(0.03)	0.13(0.02)	0.196	0.73 (0.26)	0.60 (0.15)	0.67
ET Capacity						
Oxygen flux						
Mean (s.e.)	44.60 (6.14)	32.21 (3.75)	0.136	3.11 (0.59)	6.59 (1.12)	0.017*[†]
Variance	150.83	56.38		2.42	8.76	

No flux control ratio (FCR) data are provided for the maximum electron transport capacity (ET capacity) as this is set to 1 to obtain the FCR for the routine and electron/H⁺ leak (LEAK) states. *p*-values were obtained from unpaired Student's *t*-tests, with significance: **p* < 0.05. *p*-values marked with [†] were also significant after controlling for Type I error rates (Supplementary Table 1). IMD: imidacloprid; s.e.: ±1 standard error.



The higher routine oxygen flux and low spare respiratory capacity detected in the flight muscle of bees exposed to imidacloprid may be further factors contributing to the negative effects on flight activity. The SRC is defined as the mitochondrial capacity to reach additional ATP demands that are greater than routine respiration levels, and to thus prevent ATP crisis (24). This is important in times of stress, exercise, and increased workload when the extra capacity is needed. Hence, the lower SRC in the imidacloprid-exposed bees, which was approximately half of the value for unexposed control bees, suggests a greater mitochondrial dysfunction during times of high ATP demand that is above the routine respiration rate, such as during flight. Imidacloprid increased routine respiration but did not affect the maximum ET capacity; the knock-on effect is likely a diminished SRC. We currently do not know for how long the effects of acute imidacloprid exposure on mitochondria functions last and if the resulting increases in routine respiration are transient. Therefore, the duration of lowered SRC will need to be determined in future follow-up studies.

In the brain, we detected no significant difference in routine and LEAK oxygen consumption between imidacloprid-exposed and control bees, however the maximum ET capacity in imidacloprid-exposed group showed a greater oxygen consumption. This indicates that acute imidacloprid-exposure results in elevated levels of oxygen consumption and thus, in higher maximum respiration rates. However, there was no detected difference in routine or LEAK FCRs, which are likely to be associated with the higher mean routine and LEAK oxygen consumption values. In addition, the FCR values suggest that the routine and LEAK values are in a similar ratio to the maximum ET capacity for both treatment groups. The similarity in the SRC values (0.011 for the control treatment and 0.009 for imidacloprid exposed bees) also illustrates this; these low values indicate that the routine respiration rate in the brain is at maximum ET capacity, as a SRC value of 0.01 represents that the oxygen consumption during routine and maximum ET capacity are equal. Therefore, it appears that imidacloprid increases overall oxygen consumption in the brain; however, due to high inter-individual variance, our results remain inconclusive. This high variability could be associated with the quantity of nAChRs in the brain, which is much greater than in the flight muscle and may therefore explain why there is lower variance among the flight muscles of imidacloprid-exposed bees and among control groups. Cognitive ability such as learning and memory varies among individual bees; this suggests that the brain is particularly susceptible to interspecific variation (30, 31). Hence, a possible variation in the number of neonicotinoid target sites among individual bees may lead to more variability in the response of imidacloprid on mitochondrial function.

Bees are exposed to imidacloprid both acutely and chronically, hence it is important to elucidate the mitochondrial function of both acute and chronic exposure. Chronic exposure of imidacloprid can lead to neuronal dysfunction (13). Neurons require high levels of ATP, and a stable mitochondrial membrane potential is crucial to sustain ATP production at this level

and thereby normal neuronal function (13). Moffat et al. (13) also reported that acute exposure of 10 nM (2.5 ppb) imidacloprid did not reach concentrations in the brain to induce membrane depolarisation. In contrast, 2 day chronic exposure of 1 nM led to an increased sensitivity of bumblebee neurones to acetylcholine, and elicited mitochondrial depolarisation of neurons (13). Our findings are in agreement with Nicodemo and colleagues (12), who reported tissue-specific sensitivity to acute imidacloprid exposure in the head and thorax of honeybees and negative effects on ATP synthesis. Our results will help resolve how acute imidacloprid exposure causes a different signature of mitochondrial activity compared to effects of chronic exposure. Moreover, it may now be possible to delineate where acute exposure ends and chronic effects begin.

An aspect of our study is that the worker bees were kept in an enclosed colony and were unable to fly. While this allowed a great deal of experimental control, it should be considered that exercise, such as foraging flights, can increase maximum ET capacity. Hence, mitochondrial function from laboratory-reared non-flying worker bees may vary from that of wild, free-flying and foraging worker bees. Another consideration is that social interactions may also impact brain energetics and mitochondrial function; all bees were kept in small cohorts within small containers from the time of adult eclosion, thus removing them from wider colony interactions and pheromones (7). This may impact brain mitochondrial respiration and behaviour, which may in turn not provide a true representation of *B. terrestris* in a natural or agricultural environment. The effects of acute imidacloprid were only analysed at one time point, shortly after exposure; potential long-term implications of acute exposure on the mitochondrial function in bees remain to be determined. For example, whether mitochondria exposed to imidacloprid return to pre-exposure functioning and, if so, how long such recovery may take.

In this study we have further scrutinised how imidacloprid affects the different mitochondrial respiration states. We found that imidacloprid did not affect LEAK, indicating an increase in oxygen consumption is not associated with proton leak and thus imidacloprid does not seem to have an inhibitory effect on ATP synthase. The increase in routine respiration in the muscle suggests imidacloprid may affect the rate at which electrons flow through the electron transport chain, and thereby increases the rate closer to the maximum ET capacity *via* utilisation of the SRC. A similar mechanism of increased electron transport rate could also explain how imidacloprid increased the maximum ET capacity in the brain. The SRC values were much lower in the brain compared to the flight muscle. This difference between the two tissues could explain how imidacloprid was able to increase the maximum ET capacity in the brain and not the flight muscle as only a slight increase in the rate would be capable of surpassing the maximum ET capacity. The exact mechanism by which imidacloprid affects the electron transport chain requires further investigation.

Overall, we found that acute exposure to imidacloprid, *via* feeding, increased the routine respiration rate in the flight muscle and showed a similar, marginally non-significant, trend towards higher respiration rates in the brain of workers. The higher routine respiration rates in the flight muscle may lead to short-term increases in oxygen consumption and potentially increase ATP production equating to a longer-term deprivation of energy. These results accord with those of previous studies which have shown flight duration and distance to be shorter in bumblebees exposed to imidacloprid. Negative impacts of flight activity will increase exhaustion and mortality of individual bees during flight and this reduced foraging capacity will lead to a reduction in pollination services within natural and agricultural landscapes. The broad range of responses in the brain by individual bees makes it difficult to draw clear conclusions from the measurements. Hence, more research is required to establish the impact of imidacloprid on the brain and elucidate possible explanations for the inconsistencies observed.

These results further contribute to our understanding of how sub-lethal levels of imidacloprid mediate tissue-specific effects on respiration and energy production and thereby impacts pollinator fitness. This study will provide a foundation for further research into the link between mitochondrial function and neonicotinoid exposure, as well as assist in the development and assessment of novel pesticides that minimise harmful effects on non-target species.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author.

REFERENCES

- Kremen C, Williams NM, Thorp RW. Crop pollination from native bees at risk from agricultural intensification. *Proc Natl Acad Sci U S A*. (2002) 99:16812–6. doi: 10.1073/pnas.262413599
- Gonzalez-Varo JB, Biesmeijer JC, Bommarco R, Potts SG, Schweiger O, Smith HG, et al. Combined effects of global change pressures on animal-mediated pollination. *Trends Ecol Evol*. (2013) 28:524–530. doi: 10.1016/j.tree.2013.05.008
- Hedges CP, Wilkinson RT, Devaux JBL, Hickey AJR. Hymenoptera flight muscle mitochondrial function: Increasing metabolic power increases oxidative stress. *Comp Biochem Physiol A Mol Integr Physiol*. (2019) 230:115–21. doi: 10.1016/j.cbpa.2019.01.002
- Chaban Y, Boekema EJ, Dudkina NV. Structures of mitochondrial oxidative phosphorylation supercomplexes and mechanisms for their stabilisation. *Biochim Biophys Acta Bioenerg*. (2014) 1837:418–26. doi: 10.1016/j.bbabi.2013.10.004
- Suarez RK, Darveau CA, Welch KC, O'Brien DM, Roubik DW, Hochachka PW. Energy metabolism in orchid bee flight muscles: carbohydrate fuels all. *J Exp Biol*. (2005) 208:3573–79. doi: 10.1242/jeb.01775
- Rittschof CC, Schirmeier S. Insect models of central nervous system energy metabolism and its links to behavior. *Glia*. (2018) 66:1160–75. doi: 10.1002/glia.23235
- Rittschof CC, Vekaria HJ, Palmer JH, Sullivan PG. Brain mitochondrial bioenergetics change with rapid and prolonged shifts in aggression in the honey bee, *Apis mellifera*. *J Exp Biol*. (2018) 221:176917 doi: 10.1242/jeb.176917
- Gill RJ, Raine NE. Chronic impairment of bumblebee natural foraging behaviour induced by sublethal pesticide exposure. *Funct Ecol*. (2014) 28:1459–71. doi: 10.1111/1365-2435.12292
- Stanley DA, Smith KE, Raine NE. Bumblebee learning and memory is impaired by chronic exposure to a neonicotinoid pesticide. *Sci Rep*. (2015) 5:10. doi: 10.1038/srep16508
- Buckingham SD, Lapiéd B, LeCorronc H, Grolleau F, Sattelle DB. Imidacloprid actions on insect neuronal acetylcholine receptors. *J Exp Biol*. (1997) 21:2685–92. doi: 10.1242/jeb.200.21.2685
- Colgan TJ, Fletcher IK, Arce AN, Gill RJ, Ramos Rodrigues A, Stolle E, et al. Caste- and pesticide-specific effects of neonicotinoid pesticide exposure on gene expression in bumblebees. *Mol Ecol*. (2019) 28:1964–74. doi: 10.1111/mec.15047
- Nicodemo D, Maioli MA, Medeiros HCD, Guelfi M, Balieira KVB, De Jong D, et al. Fipronil and imidacloprid reduce honeybee mitochondrial activity. *Environ Toxicol Chem*. (2014) 33:2070–5. doi: 10.1002/etc.2655
- Moffat C, Pacheco JG, Sharp S, Samson AJ, Bolland KA, Huang J, et al. Chronic exposure to neonicotinoids increases neuronal vulnerability to mitochondrial dysfunction in the bumblebee (*Bombus terrestris*). (2015) *Faseb J*. 29:2112–9. doi: 10.1096/fj.14-267179
- Friol PS, Catae AF, Tavares DA, Malaspina O, Roat TC. Can the exposure of *Apis mellifera* (Hymenoptera, Apidae) larvae to a field concentration of thiamethoxam affect newly emerged bees? *Chemosphere*. (2017) 185:56–66. doi: 10.1016/j.chemosphere.2017.06.113

AUTHOR CONTRIBUTIONS

CS and RS designed the experiment with the additional support of LC and BE. Bee husbandry, laboratory work, and data collection were performed by CS. HRR was performed by CS with the support of LC and BE. Data analysis was performed by CS with supplementary analysis by CS and IH. Figures were prepared by CS. Paper was written by CS and RS with additional editorial assistance by LC, BE, IH, and TD. All authors contributed to the article and approved the submitted version.

FUNDING

CS is a PhD candidate funded by the Future Food Beacon/Graduate Centre for International Agriculture, University of Nottingham and Rothamsted Research and BE is a PhD candidate funded by the University of Nottingham Biotechnology and Biological Sciences Doctoral Training Programme (BBSRC-DTP).

ACKNOWLEDGMENTS

The authors would like to thank Peter Graystock, Daniel Kenna, and Richard Gill at Imperial College London for their previous support to the first author and their research which helped to guide this study.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/finsc.2021.765179/full#supplementary-material>

15. Kenna D, Cooley H, Pretelli I, Ramos Rodrigues A, Gill S, Gill R. Pesticide exposure affects flight dynamics and reduces flight endurance in bumblebees. *bioRxiv*. (2018) 449280. doi: 10.1101/449280
16. Schwartz KR, Minor H, Magro C, McConnell J, Capani J, Griffin J, et al. The neonicotinoid imidacloprid alone alters the cognitive behavior in *Apis mellifera* L. and the combined exposure of imidacloprid and *Varroa destructor* mites synergistically contributes to trial attrition. *J Apic Res*. (2020) 60:431–8. doi: 10.1080/00218839.2020.1866233
17. Thompson SN, Suare RK. “Metabolism,” In: Resh VH, Cardé RT editors, *Encyclopedia of Insects*. New York, NY: Academic Press (2009), 623–627.
18. Gnaiger E. *Mitochondrial Pathways and Respiratory Control. An Introduction to OXPHOS* Analysis. 5th Ed. Bioenerg Commun (2020)
19. Reber T, Vahakainu A, Baird E, Weckstrom M, Warrant E, Dacke M. Effect of light intensity on flight control and temporal properties of photoreceptors in bumblebees. (2015) *J Exp Biol*. 218:1339–1346. doi: 10.1242/jeb.113886
20. Teulier L, Weber JM, Crevier J, Darveau CA. Proline as a fuel for insect flight: enhancing carbohydrate oxidation in hymenopterans. *Proc R Soc B*. (2016) 16:283. doi: 10.1098/rspb.2016.0333
21. Bioblast. *SUIT-003*. (2021). Available online at: <https://wiki.oroboros.at/index.php/SUIT-003>. Last. (accessed August 23, 2021).
22. R Core Team. R: A language and Environment for Statistical Computing. Version 3.6.2 (2020). Available online at: <https://www.R-project.org>.
23. Bioblast. Flux control ratio (2021). Available online at: https://wiki.oroboros.at/index.php/Flux_control_ratio. (accessed June 21, 2021).
24. Marchetti P, Fovez Q, Germain N, Khamari R, Kluza J. Mitochondrial spare respiratory capacity: Mechanisms, regulation, and significance in non-transformed and cancer cells. *FASEB J*. (2020) 34:13106–24. doi: 10.1096/fj.202000767R
25. Goulson D. REVIEW: An overview of the environmental risks posed by neonicotinoid insecticides. *J Appl Ecol*. (2013) 50:977–87. doi: 10.1111/1365-2664.12111
26. Botias C, David A, Hill EM, Goulson D. Contamination of wild plants near neonicotinoid seed-treated crops, and implications for non-target insects. *Sci Total Environ*. (2016) 566:269–78. doi: 10.1016/j.scitotenv.2016.05.065
27. Tosi S, Burgio G, Nieh JC. A common neonicotinoid pesticide, thiamethoxam, impairs honey bee flight ability. *Sci Rep*. (2017) 7:8. doi: 10.1038/s41598-017-01361-8
28. Smit WA, Becht G, Beenackers AM. Structure fatigue and enzyme activities in fast insect muscles. *J Insect Physiol*. (1967) 13:12. doi: 10.1016/0022-1910(67)90022-4
29. Usherwood PN. Insect Neuromuscular Mechanisms. *Am Zoologist*. (1967) 7:553–82. doi: 10.1093/icb/7.3.553
30. Dyer AG. The mysterious cognitive abilities of bees: why models of visual processing need to consider experience and individual differences in animal performance. *J Exp Biol*. (2012) 215:387–95. doi: 10.1242/jeb.038190
31. Chittka L. Bee Cognition. *Curr Biol* 27:19. (2017) R1049–53. doi: 10.1016/j.cub.2017.08.008
32. Image J. U. S. *National Institutes of Health, Bethesda, Maryland, USA*. (2021). Available online at: <https://imagej.nih.gov/ij/> 1997–2018

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Sargent, Ebanks, Hardy, Davies, Chakrabarti and Stöger. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.