

L-Cysteine containing vitamin supplement which prevents or alleviates alcohol-related hangover symptoms: nausea, headache, stress and anxiety.

C. J. Peter Eriksson^{1*}, Markus Metsälä^{2,§}, Tommi Möykkynen^{3,§}, Heikki Mäkisalo^{4,§}, Olli Kärkkäinen^{5,§}, Maria Palmén^{6,§}, Joonas E. Salminen^{2,§}, and Jussi Kauhanen^{7,§}

¹Department of Public Health, University of Helsinki, PO Box 20, 00014 University of Helsinki, Tukholmankatu 8 B, Finland. ²Department of Chemistry, University of Helsinki, PO Box 55, 00014 University of Helsinki, A. I. Virtasen aukio 1, Finland. ³Department of Biosciences, Molecular and Integrative Biosciences, University of Helsinki, PO Box 56, 00014 University, Viikinkaari 9, Finland. ⁴Transplantation and Liver Surgery Unit, Helsinki University Hospital, PO Box 372, 00029 HUS, 00014 University of Helsinki, Haartmaninkatu 4, Finland. ⁵School of Pharmacy, University of Eastern Finland, Yliopistonranta 1 C, Kuopio, Finland. ⁶Schildts & Söderströms, Snellmanninkatu 13, Helsinki, Finland. ⁷Institute of Public Health and Clinical Nutrition, University of Eastern Finland, PO Box 1627, 70211 Kuopio, Finland

*Corresponding author: Department of Public Health, University of Helsinki, PO Box 20, 00014 University of Helsinki, Tukholmankatu 8 B, Finland. Tel.; +358407549875; E-mail: peter.cj.eriksson@helsinki.fi

§These authors have equally contributed to this work

Key words: L-cysteine, hangover, nausea, headache, stress, anxiety

Abstract

Aims: Alcohol-related hangover symptoms: nausea, headache, stress and anxiety cause globally considerable amount of health problems and economic losses. Many of these harmful effects are produced by alcohol and its metabolite, acetaldehyde, which also is a common ingredient in alcohol beverages. The aim of the present study is to investigate the effect of the amino acid L-cysteine on the alcohol/acetaldehyde related aftereffects.

Methods: Voluntary healthy participants were recruited through advertisements. Volunteers had to have experience of hangover and/or headache. The hangover study was randomized, double-blind and placebo controlled. Nineteen males randomly swallowed placebo and L-cysteine tablets. The alcohol dose was 1.5 g/kg, which was consumed during 3 hours.

Results: The primary results based on correlational analysis showed that L-cysteine prevents or alleviates hangover, nausea, headache, stress and anxiety. For hangover, nausea and headache the results were apparent with the L-cysteine dose of 1200 mg and for stress and anxiety already with the dose of 600 mg.

Conclusions: L-cysteine would reduce the need of drinking the next day with no or less hangover symptoms: nausea, headache, stress and anxiety. Altogether, these effects of L-cysteine are unique and seems to have a future in preventing or alleviating these harmful symptoms as well as reducing the risk of alcohol addiction.

INTRODUCTION

Alcohol drinking and drunkenness commonly lead to hangover symptoms: nausea, headache, stress and anxiety which are unpleasant experiences of various physical, physiological and psychological effects. These aftereffects may last and vary from less than one hour up to several days, depending on numerous intrinsic and extrinsic factors. Globally, the alcohol-related aftereffects cause, e.g., impaired cognitive functions and performance of everyday tasks, e.g., such as driving (Gunn et al. 2018), as well as impaired work performance (presenteeism) and absence from work (absenteeism) (Thørrisen et al. 2019). Such impairments create considerable amount of economic loss worldwide (Wiese et al. 2004).

In January 2020, there were more than 700 publications in PubMed in which hangover have been studied in different contexts. Of these publications only 10 relevant studies have investigated various food supplements as possible cures for the aftereffects of alcohol drinking (Bang et al. 2015, Chauhan and Kulkarni 1991, George et al. 2019, Kim et al. 2017, Kim et al. 2018, Lee et al. 2013, Lee et al. 2014, Mammen et al. 2018, Takahashi 2010 et al, Wiese et al. 2004). In all of these 10 studies, plants, fruits and/or their extracts have been used, which complicates the evaluation of the real effective agents. Stress was not investigated in any of these studies. Anxiety was mentioned in four studies, two without response (Lee et al. 2014, Takahashi et al. 2010) and two with response (George et al. 2019, Mammen et al. 2018). Hangover, nausea and headache were the primary target for these 10 studies. Three of the studies displayed lower blood alcohol curves with the food supplements compared to placebo (Lee et al. 2013, Lee et al. 2014, Mammen et al. 2018). These effects may relate to slower gastric absorption. It is commonly known that virtually any sort of food may reduce the absorption of alcohol (Gentry 2000).

Alcohol hangover is a complex phenomenon affecting biochemical, inflammatory and neurochemical mechanisms (Palmer et al. 2019, 2020). Generally, alcohol is considered to be the agent for causing hangover and related symptoms. However, the truth is that many of these harmful effects are produced by alcohol and its metabolite, acetaldehyde (Eriksson 2001), which also is a common ingredient in alcohol beverages (Lachenmeier and Sohnius 2008). In addition, also cigarette smoking produces acetaldehyde (Seeman et al. 2002), which may explain why alcohol drinking and smoking are closely associated with each other. A study published 1967 showed that acetaldehyde in cigarette smoke binds to L-cysteine and forms a 2-methyl-L-thiazolidine-4-carboxylic acid (MTCA) derivative (Braven et al. 1967). The hypothesis was that the protective strength of amino thiols (especially L-cysteine) would be reduced, due to the acetaldehyde, which could increase the possibility of carcinogenesis. Later studies have shown that alcohol-related acetaldehyde is carcinogenic (Secretan et al. 2009) and in addition, also causes harmful symptoms during hangover (Eriksson 2001).

In our present study we investigated the effect of the semi essential amino acid, L-cysteine, on hangover symptoms: nausea, headache, stress and anxiety. L-cysteine, representing a single chemical molecule, is produced in the human body and is also an ingredient in the diet. This makes L-cysteine different to the other studied food supplements (plants, fruits and/or their extracts), in which there are a number of ingredients of unspecific nature. Our main hypothesis for the present study is that the oral administration of the L-cysteine leads to elimination of acetaldehyde in the body, which in turn prevents or alleviates alcohol aftereffects. To our knowledge, our study is the first to prove the prevention and the alleviation of alcohol-related hangover symptoms: nausea, headache, stress and anxiety with L-cysteine as a food supplement.

MATERIALS AND METHODS

The recruitment and selection of volunteers

Voluntary healthy participants were recruited through advertisements (media, internet, message boards from the University of Helsinki). To all participants, which were interested in the study, we sent the first questionnaires, which were completed and sent back to us by mail before the first session. Altogether, 27 participants were enrolled, 8 women and 19 men, aged between 21 - 50 (women) and 21 - 60 (men). All volunteers had to have some experience of hangover and/or headache. The exclusion criteria also included pregnancy (tested before sessions), menopause, alcoholism, especially low tolerance to alcohol (highlighted intoxication and/or hangover), more than 6 hours' fasting, BMI more than 30 kg/m² and less than 18 kg/m², alcohol related and non-related diseases and alcohol drinking related aggression. Also, all volunteers received a letter, in which the study was described, together with an informed consent document (to be signed before the first session).

A main problem was that only 8 women participated; one attending one session. Three women had zero response to hangover symptoms in all 3 sessions. We were left with 4 women; one unfortunately participating at different stages of the normal menstrual cycle. One of the remaining 3 women using long term steroids vomited and had nausea during all three sessions. These confounding factors prevent us from drawing reliable conclusions from the present women's data. Thus, the results of the present study are based solely on the men's data.

Study design

The study protocol has been approved by the research ethics committee of UEF (University of Eastern Finland) and by the coordinating ethics committee of HUH (Helsinki university hospital). The present hangover study was randomized, double-blind and placebo controlled. In all categories (hangover, nausea, headache, stress and anxiety) only the data of participants with a response were included. Two volunteers quit after the first session, one quit after 2 sessions, one could not finish the drinking in one session, one could not finish the drinking in 2 sessions and two volunteers did not respond in any of their 3 sessions. Twelve volunteers responded in all 3 sessions. Altogether, data was obtained from 43 participations in the 5 categories (hangover, nausea, headache, stress and anxiety). In addition, alcohol and acetaldehyde levels were also measured for two volunteers, who did not respond in any of the 5 categories, raising the number of participations to 49. All volunteers randomly swallowed during Friday evenings (August-September 2018) 6 placebo tablets, 3 placebos plus 3 L-cysteine tablets and/or 6 L-cysteine tablets (L-cysteine tablets containing 200 mg L-cysteine), every time about 15 min before 7, 8, 9, 10, 11 and 12 pm.

All tablets, excluding placebo tablets, contained also vitamin B1 (thiamine, 50 mg), B2 riboflavin, 1.4 mg), B3 (niacin, 16 mg), B6 (pyridoxine, 1.4 mg), B7 (biotin, 50 µg), B9 (folic acid, 200µg), B12 (cobalamin 2.5 µg) and vitamin C (ascorbic acid, 80 mg). This L-cysteine tablet is a fast releasing food supplement (Catapult Cat™ L-cysteine vitamin tablet), which cannot be distinguished with the placebo. The composition of all tablets also consists of filling agent (microcrystalline cellulose), anti-caking agent (magnesium salts of fatty acids and silicon dioxide), titanium dioxide, copper complexes of chlorophyll and chlorophyllin, glycerol, hydroxypropyl methylcellulose and flavor (menthol).

All sessions were done at a private hotel, during 6 subsequent Fridays, started at 6 pm with an informal event during which the informed consents were signed and second questionnaires completed. First saliva samples were taken at about 6:30 pm. Alcohol drinking started just before 7 pm and was consumed evenly in 3 doses in the period of 3 hours. The overall alcohol dose was 1.5 g/kg, which was served as 10%, v/v, mixed with lingonberry/blackcurrant juice (sucrose, 9 g/L, energy content 160 kJ). Second saliva samples were taken about half an hour after finishing drinking ca 10:30 pm, during which an intoxication questionnaire was completed. Sessions were closed around midnight after which participants went to sleep. No smoking nor eating were allowed during the evening and night. Next morning, around 8 – 9 am, the participants gathered in the same room as in the previous evening. After the third saliva sampling and completing the rest of the questionnaires the participants were free to go for breakfast and other activities.

With the first questionnaire for the present study, we could assess the inclusion and exclusion criteria for the participation, and also using the Alcohol Use Disorders Identification Test (AUDIT) (Saunders et al. 1993). The main morning questionnaires, with which the participants expressed their degree of hangover, nausea, headache and anxiety, were done with Likert Scales from 0 to 10. In addition, we also used a Likert Scale 1–4 in interviewing the degree of volunteers' stress.

Analytical procedures

Alcohol and acetaldehyde breath tests were performed before the session, at the end of sessions and in the morning using Proton Transfer Reaction Time-of-Flight Mass Spectrometry (PTR-TOF-MS) (PTR-TOF 1000, Ionicon Analytik, Innsbruck, Austria). Proton transfer reaction mass spectrometry is a very sensitive analytical methodology that can be used for online analysis of various volatile organic compounds in the gas phase (Blake et al. 2009). The sensitivity is high enough to enable the analysis of endogenous ethanol and acetaldehyde levels in human exhaled breath. A closely related technique, selected-ion flow-tube mass spectrometry (SIFT-MS), has previously been used to analyse acetaldehyde and ethanol from exhaled breath (Turner et al. 2006). The PTR-TOF-MS operating conditions were as follows: field density ratio (E/N) of 127 Td; drift tube pressure of 2.20 mbar, H₂O flow of 5.0 standard cubic centimetres per minute (sccm); ion source current of 3.0 mA; inlet flow of 50 sccm and drift tube voltage of 551 V. Drift tube and inlet temperatures were kept at 70 °C. Sampling frequency was 1 Hz, meaning one spectrum was recorded every second. Breath samples were obtained using Ionicon BET-med breath sampler. The breath sampler was connected to a device (Multiple Breath Gas Sampler, Loccioni) that measured the exhaled volume of air and CO₂ concentration. Breath samples were collected such that the volunteers exhaled a total of 1300 ml of air for each sample.

The acetaldehyde concentration was quantified at the protonated mass-to-charge ratio (m/z) signal of 46 that corresponds to the less abundant isotopologue of acetaldehyde. This was done to reduce the effect of signal saturation that takes place due to the high concentration of acetaldehyde after alcohol drinking. The signal at m/z 46 was multiplied by a factor of 47 to get the concentration of the most abundant acetaldehyde isotopologue. Regardless, the very high ethanol concentration in breath caused problems for the operation of the instrument and the acetaldehyde analysis. In practice, we observed that the mean acetaldehyde levels had significant differences between the six sessions. For this reason, the individual acetaldehyde measurements were normalized by multiplying with the following ratio: mean acetaldehyde concentration of all sessions / mean acetaldehyde concentration of session.

In addition, alcohol breath tests were also measured by the Dräger Alcometer, Alcotest 3000 (Drägerwerk AG & Co. KGaA, Lübeck, Germany). Breath alcohol and acetaldehyde concentrations

were converted by the blood/breath ratio (2300:1) for alcohol (Jones 2000) and (190:1) acetaldehyde (Jones 1995), respectively. Cortisol and testosterone were measured from saliva samples before alcohol drinking, after the drinking and next morning. Salivary cortisol and testosterone determinations were made by the enzyme-linked immunosorbent assay (ELISA) analysis (see Acknowledgements).

Statistical methods

The statistics were performed with the (IBM) SPSS software version 24. In the present study all statistical significances associated with L-cysteine were 1-tailed. This presumption is reasonable, since it is known that L-cysteine binds to acetaldehyde and consequently diminishes alcohol-related hangover symptoms: nausea, headache, stress and anxiety. For the other correlations without L-cysteine, the significances were 2-tailed. Spearman's rho coefficients are used for all correlation analyses (all Figures). Related-Samples Wilcoxon Sign Rank Test (for median differences), Related-Samples Friedman's Two-Way Analysis of Variance by Ranks and Paired Samples and T-Test were used. (See Figures). Standard error of the mean (SEM) was used in alcohol and acetaldehyde calculations.

RESULTS

Alcohol and acetaldehyde concentrations

In the present study the overall mean alcohol concentration after 3 hours' evening drinking was 28.35 ± 0.69 mM (1.31 ± 0.03 %) (N = 49) without any significant differences between the placebo and L-cysteine groups. Neither were there any separate significant differences between evening alcohol concentration and the 5 categories hangover symptoms: nausea, headache, stress and anxiety) in placebo and L-cysteine groups. Similarly, the overall average morning alcohol concentration was 3.45 ± 0.83 mM without any significant differences between the placebo and L-cysteine groups. Thus, in the present study, L-cysteine did not affect the evening absorption or the rate of alcohol elimination.

The evening mean acetaldehyde level was 6.31 ± 0.26 μ M (N = 49) without any significant differences between the placebo and L-cysteine groups. However, in the category nausea and stress, ingestion of L-cysteine (1200 mg) resulted in significant effects on evening acetaldehyde (Fig. 1). Morning acetaldehyde concentration was 4.47 ± 0.60 μ M without any significant differences between placebo and L-cysteine.




Fig. 1. The effect of L-cysteine on evening acetaldehyde in the categories of morning nausea (A) and stress (B). Individual data of 9 men attending all 3 sessions are shown (A) and 6 men attending all 3 sessions plus 3 single cases (B). Friedman related-samples analysis of variance by ranks shows differences between placebo (red dots) and L-cysteine (1200 mg, blue dots) with the significance of $P = 0.048$ (A) and $P = 0.051$ (B).

Correlational analyses

Our correlational study is designed so that all the data points in the figures represent original data. Although the number of volunteers was rather low, several significant results were obtained by L-cysteine when hangover symptoms: nausea, headache, stress and anxiety were examined the next morning, ca. 10 hours after evening drinking was finished. One main hangover symptom is nausea, which was prevented or alleviated by L-cysteine (Fig. 2A). Nausea was also significantly positively correlated with hangover (34 participations, with 9 subjects attending all 3 sessions, 0.437 , $P = 0.010$). However, headache was more strongly correlated with hangover than with nausea (participations were 43 with 12 subjects attending all 3 sessions, 0.690 , $P = 0.000$). Nausea and headache symptoms may thus share some affecting parameters, but headache also seems to have an additional parameter in common with hangover (participations were 34 with 9 subjects attending all 3 sessions, 0.370 , $P = 0.031$).




Fig. 2. The effect of evening L-cysteine on alcohol-related morning nausea (A), headache (B) and hangover (C). Individual data of men are shown, all of which displayed at least a hint of nausea (A), headache (B) and/or hangover (C) at some of the sessions. Altogether the participations were 34 with 9 subjects attending all 3 sessions (A), 43 with 12 subjects attending all 3 sessions (B, C). Correlations analysis displayed significance with -0.287 , $P = 0.050$ (A), negligible with -0.165 , $P = 0.145$ (B) and significance with -0.258 , $P = 0.047$ (C). Moreover, related-samples Wilcoxon sign rank test tests (median differences) displayed significant differences between placebo (red dots) and L-cysteine (1200 mg, blue dots), $P = 0.013$ (A), $P = 0.010$ (B) and $P = 0.043$.

From the correlations between L-cysteine and hangover symptoms: nausea and headache, it could be seen that the dose of L-cysteine, 600 mg, had little effect on these symptoms, as compared to the dose of 1200 mg, (Fig. 2A, B and C). However, for stress and anxiety, a correlation with L-cysteine is present already at a dose level of 600 mg (Fig. 3). Moreover, both stress (headache correlation

0.574, $P = 0.005$; hangover correlation 0.503, $P = 0.017$) and anxiety (headache correlation 0.594, $P = 0.001$; hangover correlation 0.586, $P = 0.001$) were significantly associated with headache and hangover, as also with each other (stress vs anxiety correlation 0.806, $P = 0.000$).

We also studied the possible inter-relations between the hangover and stress and anxiety symptoms, and the steroids cortisol and testosterone. The only trend effect was a positive correlation (0.304, $P = 0.086$) between cortisol and nausea. In fact, the positive cortisol/nausea correlation has been recognized in an earlier Finnish study by Ylikahri et al. (1974).




Fig. 3. The effect of L-cysteine on alcohol-related morning stress (A) and anxiety (B). Individual data of men are shown, all of which displayed at least a hint of stress (A) and anxiety (B) at some of the sessions. Altogether the participations were 22 with 6 subjects (A) and 27 with 7 subjects, attending all 3 sessions. Correlations analysis displayed significance with -0.382 , $P = 0.039$ (A) and negligible with -0.250 , $P = 0.104$. (B). In addition, paired samples t-test (with 7 subjects) displayed a significance of $P = 0.039$ between placebo (red dots) and L-cysteine (600 mg, purple dots) (A). Moreover, related-samples Wilcoxon sign rank test (median differences) displayed a significance of $P = 0.052$ (B).

Discussion

Alcohol and acetaldehyde

No differences in overall evening and morning alcohol concentrations were observed between placebo and L-cysteine groups. This is plausibly due to the strict control imposed on the volunteers' evening drinking which led to a very low variation in the blood alcohol levels. The lack of differences in morning alcohol levels showed that the L-cysteine did not change the rate of alcohol elimination. Altogether, it was impossible to detect any significant alcohol correlations within the different categories (hangover symptoms: nausea, headache, stress and anxiety) between the placebo and L-cysteine groups. Regarding the pulmonary acetaldehyde, the difficulty is that the exhaled breath sample may be contaminated by acetaldehyde derived by the oral microflora. Most

likely, the real pulmonary acetaldehyde concentration is a bit lower than our measured concentrations. In addition, technical problems caused by the high ethanol content of exhaled breath during the sessions, most likely introduced additional noise into the acetaldehyde data. However, in spite of these problems, with acetaldehyde and L-cysteine, two significant differences emerged between placebo and L-cysteine (see result section, Fig. 1A and B).

Hangover, nausea, headache, stress and anxiety

In our present study, hangover represents disagreeable aftereffects of drunkenness involving physical, physiological and psychological experience. Instead of using a hangover scale that includes a number of hangover symptoms such as HSS Hangover Symptoms Scale (Slutske et al. 2003), AHS Acute Hangover Scale (Rohsenow et al. 2007) or AHSS Alcohol Hangover Severity Scale (Penning et al. 2013), we just used a single Likert scales for hangover, stress and anxiety symptoms. Earlier 6 food supplement studies demonstrate some success treating nausea (Bang et al. 2015, George et al. 2019, Kim et al. 2017, Mammen et al. 2018, Takahashi et al. 2010, Wiese et al. 2004), headache (Bang et al. 2015, George et al. 2019, Kim et al. 2017, Kim et al. 2018, Mammen et al. 2018) and hangover (Bang et al. 2015, Chauhan and Kulkarni 1991, Kim et al. 2017, Mammen et al. 2018). Anxiety has been treated only in two studies (George et al. 2019, Mammen et al. 2018). Interestingly, stress has not been treated in any of the earlier studies. In our present study we observed statistically significant effects in all of the five categories.

Vitamins

The vitamins included in the L-cysteine tablets may or may not had an effect on the hangover symptoms. The neutralizing effect of L-cysteine on acetaldehyde (Fig. 1), which is a major factor promoting hangover symptoms, indicates that L-cysteine has a main impact in diminishing the alcohol-related hangover symptoms. Altogether, the present study was focused only on the effect of L-cysteine on hangover and its symptoms. A separate study with alcohol-related hangover symptoms with all different vitamins would certainly be an interesting topic for further research.

Ethics and L-cysteine

In many parts of the world where alcohol drinking is legal, a widely held belief still exists, that hangover should not be cured because that would increase the risk of more alcohol consumption. However, this belief may be misinformed and not entirely correct. The fact of the matter is that higher degree of alcohol-related hangover and stress symptoms lead to more effort “curing” the aftereffects by drinking alcohol. This is especially the case with individuals with predisposition for developing alcohol addiction (Earleywine 1999, Newling and Pretorius 1990). Moreover, it seems that there is no evidence for why preventing or alleviating hangover would lead to increased alcohol consumption (Wiese et al. 2000). Finally, it has to be recognized that not all people who experience hangovers continue to drink.

Some of the previously studied food supplements, plants, fruits and/or their extracts may have some similar effects as L-cysteine, but possibly also various other unwanted effects as well. The advantage with L-cysteine is that this semi-essential amino acid already is present in the human body. This means that the addition of L-cysteine can directly and dose dependently enter into the normal metabolism and help to neutralize the detrimental acetaldehyde, as well as prevent or alleviate the alcohol aftereffects.

Moreover, since acetaldehyde is carcinogenic, binding it by L-cysteine may also reduce the probability for the development of cancer (Hellström et al. 2017, Juliano et al. 2011). Thus, all these actions of L-cysteine are ethically sound.

Conclusion

The final conclusion is that L-cysteine reduces alcohol-related hangover symptoms: nausea, headache, stress, and anxiety. Altogether, the effects of L-cysteine on hangover symptoms: nausea, headache, stress and anxiety are unique and seems to have a future in preventing or alleviating these harmful symptoms as well as reducing the risk of alcohol addiction.

REFERENCES

- Bang JS, Chung YH, Chung SJ, *et al.* (2015) Clinical effect of a polysaccharide-rich extract of *Acanthopanax senticosus* on alcohol hangover. *Pharmazie* **70**:269–73.
- Blake RS, Monks PS, Ellis AM (2009) Proton-transfer reaction mass spectrometry. *Chemical Reviews* **109**:861–96.
- Braven J, Bonker GJ, Fenner ML, *et al.* (1967) The mechanism of carcinogenesis by tobacco smoke. Some experimental observations and a hypothesis. *Br J Cancer* **21**:623–33.
- Chauhan BL, Kulkarni RD (1991) Alcohol hangover and Liv.52. *Eur J Clin Pharmacol* **40**:187–8.
- Earleywine M (1999) Personality risk for alcoholism covaries with hangover symptoms. *Addict Behav* **24**:121–5.
- Eriksson CJP (2001) The role of acetaldehyde in the actions of alcohol (update 2000). *Alcohol Clin Exp Res* (**5 Suppl ISBRA**):15S–32S.
- Gentry RT (2000) Effect of food on the pharmacokinetics of alcohol absorption. *Alcohol Clin Exp Res* **24**:403–4.
- George A, Udani JK, Yusof A (2019) Effects of *Phyllanthus amarus* PHYLLPRO™ leaves on hangover symptoms: a randomized, double-blind, placebo-controlled crossover study. *Psychol Addict Behav*. **33**:685–96.
- Gunn C, Mackus M, Griffin C, *et al.* (2018) A systematic review of the next-day effects of heavy alcohol consumption on cognitive performance. *Addiction* **113**:2182–93.
- Hellström PM, Hendolin PM, Kaihovaara P, *et al.* (2017) Slow-release L-cysteine capsule prevents gastric mucosa exposure to carcinogenic acetaldehyde: results of a randomised single-blinded, cross-over study of *Helicobacter*-associated atrophic gastritis. *Scand J Gastroenterol* **52**:230–7.
- Jones AW (1995) Measuring and reporting the concentration of acetaldehyde in human breath. *Alcohol Alcohol* **30**:271-85.
- Jones AW (2000) Medicolegal alcohol determination - blood- or breath-alcohol concentration? *Forensic Sci Rev* **12**:23-47.
- Juliano C, Cossu M, Rota MT, *et al.* (2011) Buccal tablets containing cysteine and chlorhexidine for the reduction of acetaldehyde levels in the oral cavity. *Drug Dev Ind Pharm* **37**:1192–9.

- Kim H, Kim YJ, Jeong HY, *et al.* (2017) A standardized extract of the fruit of *Hovenia dulcis* alleviated alcohol-induced hangover in healthy subjects with heterozygous ALDH2: A randomized, controlled, crossover trial. *J Ethnopharmacol* **209**:167–74.
- Kim MJ, Lim SW, Kim JH, *et al.* (2018) Effect of mixed fruit and vegetable juice on alcohol hangovers in healthy adults. *Prev Nutr Food Sci* **23**:1–7.
- Lachenmeier DW, Sohnius EM (2008) The role of acetaldehyde outside ethanol metabolism in the carcinogenicity of alcoholic beverages: evidence from a large chemical survey. *Food Chem Toxicol* **46**:2903–11.
- Lee H, Isse T, Kawamoto T, *et al.* (2013) Effect of Korean pear (*Pyrus pyrifolia* cv. Shingo) juice on hangover severity following alcohol consumption. *Food Chem Toxicol* **58**:101–6.
- Lee MH, Kwak JH, Jeon G, *et al.* (2014) Red ginseng relieves the effects of alcohol consumption and hangover symptoms in healthy men: a randomized crossover study. *Food Funct* **5**:528–34.
- Mammen RR, Natinga MJ, Mohanan R, *et al.* (2018). Clove bud polyphenols alleviate alterations in inflammation and oxidative stress markers associated with binge drinking: A randomized double-blinded placebo-controlled crossover study. *J. Med Food* **21**:1188–96.
- Newlin DB, Pretorius MB (1990) Sons of alcoholics report greater hangover symptoms than sons of nonalcoholics: a pilot study. *Alcohol Clin Ex. Res* **14**:713–6.
- Palmer E, Tyacke R, Sastre M, *et al.* (2019) Alcohol hangover: Underlying biochemical, inflammatory and neurochemical mechanisms. *Alcohol Alcohol* **54**:196–203.
- Palmer EOC, Arnoldy L, Ayre E, *et al.* (2020) Proceedings of the 11th Alcohol Hangover Research Group Meeting, in Nadi, Fiji. *Proceedings* **43**, 1; doi:10.3390/proceedings2020043001.
- Penning R, McKinney A, Bus LD, *et al.* (2013) Measurement of alcohol hangover severity: development of the Alcohol Hangover Severity Scale (AHSS). *Psychopharmacology (Berl)*. **225**:803–10.
- Rohsenow DJ, Howland J, Minsky SJ, *et al.* (2007) The Acute Hangover Scale: A new measure of immediate hangover symptoms. *Addict Behav* **32**:1314–20.
- Saunders JB, Aasland OG, Babor TF, *et al.* (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption--II. *Addiction* **88**:791–804.
- Secretan B, Straif K, Baan R, *et al.* (2009) A review of human carcinogens--Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol* **10**:1033–4. Review.
- Seeman JI, Dixon M, Haussmann HJ (2002) Acetaldehyde in mainstream tobacco smoke: formation and occurrence in smoke and bioavailability in the smoker. *Chem Res Toxicol* **15**:1331–50. Review.
- Slutske WS, Piasecki TM, Hunt-Carter EE (2003) Development and initial validation of the Hangover Symptoms Scale: prevalence and correlates of hangover symptoms in college students. *Alcohol Clin Exp Res* **27**:1442–50.
- Takahashi M, Li W, Koike K, *et al.* (2010) Clinical effectiveness of KSS formula, a traditional folk remedy for alcohol hangover symptoms. *J Nat Med* **64**:487–41.
- Thørrisen MM, Bonsaksen T, Hashemi N, *et al.* (2019) Association between alcohol consumption and impaired work performance (presenteeism): a systematic review. *BMJ Open* **9**:e029184.
- Turner C, Španěl P, Smith S (2006) A longitudinal study of ethanol and acetaldehyde in the exhaled breath of healthy volunteers using selected -ion flow-tube mass spectrometry. *Rapid Commun Mass Spectrom* **20**: 61–8 .
- Wiese JG, Shlipak MG, Browner WS (2000) The alcohol hangover. *Ann Intern Med* **132**:897–902. Review.
- Wiese J, McPherson S, Odden MC, *et al.* (2004) Effect of *Opuntia ficus indica* on symptoms of the alcohol hangover. *Arch Intern Med* **164**:1334–40.

Ylikahri RH, Huttunen MO, Eriksson CJP, *et al.* (1974) Metabolic studies on the pathogenesis of hangover. *Eur J Clin Invest* **4**:93–100.

ACKNOWLEDGEMENTS

This work was supported by Catapult Cat Pty Ltd, Espoo, Finland; and also supported by the University of Eastern Finland. We thank our volunteers that participated in the sessions. We also thank Karin von Wachenfeldt and Peter Blom for steroid analyses (Truly Labs AB, Medicin Village, Lund, Sweden).

CONFLICTS OF INTEREST STATEMENT

Authors have no conflict of interest. The authors were fully responsible for the study design and the funding agency had no role in the collection, analysis, interpretation of data and in writing of the manuscript.