

<https://helda.helsinki.fi>

EAACI Biologicals Guidelines-Recommendations for severe asthma

Agache, Ioana

2021-01

Agache , I , Akdis , C , Akdis , M , Canonica , G W , Casale , T , Chivato , T , Corren , J , Chu , D K , Del Giacco , S , Eiwegger , T , Flood , B , Firinu , D , Gern , J E , Hamelmann , E , Hanania , N , Hernandez-Martin , I , Knibb , R , Mäkelä , M , Nair , P , O'Mahony , L , Papadopoulos , N G , Papi , A , Park , H-S , Perez de Llano , L , Pfaar , O , Quirce , S , Sastre , J , Shamji , M , Schwarze , J , Palomares , O & Jutel , M 2021 , ' EAACI Biologicals Guidelines-Recommendations for severe asthma ' , Allergy : European journal of allergy and clinical immunology , vol. 76 , no. 1 , pp. 14-44 . <https://doi.org/10.1111/all.14425>

<http://hdl.handle.net/10138/339440>

<https://doi.org/10.1111/all.14425>

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

GUIDELINES



WILEY

EAACI Biologicals Guidelines—Recommendations for severe asthma

Ioana Agache¹ | Cezmi A. Akdis^{2,3} | Mubeccel Akdis² | Giorgio Walter Canonica⁴ | Thomas Casale⁵ | Tomas Chivato⁶ | Jonathan Corren⁷ | Derek K. Chu⁸ | Stefano Del Giacco⁹ | Thomas Eiwegger^{10,11,12} | Breda Flood¹³ | Davide Firinu⁹ | James E. Gern¹⁴ | Eckard Hamelmann¹⁵ | Nicola Hanania¹⁶ | Irene Hernández-Martín¹⁷ | Rebeca Knibb¹⁸ | Mika Mäkelä¹⁹ | Parameswaran Nair^{20,21} | Liam O'Mahony²² | Nikolaos G. Papadopoulos^{23,24} | Alberto Papi²⁵ | Hae-Sim Park²⁶ | Luis Pérez de Llano²⁷ | Oliver Pfaar²⁸ | Santiago Quirce²⁹ | Joaquin Sastre³⁰ | Mohamed Shamji^{31,32} | Jurgen Schwarze³³ | Oscar Palomares³⁴ | Marek Jutel^{35,36}

¹Faculty of Medicine, Transylvania University, Brasov, Romania

²Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland

³Christine-Kühne-Center for Allergy Research and Education (CK-CARE), Davos, Switzerland

⁴Personalized Medicine, Asthma and Allergy, Humanitas Clinical and Research Center, IRCCS, Rozzano, Italy

⁵Division of Allergy and Immunology, University of South Florida Morsani College of Medicine, Tampa, FL, USA

⁶School of Medicine, University CEU San Pablo, Madrid, Spain

⁷David Geffen School of Medicine at UCLA, Los Angeles, CA, USA

⁸Department of Health Research Methods, Evidence and Impact, Division of Immunology and Allergy, and Department of Medicine, McMaster University, Hamilton, ON, Canada

⁹Department of Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy

¹⁰Translational Medicine Program, Research Institute, Hospital for Sick Children, Toronto, ON, Canada

EXTERNAL PEER REVIEW PANEL: Leonard B. Bacharier¹, Eugene R. Bleeker², Christopher Brightling³, William W. Busse⁴, Kian Fan Chung⁵, Borja G. Cosío⁶, Alvaro Cruz⁷, Christian Domingo Ribas⁸, Christer Jansen⁹, Stephen Holgate¹⁰, Sejal Saglani¹¹, Stanley J. Szefler¹².

¹Division of Allergy, Immunology and Pulmonary Medicine, Department of Pediatrics, Washington University, St Louis, Mo; ²Department of Medicine, Division of Genetics, Genomics and Precision Medicine, University of Arizona Health Sciences, Tucson, Arizona, USA; ³Institute for Lung Health, Department of Respiratory Sciences, University of Leicester, Leicester, UK; ⁴Division of Allergy, Pulmonary and Critical Care Medicine, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; ⁵Experimental Studies Medicine at National Heart & Lung Institute, Imperial College London & Royal Brompton & Harefield NHS Trust, London, UK; ⁶Institut d'Investigació Sanitària Illes Balears (IdISBa), Palma de Mallorca, Spain; CIBER de Enfermedades Respiratorias, CIBERES, Spain; Department of Respiratory Medicine, Hospital Universitario Son Espases, Palma de Mallorca, Spain; Clínica Rotger Quiron, Palma de Mallorca, Spain; ⁷Fundação ProAR, Federal University of Bahia and GARD/WHO Planning Group, Salvador, Brazil; ⁸Departament de Medicina, Universitat Autònoma de Barcelona (UAB), Barcelona, Spain; Servei de Pneumologia, Corporació Sanitària Parc Taulí, Parc Taulí s/n, 08208, Sabadell (Barcelona), Spain; ⁹Department of Medical Sciences: Respiratory, Allergy and Sleep Research, Uppsala University, Uppsala, Sweden; ¹⁰Faculty of Medicine, University of Southampton, Southampton, UK; ¹¹Department of National Heart and Lung Institute, Imperial College London, UK; Department of Respiratory Paediatrics, Royal Brompton Hospital, London, UK; ¹²Section of Pulmonary and Sleep Medicine, Department of Pediatrics, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, Colorado.

Abbreviations: ACQ, asthma control questionnaire; ACT, asthma control test; AD, atopic dermatitis; ADA, anti-drug antibodies; AQLQ, asthma quality of life questionnaire; AR, allergic rhinitis; ATS, American Thoracic Society; COI, conflict of interest; CRSwNP, chronic rhinosinusitis with nasal polyps; DCs, dendritic cells; EAACI, European Academy of Allergy and Clinical Immunology; ED, emergency department; EMA, European Medicines Agency; ERS, European Respiratory Society; EtD, evidence to decision; Fab, antigen-binding fragment; Fc, fragment crystallizable; FcεRI, IgE high-affinity receptor; FDA, Food and Drug Administration; FeNO, fractional exhaled nitric oxide; FEV1, forced expiratory volume at the end of the first second of forced expiration; GDG, guidelines development group; GETE, global evaluation of treatment effectiveness; GINA, Global Initiative for Asthma; GRADE, Grading of Recommendations Assessment, Development, and Evaluation; HCP, healthcare professional; ICERs, incremental cost-effectiveness ratios; ICS, inhaled corticosteroids; Ig, immunoglobulin; IL, interleukin; IL-4Rα, the α subunit of the IL-4 receptor; IL-5Rα, the α subunit of the IL-5 receptor; ISAR, International Severe Asthma Registry; IV, intravenous; JAK, Janus kinase; Mab, monoclonal antibody; NICE, National Institute for Health and Care Excellence; NK, natural killer; OCS, oral corticosteroids; PEF, peak expiratory flow; PI3K, phosphoinositide 3 kinase; PICO, population, intervention, comparator and outcomes; PROs, patient-reported outcomes; QALY, quality-adjusted life-years; QoL, quality of life; ROB, risk of bias; SANI, Severe Asthma Network in Italy; SMART, Standardized Measure to Assess Response to Therapy; SMD, small-molecule drug; SOF, summary of findings; SPACE, Severe Paediatric Asthma Collaborative in Europe; SRs, systematic literature reviews; STAT, signal transducer and activator of transcription; T2, type 2; TH, T helper; TNF, tumour necrosis factor; TSLP, thymic stromal lymphopoietin.

Oscar Palomares and Marek Jutel are joint last authorship.

¹¹Department of Immunology, University of Toronto, Toronto, ON, Canada

¹²Division of Immunology and Allergy, Food Allergy and Anaphylaxis Program, The Hospital for Sick Children, Departments of Paediatrics and Immunology, University of Toronto, Toronto, ON, Canada

¹³European Federation of Allergy and Airway Diseases, Brussels, Belgium

¹⁴Department of Pediatrics, School of Medicine and Public Health, University of Wisconsin, Madison, WI, USA

¹⁵Children's Center Bethel, Evangelical Hospital Bethel, University of Bielefeld, Bielefeld, Germany

¹⁶Section of Pulmonary, Critical Care and Sleep Medicine, Baylor College of Medicine, Houston, TX, USA

¹⁷Department of Allergy, Hospital Universitario La Paz, Madrid, Spain

¹⁸Department of Psychology, School of Life and Health Sciences, Aston University, Birmingham, UK

¹⁹Skin and Allergy Hospital, Helsinki University Hospital and University of Helsinki, Helsinki, Finland

²⁰Division of Respiratory, Department of Medicine, McMaster University, Hamilton, ON, Canada

²¹Firestone Institute for Respiratory Health, St Joseph's Healthcare, Hamilton, ON, Canada

²²Departments of Medicine and Microbiology, APC Microbiome Ireland, University College Cork, Cork, Ireland

²³Division of Infection, Immunity and Respiratory Medicine, University of Manchester, Manchester, UK

²⁴Allergy Department, 2nd Pediatric Clinic, National Kapodistrian University of Athens, Athens, Greece

²⁵Research Center on Asthma and COPD, Department of Medical Sciences, University of Ferrara, Ferrara, Italy

²⁶Department of Allergy and Clinical Immunology, Ajou University, Ajou, Korea

²⁷Department of Respiratory Medicine, Hospital Lucus Augusti, Lugo, Spain

²⁸Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Marburg, Germany

²⁹Department of Allergy, La Paz University Hospital, IdiPAZ, CIBER of Respiratory Diseases (CIBERES), Universidad Autónoma de Madrid, Madrid, Spain

³⁰Facultad de Medicina, Universidad Autónoma de Madrid, Madrid, Spain

³¹Immunomodulation and Tolerance Group, Allergy and Clinical Immunology, Inflammation, Repair, Development, National Heart and Lung Institute, London, UK

³²Imperial College NIHR Biomedical Research Centre, Asthma UK Centre in Allergic Mechanisms of Asthma, London, UK

³³Centre for Inflammation Research, Child Life and Health, The University of Edinburgh, Edinburgh, UK

³⁴Department of Biochemistry and Molecular Biology, Chemistry School, Complutense University of Madrid, Madrid, Spain

³⁵Department of Clinical Immunology, University of Wrocław, Wrocław, Poland

³⁶ALL-MED[™] Medical Research Institute, Wrocław, Poland

Correspondence

Ioana Agache, Faculty of Medicine,
Transylvania University, 2A, Pictor Ion
Andreescu, Brasov 500051, Romania.
Email: ibrumaru@unitbv.ro

Abstract

Severe asthma imposes a significant burden on patients, families and healthcare systems. Management is difficult, due to disease heterogeneity, co-morbidities, complexity in care pathways and differences between national or regional healthcare systems. Better understanding of the mechanisms has enabled a stratified approach to the management of severe asthma, supporting the use of targeted treatments with biologicals. However, there are still many issues that require further clarification. These include selection of a certain biological (as they all target overlapping disease phenotypes), the definition of response, strategies to enhance the responder rate, the duration of treatment and its regimen (in the clinic or home-based) and its cost-effectiveness. The EAACI Guidelines on the use of biologicals in severe asthma follow the GRADE approach in formulating recommendations for each biological and each outcome. In addition, a management algorithm for the use of biologicals in the clinic is proposed, together with future approaches and research priorities.

KEYWORDS

biologicals, cost-effectiveness, GRADE, guidelines, severe asthma

1 | INTRODUCTION

1.1 | The current landscape of severe asthma

1.1.1 | Definitions and burden

Asthma is a heterogeneous chronic inflammatory disease of the lower airways that affects nearly 400 million people worldwide.¹ It is the most frequent chronic disease in children. The current estimate is that 3 to 10% of adult asthmatics suffer from severe asthma.² The real prevalence might be lower, as these figures include cases with poor adherence, untreated co-morbidities or misdiagnosis.³⁻⁷ Although severe asthma comprises a small proportion of asthma cases, it is associated with increased mortality and hospitalization, reduced quality of life (QoL) and increased healthcare costs.⁸⁻¹³ Extrapolating adult severity classifications to children is difficult for a number of reasons ranging from the clinical phenotypes to lung function parameters and response to treatment. Therefore, the incidence of paediatric severe asthma affecting up to 2.5% of all children with asthma¹⁵ is not directly comparable with the incidence of severe asthma in adults. In children, severe asthma accounts approximately for half of all healthcare resources for paediatric asthma and it is associated with increased risk of mortality, and of development of chronic obstructive pulmonary disease in adulthood.¹⁵⁻¹⁹

The European Respiratory Society (ERS)/American Thoracic Society (ATS) guidelines set out a framework for the diagnosis and management of severe asthma in adults and children.^{20,21} The Global Initiative for Asthma (GINA) has produced a useful guide for the diagnosis and management of “Difficult-to-treat” and Severe Asthma in adolescents and adults.²² Patients with a confirmed diagnosis of asthma who have had modifiable factors addressed, poorly controlled while receiving high-dose inhaled corticosteroid (ICS) treatment, or with requirement of a high level of treatment to maintain control are classified as severe asthma.²⁰

1.1.2 | Severe asthma phenotypes and endotypes—practical implications for management

Following recent advances in our understanding of asthma mechanisms and therapeutic responsiveness, the concept of asthma as a single entity has been replaced with a model of a complex biological network with distinct, but interrelating immune–inflammatory pathways continuously modified by multiple external and internal factors.²³⁻²⁸ Several visible properties underpin the definition of severe adult asthma phenotypes: clinical, functional, morphological, inflammatory, molecular and microbiome-related.^{25,29-31} However, severe asthma phenotypes do not necessarily relate to or give insights into the underlying pathogenetic mechanisms; these are best described by disease endotypes. The major immune-inflammatory pathways for severe asthma include type-2 (T2) high, T2 low and mixed endotypes which may share certain genetic and epigenetic, metabolic, neurogenic and remodelling characteristics.^{24,31,32}

Severe paediatric asthma has been associated in most cases with severe and multiple aeroallergen sensitization, allergic rhinitis, food

allergy, eosinophilic airway inflammation, exposure to environmental tobacco smoke and airway remodelling.³³⁻³⁵ Similar to adults, it is quite heterogeneous.³⁶⁻³⁸ Unfortunately, less is known about its endotypes and related biomarkers. Mechanisms of adult severe asthma cannot be extrapolated to children. Children with eosinophilic asthma may not have increases in other T2 markers.³⁵ Unlike adults, intraepithelial neutrophils were associated with better lung function.³⁹ The potential roles of the innate epithelial cytokine interleukin (IL)-33 and of lineage negative innate lymphoid cells (ILCs) are also described.^{40,41}

The advent of biologicals for asthma represents a giant leap forward for severe asthma management. Considering the availability of several specific targeted therapies for T2 asthma, the management approach to severe asthma currently includes a phenotyping step for the identification of allergic and eosinophilic and non-T2 phenotypes.^{2,20,21,42} This stratified approach is based on measurement of biomarkers such as eosinophils in blood or sputum, fractional exhaled nitric oxide (FeNO) and specific immunoglobulin (Ig) E of clinical relevance.⁴²⁻⁴⁶ Whether this stratified approach will improve the burden of severe asthma remains to be proven by real-life studies and registry data.^{43,44,47} While targeted treatment strategies and mathematical models such as multidimensional endotyping are currently being tested, they will ultimately require validated guidelines for optimal implementation.^{30,43,48-50} In addition, no specific investigation into tailored treatment for young children with severe asthma exists and the efficacy of biologicals is less frequently tested in children. The same holds true for treatment of T2-low severe asthma. Given the relatively high costs of biological therapies, cost-effectiveness analyses are a prerequisite for coverage and reimbursement. Carefully targeting biological therapy to specific populations, such as high-responders, or discounting the acquisition price in order to further improve value are currently advocated.⁵¹

1.1.3 | Biologicals

Biologic products (biologicals) include a wide range of products such as vaccines, blood and blood components, allergenics, somatic cells, gene therapy, tissues and recombinant therapeutic proteins. They are isolated from a variety of natural sources—human, animal or microorganism—and may be produced by biotechnology methods and other cutting-edge technologies. For the purpose of this guideline, we refer to monoclonal antibodies (mAb) as biologicals. In contrast to chemical compounds and small-molecule agonists or antagonists, biologicals bind a specific determinant, for example a cytokine or receptor. Owing to this selectivity, biologicals are ideal for “personalized” or “precision” medicine.⁵²

1.1.4 | Current management of severe asthma

The GINA 2020 asthma guideline² recommends a decision tree for the diagnosis and management of severe asthma with distinct tasks for primary and specialized care while reinforcing the value of the patient perspective and collaborative care by primary care physicians, specialists

and other healthcare professionals (HCP). At the specialist level, assessment of the severe asthma phenotype during high-dose ICS treatment or lowest possible dose of oral corticosteroids (OCS) and of the factors contributing to symptoms, QoL and exacerbations are advocated. T2 inflammation is defined by the presence of blood eosinophils $\geq 150/\mu\text{L}$ and/or FeNO ≥ 20 ppb and/or sputum eosinophils $\geq 2\%$, and/or asthma that is clinically allergen-driven and/or that require maintenance OCS. A tentative approach for first-line biological targeting the T2 pathway is based on biomarkers and predictors of response, while taking into consideration local payer eligibility criteria, cost, dosing frequency, route (subcutaneous or intravenous) and patient preference. Response to the biological is evaluated after 4 months and if favourable the biological is continued with re-evaluation every 3–6 months. If the patient fails to respond to the initial agent but is still eligible for T2 targeted treatment, GINA 2020 recommends switching to another biological. GINA 2020 acknowledges that at present, there are no well-defined criteria for a good response, but recommends considering exacerbation frequency, symptom control, lung function, medication side-effects, treatment intensity (including OCS dose) and patient satisfaction as important factors.

The 2014 ERS/ ATS guidelines on severe asthma in adults and school age children introduced the definition of severe asthma as it is

used today, and highlighted the necessity of always confirming the diagnosis of asthma and excluding other conditions that may mimic asthma. This guideline also provided specific recommendations for the use of sputum eosinophil count and FeNO to guide the phenotypic-driven approach.²⁰ The 2019 revision of the ERS/ATS guidelines provided some specific recommendations for the use of biologicals targeting the IL-5, IgE and IL-4/IL-13 pathways based on the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) methodology.²¹ Recommendations are formulated for all anti-IL-5 interventions together, receiving a conditional recommendation for use as add-on treatment in severe eosinophilic asthma. Dupilumab received a conditional recommendation for adult patients with severe eosinophilic asthma and for those with severe corticosteroid-dependent asthma regardless of blood eosinophil levels. A blood eosinophil cut-point of $\geq 150/\mu\text{L}$ received a conditional recommendation to guide anti-IL-5 initiation in adult patients with severe asthma and specific eosinophil ($\geq 260/\mu\text{L}$) and FeNO (≥ 19.5 ppb) cut-offs are suggested to identify adolescents or adults with the greatest likelihood or response to anti-IgE therapy.

Currently, five biologicals are approved for severe asthma (in alphabetical order): benralizumab^{53,54}; dupilumab^{55,56}; mepolizumab^{57,58}; omalizumab^{59,60}; and reslizumab^{61,62} (Tables 1–3).

TABLE 1 Monoclonal antibodies approved for severe asthma—targets and mechanism of action

Biological	Target and mechanism of action
Benralizumab	IgG1 kappa humanised monoclonal antibody. Binds to the α subunit of the IL-5 receptor (IL-5R α) with inhibition of the hetero-oligomerization of α and β subunits and thus no signal transduction occurs. The afucosylated site of benralizumab enhances its binding to Fc γ RIIIa leading to antibody-dependent cell-mediated cytotoxicity by NK cells and macrophages. Depletes the eosinophils and reduces basophil levels.
Dupilumab	IgG4 human monoclonal antibody. Binds to the α subunit of the IL-4 receptor (IL-4R α) shared by IL-4 and IL-13 receptor complexes, thus simultaneously inhibiting both IL-4- and IL-13-mediated signalling pathways. IL-4 and IL-13 are key cytokines for orchestration of type 2 immune responses. Simultaneous blocking of Type I receptor (IL-4R α/γ c) and Type 2 receptor (IL-4R α /IL-13R α) inhibit at the same time type 2 responses depending on IL-4 and IL-4/IL-13, respectively, in hematopoietic and non-hematopoietic cells.
Mepolizumab	IgG1 kappa humanised monoclonal antibody specific for IL-5. Binds to a specific epitope of IL-5 and prevents it from binding to IL-5R α . Inhibits the maturation, activation, proliferation and recruitment of eosinophils.
Omalizumab	IgG1 kappa humanised monoclonal antibody. Binds to free IgE and inhibits the binding of IgE to both the high- and low-affinity IgE receptors (Fc ϵ RI and CD23, respectively). The reduction in surface bound IgE on Fc ϵ RI-bearing cells reduces the expression of this receptor in mast cells, basophils and dendritic cells, thus blocking the degree of release of cytokines and mediators of the allergic response and IgE-mediated presentation of Th2 cells. It also enhance the production of IFN- α by pDCs, thus reducing viral-induced exacerbations.
Reslizumab	IgG4 kappa humanised monoclonal antibody. Binds to a specific epitope IL-5 and prevents it from binding to IL-5R α . The <i>in vitro</i> affinity of reslizumab for IL-5 is higher than mepolizumab and the capacity to suppress IL-5-dependent proliferation <i>in vitro</i> is also superior. Inhibits the maturation, activation, proliferation and recruitment of eosinophils.

TABLE 2 Summary of regulatory approvals for biologicals in severe asthma—European Medical Agency

Product	Population	Posology	Remarks
Benralizumab (Fasenra®)	Add-on maintenance treatment in adult and adolescent patients with severe eosinophilic asthma inadequately controlled despite high-dose ICS corticosteroids plus LABA.	30mg SC every 4 weeks for the first 3 doses, and then every 8 weeks thereafter.	Intended for long-term treatment. The need for continued therapy should be considered at least on an annual basis as determined by physician assessment of the patient's disease severity and level of control of exacerbations. Special caution for helminth infections.
Dupilumab (Dupixent®)	Adults and adolescents 12 years and older as add-on maintenance treatment for severe asthma with type 2 inflammation characterised by raised blood eosinophils (> 150) and/or raised FeNO >20), inadequately controlled with high dose ICS plus another maintenance treatment.	On OCS or with co-morbid AD, initial dose of 600 mg followed by 300 mg every other week SC. For all other patients, initial dose of 400 mg followed by 200 mg every other week SC.	Intended for long-term treatment. The need for continued therapy should be considered at least on an annual basis based on level of asthma control.
Mepolizumab (Nucala®)	Severe refractory eosinophilic asthma in adults, adolescents and children aged 6 years and older.	> 12 years old 100 mg SC (pre-filled pen) once every 4 weeks. Children 6-11 years old 40 mg SC (powder for solution) once every 4 weeks.	Intended for long-term treatment. The need for continued therapy should be considered at least on an annual basis as determined by physician assessment of the patient's disease severity and level of control of exacerbations.
Omalizumab (Xolair®)	>12 years old: add-on therapy to improve asthma control in patients with severe persistent allergic asthma who have a positive skin test or in vitro reactivity to a perennial aeroallergen and who have reduced lung function (FEV1<80%) as well as frequent daytime symptoms or night-time awakenings, and who have had multiple documented severe asthma exacerbations despite daily high-dose ICS + LABA. 6-12 years old: add-on therapy to improve asthma control in patients with severe persistent allergic asthma who have a positive skin test or in vitro reactivity to a perennial aeroallergen and frequent daytime symptoms or night-time awakenings, and who have had multiple documented severe asthma exacerbations despite daily ICS+LABA.	SC every 2-4 weeks based on total IgE level and body weight.	Intended for long-term treatment. Assessed for effectiveness at 16 weeks (GETE) before further injections are administered.
Reslizumab (Cinqero®)	Adult patients with severe eosinophilic asthma inadequately controlled despite high-dose ICS plus another maintenance treatment.	iv per body weight (see table)	Intended for long-term treatment. The need for continued therapy should be considered at least on an annual basis as determined by physician assessment of the patient's disease severity and level of control of exacerbations.

1.2 | Purpose of the EAACI Guidelines for the use of biologicals in severe asthma

Delivering high-quality clinical care is a central priority for allergists, pneumologists, paediatricians and other specialities caring for patients with allergic diseases and asthma. The European Academy of Allergy and Clinical Immunology (EAACI) develops and updates each year resources to help HCP and researchers to design the best interventions, deliver high standard care and to assess their actions and decisions for purposes of quality improvement and/or reporting.

EAACI Guidelines include recommendations for the management of patients with particular conditions or diseases. Guidelines are developed using a systematic process and are based on available evidence and the clinical experience and expertise of all interested stakeholders.

EAACI developed a Position Paper on Biologicals in 2015.⁵² Due to the rapid accrual of evidence and new therapies, advancement of guideline development methodologies, and the need to broaden the scope of the 2015 recommendations a new guideline was therefore needed.

The current guidelines address only treatment with biologicals of severe asthma and do not address any topics related to severe asthma correct diagnosis, background controller treatment, achieving asthma control, monitoring adherence or treating its co-morbidities.

The EAACI Guidelines for the use of biologicals in severe asthma are not intended to impose a standard of care. Instead, they provide the framework for rational decisions for the use of biologicals in severe asthma by HCPs, patients, third-party payers, institutional review committees and other stakeholders. Statements regarding the underlying values and preferences as well as qualifying remarks accompanying

TABLE 3 Summary of regulatory approvals for biologicals in severe asthma—Food and Drug Administration

Product	Population	Posology	Remarks
Benralizumab (Fasenra®)	Add-on maintenance treatment of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype (blood eosinophil counts greater than or equal to 150 cells/ μ L).	30 mg administered SC once every 4 weeks for the first 3 doses, and then once every 8 weeks thereafter.	Caution for: 1. Hypersensitivity reactions 2. Helminth infections 3. Abrupt discontinuation of OCS or ICS
Dupilumab (Dupixent®)	Add-on maintenance treatment in patients with moderate-to-severe asthma aged 12 years and older with an eosinophilic phenotype or with oral corticosteroid dependent asthma.	1. An initial dose of 400 mg followed by 200 mg given every other week. 2. An initial dose of 600 mg followed by 300 mg given every other week for patients requiring concomitant oral corticosteroids or with co-morbid moderate-to-severe atopic dermatitis start with an initial dose of 600 mg followed by 300 mg given every other week.	Caution for: 1. Hypersensitivity reactions 2. Helminth infections 3. Abrupt discontinuation of OCS or ICS 4. Eosinophilic conditions
Mepolizumab (Nucala®)	Add-on maintenance treatment of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype (blood eosinophils of \geq 150 cells/mcL at screening (within 6 weeks of dosing) or blood eosinophils of \geq 300 cells/mcL within 12 months of enrollment).	100 mg administered subcutaneously once every 4 weeks.	Caution for: 1. Hypersensitivity reactions 2. Helminth infections 3. Herpes zoster infections 4. Abrupt discontinuation of OCS or ICS
Omalizumab (Xolair®)	Moderate to severe persistent asthma in patients 6 years of age and older with a positive skin test or <i>in vitro</i> reactivity to a perennial aeroallergen and symptoms that are inadequately controlled with inhaled corticosteroids.	75 to 375 mg SC every 2 or 4 weeks. Determine dose (mg) and dosing frequency by serum total IgE level (IU/mL), measured before the start of treatment, and body weight (kg). See the dose determination charts.	Black box warning for anaphylaxis Caution for: 1. Abrupt discontinuation of OCS or ICS 2. Eosinophilic conditions 3. Serum sickness
Reslizumab (Cinqair®)	Add-on maintenance treatment of patients with severe asthma aged 18 years and older with an eosinophilic phenotype (blood eosinophil count of at least 400 cells/mcL within 3 to 4 weeks of dosing).	3 mg/kg once every 4 weeks administered by intravenous infusion over 20-50 minutes.	Black-box warning for anaphylaxis Caution for: 1. Helminth infections 2. Abrupt discontinuation of OCS or ICS

each recommendation are an integral part of the Guidelines and aim to facilitate more accurate interpretation. They should never be omitted or ignored when quoting Guidelines recommendations.

1.2.1 | Target audience

The target audience includes all HCPs involved in the management of severe asthma, patients and caregivers, basic scientists involved in biologicals development, regulatory authorities and policymakers.

1.2.2 | Biologicals included—rationale for choosing

The EAACI Guidelines provide recommendations for the use of biologicals with current regulatory approval in patients with severe asthma: benralizumab, dupilumab, mepolizumab, omalizumab, reslizumab (in alphabetical order).

Additional comments are provided for the biologicals currently tested and not yet approved and for doses/routes not approved by regulatory authorities.

2 | METHODS

The EAACI Guidelines followed the GRADE methodology (available at www.gradeworkinggroup.org). Training was conducted with all members of the guidelines development group (GDG) to prepare them for their roles, including specific sessions on the GRADE methodology.

2.1 | The Guidelines Development Group

A Core Leadership Team (Table S1) supervised the project and was responsible for defining the project scope, drafting the clinical questions to be addressed by the guideline, coordinating the search and drafting the manuscript together with the Voting Panel (Table S1). It was led by three chairs with both content and methodologic expertise. The Core Leadership Team received support from a methodologist team, who advised on the process and provide input on the GRADE summary of findings (SOF) tables and from experts in guideline development.

The methodologist team conducted the systematic literature reviews (SRs) for each of the clinical questions, graded the quality

TABLE 4 Structured questions for the systematic reviews

“Is the treatment with biologicals (i.e., benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) efficacious and safe for patients with uncontrolled severe eosinophilic asthma?”

“Is treatment with benralizumab, dupilumab and omalizumab efficacious and safe for patients with allergic asthma?”

“Is the treatment with dupilumab efficacious and safe for patients with severe asthma?”

of evidence, developed the SOF tables and provided the evidence reports. Narrative reviews were conducted by different content specialist subgroups for each topic to be covered to complement the SRs.

The Voting Panel, composed of content experts, decided which clinical questions are to be asked and which outcomes are critical, important and of low importance, and voted for the final recommendations after reviewing the evidence provided by the methodology team and the narrative reviews. It included specialists with expertise and clinical experience in treating severe asthma, biologists and clinical immunology experts, as well as patient representatives.

In accordance with EAACI policy, everyone who was intellectually involved in the project (ie considered for guideline authorship) disclosed all potential conflict of interest (COIs) in writing at the beginning, middle and end of the project. The Guideline Oversight Committee (Table S1) was responsible for developing and implementing rules related to COIs.

2.2 | Definitions

For the purpose of the three SRs^{63–65} that informed the recommendations, severe asthma populations were defined as follows:

- Eosinophilic asthma: subjects with any of the following: a sputum eosinophil count of >1% or an asthma-related peripheral blood eosinophil count of ≥ 150 cells/ μ L, or a fractional exhaled nitric oxide (FeNO) of ≥ 20 ppb⁶⁶
- Allergic asthma: subjects diagnosed with moderate-to-severe allergic asthma with asthma symptoms due to exposure to a perennial aeroallergen and serum total IgE levels 30–1300 IU/mL not adequately controlled on ICS and/or other background controllers.
- For dupilumab SR, the severe T2 asthma population was defined as subjects with confirmed diagnosis of asthma inadequately controlled on ICS and additional controllers

For the recommendations, the population was defined as in the clinical trials that informed the regulatory approval.

2.3 | TF questions and prioritizations of key outcomes

Clinically relevant interventions and comparators were developed balancing comprehensiveness with feasibility (Table 4). The most

challenging decision in framing the question was how broadly the patients and intervention should be defined. The underlying biology of asthma suggested that across the range of patients and interventions, it is plausible that the magnitude of effect on the key outcomes is different; thus, the GDG defined subpopulations based on age (6–11 years old, 12–18 years old, >18 years old), co-morbidities (chronic rhinosinusitis with nasal polyps (CRSwNP) for anti-IL-5 interventions and dupilumab; atopic dermatitis (AD) for dupilumab; chronic urticaria (CU), allergic rhinitis (AR) and food allergy for omalizumab, biomarkers (blood eosinophils, atopy, specific and total IgE, FeNO), dose, etc. The outcomes were evaluated per product and subgroup.

As required by the GRADE approach, asthma-related outcomes were prioritized in a first step by the GDG using a 1 to 9 scale (7 to 9 critical; 4 to 6 important; 1 to 3 of limited importance). The critical outcomes were as follows: severe asthma exacerbations, asthma control measured by the asthma control questionnaire (ACQ) and asthma control test (ACT), QoL measured by asthma quality of life questionnaire (AQLQ) and safety. The important outcomes were as follows: lung function measured by the forced expiratory volume at first second (FEV₁), decrease in ICS and OCS dose and in rescue medication use. FeNO, sputum and blood eosinophils were scored as low importance (Table 5). After reviewing the evidence, the prioritization of the outcomes was reassessed to ensure that important outcomes that were not initially considered are included and to reconsider the relative importance of outcomes in light of the available evidence. All asthma-related relevant outcomes were addresses simultaneously. As per GRADE methodology when surrogate outcomes were used (FeNO, FEV₁), the quality of the evidence was down-graded.

The GDG framed a separate cost-effectiveness question to assess the economic impact of the biologicals versus standard of care. The outcomes of interest were costs and resources use, the incremental cost-effectiveness ratios (ICERs) per both quality-adjusted life-years (QALY) and asthma-related outcomes.

The GDG also defined and addressed clinical questions not covered by the systematic reviews (Table 6).

2.4 | The minimal important difference

To evaluate the imprecision for each outcome, the minimal important difference (MID) thresholds were considered: St George's Respiratory Questionnaire (SGRQ) score change of 4 units, Asthma Control Questionnaire (ACQ-5, ACQ-6 and ACQ-7) score change

TABLE 5 Asthma-related outcomes; grading importance for the systematic reviews

Outcome	Importance
Severe asthma exacerbations Asthma control Quality of life Safety (adverse events)	Critical
Lung function (FEV ₁) Decrease in ICS dose and OCS dose Rescue medication use	Important
FeNO, sputum and blood eosinophils	Low importance

of 0.5 units, Asthma Quality of Life Questionnaire (AQLQ) score change of 0.5 units (with disclaimer as calculated pre-/post-treatment), FEV₁ change in litres 0.20, OCS reduction by 50%, rescue medication reduction by 0.81 puff/day and a reduction of at least 20% in FeNO for values over 50 ppb or more than 10 ppb for values lower than 50 ppb.⁶⁷⁻⁷⁰

2.5 | The GRADE approach (search, appraisal of the evidence)

Key principles and provisions, key terms, descriptions, drug categories, PICO (population, intervention, comparator and outcomes) questions, search methodology and evidence reporting used in the guideline development process were predefined.

Separate systematic reviews on eosinophilic asthma (benralizumab, dupilumab, mepolizumab, omalizumab, reslizumab), allergic asthma (benralizumab, dupilumab, omalizumab) and severe T2 asthma (dupilumab) were conducted to inform the recommendations.⁶³⁻⁶⁵ A GRADE SOF table was provided for each PICO question. The quality of evidence was evaluated based on GRADE quality assessment criteria by two independent reviewers and discordance resolved by consensus. Quality assessment includes the risk of bias (ROB) in included trials, the likelihood of publication bias, inconsistency between trial results, indirectness of the evidence (eg differences between populations, interventions or outcomes of interest in the group to whom the recommendation applies versus those who were included in the studies referenced) and imprecision (wide confidence intervals, usually due to a small number of patients or events, or those situations where clinical decision-making would differ at the extremes of the confidence interval).^{71,72} The quality of evidence for each outcome was rated as high, moderate, low or very low. In the absence of any data, the level of evidence was rated as very low, based on clinical experience only. Search results were pooled in an evidence report as SOF tables and accompanied by a qualitative summary of the evidence for each PICO question. The Content Panel reviewed

the drafted evidence report to address evidence gaps prior to presentation to the Voting Panel.

2.6 | Additional evidence

In support of formulated recommendations, the GDG performed narrative reviews collecting evidence on phase IV, observational and real-world trials and on clinical questions not addressed by the SRs (Table S2).

2.7 | Consensus building and formulating recommendations

After reviewing the evidence report and the additional evidence, the Voting Panel decided in a face-to-face meeting followed by subsequent emails regarding the final recommendations. For each PICO question, the Voting Panel heard an oral summary of the evidence and provided votes on the direction and strength of the related recommendation. A 70% consensus threshold was reached for all recommendation presented below. The recommendations follow the data included in the evidence-to-decision (EtD) tables and take into consideration the balance of desirable and undesirable consequences, quality of evidence, patients' values and preferences, feasibility, and acceptability of various interventions, use of resources paid for by third parties, equity considerations, impacts on those who care for patients and public health impact.^{71,72} A strong recommendation was made in favour of an intervention when the GDG was certain that the desirable consequences outweighed the undesirable consequences. A conditional recommendation was provided if there were reasons for uncertainty on the benefit-risk profile, especially for low or very low quality of evidence. The underlying values and preferences played a key role in formulating recommendations. As the key target audience of the EAACI Guidelines is HCPs and the patients they treat, the perspective chosen when formulating recommendation was mainly that of the HCPs and of

the patient, although the health system perspective was also evaluated.⁷³ Recommendation is provided per product, per subgroups and per outcomes. The recommendations formulated in these guidelines should be used following the GRADE interpretation (Table 7). These recommendations should be reconsidered when new evidence becomes available and an update of these guidelines is planned for 2024.

Where no evidence was available, the GDG formulated expert-based recommendations.

The Guidelines were available on the EAACI website for two weeks (20 April- 4 May) for public comment and were external peer-reviewed. All comments received were carefully revised by the GDG and where applicable incorporated.

2.8 | Final review and approval of the guideline by EAACI

In addition to journal and external peer review, the EAACI Scientific Committee and Executive Committee reviewed the manuscript. These EAACI oversight groups did not mandate that certain recommendations be made within the guideline, but rather serve as peer reviewers.

3 | KEY RECOMMENDATIONS (BIOLOGICALS ARE MENTIONED IN ALPHABETICAL ORDER)

3.1 | Benralizumab—severe eosinophilic asthma—adults and paediatric population 12- to 17- years old

IL-5 is one of the major cytokines responsible for the growth, differentiation, recruitment, activation and survival of eosinophils. Benralizumab is a humanized Mab that binds to the α subunit of the IL-5 receptor (IL-5R α) via its antigen-binding fragment (Fab) domain, blocking the binding of IL-5 to its receptor and resulting in inhibition of eosinophil differentiation and maturation in bone marrow. In addition, this antibody is able to bind through its afucosylated fragment crystallizable (Fc) domain of the IgG receptor Fc γ RIIIa on natural killer (NK) cells, macrophages and neutrophils, thus strongly inducing antibody-dependent, cell-mediated cytotoxicity in both circulating and tissue-resident eosinophils.⁷⁴ This double function of benralizumab rapidly induces and maintains depletion of eosinophils that is much greater than that induced by other monoclonal antibodies targeting the IL-5 pathway.^{75,76} This effect translates clinically into a rapid and significant improvement of patient-reported outcomes (PROs) and of lung function (peak expiratory flow (PEF), together with a significant decrease in asthma exacerbations and OCS use.⁷⁷⁻⁸¹

The summary of the supportive evidence is presented in Tables S3 and S4. Recommendations are based on the evidence-to-decision Tables S5-S7.

TABLE 6 Clinical questions not covered by the systematic reviews

1. Dose not approved by FDA/EMA
2. Route not approved by FDA/EMA
3. Biological not approved by EMA/FDA
4. Relevance of clinical trial population for real-world patients
5. Efficacy in the paediatric population, especially 6-11 years old
6. Safety long term (> 5 years)
7. Safety in the paediatric population
8. Immunogenicity
9. Best method to monitor anti-drug antibodies (ADA)
10. Joint treatment of co-morbidities (asthma and CRSwNP or AR or AD)
11. Biomarkers
12. Continuation/discontinuation criteria
13. Switching rules
14. Defining efficacy; definition of responder, partial responder (dissociated outcome), non-responder
15. Time to achieve efficacy
16. Treatment duration
17. Combinations between biologicals
18. Effects after administration in the emergency department
19. Health economics data not included in the SR

Recommendations are formulated separately for the adult population (Box 1) and the 12- to 17-year-old population (Box 2).

3.1.1 | Justification

There is high certainty for adults for decreasing asthma exacerbations in the overall group and for OCS decrease in the subgroup with >150 eosinophils/ μ L. Although there is high certainty for improving asthma control and QoL, the effect of benralizumab did not reach the MID; thus, a conditional recommendation was formulated.

The GDG formulated conditional recommendations for the paediatric population due to the low numbers of subjects 12-17 years old included in clinical trials.

3.1.2 | Subgroups: stratified by biomarkers and co-morbidities

The higher the blood eosinophils, the higher the expected impact of benralizumab on exacerbations, asthma control, QoL and lung function (Table S8) (conditional recommendation).

TABLE 7 Interpretation of GRADE recommendations

Implications	Strong recommendation	Conditional (weak) recommendation
For patients	Most individuals in this situation would want the recommended course of action and only a small proportion would not. Formal decision aids are not likely to be needed to help individuals make decisions consistent with their values and preferences.	The majority of individuals in this situation would want the suggested course of action but many would not.
For clinicians	Most individuals should receive the intervention. Adherence to this recommendation according to the guideline could be used as a quality criterion or performance indicator.	Recognise that different choices will be appropriate for individual patients and that you must help each patient arrive at a management decision consistent with his or her values and preferences. Decision aids may be useful helping individuals making decisions consistent with their values and preferences.
For policy makers	The recommendation can be adapted as policy or performance measure in most situations	Policy making will require substantial debate and involvement of various stakeholders. Documentation of appropriate (e.g. shared) decision-making processes can serve as performance measure.

BOX 1 Recommendations for benralizumab as add-on treatment in adults with uncontrolled severe eosinophilic asthma

1. Benralizumab is recommended in adults with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Decrease or withdraw oral corticosteroids for blood eosinophils > 150/ μ L	Strong recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation**
2. Benralizumab demonstrated a good safety profile; however, patients should be regularly screened for parasitic infections in endemic areas		Conditional recommendation

*population: severe asthma uncontrolled by high-dosage ICS + LABA with baseline blood eosinophil cell counts >300 cells/ μ L or >150 cells/ μ L for OCS-dependent patients.

**Although the effect size is small, it might be beneficial in patients with very low lung function.

Neither the atopic status nor total IgE predicts the magnitude of effect of benralizumab (conditional recommendation) (Table S9).

Benralizumab can be recommended in patients with severe eosinophilic asthma and CRSwNP uncontrolled despite optimal treatment to:

1. Decrease asthma exacerbations (conditional recommendation)

2. Improve lung function (conditional recommendation) (Table S10)

3.2 | Benralizumab—severe allergic asthma—adults

In humans, IL-5 is often co-expressed with other T2 cytokines including IL-4 and IL-13 and associated in atopic individuals with increased

BOX 2 Recommendations for benralizumab as add-on treatment in the paediatric population 12- to 17- years old with uncontrolled severe eosinophilic asthma

1. Benralizumab is recommended in the paediatric population 12- to 17- years old with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation**
2. Benralizumab demonstrated a good safety profile; however, patients should be regularly screened for parasitic infections in endemic areas		Conditional recommendation

*Population: severe asthma uncontrolled by high-dosage ICS + LABA with baseline blood eosinophil cell counts >300 cells/ μ L or >150 cells/ μ L for OCS-dependent patients.

**Although the effect size is small, it might be beneficial in patients with very low lung function.

IgE production.^{28,30,31,82,83} Thus, targeting the IL-5 pathway might be beneficial in allergic asthma.

From all anti-IL-5 biologicals, only benralizumab was evaluated in the regulatory approved dose in a post hoc analysis of a subpopulation of adults with severe allergic asthma.⁸⁴

The summary of the supportive evidence is presented in Tables S11 and S12. Recommendations are based on the evidence-to-decision Tables S13-S15 and are presented in Box 3

3.2.1 | Justification

There is high certainty for adults for decreasing asthma exacerbations; however, data are derived from a post hoc analysis. Although there is high certainty for improving asthma control and QoL, the effect of benralizumab did not reach the MID; thus, a conditional recommendation was formulated.

3.3 | Dupilumab—severe eosinophilic asthma—adults and paediatric population 12-17 years old

IL-4 and IL-13 are key cytokines in driving the initiation and the chronicity of T2 inflammation, where IL-4 is considered an initiator of T2 immune responses and IL-13 an effector molecule.⁸⁵⁻⁸⁹

Dupilumab is a human IgG4 Mab that targets the IL-4 receptor alpha chain (IL-4R α), common to both IL-4R complexes: type 1 (IL-4R α / γ c; IL-4 specific) and type 2 (IL-4R α /IL-13R α 1; IL-4 and IL-13 specific).^{90,91} In mice models, dual IL-4/IL-13 blockade prevents eosinophil infiltration into lung tissue without affecting circulating eosinophils.⁹⁰ Supported by a strategic mechanism of action inhibiting T2-mediated inflammation, dupilumab's efficacy has been assessed across a range of atopic diseases, such as AD, asthma and chronic rhinosinusitis with nasal polyps (CRSwNP) which often occur together as co-morbidities.⁹¹ Previous to being approved for T2 asthma, dupilumab has been approved in the United States for

BOX 3 Recommendations for benralizumab as add-on treatment in adults with uncontrolled severe allergic asthma

1. Benralizumab is recommended in adults with uncontrolled severe allergic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation**
2. Benralizumab demonstrated a good safety profile; however, patients should be regularly screened for parasitic infections in endemic areas		Conditional recommendation

*Population: severe asthma uncontrolled by high-dosage ICS + LABA with baseline blood eosinophil cell counts >300 cells/ μ L or >150 cells/ μ L meeting the criteria for atopy (by Phadiatop test) and serum IgE concentration of 30 to 700 kU/L (US) and 30 to 1500 kU/L (EU).

**Although the effect size is small, it might be beneficial in patients with very low lung function.

the treatment of atopic dermatitis (AD) and CRSwNP and in Europe for AD. Convincing clinical results were recently published for severe uncontrolled asthma⁹²⁻⁹⁴ and for CRSwNP.⁹⁵

The summary of the supportive evidence is presented in Tables S16 and S17. Recommendations follow the evidence-to-decision Tables S18 and S19.

Recommendations are formulated for the adult and adolescent population altogether (Box 4) as we did not perform a separate analysis for the 12- to 17-year-old subgroup.

3.3.1 | Justification

There is high certainty for adults for decreasing asthma exacerbations and moderate certainty for improving asthma control and QoL and the decrease in rescue medication use; thus, a conditional recommendation was formulated. The increase in FEV₁ is significant (above the MID) for the adult subgroup with blood eosinophils >300 cells/ μ L or with FeNO levels >50 ppb.⁶⁴ More efficacy and safety data are needed in the paediatric population.

3.3.2 | Subgroups: stratified by biomarkers and comorbidities

The higher the blood eosinophils and FeNO the higher, the clinical efficacy of dupilumab, both in reduced exacerbations and in improving lung function (conditional recommendation) (Tables S20 and S21).

Dupilumab can be recommended in adults and adolescents with severe eosinophilic asthma with chronic rhinosinusitis with nasal polyps who are uncontrolled despite optimal treatment to:

1. Decrease exacerbations (conditional recommendation)
2. Improve asthma control (conditional recommendation)
3. Improve quality of life (conditional recommendation)
4. Improve lung function (conditional recommendation)

3.4 | Dupilumab—severe allergic asthma—adults and paediatric population 12- to 17- years old

Ligand binding of the type 1 (IL-4R α / γ c; IL-4 specific) and type 2 (IL-4R α /IL-13R α 1; IL-4 and IL-13 specific) receptors activates a signal transduction cascade that mainly leads to the modulation of expression of genes involved in IgE class switching, T helper (TH) 2 cell differentiation and M2 macrophage polarization.^{31,82,86-88} IL-4 has been shown to stimulate IgE production from B cells. In asthma, IL-4 plays a major role in TH2 cell proliferation, cytokine production and IgE synthesis. Combined analyses of genetic alterations in the IL-4/IL-13 pathway showed a profound influence on serum IgE levels and the risk of childhood asthma.⁹⁶ Dupilumab was specifically evaluated in a

post hoc analysis of patients with moderate-to-severe allergic asthma enrolled in QUEST trial. Allergic asthma was defined by total serum IgE \geq 30 IU/mL and \geq 1 perennial aeroallergen-specific IgE \geq 0.35 kU/L at baseline.⁹⁷

The summary of the supportive evidence is presented in Tables S22 and S23. Recommendations follow the evidence-to-decision Tables S24 and S25.

Recommendations are formulated for the adult and adolescent population altogether (Box 5) as we did not perform a separate analysis for the 12- to 17-year-old subgroup.

3.4.1 | Justification

There is high certainty for adults for decreasing asthma exacerbations and for improving asthma control. However, because the improvement in asthma control did not reach the MID, a conditional recommendation was formulated. The increase in FEV₁ is significant (above the MID) for the adult subgroup with blood eosinophils >300 cells/ μ L or with FeNO levels > 50 ppb.⁶⁴ More efficacy and safety data are needed in the paediatric population. As data are derived from a post hoc analysis, a conditional recommendation was formulated.

3.5 | Dupilumab—severe T2 asthma—adults and paediatric population 12- to 17- years old

T2 asthma encompasses several pathways that are concomitantly activated, including cells of the innate immune system such as innate lymphoid cells, macrophages, neutrophils and natural killer T cells which are effective producers of a variety of cytokines and seem to play important roles in the development of nonallergic asthma.^{27,30,32} IL-13 cytokine serum levels were significantly high in atopic and nonatopic asthma patients compared to healthy controls. IL-4 and IL-13 can curtail chemotaxis and several effector functions of neutrophils in humans.^{98,99} In a recent analysis of the QUEST trial, dupilumab reduced severe exacerbation rates, improved FEV₁ and asthma control, and suppressed T2 inflammatory biomarkers in patients with uncontrolled, moderate-to-severe asthma with or without evidence of allergic asthma.⁹⁷ Following this rationale, the GDG decided to perform a SR on the efficacy and safety of dupilumab without using the criteria for eosinophilic or allergic asthma.⁶⁵ This SR is the basis for the recommendations for dupilumab as add-on treatment in patients with T2 asthma uncontrolled under medium- to high-dose ICS plus up to two additional controllers.

The summary of the supportive evidence is presented in Tables S26 and S27. Recommendations follow the evidence-to-decision Tables S28-S30.

Recommendations are formulated for the adult and adolescent population altogether (Box 6) as there were no differences between the adults and the 12- to 17-year-old subgroup.

BOX 4 Recommendations for dupilumab as add-on treatment in adults and paediatric population 12- to 17- years old with uncontrolled severe eosinophilic asthma

1. Dupilumab is recommended in adults and paediatric population 12- to 17- years old with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function**	Strong recommendation
	Decrease rescue medication use***	Conditional recommendation
2. Dupilumab demonstrated a good safety profile; however, longer term data (up to 2 years) are extrapolated from atopic dermatitis studies and careful reporting of all drug-related adverse events is recommended		Conditional recommendation

*Population: severe asthma uncontrolled by medium-/high-dose ICS plus up to 2 additional controllers including OCS; T2 inflammation (EMA requirements) characterized by raised blood eosinophils (>150) and/or raised FeNO > 20.

**Population: adult subgroup with blood eosinophils >300 cells/ μ L or with FeNO levels >50 ppb.⁶⁴

***Although the effect size is small, it might be beneficial in patients with increased risk due to excessive use of rescue medication.

BOX 5 Recommendations for dupilumab as add-on treatment in adults and in the paediatric population 12-17 years old with uncontrolled severe allergic asthma

1. Dupilumab is recommended in adults and in the paediatric population 12- to 17- years old with uncontrolled severe allergic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function**	Conditional recommendation
2. Dupilumab demonstrated a good safety profile; however, longer term data (up to 2 years) are extrapolated from atopic dermatitis studies and careful reporting of all drug-related adverse events is recommended		Conditional recommendation

*Population: severe asthma uncontrolled by medium-/high-dose ICS plus up to 2 additional controllers including OCS; T2 inflammation (EMA requirements) characterized by raised blood eosinophils (>150) and/or raised FeNO > 20.

**Population: adults subgroup with blood eosinophils >300 cells/ μ L or with FeNO levels > 50 ppb.⁶⁴

3.5.1 | Justification

There is high certainty for adults for decreasing asthma exacerbations and for OCS reduction. Although there is high certainty for improving asthma control and QoL, the effect does not reach above the MID; thus, a conditional recommendation was formulated. The increase in FEV₁ is significant (above the MID) for the adult subgroup with blood eosinophils >300 cells/ μ L and/or with FeNO levels >50 ppb⁶⁵ and for OCS-dependent patients.⁹⁴ More efficacy and safety data are needed in the paediatric population.

3.5.2 | Subgroups

Dupilumab is effective in both allergic* and nonallergic asthma in a T2 asthma context (conditional recommendation).

*a total serum IgE \geq 30 IU/mL and \geq 1 positive perennial aeroallergen-specific IgE value (\geq 0.35 kU/L) at baseline (Table S31A,B).

3.6 | Mepolizumab—eosinophilic severe asthma—adults and paediatric population 12-17 years old

Mepolizumab is humanized Mab of IgG1 κ type, which targets human IL-5 with high-affinity and specificity. Mepolizumab inhibits the bioactivity of IL-5 with nanomolar potency by blocking the binding of IL-5 to the alpha chain of the IL-5 receptor complex expressed on the eosinophil cell surface, thereby inhibiting IL-5 signalling and reducing the production and survival of eosinophils.¹⁰⁰ It was approved for severe asthma with peripheral eosinophilia. Good control of peripheral eosinophilia has been readily demonstrated together with translation into a significant reduction in asthma exacerbations and an OCS sparing effect.¹⁰¹⁻¹⁰³ Its effect on asthma control, QoL and lung function is less clear.

The summary of the supportive evidence is presented in Tables S32 and S33. Recommendations follow the evidence-to-decision Tables S34-S37.

Recommendations are formulated separate for the adult population (Box 7) and the 12- to 17-year-old population (Box 8).

BOX 6 Recommendations for dupilumab as add-on treatment in adults and paediatric population 12- to 17- years old with uncontrolled severe T2 asthma

1. Dupilumab is recommended in adults and paediatric population 12- to 17- years old with uncontrolled severe T2 asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Decrease or withdraw oral corticosteroids**	Strong recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function***	Strong recommendation
	Decrease rescue medication****	Conditional recommendation
2. Dupilumab demonstrated a good safety profile; however, longer term data (up to 2 years) are extrapolated from atopic dermatitis studies and careful reporting of all drug-related adverse events is recommended		Conditional recommendation

*Population: severe asthma uncontrolled by medium-/high-dose ICS plus up to 2 additional controllers including OCS; T2 inflammation (EMA requirements) characterized by raised blood eosinophils (>150) and/or raised FeNO > 20.

**Population: Patients with severe asthma on maintenance OCS and high-dose ICS in combination with a second controller.

***Population: adult subgroup with blood eosinophils >300 cells/ μ L or with FeNO levels >50 ppb⁶⁵ or OCS-dependent patients.⁹⁴

****Although the effect size is small, it might be beneficial in patients with increased risk due to excessive use of rescue medication.

BOX 7 Recommendations for mepolizumab as add-on treatment in adults with uncontrolled severe eosinophilic asthma

1. Mepolizumab is recommended in adults with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Decrease or withdraw oral corticosteroids	Strong recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation**
2. No recommendation can be made for reducing rescue medication use as the effect size is small		
3. Mepolizumab demonstrated a good safety profile (long-term safety data up to 5 years); however, patients should be regularly screened for parasitic infections in endemic areas		Conditional recommendation

*Population: Severe eosinophilic asthma defined as presence of eosinophilic inflammation determined by a blood eosinophil level of either 300 cells or more per μ L in the past 12 months or 150 cells or more per μ L at initiation.

**Although the effect size is small, the improvement might be relevant in severe asthma with very low lung function.

No recommendations can be formulated for children 6- to 11- years old as per extrapolated data from mepolizumab 12- to 17- years old population.

3.6.1 | Justification

There is high certainty for the decrease of moderate and severe asthma exacerbations and for the decrease in OCS and moderate or low certainty for improving asthma control, quality of life and lung function and for monitoring for parasitic infections. Only conditional recommendations can be formulated for the paediatric group 12- to 17- years old as there were low numbers enrolled in clinical trials.

3.6.2 | Subgroups: stratified by biomarkers and co-morbidities

The higher the blood eosinophils, the higher the expected impact of mepolizumab on exacerbations (Table S38) (conditional recommendation).

Mepolizumab can be recommended in adult patients with severe eosinophilic asthma and CRSwNP uncontrolled despite optimal treatment to:

- decrease asthma exacerbations (conditional recommendation)
- decrease or withdraw OCS (conditional recommendation)
- Improve lung function (conditional recommendation)
- Improve quality of life (conditional recommendation)

For the use of mepolizumab in allergic asthma in a licensed dose, an open multicentre, open-label, single-arm study showed clinically significant improvements in asthma control, health status and exacerbation rate.¹⁰⁴ The post hoc meta-analysis of two phase 3 studies showed efficacy of mepolizumab regardless of allergic characteristics or omalizumab eligibility.¹⁰⁵ Recent reports show that both mepolizumab and reslizumab can control severe asthma patients that failed to respond to omalizumab.¹⁰⁶ The GDG considered that there are not enough data to conduct a SR on the licenced dose of mepolizumab in severe allergic asthma and a recommendation cannot be currently formulated.

3.7 | Omalizumab—severe eosinophilic asthma—adults

While immunoglobulin E is a prominent biomarker for early-onset asthma, its serum levels are often also elevated in nonallergic late-onset asthma. The pattern of IgE expression in the latter is mostly polyclonal and frequently associated with high blood eosinophils.^{107,108} The innate immune response also significantly contributes to this asthma phenotype. Release of IL-33 and TSLP from respiratory epithelium and activation of ILC2s via its receptor ST2 followed by T2 cytokine release from ILC2s and TH 2 cells drives further massive local B-cell activation and IgE formation, mast cell degranulation and finally eosinophil attraction.^{99,109,110} Omalizumab indirectly downregulates the IgE high-affinity receptor (FcεRI) expression on basophils, mast cells and dendritic cells (DCs), decreasing T2 cytokine production and inhibiting the eosinophilic inflammation.¹¹¹⁻¹¹⁷ Through a different mechanism, long-term omalizumab treatment dampens T2 inflammation acting on different cell types that play a pivotal role in the pathogenesis of asthma such as plasmacytoid DCs (pDCs) and CD-4 T helper cells.¹¹⁸ Several publications of case reports or short series describe the use of omalizumab

adults and children suffering from severe nonallergic asthma, sometimes reporting the successful withdrawal of systemic corticosteroids under treatment.¹¹⁹⁻¹²¹ Better response to omalizumab for eosinophil counts >300 cells/μL are shown in a prospective trial and in a recent pooled analysis of the two pivotal trials of omalizumab in allergic asthma.¹²²⁻¹²⁴ However, in two real-life studies omalizumab proved its efficacy regardless of the blood eosinophils status.^{125,126}

The summary of the supportive evidence is presented in Table S39. Recommendations follow the evidence-to-decision Tables S40 and S41. Recommendations are formulated only for the adult population (Box 9).

3.8 | Omalizumab—moderate-to-severe allergic asthma—adults and paediatric population 12- to 17- years old

The first in a new class of biological therapies for allergic asthma, omalizumab works by binding to free IgE thereby, reducing the amount available to bind the IgE high-affinity receptor (FcεRI) on mast cells, basophils and antigen presenting cells.⁸⁰ The ability to detach IgE from its receptor following omalizumab was recently described.¹¹⁸ The pDCs act at the crossroads between innate and adaptive immunity. IgE receptor cross-linking on pDCs suppresses their anti-viral activity. Thus, a reduced expression of IgE receptors on pDCs and a reduced amount of circulating IgE might generally strengthen anti-viral immune responses following omalizumab administration.^{128,129} Preclinical and clinical evidence supports the existence of a close counter-regulation of the high-affinity IgE receptor and interferon (IFN) pathways, and a potential dual mechanism of action and therapeutic benefit for omalizumab, which may enhance the prevention and treatment of virally induced asthma exacerbations.^{128,130-132} Extensive experience with omalizumab

BOX 8 Recommendations for mepolizumab as add-on treatment in paediatric population 12- to 17- years old with uncontrolled severe eosinophilic asthma

1. Mepolizumab is recommended in children 12- to 17- years old with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Conditional recommendation
	Decrease or withdraw oral corticosteroids	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation
2. Mepolizumab demonstrated a good safety profile (long-term safety data up to 5 years); however, patients should be regularly screened for parasitic infections in endemic areas		Conditional recommendation

*Population: Severe eosinophilic asthma defined as presence of eosinophilic inflammation determined by a blood eosinophil level of either 300 cells or more per μL in the past 12 months or 150 cells or more per μL at initiation.

BOX 9 Recommendations for omalizumab as add-on treatment in adults with uncontrolled severe eosinophilic asthma (both allergic and nonallergic)

1. Omalizumab is recommended in adults with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Improve quality of life	Conditional recommendation
	Improve lung function	Conditional recommendation**
	Decrease the use of rescue medication	Conditional recommendation***
2. Omalizumab has demonstrated a long-term (>10 years) good safety profile; however, 60 minutes monitoring for anaphylaxis is recommended for the first 3 administrations****		Conditional recommendation

*Population: adults with uncontrolled severe persistent allergic asthma with FENO \geq 24ppb, and blood eosinophil counts \geq 260/ μ L.

**Although the effect size is small, the improvement might be relevant in severe asthma with very low lung function.

***Although the effect size is small, it might be beneficial in patients with increased risk due to excessive use of rescue medication.

****The recommendations on the duration of monitoring after omalizumab injection and for the first 3 administrations are expert opinion-based following the data reported in the literature¹²⁷; of note, anaphylaxis has also been documented two hours after administration and beyond 1 year after beginning regularly administered treatment; thus, it is up to the clinician to decide based on the personal history of the patient.

treatment for severe allergic asthma confirmed its effectiveness and safety, reducing symptoms, frequency of reliever use and severe exacerbations.^{133–139}

The summary of the supportive evidence is presented in Tables S42 and S43. Recommendations follow the evidence-to-decision Tables S44 and S45. As we did not perform a separate analysis for the 12–17 population, recommendations are formulated altogether for the adult and adolescent^{12–17} population (Box 10).

3.8.1 | Justification

There is high certainty for the decrease in asthma exacerbations. Although there is high certainty for improving quality of life and for decreasing ICS and rescue medication use, the GDG formulated conditional recommendations as the effect size did not reach the MID (where applicable) or was very small or there was a large confidence interval. There was moderate certainty for improving asthma control, not reaching the MID; however, there was high certainty for the improvement in the global evaluation of treatment effectiveness (GETE).

3.8.2 | Subgroups: stratified by biomarkers and co-morbidities

Serum IgE thresholds (within regulatory limits) do not influence response (conditional recommendation).

The effect of omalizumab on exacerbations does not depend on blood eosinophils (conditional recommendation) (Table S46).

Omalizumab may control associated co-morbidities such as allergic rhinitis, chronic urticaria food allergy, CRSwNP but

asthma-related recommendations and posology should be primarily used (conditional recommendation, expert opinion-based).

3.9 | Omalizumab—moderate-to-severe allergic asthma—paediatric population 6 –to 11- years old

As there is good evidence with omalizumab for the paediatric asthma subgroup 6–11 years old, a separate analysis was performed (see evidence-to-decision Table S47) and the GDG formulated separate recommendations (Box 11).

3.10 | Reslizumab—severe eosinophilic asthma—adults

Reslizumab is an IL-5 antagonist (IgG4-kappa) binding to the IL-5 using a different epitope as compared to mepolizumab and in vitro with greater potency.¹⁴⁰ It is administered via intravenous (iv) infusion, and dosing is based on patients' weight.¹⁴¹ Administration is recommended immediately after preparation. In clinical trials, rare cases of anaphylaxis were observed within 20 minutes of infusion and were reported as early as the second dose. The pooled analysis of six clinical trials reports only three cases of anaphylaxis related to reslizumab, successfully managed with standard therapies.¹⁴² By week 52, patients receiving reslizumab showed a 92% reduction in mean blood eosinophil counts, and this effect translates into a significant decrease in asthma exacerbations.^{143–146}

The summary of the supportive evidence is presented in Tables S48 and S49. Recommendations (Box 12) follow the evidence-to-decision Tables S50–S52.

BOX 10 Recommendations for omalizumab as add-on treatment in adults and the paediatric population 12-17 years old with uncontrolled moderate-to-severe allergic asthma

1. Omalizumab is recommended in adults and the paediatric population 12- to 17- years old with uncontrolled severe allergic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Improve asthma control	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Decrease in the use of ICS	Conditional recommendation**
	Decrease the use of rescue medication	Conditional recommendation***
2. Omalizumab has demonstrated a long-term (>10 years) good safety profile; however, 60 minutes monitoring for anaphylaxis is recommended for the first 3 administrations ****		Conditional recommendation

*Population: moderate-to-severe asthma, total IgE level of 30-700 IU/mL (US) and 30-1500 IU/mL (EU) \pm one perennial aeroallergen.

**Of particular importance for the paediatric population.

***Although the effect size is small, it might be beneficial in patients with increased risk due to excessive use of rescue medication.

****The recommendations on the duration of monitoring after omalizumab injection and for the first 3 administrations are expert opinion-based¹²⁷; of note, anaphylaxis has also been documented two hours after administration and beyond 1 year after beginning regularly administered treatment; thus, it is up to the clinician to decide based on the personal history of the patient.

3.10.1 | Justification

Although there is high certainty for improving asthma control and QoL, the effect does not reach the minimal important difference; thus, a conditional recommendation was formulated. There is moderate certainty for improving lung function and low certainty for monitoring for parasitic infections and the iv administration.

3.10.2 | Subgroups: stratified by biomarkers and co-morbidities

The higher the blood eosinophils, the higher the expected impact of reslizumab on lung function and on asthma control (conditional recommendation) (Table S53).

Reslizumab can be recommended in patients with severe eosinophilic asthma with CRSwNP uncontrolled despite optimal treatment to:

1. decrease moderate and severe asthma exacerbations (conditional recommendation)
2. Improve asthma control (conditional recommendation) (Table S54)

BOX 11 Recommendations for omalizumab as add-on treatment for children 6- to 11- years old with uncontrolled moderate-to-severe allergic asthma

1. Omalizumab is recommended for children 6- to 11- years old with uncontrolled allergic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Conditional recommendation
	Improve asthma control	Conditional recommendation
	Improve quality of life	Conditional recommendation
	Decrease in the use of ICS	Conditional recommendation**
2. Omalizumab has demonstrated a long-term (> 10 years) good safety profile; however, 60 minutes monitoring for anaphylaxis is recommended for the first 3 administrations***		Conditional recommendation
3. Serum IgE thresholds (within regulatory limits) do not influence response		Conditional recommendation
4. Omalizumab might reduce viral-induced exacerbations in this particular population		Conditional recommendation

*Population: moderate-to-severe asthma, total IgE level of 30-700 IU/mL (US) 30-1500 IU/mL (EU) \pm one perennial aeroallergen.

**Of particular importance for the paediatric population.

***The recommendations on the duration of monitoring after omalizumab injection and for the first 3 administrations are expert opinion-based¹²⁷; of note, anaphylaxis has also been documented two hours after administration and beyond 1 year after beginning regularly administered treatment; thus, it is up to the clinician to decide based on the personal history of the patient.

BOX 12 Recommendations for reslizumab as add-on treatment in adults with uncontrolled severe eosinophilic asthma

1. Reslizumab is recommended in adults with uncontrolled severe eosinophilic asthma* in spite of optimal controller treatment to:	Decrease severe asthma exacerbations	Strong recommendation
	Improve quality of life	Conditional Recommendation
	Improve asthma control	Conditional recommendation
	Improve lung function	Conditional recommendation**
2. No recommendation can be made for reducing rescue medication use as the effect size is small; OCS reduction is not reported		
3. Reslizumab demonstrated a good safety profile; however: patients should be regularly screened for parasitic infections in endemic areas patients should be carefully monitored 30 minutes after the iv administration for the risk of anaphylaxis		Conditional recommendation

*Population: at least one blood eosinophil count of 400 cells per μL or higher during a 2-4 weeks of screening period and inadequately controlled asthma, receiving at least a medium dose of ICS with or without another controller drug including OCS.

**Although the effect size is small, the improvement might be relevant in severe asthma with very low lung function.

3.10.3 | Additional considerations

Reslizumab's flexibility in dosage based on body weight might benefit patients with uncontrolled severe eosinophilic asthma not responding to other anti-IL-5 interventions (conditional recommendation, expert opinion-based).

As there are no paediatric data available, no recommendations can be formulated for the use of reslizumab in children with uncontrolled severe asthma.

3.11 | Comparison between biologicals

No comparison can be made between the efficacy and safety of different biologics (strong recommendation, expert opinion-based). Baseline asthma severity, atopic status definition, lung function, eosinophil cut-offs or exacerbation history and asthma duration are all important modulators of treatment efficacy. The rate of background exacerbations (in year prior to trial) or the placebo exacerbations rate (during the trial) should also be considered. These differ across trials because of different inclusion or exclusion criteria; thus, the indirect treatment comparisons may be erroneous or biased.

3.12 | Implementation consideration (for all biologicals)

The GDG formulated strong recommendations for the reduction in asthma exacerbations and for the reduction in OCS dose and conditional recommendations for the other asthma-related outcomes. According to GRADE for strong recommendations, most individuals should receive the intervention and the recommendation can be adapted as policy or performance measure in most situations

(Table 6). However, the GDG cautions on several unsolved key pillars supporting the implementation of these recommendations, such as independent high-quality cost-effectiveness studies, selection of responders, documentation of the disease-modifying effect together with long-term safety data, studies addressing a priori severe asthma together with its co-morbidities. Additional considerations are acceptability of the iv administration for reslizumab or the possibility of self-administration of biologicals at home and at longer intervals.¹⁴⁷ The cost-effectiveness of biologicals based on real-world treatment patterns is unknown. Including broader evidence on treatment discontinuation, caregiver burden and OCS reduction from real-world studies and severe asthma registries may better reflect the effects and value of biologicals for all healthcare stakeholders.⁴⁸ Last but not least, the value of the recommendations depends also on the setting in which the current guideline will be implemented, as recommendation suitable for resource-rich environments might change from strong to conditional in resource-poor environments.

4 | OTHER BIOLOGICALS CURRENTLY TESTED FOR SEVERE ASTHMA

Thymic stromal lymphopoietin (TSLP) is a critical upstream epithelial-derived cytokine inducing T2 inflammation. TSLP activates distinct immune cell cascades in the context of innate and adaptive immune-mediated T2 inflammation.¹⁴⁸ TSLP's importance in human asthma has been repeatedly documented.¹⁴⁹⁻¹⁵¹ Targeting of TSLP-mediated signalling is a potential therapeutic strategy for severe asthma.

Tezepelumab, which is the first-in-class anti-TSLP mAb, is a fully human IgG2 λ mAb that binds human TSLP and prevents interaction with its receptor. Because of the excellent results in the phase II trial which showed a notable reduction in the annual

asthma exacerbation rate in a large population of severe asthma patients,¹⁵² FDA granted tezepelumab a “breakthrough therapy designation” for the treatment of severe asthma without an eosinophilic phenotype.¹⁵³ The drug is currently in phase III clinical trials. In order to formulate a preliminary opinion, a SR was performed for tezepelumab following the same methodology as for the currently approved biologicals for asthma (Table S55). There was moderate certainty that tezepelumab decreases asthma exacerbations, low certainty for improving asthma control and QoL and very low certainty for improving FEV₁ and for decreasing FeNO. There was low certainty for an increase in treatment-related AEs and SAEs. The subgroup analysis by eosinophil and FeNO levels found no difference for the exacerbation rate and FEV₁. For FEV₁, another subgroup analysis by IgE levels found no difference either.

MSTT1041A, an anti-ST2 (IgG2) human monoclonal antibody, has recently been tested in a phase IIb trial, and results are soon to be published.¹⁴⁹ Several other anti-IL-33 Mabs are currently being tested.¹⁵⁴ The anti-IL-33 antibody REGN3500 (SAR440340) met the primary end point of improvement in loss of asthma control when comparing REGN3500 monotherapy to placebo. The greatest improvement was observed in patients with blood eosinophil levels ≥ 300 cells/ μ L.¹⁵⁵ Other anti-IL-33 antibodies, GSK3772847 (NCT03207243) and etokimab, are currently being tested in phase II trials.

5 | BIOLOGICALS EVALUATED FOR SEVERE ASTHMA CURRENTLY DISCONTINUED

Despite promising findings in several experimental models of allergic inflammation, the results of multicentre studies evaluating the efficacy of anti-IL4 and anti-IL-13 mAbs in patients with asthma were negative (Table S56). It appears that individual blockade of IL-4 or IL-13 alone is insufficient to inhibit the complex allergic inflammation. For example, blocking IL-13 does not impact significantly tissue eosinophilia. Other explanations could be that the biomarkers used to identify responders to anti-IL-13 therapy (eg periostin, DPP-4, peripheral eosinophil count) were not optimal or non-T2 asthma patients were included in phase III trials. Finally, systemic administration could not achieve optimal lung concentrations. The preliminary efficacy of a nebulized inhaled anti-IL-13 mAb antigen-binding fragment in macaque model of asthma was recently reported.¹⁵⁶

Targeting the IgE pathway via depletion of IgE-switched and memory B cells was not sufficient for a clinically meaningful benefit for adults with allergic asthma uncontrolled by standard therapy (Table S56), indicating that there are major pathological mechanisms that extend beyond the new, local production of IgE in the lung.¹⁵⁷

The results of clinical trials of biological agents targeting mediators associated with noneosinophilic inflammation, such as IL-17 and tumour necrosis factor (TNF)- α , were disappointing (Table S56). A

better understanding of the mechanisms of noneosinophilic inflammation in asthma should lead to improved therapies, with potential targeted treatment for.²⁹⁻³¹

6 | CLINICAL DECISION ALGORITHM FOR THE INITIATION AND FOLLOW-UP OF BIOLOGICALS FOR THE TREATMENT OF PATIENTS WITH UNCONTROLLED SEVERE ASTHMA

The algorithm developed by the GDG assumes that the diagnosis of severe asthma was correctly performed according to the current guidelines (GINA 2020, ERS/ATS), all co-morbidities and factors influencing asthma control were correctly addressed and patients are still symptomatic on high-intensity controller medication or deteriorate upon step-down.^{2,20,21}

The GDG developed a three-pillar decision tree aiming to help the clinician in reaching the decision to start on a particular biological (Figure 1) based on the combination between phenotypic traits, biomarkers and clinically relevant asthma-related end points (exacerbations, symptoms, lung function, QoL, etc), including safety (conditional recommendation, expert opinion-based). Cost-related and regulatory aspects should also be considered in reaching the decision to start with a particular biological.

The GDG recommends re-evaluation of response after 4-6 months (conditional recommendation, expert opinion-based). Of note, in some patients, exacerbations might not decrease significantly in this short period of time and a composite end point should be used to define response.^{158,159} The definition of a suboptimal response relies on individualized predefined cut-offs of the selected outcomes (conditional recommendation, expert opinion-based). As there is no consensus or validated criteria to define response, the GDG recommends individualized predefined targets established by informed shared decision focused on the patient's goals to control their asthma, in alignment with the principles of personalized treatment. Cost-related, administration at home or in the clinic and regulatory aspects should also be considered in establishing the predefined goals.

For suboptimal response, the GDG recommends to re-assess airway inflammation and airway hyperresponsiveness (conditional recommendation, expert opinion-based). Induced sputum is the preferred option to re-assess airway inflammation (conditional recommendation, expert opinion-based) as a noninvasive validated tool. The GDG strongly recommends joint efforts from academia, industry and healthcare systems to develop both educational tools and resources supporting a wider use of induced sputum evaluation in severe asthma.

If airway eosinophilia is not controlled, the clinician is advised to address the following possibilities:

1. Patient is not adherent to the background controller treatment or to the general management plan installed to ensure optimal asthma control. The GDG considers this as the first step to be




FIGURE 1 Decision tree guiding biological use for patients with uncontrolled T2 severe asthma at the point of care. This algorithm is based on a precise diagnosis of severe asthma uncontrolled under high-intensity treatment after all other measures for reaching asthma control were implemented, as described in the GINA 2020 and the ERS/ATS guidelines. A three-pillar decision tree supports the clinician in reaching the decision to start on a particular biological based on the combination between phenotypic traits, biomarkers and clinically relevant asthma-related end points, including safety (conditional recommendation, expert opinion-based). Cost-related and regulatory aspects should also be considered in reaching the decision to start with a particular biological. Re-evaluation of response should be done after 4–6 mo (conditional recommendation, expert opinion-based). As there is no consensus or validated criteria to define response, the definition of a suboptimal response relies on individualized predefined cut-offs of the selected outcomes established by informed shared decision focused on the patient's goals to control its asthma, in alignment with the principles of personalized treatment (conditional recommendation, expert opinion-based). Cost-related and regulatory aspects should also be considered in establishing the predefined goals. I. For suboptimal response, re-assessment of airway inflammation using induced sputum and airway hyperresponsiveness is recommended (conditional recommendation, expert opinion-based). A. If lung eosinophilia is not controlled, the clinician is advised to address the following possibilities: (1) Patient is not adherent to the background controller treatment or to the general management plan installed to ensure optimal asthma control. The asthma management plan should be re-discussed with the patient. (2) Eosinophilia is not driven by the pathway targeted by the biological used. In this case, consider switching to biologicals targeting a different pathway. (3) Inadequate dosing. In this case, consider switching to a biological targeting the same pathway but with different mechanism of action or route of administration. (4) Development of neutralizing anti-drug antibodies (ADA). The titre, affinity, isotype and epitope mapping are important steps in characterizing ADA. In this case, consider switching to biologicals targeting a different pathway or to a biological targeting the same pathway but with different mechanism of action or route of administration. (5) Other immune dysfunctions such as predominant ILC2 activation, autoimmune mechanisms or IL-5-anti-IL-5 complement activating immune complexes are driving the treatment-resistant lung eosinophilia. As these cases are very rare, referral to a specialized centre after all other causes were excluded is recommended. B. If at re-evaluation for suboptimal response, there is no lung eosinophilia and neutrophilic inflammation is present; the biological should be interrupted and measures addressing non-T2 asthma should be considered (conditional recommendation, expert opinion-based). In case of no inflammation, addressing airway hyperresponsiveness (LABA/LAMA combinations or bifunctional drugs) or airway remodelling (bronchial thermoplasty in selected cases) is recommended (conditional recommendation, expert opinion-based). II. In a patient with good response according to the individualized predefined targets, treatment should be continued, pending on the cost-efficacy evaluation and local regulatory status, while continuously monitoring for efficacy and safety (conditional recommendation, expert opinion-based)

considered in case of suboptimal response. Many patients stop their background controller treatment following the first 3–4 administrations of the biological (Table S2) without consulting with their HCP.¹⁶⁰ In this case, the management plan should be re-discussed with the patient. Adherence could be monitored more closely by medication refills or by using smart inhalers with dose counting. The GDG considers that the shared decision process at the start of the treatment and establishment of common goals following the administration of the biological might mitigate the nonadherence to the overall management plan.

2. Airway eosinophilia is not driven by the pathway targeted by the biological used.^{106,160,161} In this case, consider switching to biologicals targeting a different pathway.

3. Inadequate dosing. In this case, consider switching to a biological targeting the same pathway but with different mechanism of action or route of administration.¹⁶³

4. Development of neutralizing anti-drug antibodies (ADA). Following the ABIRISK consortium recommendations, the titre, affinity, isotype and epitope mapping are important steps in characterizing ADA.¹⁶⁴ In this case, consider switching to biologicals targeting a different pathway or to a biological targeting the same pathway but with different mechanism of action or route of administration.

5. Other immune dysfunctions such as predominant ILC2 activation, autoimmune mechanisms (eg driven by Charcot-Leyden crystals) or IL-5-anti-IL-5 complement activating immune complexes are driving the treatment-resistant lung eosinophilia.^{165–167} As these

cases are very rare, referral to a specialized centre after all other caused were excluded is recommended.

If at re-evaluation for suboptimal response, there is no airway eosinophilia and neutrophilic inflammation is present; the biological should be interrupted and measures addressing non-T2 asthma such as macrolides¹⁶⁸ should be considered (conditional recommendation, expert opinion-based). In cases with no airway inflammation, addressing airway hyperresponsiveness (LABA/LAMA combinations or bifunctional drugs)¹⁷⁰ or airway remodelling (bronchial thermoplasty¹⁷⁰ in selected cases) is recommended (conditional recommendation, expert opinion-based).

For the duration of a biological treatment in a patient with good response according to the individualized predefined targets, the GDG recommends continuing treatment, pending on the cost-efficacy evaluation and local regulatory status, while continuously monitoring for efficacy and safety (conditional recommendation, expert opinion-based). The rationale behind this recommendation is the evidence that upon interruption of the biological, as for any other background controller of asthma, all the beneficial effect is lost. The longer term administration informs as well on the long-term safety profile and might facilitate a disease-modifying effect.

7 | DISCUSSION

7.1 | Relevance of the EAACI Guidelines compared to GINA and ERS/ATS recommendations

The EAACI Guidelines recommendations for the use of currently regulatory approved biologicals in uncontrolled severe asthma are formulated per product and with a careful description of the population where the recommendation is applicable. In comparison with the current guidelines (GINA 2020 and ERS/ATS), the EAACI GDG considered this approach pertinent with the precision medicine framework, where an exact characterization of the target population is advocated. Until real-world evidence accumulates, the only detailed description of the phenotype for each biological are the inclusion/exclusion criteria for the pivotal trials that allowed the regulatory approval. Although not immediately very helpful for the clinician, the GDG considered important to acknowledge in the recommendations the heterogeneity of the patients with severe asthma and tried to be as precise as possible in the description of the phenotype. Another key difference is the formulation of recommendations per product as there is no proven "class effect" of the biologicals even if they target the same pathway (eg IL-5) or the same cell (eosinophils). This aspect is evident in the case of response to biologicals from the same "class" after switching for lack of response.

The National Institute for Health and Care Excellence (NICE) in the United Kingdom suggests re-evaluating patients on mepolizumab-based therapy after 12 months to verify whether the frequency of asthmatic exacerbations has been reduced by at least 50%.¹⁷¹ Similar to GINA 2019 and ERS/ATS guidelines, the clinical decision algorithm proposed by the EAACI Guidelines proposes re-evaluation after 4-6 months. This

arbitrary cut-off was chosen based on the high-cost of these drugs with the assumption that the duration is long enough to select responders from suboptimal response. In addition, as no validated criteria exist for defining optimal response, the EAACI Guidelines advocate for individual predefined targets established by informed shared decision focused on the patient's goals to control their asthma.⁴³

7.2 | Future perspectives: barriers and facilitators

7.2.1 | Precision medicine using multiple or upstream targets

Focusing on antagonizing one cytokine alone (anti-IL-4, anti-IL-13 interventions) was either unsuccessful on major clinical end point or provided a dissociated effect,^{26,63-65,172} with impact mainly on exacerbations and less on lung function or asthma symptoms (anti-IL-5, anti-IgE interventions). This dissociated effect seems to be less pronounced for dupilumab, which binds to IL-4R α and consequently blocks both IL-4 and IL-13 signalling. These observations suggest that only the effective simultaneous blockade of two or more main pathogenic pathways in asthma (IL-5, IL-4/IL-13, IgE) would be more effective in the treatment of severe T2 asthma.²⁸ The combined blockade of the IL-13 and IL-33 pathways leads to a greater inhibition of T2 inflammation over inhibition of either pathway alone.¹⁷³ Similarly, co-blockade of IL-13 and IL-25 attenuated airway hyperreactivity, eosinophil infiltration in the lung and mucus hyperproduction in a mouse model of allergic asthma.¹⁷⁴ A novel dual antagonist anti-TSLP/IL-13 bispecific antibody has entered preclinical testing.¹⁷⁵

Small-molecule drug (SMD)-based therapies represent an active field in pharmaceutical research and development. SMDs expand biologicals' therapeutic targets by reaching the intracellular compartment by delivery as either an oral or topically based formulation, offering both convenience and lower costs.¹⁷⁶ A randomized, double-blind, placebo-controlled, crossover study investigating the efficacy, safety, tolerability and pharmacokinetics of a phosphoinositide 3 kinase (PI3K) δ inhibitor, nemiralisib, in patients with persistent, uncontrolled asthma did not translate into meaningful clinical improvement in spite of local inhibition of PI3K δ . Further studies will investigate its potential efficacy in more specific phenotypes, including those colonized with bacteria or frequent exacerbators.¹⁷⁷ Being pivotal for signalling for multiple asthma-relevant cytokines, including IL-4, IL-5, IL-13 and TSLP, Janus kinase (JAK)/ signal transducer and activator of transcription (STAT) inhibition, especially delivered through the inhaled route, might be a novel intervention strategy for severe asthma.^{178,179}

7.2.2 | Targeted treatment of non-T2 asthma

No targeted therapies exist at present for the non-T2 severe asthma. It remains to be seen whether such an endotype really exists as many data derive from cross-sectional studies or are confounded by the

concomitant use of high-dose ICS or OCS or by the co-existence of infection.¹⁸⁰ Nevertheless, the identification of a non-T2 profile is essential as it excludes treatment with the current biologicals available for severe asthma. The NHLBI PrecISE network is currently exploring therapeutic approaches for non-T2 asthma.

7.2.3 | Multidimensional endotyping

Current understanding of the heterogeneity of severe asthma requires a shift in the methodological approach from investigator-imposed hypothesis-driven clusters to the unbiased approach of data-driven models leading to the discovery of new pathogenetic pathways. At present, large amount of multi-omics, imaging, information from medical devices, Apps and electronic health records data are available and require effective analytic tools. Advanced machine learning methods such as deep learning and platforms for cognitive computing represent the future toolbox for the data-driven analysis of big data.^{31,44,49,50}

7.2.4 | The disease-modifying effect

The “holy Grail” for the use of biologicals in severe asthma is to validate their disease-modifying potential.¹⁸¹ If this proves to be true, we might consider a future potential role for biologicals in mild asthma to prevent the evolution towards severe cases or for the primary prevention of asthma in high-risk individuals. Several pathways for immune modulation in T2-driven inflammatory diseases are described, from trained immunity, epigenetic reprogramming, B and T regulatory cells, to microbiome, and novel metabolic pathways.^{109,182-187} The induction of immune tolerance involves molecular mechanisms of anergy, deletion, suppression, immune privilege and ignorance. Most of the knowledge accumulated of T2 immune modulation followed the application of allergen immunotherapy (AIT) in asthma.¹⁸⁸⁻¹⁹⁰ Currently, none of the biologicals approved for the treatment of severe asthma demonstrated any disease-modifying effect, as the efficacy is lost a few weeks or months after the treatment is stopped. Data from mechanistic studies on biologicals are very scarce and frequently contradictory.

7.2.5 | Long-term safety

In healthy individuals, eosinophils contribute to protective immune responses directed against parasites, viral, bacterial and fungal pathogens, are crucial for the survival of long-lived plasma cells and are critical regulators of local immunity and remodelling/repair in both health and disease.^{191,192} Homeostatic eosinophils present in healthy individuals in various tissues are related to the control of glucose homeostasis, protection against obesity, regulation of mammary gland development, preparation of the uterus for pregnancy and the maintenance of the intestinal homeostasis in collaboration

with the local microbiota.¹⁹³⁻¹⁹⁶ In the lung, homeostatic eosinophils have been shown to suppress T2-driven airway responses.¹⁹⁷

Besides its role against parasites, IgE can exert anti-neoplastic surveillance via mast cell and eosinophil-mediated cytotoxicity or by engaging and re-educating alternatively activated macrophages towards pro-inflammatory phenotypes and by priming all subsets to mediate anti-tumour functions.¹⁹⁸⁻²⁰⁰ IgE deficiency was associated with a higher rate of prior diagnosis of malignancies compared with individuals with high or very high IgE levels.²⁰¹ However, prospective studies are essential to better evaluate the association between IgE levels and risk of cancer.

The T2 cytokines IL-4 and IL-13, which signal through IL-4R α , trigger a specialized macrophage phenotype (M(IL-4)) that promotes control of helminth infection and tissue repair in the lung and in the liver.²⁰²⁻²⁰⁴ IL-4 drives the production of defence collagens, SP-A and C1q, and the expression of their receptor, myosin 18A.²⁰⁴ By controlling complement activation, IL-4 regulates the induction of IL-6, thereby influencing a key pathway involved in regenerating liver cell proliferation and survival.²⁰⁵

Omalizumab and mepolizumab have evidence for long-term safety above 5 years. For all the other biologicals, there is evidence for long-term safety up until 2 years (Table S2). Thus, postmarketing surveillance, especially collected through structured registries such as International Severe Asthma Registry (ISAR) or Severe Asthma Network in Italy (SANI), is of utmost importance.^{206,207}

7.2.6 | Efficacy versus effectiveness in a real-world setting

Several reports show that a considerable proportion of patients with severe asthma remain uncontrolled and are not eligible for any of the available biological treatments.¹⁹⁹ Of note, patient selection for biologicals in real life might not be optimal: many omalizumab users have low or very low adherence rates for ICSs and/or ICS-LABA in the 12 months before omalizumab initiation compared the matched cohort of nonusers.²⁰⁸ In addition, there might be a selection bias as patients prescribed mepolizumab had a different prevalence of certain co-morbidities such as CRSwNP, higher disease burden, higher healthcare resource utilization and costs compared with patients prescribed omalizumab.^{209,210} There is inequity in access to biologicals, as higher likelihood of use was related to middle age, higher income, commercial insurance and access to a specialist.²¹¹

A validated assessment tool is needed to adequately evaluate response to biologicals in real-world settings. The Real-life Effectiveness of Omalizumab Therapy study evaluated The Standardized Measure to Assess Response to Therapy (SMART), a tool designed to define response by physician's subjective assessment of asthma symptoms and control and objective assessment of 6 parameters (exacerbations, steroid bursts, emergency department visits and hospitalizations; lung function; ACT score). True responders are defined as meeting both subjective and objective criteria.²¹²

7.2.7 | Efficacy and safety in the paediatric population with severe asthma

Omalizumab has good evidence for both for the 6- to 11- and the 12- to 17-year-old subgroups. Data on the efficacy and safety of benralizumab, dupilumab and mepolizumab in the 12- to 17-year-old patients with severe asthma subgroup are limited. Reslizumab has no paediatric data reported (Table S2). Data are frequently extrapolated from adult trials, and evidence for long-term use is lacking.²¹³ The development of new drugs for the treatment of paediatric severe asthma proves difficult as: (a) criteria to diagnose severe asthma in children are ambiguous and require extensive step-by-step assessment; (b) there is a limited availability of a very heterogeneous population to enter randomized placebo-controlled trials; (c) the requirements of the Paediatric Investigational Plan (EMA) or Paediatric Study Plan (FDA) are quite stringent. Registries like "Severe Paediatric Asthma Collaborative in Europe" (SPACE), the Severe asthma registry of the German Asthma Net or the Children's Health Foundation Paediatric Asthma Registry might prove of help. Large-scale international consortia evaluating severe paediatric asthma using unbiased methods such as multidimensional endotyping could help to overcome this major unmet need in the field of biologicals for asthma.

7.2.8 | Overall efficacy on asthma and its comorbidities

There is high incidence of T2-driven co-morbidity in severe asthma cases such as chronic rhinosinusitis with nasal polyps (CRSwNP), allergic rhinitis, atopic dermatitis, food allergy, anaphylaxis, allergic conjunctivitis.²¹⁴ A retrospective, observational study that assessed the efficacy of omalizumab in patients with asthma and other concomitant allergic diseases such as rhinosinusitis, atopic dermatitis and allergic broncho-pulmonary aspergillosis showed improvement in symptoms of these allergic diseases.²¹⁵ As all regulatory approved T2 biologicals are systemically bioavailable, improving the overall patient well-being by acting on the nonairway targets should be further explored, both in RCTs and in real-life studies. Currently, dupilumab is approved for asthma, atopic dermatitis and chronic rhinosinusitis with nasal polyps and omalizumab is approved for asthma and chronic urticaria and is being assessed in allergic rhinitis, CRSwNP and food allergies. For the EAACI Guidelines, subgroup analysis for reported co-existing co-morbidities was performed and where possible recommendations were formulated for the use of the biological for severe asthma and associated co-morbidities.

7.2.9 | Impact on small airways disease

Severe asthma almost always involves small airways where inhaled drugs have difficult access.^{216,217} For benralizumab, dupilumab and mepolizumab, there is evidence of improving small airways obstruction.^{77,218,219}

7.3 | Additional major unmet needs and research priorities

The GDG proposed several key areas of interest both for the clinician and for the basic researcher and from the healthcare point of view (Box 13). Unmet needs have been assessed from the perspectives of different stakeholders and in most past, converged.²²⁰

8 | CONCLUSION

The addition of targeted treatment for severe asthma based on phenotyping has proved of real value as is recommended by all contemporary guidelines on the management of severe asthma. This significant change in the management of severe asthma was supported by improved understanding of the contribution of immune-inflammatory mechanisms, followed by a relative fast development of biologicals and small molecules specifically targeting the innate and adaptive immune response. There are several critical points impacting the efficacy of this stratified approach, from the complexity of disease endotypes to the effectiveness in real-world settings. The EAACI Guidelines on the use of biologicals for uncontrolled severe asthma offers a desk reference tool for the healthcare providers, patients, regulators and healthcare systems based on a critical appraisal of the current evidence and a structured approach in formulating recommendations in alignment with the key principles of personalized medicine and implementation science.

ACKNOWLEDGMENTS

The GDG is grateful to all the methodology team from the Iberoamerican Cochrane Center (Biomedical Research Institute Sant Pau) who conducted the systematic reviews for efficacy, safety and economic evidence.

CONFLICT OF INTEREST

IA serves as Associate Editor of Allergy. CA and MA report grants from Allergopharma, Idorsia, Swiss National Science Foundation, Christine Kühne-Center for Allergy Research and Education, European Commission Horizon 2020 Framework Programme, Cure, Novartis Research Institutes, AstraZeneca and SciBase, and CA is also on the Sanofi/Regeneron advisory board. T Casale reports grants and/or personal fees from Genentech, Novartis, Sanofi Regeneron, GSK and Amgen. JC declares grants or personal fees from AZ, Genentech/Roche, Novartis, Optinose, Sanofi, Stallergenes and Teva. JC declares grants or personal fees from AZ, Genentech/Roche, Novartis, Optinose, Sanofi, Stallergenes and Teva. DKC is an Emerging Clinician-Scientist Research Fellow, supported by CAAIF, CSACI and AllerGen NCE Inc SG reports personal fees from AstraZeneca, GSK and Novartis. TE has received grants or other from DBV, Innovation Fund Denmark, Regeneron, the Allergy and Anaphylaxis Program SickKids; serves as associate editor for Allergy and in the local advisory board of ALK. JG reports personal fees from

BOX 13 Gaps in evidence for the use of biologicals in severe asthma and plan to address

Gaps in evidence	Plan to address	Priority
Standardizing the use of biologicals in clinical practice 1. Criteria for responders and suboptimal response (early stopping rules) 2. Switching rules 3. Duration of treatment in responders (late stopping rules) 4. Long-term treatment regimen in responders: longer interval, down-dosing, possibility of stopping treatment during the summer months, switch to strategies like topical application, etc 5. Identification of factors related to failure 6. Efficacy in cases of prior failure to other biologicals 7. Routine measurement of ADA	Prospective trials testing the clinical question followed by validation in independent population	High
Implementation of guidelines for the use of biologicals in clinical practice	In-depth education of HCPs on T2 inflammation recognizing the involvement of both the innate and the adaptive immune systems.	High
Long-term safety data	Well-structured postmarketing surveillance using severe asthma registries	High
Assess the long-term efficacy/disease-modifying effect of biologicals in severe asthma (after treatment cessation)	Identify biomarkers related to the course of asthma ²²¹ Well-designed RCT and real-life studies focusing on long-term efficacy Mechanistic studies at a single-cell level	High
Efficacy and safety data in the paediatric population	RCT and RWE trials/registries focused primarily on the paediatric population	High
Cost-effectiveness of biologicals in severe asthma	Sectoral and generalized cost-effectiveness analysis, including the real-world perspective Long-term perspective as disease-modifying intervention and thereby influence long-term cost	High
Identification of clinically relevant biomarkers in order to select responders to the current available biologicals	Proof of concept studies evaluating patient selection based on biomarkers	High
Impact of multi-morbidities (allergic rhinitis, atopic dermatitis, CRSwNP, food allergy, etc)	Studies evaluating the global effect of biologicals on multi-morbidities	High
Fair accessibility to severe asthma correct diagnosis and optimal targeted treatment	Reorganization of severe asthma care Implementation of the patients' perspective from research to models of care Implementation of management pathways/clinical decision systems	High
Comparison between biologicals	Independent head-to-head comparison between biologicals, ideally with crossover design	High
Alignment of studies (including RWE) with guidance from regulatory bodies.	Work in partnership with regulatory bodies to continuously review trial methodology and outcomes.	Medium
Correlation between biological and clinical response to biologicals	Well-designed RCT, example for personalized medicine	Medium
The impact of age on the short and the long-term effects (efficacy and safety) of treatment with biologics?	Well-designed RCT, example for personalized medicine	Medium
Does "resistance" occur as in antibiotic or anti-cancer therapy and what are the underlying molecular mechanisms?	Well-designed RCT, example for personalized medicine	Medium
Validation of different regimens: shorter or longer intervals ("pulse-wise") rather than as a chronic ("maintenance") therapy (eg to prevent resistance)?	RCTs and real-life studies testing different approaches in terms of dose, duration and route	Medium
Seasonal approach with other biologicals following the model of omalizumab	RCTs and real-life studies	Medium

Regeneron, Ena Therapeutics and MedImmune/AstraZeneca, and stock options from Meissa Vaccines Inc. In addition, JG has pending patents on Methods of Propagating Rhinovirus C in Previously Unsusceptible Cell Lines and on Adapted Rhinovirus C. NH reports funding, honoraria or personal fees from GSK, AstraZeneca, Boehringer Ingelheim, Novartis and Sanofi Genzyme, Regeneron, Genentech, Sunovion and Mylan. RK reports grants from Asthma UK, Food Standards Agency, NIH and other from Aimmune. PN reports grants and/or personal fees from AZ, Novartis, Teva, Sanofi, Roche, Novartis, Merck and Equillium. LOM reports grants from GSK and personal fees from AHL. NGP reports personal fees from Novartis, Nutricia, HAL, Menarini/Faes Farma, Sanofi, Mylan/Meda, Biomay, AstraZeneca, GSK, MSD, Asit Biotech and Boehringer Ingelheim, and grants from Gerolymatos International SA and Capricare. AP has received grants, personal fees, nonfinancial support or other from GlaxoSmithKline, AstraZeneca, Boehringer Ingelheim, Chiesi Farmaceutici, Teva, Mundipharma, Zambon, Novartis, Menarini, Sanofi/Regeneron, Roche, Fondazione Maugeri, Fondazione Chiesi and Edmond Pharma. LP reports grants, personal fees or nonfinancial support from Novartis, AstraZeneca, GSK, Teva, Boehringer Ingelheim, Chiesi, Sanofi, Menarini, Mundipharma, Esteve and ROVI. OP reports grants, personal fees or other from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding BV/HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Lofarma, Biomay, Circassia, ASIT Biotech Tools SA, Laboratorios LETI/LETI Pharma, MEDA Pharma/MYLAN, Anergis SA, Mobile Chamber Experts (a GA2LEN Partner), Indoor Biotechnologies, GSK, Astellas Pharma Global, EUFOREA, ROXALL, Novartis, Sanofi Aventis, Med Update Europe GmbH and streamedup! GmbH. SQ reports personal fees and nonfinancial support from GSK, AZ, Sanofi, Novartis, Mundipharma, Teva and Allergy Therapeutics. JSastre declares personal fees and nonfinancial support from Novartis, GSK, AstraZeneca, Sanofi and Mundipharma. MS reports personal fees or research funding from ASIT Biotech.sa, Allergy Therapeutics, ALK, Regeneron, Merck, Immune Tolerance Network, Leti Laboratorios and Allergopharma. OP received research grants from Immunotek SL and Novartis; received fees for giving scientific lectures from Allergy Therapeutics, Amgen, AstraZeneca, Diater, GSK, Immunotek S.L, Novartis, Sanofi Genzyme and Stallergenes; participated in advisory boards from Novartis and Sanofi Genzyme. MJ reports personal fees from ALK-Abello, Allergopharma, Stallergenes, Anergis, Allergy Therapeutics, Circassia, Leti, Biomay, HAL, AstraZeneca, GSK, Novartis, Teva, Vectura, UCB, Takeda, Roche, Janssen, MedImmune and Chiesi. All other authors have no conflict of interest within the scope of the submitted work.

ORCID

Ioana Agache  <https://orcid.org/0000-0001-7994-364X>
 Cezmi A. Akdis  <https://orcid.org/0000-0001-8020-019X>
 Mubeccel Akdis  <https://orcid.org/0000-0003-0554-9943>
 Thomas Casale  <https://orcid.org/0000-0002-3149-7377>
 Thomas Eiwegger  <https://orcid.org/0000-0002-2914-7829>
 Davide Firinu  <https://orcid.org/0000-0002-5768-391X>

James E. Gern  <https://orcid.org/0000-0002-6667-4708>
 Eckard Hamelmann  <https://orcid.org/0000-0002-2996-8248>
 Rebeca Knibb  <https://orcid.org/0000-0001-5561-0904>
 Parameswaran Nair  <https://orcid.org/0000-0002-1041-9492>
 Liam O'Mahony  <https://orcid.org/0000-0003-4705-3583>
 Nikolaos G. Papadopoulos  <https://orcid.org/0000-0002-4448-3468>
 Hae-Sim Park  <https://orcid.org/0000-0003-2614-0303>
 Luis Pérez de Llano  <https://orcid.org/0000-0003-2652-6847>
 Oliver Pfaar  <https://orcid.org/0000-0003-4374-9639>
 Joaquin Sastre  <https://orcid.org/0000-0003-4689-6837>
 Mohamed Shamji  <https://orcid.org/0000-0003-3425-3463>
 Oscar Palomares  <https://orcid.org/0000-0003-4516-0369>

REFERENCES

1. Global Burden of Disease 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2016;2017(390):1211-1259.
2. https://ginasthma.org/wp-content/uploads/2020/04/GINA-2020-full-report_final_wms.pdf. Accessed April 10, 2020
3. Hekking PP, Wener RR, Amelink M, Zwinderman AH, Bouvy ML, Bel EH. The prevalence of severe refractory asthma. *J Allergy Clin Immunol*. 2015;135:896-902.
4. Corren J, Yawn BP. Advancing the care of severe asthma: differential diagnosis, multidisciplinary management, and patient engagement [published online ahead of print, 2019 Feb 20]. *Am J Med*. 2019. <https://doi.org/10.1016/j.amjmed.2019.02.002>
5. Tay TR, Hew M. Comorbid, "treatable traits" in difficult asthma: current evidence and clinical evaluation. *Allergy*. 2018;73:1369-1382.
6. Bender BG. Sorting out nonadherence and airway inflammation in treatment escalation for severe asthma. *Am J Respir Crit Care Med*. 2019;199:400-402.
7. Lee J, Tay TR, Radhakrishna N, et al. Nonadherence in the era of severe asthma biologics and thermoplasty. *Eur Respir J*. 2018;51:1701836.
8. Settignano RA, Kreindler JL, Chung Y, Tkacz J. Evaluating direct costs and productivity losses of patients with asthma receiving GINA 4/5 therapy in the United States. *Ann Allergy Asthma Immunol*. 2019;123(6):564-572.
9. Ortega H, Hahn B, Tran JN, et al. Disease burden in patients with asthma before initiating biologics: a retrospective cohort database study. *Allergy Asthma Proc*. 2019;40(3):146-153.
10. Chen S, Golam S, Myers J, et al. Systematic literature review of the clinical, humanistic, and economic burden associated with asthma uncontrolled by GINA Steps 4 or 5 treatment. *Curr Med Res Opin*. 2018;34(12):2075-2088.
11. Perez de Llano L, Martinez-Moragon E, Plaza Moral V, et al. Unmet therapeutic goals and potential treatable traits in a population of patients with severe uncontrolled asthma in Spain. ENEAS study. *Respir Med*. 2019;151:49-54.
12. Canonica GW, Colombo GL, Bruno GM, et al. Shadow cost of oral corticosteroids-related adverse events: a pharmaco-economic evaluation applied to real-life data from the Severe Asthma Network in Italy (SANI) registry. *World Allergy Organ J*. 2019;12(1):100007.
13. Barry LE, Sweeney J, O'Neill C, Price D, Heaney LG. The cost of systemic corticosteroid-induced morbidity in severe asthma: a health economic analysis. *Respir Res*. 2017;18:129.

14. Bourdin A, Fabry-Vendrand C, Ostinelli J, et al. the burden of severe asthma in France: a case-control study using a medical claims database. *J Allergy Clin Immunol Pract.* 2019;7(5):1477-1487.
15. Ahmed H, Turner S. Severe asthma in children—a review of definitions, epidemiology, and treatment options in 2019. *Pediatr Pulmonol.* 2019;54(6):778-787.
16. Fleming L, Murray C, Bansal AT, et al. The burden of severe asthma in childhood and adolescence: results from the paediatric U-BIOPRED cohorts. *Eur Respir J.* 2015;46:1322-1333.
17. Selby L, Saglani S. Severe asthma in children: therapeutic considerations. *Curr Opin Allergy Clin Immunol.* 2019;19(2):132-140.
18. McGeachie MJ, Yates KP, Zhou X, et al. Patterns of growth and decline in lung function in persistent childhood asthma. *N Engl J Med.* 2016;374(19):1842-1852.
19. Bui DS, Lodge CJ, Burgess JA, et al. Childhood predictors of lung function trajectories and future COPD risk: a prospective cohort study from the first to the sixth decade of life. *Lancet Respir Med.* 2018;6(7):535-544.
20. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J.* 2014;43:343-373.
21. Holguin F, Cardet JC, Chung KF, et al. Management of severe asthma: a European Respiratory Society/American Thoracic Society guideline. *Eur Respir J.* 2020;55(1):1900588.
22. Difficult-to-treat and Severe Asthma in Adolescent and Adult Patients. Diagnosis and Management. A GINA Pocket Guide for Health Professionals. 2018. <https://ginasthma.org/wp-content/uploads/2018/11/GINA-SA-FINAL-wms.pdf>.
23. Anderson GP. Endotyping asthma: new insights into key pathogenic mechanisms in a complex, heterogeneous disease. *Lancet.* 2008;372(9643):1107-1119.
24. Lötvalld J, Akdis CA, Bacharier LB, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol.* 2011;127(2):355-360.
25. Agache I, Akdis C, Jutel M, Virchow JC. Untangling asthma phenotypes and endotypes. *Allergy.* 2012;67(7):835-846.
26. Agache IO. From phenotypes to endotypes to asthma treatment. *Curr Opin Allergy Clin Immunol.* 2013;13(3):249-256.
27. Agache I, Akdis CA. Endotypes of allergic diseases and asthma: An important step in building blocks for the future of precision medicine. *Allergol Int.* 2016;65(3):243-252.
28. Agache I, Akdis CA. Precision medicine and phenotypes, endotypes, genotypes, regiotypes, and theratypes of allergic diseases. *J Clin Invest.* 2019;130:1493-1503.
29. Gibson PG, McDonald VM. Management of severe asthma: targeting the airways, comorbidities and risk factors. *Intern Med J.* 2017;47(6):623-631.
30. Chung KF, Adcock IM. Precision Medicine for the discovery of treatable mechanisms in severe asthma. *Allergy.* 2019;74(9):1649-1659.
31. Agache I. Severe asthma phenotypes and endotypes. *Semin Immunol.* 2019;46:101301.
32. Agache I, Sugita K, Morita H, et al. The complex type 2 endotype in allergy and asthma: from laboratory to bedside. *Curr Allergy Asthma Rep.* 2015;15(6):29.
33. Liu AH, Babineau DC, Krouse RZ, et al. Pathways through which asthma risk factors contribute to asthma severity in inner-city children. *J Allergy Clin Immunol.* 2016;138(4):1042-1050.
34. Pongracic JA, Krouse RZ, Babineau DC, et al. Distinguishing characteristics of difficult-to-control asthma in inner-city children and adolescents. *J Allergy Clin Immunol.* 2016;138(4):1030-1041.
35. Bossley CJ, Fleming L, Gupta A, et al. Pediatric severe asthma is characterized by eosinophilia and remodeling without T(H)2 cytokines. *J Allergy Clin Immunol.* 2012;129:974-978.
36. Saglani S. Childhood severe asthma: New insights on remodelling and biomarkers. *Paediatr Respir Rev.* 2017;24:11-13.
37. Teague WG, Phillips BR, Fahy JV, et al. Baseline features of the severe asthma research program (SARP III) cohort: differences with age. *J Allergy Clin Immunol Pract.* 2018;6:545-554.
38. Nayeem SS, Bush A, Silveira LP, et al. Clinical and pathological characteristics of severely asthmatic children with persistent airflow limitation. *Thorax.* 2017;72:A45-A46.
39. Andersson CK, Adams A, Nagakumar P, et al. Intraepithelial neutrophils in pediatric severe asthma are associated with better lung function. *J Allergy Clin Immunol.* 2017;139:1819-1829.
40. Saglani S, Lui S, Ullmann N, et al. IL-33 promotes airway remodeling in pediatric patients with severe steroid-resistant asthma. *J Allergy Clin Immunol.* 2013;132:676-685.
41. Nagakumar P, Denney L, Fleming L, et al. Type 2 innate lymphoid cells in induced sputum from children with severe asthma. *J Allergy Clin Immunol.* 2016;137:624-626.
42. Bousquet J, Brusselle G, Buhl R, et al. Care pathways for the selection of a biologic in severe asthma. *Eur Respir J.* 2017;50(6):1701782.
43. Papadopoulos NG, Barnes P, Canonica GW, et al. The evolving algorithm of biological selection in severe asthma *Allergy.* 2020;75: 1555-1563. <https://doi.org/10.1111/all.14256>
44. Seys SF, Quirce S, Agache I, et al. Severe asthma: Entering an era of new concepts and emerging therapies: Highlights of the 4th international severe asthma forum, Madrid, 2018. *Allergy.* 2019;74(11):2244-2248.
45. Agache I, Cojanu C, Laculiceanu A, Rogozea L. Critical Points on the Use of Biologicals in Allergic Diseases and Asthma. *Allergy Asthma Immunol Res.* 2020;12(1):24-41.
46. Diamant Z, Vijverberg S, Alving K, et al. Towards clinically applicable biomarkers for asthma – an EAACI position paper. *Allergy.* 2019;74(10):1835-1851.
47. Agache I, Rogozea L. Asthma biomarkers: do they bring precision medicine closer to the clinic? *Allergy Asthma Immunol Res.* 2017;9(6):466-476.
48. Agache I, Annesi-Maesano I, Bonertz A, et al. Prioritizing research challenges and funding for allergy and asthma and the need for translational research-The European Strategic Forum on Allergic Diseases. *Allergy.* 2019;74(11):2064-2076.
49. Hinks TSC, Brown T, Lau LCK, et al. Multidimensional endotyping in patients with severe asthma reveals inflammatory heterogeneity in matrix metalloproteinases and chitinase 3-like protein 1. *J Allergy Clin Immunol.* 2016;138(1):61-75.
50. Agache I, Strasser DS, Pierlot GM, et al. Monitoring inflammatory heterogeneity with multiple biomarkers for multidimensional endotyping of asthma. *J Allergy Clin Immunol.* 2018;141(1):442-445.
51. Anderson WC 3rd, Szeffler SJ. Cost-effectiveness and comparative effectiveness of biologic therapy for asthma: to biologic or not to biologic? *Ann Allergy Asthma Immunol.* 2019;122(4):367-372.
52. Boyman O, Kaegi C, Akdis M, et al. EAACI IG Biologicals task force paper on the use of biologic agents in allergic disorders. *Allergy.* 2015;70(7):727-754.
53. <https://www.ema.europa.eu/en/medicines/human/EPAR/fasenra>. Accessed December 16, 2019.
54. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2017/761070Orig1s000Approv.pdf. Accessed December 16, 2019.
55. <https://www.ema.europa.eu/en/medicines/human/summaries-opinion/dupixent-0>. Accessed December 16, 2019.
56. https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/761055s007lbl.pdf. Accessed December 16, 2019.
57. <https://www.ema.europa.eu/en/medicines/human/EPAR/nucala>. Accessed December 16, 2019.
58. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761122s000lbl.pdf. Accessed December 16, 2019.
59. <https://www.ema.europa.eu/en/medicines/human/EPAR/xolair>. Accessed December 16, 2019.

60. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/103976s5225lbl.pdf. Accessed December 16, 2019.
61. <https://www.ema.europa.eu/en/medicines/human/EPAR/cinquaero>. Accessed December 16, 2019.
62. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/761033lbl.pdf. Accessed December 16, 2019.
63. Agache I, Beltran J, Akdis C, et al. Efficacy and safety of treatment with biologicals (benralizumab, dupilumab, mepolizumab, omalizumab and reslizumab) for severe eosinophilic asthma. A systematic review for the EAACI Guidelines - recommendations on the use of biologicals in severe asthma. *Allergy*. 2020; May;75(5):1023–1042. <https://doi.org/10.1111/all.14221>
64. Agache I, Rocha C, Beltran J, et al. Efficacy and safety of treatment with biologicals (benralizumab, dupilumab and omalizumab) for severe allergic asthma: A systematic review for the EAACI Guidelines - recommendations on the use of biologicals in severe asthma. *Allergy*. 2020;75:1043–1057. <https://doi.org/10.1111/all.14235>
65. Agache I, Song Y, Rocha C, et al. Efficacy and safety of treatment with dupilumab for severe asthma: A systematic review of the EAACI guidelines-Recommendations on the use of biologicals in severe asthma. *Allergy*. 2020 May;75(5):1058-1068. <https://doi.org/10.1111/all.14268>
66. Walford HH, Doherty TA. Diagnosis and management of eosinophilic asthma: a US perspective. *J Asthma Allergy*. 2014;7:53–65.
67. Santanello NC, Zhang J, Seidenberg B, Reiss TF, Barber BL. What are minimal important changes for asthma measures in a clinical trial? *Eur Respir J*. 1999;14(1):23–27.
68. Jones PW. Interpreting thresholds for a clinically significant change in health status in asthma and COPD. *Eur Respir J*. 2002;19(3):398–404.
69. Juniper EF, Svensson K, Mörk AC, Ståhl E. Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med*. 2005;99(5):553–558.
70. Dweik RA, Boggs PB, Erzurum SC, et al. American Thoracic Society Committee on Interpretation of Exhaled Nitric Oxide Levels (FENO) for Clinical Applications. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *Am J Respir Crit Care Med*. 2011;184(5):602–615.
71. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336(7650):924.
72. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383–394.
73. https://apps.who.int/iris/bitstream/handle/10665/75146/9789241548441_eng.pdf;jsessionid=CA74A1F992AE5574F7B899567C721BC1?sequence=1. Accessed February 2, 2020.
74. Kolbeck R, Kozhich A, Koike M, et al. MEDI-563, a humanized anti-IL-5 receptor alpha mAb with enhanced antibody-dependent cell-mediated cytotoxicity function. *J Allergy Clin Immunol*. 2010;125:1344–1353.
75. Dávila González I, Moreno Benítez F, Quirce S. Benralizumab: a new approach for the treatment of severe eosinophilic asthma. *J Investig Allergol Clin Immunol*. 2019;29(2):84–93.
76. Laviolette M, Gossage DL, Gauvreau G, et al. Effects of benralizumab on airway eosinophils in asthmatic patients with sputum eosinophilia [published correction appears in *J Allergy Clin Immunol*. 2014;133(4): 1232]. *J Allergy Clin Immunol*. 2013;132(5):1086–1096.e5.
77. Panettieri RA Jr, Welte T, Shenoy KV, et al. Onset of effect, changes in airflow obstruction and lung volume, and health-related quality of life improvements with benralizumab for patients with severe eosinophilic asthma: phase IIIb randomized, controlled trial (SOLANA). *J Asthma Allergy*. 2020;13:115–126.
78. Chupp G, Lugogo NL, Kline JN, et al. Rapid onset of effect of benralizumab on morning peak expiratory flow in severe, uncontrolled asthma. *Ann Allergy Asthma Immunol*. 2019;122(5):478–485.
79. Bleecker ER, FitzGerald JM, Chaney P, et al. Efficacy and safety of benralizumab for patients with severe asthma uncontrolled with high-dosage inhaled corticosteroids and long-acting β 2-agonists (SIROCCO): a randomised, multicentre, placebo-controlled phase 3 trial. *Lancet*. 2016;388(10056):2115–2127.
80. FitzGerald JM, Bleecker ER, Nair P, et al. Benralizumab, an anti-interleukin-5 receptor α monoclonal antibody, as add-on treatment for patients with severe, uncontrolled, eosinophilic asthma (CALIMA): a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet*. 2016;388(10056):2128–2141.
81. Nair P, Wenzel S, Rabe KF, et al. Oral glucocorticoid-sparing effect of benralizumab in severe asthma. *N Engl J Med*. 2017;376(25):2448–2458.
82. Palomares Ó, Sánchez-Ramón S, Dávila I, et al. dIvergEnt: how IgE axis contributes to the continuum of allergic asthma and anti-IgE therapies. *Int J Mol Sci*. 2017;18(6):1328.
83. Agache I, Strasser DS, Klenk A, et al. Serum IL-5 and IL-13 consistently serve as the best predictors for the blood eosinophilia phenotype in adult asthmatics. *Allergy*. 2016;71(8):1192–1202.
84. Chipps BE, Newbold P, Hirsch I, Trudo F, Goldman M. Benralizumab efficacy by atopy status and serum immunoglobulin E for patients with severe, uncontrolled asthma. *Ann Allergy Asthma Immunol*. 2018;120(5):504–511.
85. Gour N, Wills-Karp M. IL-4 and IL-13 signaling in allergic airway disease. *Cytokine*. 2015;75:68–78.
86. Marone G, Granata F, Pucino V, et al. The intriguing role of interleukin 13 in the pathophysiology of asthma. *Front Pharmacol*. 2019;10:1387.
87. Alasandagutti ML, Ansari MS, Sagurthi SR, Valluri V, Gaddam S. Role of IL-13 genetic variants in signalling of asthma. *Inflammation*. 2017;40(2):566–577.
88. McDowell PJ, Heaney LG. Different endotypes and phenotypes drive the heterogeneity in severe asthma. *Allergy*. 2020;75(2):302–310.
89. Junttila IS. Tuning the cytokine responses: an update on interleukin (IL)-4 and IL-13 receptor complexes. *Front Immunol*. 2018;9:888.
90. Le Floch A, Allinne J, Nagashima K, et al. Dual blockade of IL-4 and IL-13 with dupilumab, an IL-4R α antibody, is required to broadly inhibit type 2 inflammation. *Allergy*. 2020;75:1188–1204. <https://doi.org/10.1111/all.14151>
91. Matsunaga K, Katoh N, Fujieda S, Izuhara K, Oishi K. Dupilumab: Basic aspects and applications to allergic diseases [published online ahead of print, 2020 Jan 29]. *Allergol Int*. 2020;S1323–8930(20):30008–30013.
92. Wenzel S, Castro M, Corren J, et al. Dupilumab efficacy and safety in adults with uncontrolled persistent asthma despite use of medium-to-high-dose inhaled corticosteroids plus a long-acting β 2 agonist: a randomised double-blind placebo-controlled pivotal phase 2b dose-ranging trial. *Lancet*. 2016;388(10039):31–44.
93. Castro M, Corren J, Pavord ID, et al. Dupilumab efficacy and safety in moderate-to-severe uncontrolled asthma. *N Engl J Med*. 2018;378(26):2486–2496.
94. Rabe KF, Nair P, Brusselle G, et al. Efficacy and safety of dupilumab in glucocorticoid-dependent severe asthma. *N Engl J Med*. 2018;378(26):2475–2485.
95. Bachert C, Han JK, Desrosiers M, et al. Efficacy and safety of dupilumab in patients with severe chronic rhinosinusitis with nasal polyps (LIBERTY NP SINUS-24 and LIBERTY NP SINUS-52): results from two multicentre, randomised, double-blind, placebo-controlled, parallel-group phase 3 trials. *Lancet*. 2019;394(10209):1638–1650.

96. Kabesch M, Schedel M, Carr D, et al. IL-4/IL-13 pathway genetics strongly influence serum IgE levels and childhood asthma. *J Allergy Clin Immunol*. 2006;117(2):269-274.
97. Corren J, Castro M, O'Riordan T, et al. Dupilumab efficacy in patients with uncontrolled, moderate-to-severe allergic asthma. *J Allergy Clin Immunol Pract*. 2020;8(2):516-526.
98. Heeb LEM, Egholm C, Boyman O. Evolution and function of interleukin-4 receptor signaling in adaptive immunity and neutrophils. *Genes Immun*. 2020;21(3):143-149.
99. Kubo M. Innate and adaptive type 2 immunity in lung allergic inflammation. *Immunol Rev*. 2017;278(1):162-172.
100. Emma R, Morjaria JB, Fuochi V, Polosa R, Caruso M. Mepolizumab in the management of severe eosinophilic asthma in adults: current evidence and practical experience. *Ther Adv Respir Dis*. 2018;12:1753466618808490. <https://doi.org/10.1177/1753466618808490>
101. Chupp GL, Bradford ES, Albers FC, et al. Efficacy of mepolizumab add-on therapy on health-related quality of life and markers of asthma control in severe eosinophilic asthma (MUSCA): a randomised, double-blind, placebo-controlled, parallel-group, multicentre, phase 3b trial. *Lancet Respir Med*. 2017;5(5):390-400.
102. Bel EH, Wenzel SE, Thompson PJ, et al. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med*. 2014;371(13):1189-1197.
103. Ortega HG, Liu MC, Pavord ID, et al. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med*. 2014;371(13):1198-1207.
104. Chapman KR, Albers FC, Chipps B, et al. The clinical benefit of mepolizumab replacing omalizumab in uncontrolled severe eosinophilic asthma. *Allergy*. 2019;74(9):1716-1726.
105. Humbert M, Albers FC, Bratton DJ, et al. Effect of mepolizumab in severe eosinophilic asthma according to omalizumab eligibility. *Respir Med*. 2019;154:69-75.
106. Pérez de Llano LA, Dacal Rivas D, Cosío BG. Mepolizumab and reslizumab, two different options for severe asthma patients with prior failure to omalizumab. *Allergy*. 2020;75(4):940-942.
107. Moore WC, Meyers DA, Wenzel SE, et al. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med*. 2010;181(4):315-323.
108. Bachert C, Humbert M, Hanania NA, et al. *Staphylococcus aureus* and its IgE-inducing enterotoxins in asthma: current knowledge [published online ahead of print, 2020 Jan 24]. *Eur Respir J*. 2020;1901592: <https://doi.org/10.1183/13993003.01592-2019>
109. Boonpiyathad T, Sözen ZC, Satitsuksanoa P, Akdis CA. Immunologic mechanisms in asthma. *Semin Immunol*. 2019;46:101333.
110. Teufelberger AR, Nordengrün M, Braun H, et al. The IL-33/ST2 axis is crucial in type 2 airway responses induced by *Staphylococcus aureus*-derived serine protease-like protein D. *J Allergy Clin Immunol*. 2018;141(2):549-559.e7.
111. Chanez P, Contain-Bordes C, Garcia G, et al. Omalizumab-induced decrease of Fc ϵ RI expression in patients with severe allergic asthma. *Respir Med*. 2010;104:1608-1617.
112. Takaku Y, Soma T, Nishihara F, et al. Omalizumab attenuates airway inflammation and interleukin-5 production by mononuclear cells in patients with severe allergic asthma. *Int Arch Allergy Immunol*. 2013;161:S107-S117.
113. Noga O, Hanf G, Brachmann I, et al. Effect of omalizumab treatment on peripheral eosinophil and T-lymphocyte function in patients with allergic asthma. *J Allergy Clin Immunol*. 2006;117(6):1493-1499.
114. Roth M, Tamm M. The effects of omalizumab on IgE-induced cytokine synthesis by asthmatic airway smooth muscle cells. *Ann Allergy Asthma Immunol*. 2010;104:152-160.
115. van Rensen ELJ, Evertse CE, van Schadewijk W, et al. Eosinophils in bronchial mucosa of asthmatics after allergen challenge: effect of anti-IgE treatment. *Allergy*. 2009;64(1):72-80.
116. Djukanović R, Wilson SJ, Kraft M, et al. Effects of treatment with anti-immunoglobulin E antibody omalizumab on airway inflammation in allergic asthma. *Am J Respir Crit Care Med*. 2004;170(6):583-593.
117. Massanari M, Holgate ST, Busse WW, Jimenez P, Kianifard F, Zeldin R. Effect of omalizumab on peripheral blood eosinophilia in allergic asthma. *Respir Med*. 2010;104(2):188-196.
118. Maggi L, Rossetini B, Montaini G, et al. Omalizumab dampens type 2 inflammation in a group of long-term treated asthma patients and detaches IgE from Fc ϵ RI. *Eur J Immunol*. 2018;48(12):2005-2014.
119. Domingo C, Pomares X, Angril N, Rudi N, Amengual MJ, Mirapeix RM. Effectiveness of omalizumab in non-allergic severe asthma. *J Biol Regul Homeost Agents*. 2013;27(1):45-53.
120. de Llano LP, Vennera Mdel C, Álvarez FJ, et al. Effects of omalizumab in non-atopic asthma: results from a Spanish multicenter registry. *J Asthma*. 2013;50(3):296-301.
121. Bourgoin-Heck M, Amat F, Trouvé C, et al. Omalizumab could be effective in children with severe eosinophilic non-allergic asthma. *Pediatr Allergy Immunol*. 2018;29(1):90-99.
122. Busse W, Spector S, Rosén K, Wang Y, Alpan O. High eosinophil count: a potential biomarker for assessing successful omalizumab treatment effects. *J Allergy Clin Immunol*. 2013;132(2):485-486.
123. Hanania NA, Wenzel S, Rosén K, et al. Exploring the effects of omalizumab in allergic asthma: an analysis of biomarkers in the EXTRA study. *Am J Respir Crit Care Med*. 2013;187(8):804-811.
124. Casale TB, Chipps BE, Rosén K, et al. Response to omalizumab using patient enrichment criteria from trials of novel biologics in asthma. *Allergy*. 2018;73(2):490-549.
125. Casale TB, Luskin AT, Busse W, et al. Omalizumab effectiveness by biomarker status in patients with asthma: evidence from PROSPERO, a prospective real-world study. *J Allergy Clin Immunol Pract*. 2019;7(1):156-164.
126. Humbert M, Taillé C, Mala L, et al. Omalizumab effectiveness in patients with severe allergic asthma according to blood eosinophil count: the STELLAIR study. *Eur Respir J*. 2018;51(5):1702523.
127. Lieberman PL, Jones I, Rajwanshi R, Rosén K, Umetsu DT. Anaphylaxis associated with omalizumab administration: Risk factors and patient characteristics. *J Allergy Clin Immunol*. 2017;140(6):1734-1736.e4.
128. Gill MA, Bajwa G, George TA, et al. Counter-regulation between the Fc ϵ RI pathway and antiviral responses in human plasmacytoid dendritic cells. *J Immunol*. 2010;184:5999-6006.
129. Gill MA, Liu AH, Calatroni A, et al. Enhanced plasmacytoid dendritic cell antiviral responses after omalizumab. *J Allergy Clin Immunol*. 2018;141(5):1735-1743.
130. Efthimiou J, Poll C, Barnes PJ. Dual mechanism of action of T2 inhibitor therapies in virally induced exacerbations of asthma: evidence for a beneficial counter-regulation. *Eur Respir J*. 2019;54(1):1802390.
131. Teach SJ, Gill MA, Togias A, et al. Preseasonal treatment with either omalizumab or an inhaled corticosteroid boost to prevent fall asthma exacerbations. *J Allergy Clin Immunol*. 2015;136(6):1476-1485.
132. Busse WW, Morgan WJ, Gergen PJ, et al. Randomized trial of omalizumab (anti-IgE) for asthma in inner-city children. *N Engl J Med*. 2011;364(11):1005-1015.
133. Chipps BE, Figliomeni M, Spector S. Omalizumab: an update on efficacy and safety in moderate-to-severe allergic asthma. *Allergy Asthma Proc*. 2012;33(5):377-385.

134. Hanania NA, Alpan O, Hamilos DL, et al. Omalizumab in severe allergic asthma inadequately controlled with standard therapy: a randomized trial. *Ann Intern Med.* 2011;154(9):573-582.
135. Lanier BQ, Corren J, Lumry W, Liu J, Fowler-Taylor A, Gupta N. Omalizumab is effective in the long-term control of severe allergic asthma. *Ann Allergy Asthma Immunol.* 2003;91(2):154-159.
136. Ayres J, Higgins B, Chilvers ER, Ayre G, Blogg M, Fox H. Efficacy and tolerability of anti-immunoglobulin E therapy with omalizumab in patients with poorly controlled (moderate-to-severe) allergic asthma. *Allergy.* 2004;59(7):701-708.
137. Vignola AM, Humbert M, Bousquet J, et al. Efficacy and tolerability of anti-immunoglobulin E therapy with omalizumab in patients with concomitant allergic asthma and persistent allergic rhinitis: SOLAR. *Allergy.* 2004;59(7):709-717.
138. Humbert M, Beasley R, Ayres J, et al. Benefits of omalizumab as add-on therapy in patients with severe persistent asthma who are inadequately controlled despite best available therapy (GINA 2002 step 4 treatment): INNOVATE. *Allergy.* 2005;60(3):309-316.
139. Lanier B, Bridges T, Kulus M, Taylor AF, Berhane I, Vidaurre CF. Omalizumab for the treatment of exacerbations in children with inadequately controlled allergic (IgE-mediated) asthma. *J Allergy Clin Immunol.* 2009;124(6):1210-1216.
140. Liddament M, Husten J, Estephan T, et al. Higher Binding Affinity and in vitro Potency of Reslizumab for Interleukin-5 Compared With Mepolizumab. *Allergy Asthma Immunol Res.* 2019;11(2):291-298.
141. Matera MG, Rogliani P, Calzetta L, Cazzola M. Pharmacokinetic/pharmacodynamic profile of reslizumab in asthma. *Expert Opin Drug Metab Toxicol.* 2018;14(2):239-245.
142. Virchow JC, Katial R, Brusselle GG, et al. Safety of reslizumab in uncontrolled asthma with eosinophilia: a pooled analysis from 6 trials. *J Allergy Clin Immunol Pract.* 2020;8(2):540-548.
143. Castro M, Mathur S, Hargreave F, et al. Reslizumab for poorly controlled, eosinophilic asthma: a randomized, placebo-controlled study. *Am J Respir Crit Care Med.* 2011;184(10):1125-1132.
144. Castro M, Zangrilli J, Wechsler ME, et al. Reslizumab for inadequately controlled asthma with elevated blood eosinophil counts: results from two multicentre, parallel, double-blind, randomised, placebo-controlled, phase 3 trials. *Lancet Respir Med.* 2015;3(5):355-366.
145. Corren J, Weinstein S, Janka L, Zangrilli J, Garin M. Phase 3 study of reslizumab in patients with poorly controlled asthma: effects across a broad range of eosinophil counts. *Chest.* 2016;150(4):799-810.
146. Bjermer L, Lemiere C, Maspero J, Weiss S, Zangrilli J, Germinaro M. Reslizumab for inadequately controlled asthma with elevated blood eosinophil levels: a randomized phase 3 study. *Chest.* 2016;150(4):789-798.
147. Shaker M, Briggs A, Dbouk A, Dutille E, Oppenheimer J, Greenhawt M. Estimation of health and economic benefits of clinic versus home administration of omalizumab and mepolizumab. *J Allergy Clin Immunol Pract.* 2020;8(2):565-572.
148. Kabata H, Flamar AL, Mahlaköiv T, et al. Targeted deletion of the TSLP receptor reveals cellular mechanisms that promote type 2 airway inflammation [published online ahead of print, 2020 Feb 17]. *Mucosal Immunol.* 2020. <https://doi.org/10.1038/s41385-020-0266-x>
149. Wang W, Li Y, Lv Z, et al. Bronchial allergen challenge of patients with atopic asthma triggers an alarmin (IL-33, TSLP, and IL-25) response in the airways epithelium and submucosa. *J Immunol.* 2018;201(8):2221-2231.
150. Li Y, Wang W, Lv Z, et al. Elevated expression of IL-33 and TSLP in the airways of human asthmatics in vivo: a potential biomarker of severe refractory disease. *J Immunol.* 2018;200(7):2253-2262.
151. Huang YC, Weng CM, Lee MJ, Lin SM, Wang CH, Kuo HP. Endotypes of severe allergic asthma patients who clinically benefit from anti-IgE therapy. *Clin Exp Allergy.* 2019;49(1):44-53.
152. Corren J, Parnes JR, Wang L, et al. Tezepelumab in adults with uncontrolled asthma. *N Engl J Med.* 2017;377(10):936-946.
153. <https://www.fdanews.com/articles/188355-fda-awards-astra-zeneca-and-amgens-tezepelumab-breakthrough-designation>.
154. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2016-001549-13/>.
155. <https://www.sanofi.com/en/media-room/press-releases/2019/2019-06-21-07-00-00>.
156. Lightwood D, Tservistas M, Zehentleitner M, et al. Efficacy of an inhaled IL-13 antibody fragment in a model of chronic asthma. *Am J Respir Crit Care Med.* 2018;198(5):610-619.
157. Harris JM, Maciuga R, Bradley MS, et al. A randomized trial of the efficacy and safety of quilizumab in adults with inadequately controlled allergic asthma. *Respir Res.* 2016;17:29.
158. Wildfire JJ, Gergen PJ, Sorkness CA, et al. Development and validation of the Composite Asthma Severity Index—an outcome measure for use in children and adolescents. *J Allergy Clin Immunol.* 2012;129(3):694-701.
159. Fitzpatrick AM, Szeffler SJ, Mauger DT, et al. Development and initial validation of the Asthma Severity Scoring System (ASSESS). *J Allergy Clin Immunol.* 2020;145(1):127-139.
160. d'Ancona G, Kavanagh J, Roxas C, et al. Adherence to inhaled corticosteroids and clinical outcomes in mepolizumab therapy for severe asthma [published online ahead of print, 2020 Feb 20]. *Eur Respir J.* 2020;1902259.
161. Bagnasco D, Menzella F, Caminati M, et al. Efficacy of mepolizumab in patients with previous omalizumab treatment failure: Real-life observation. *Allergy.* 2019;74(12):2539-2541.
162. Matsuno O, Minamoto S. Eosinophils depletion therapy for severe asthma management following favorable response to mepolizumab. *Respir Med Case Rep.* 2019;28:100899.
163. Mukherjee M, Aleman Paramo F, Kjarsgaard M, et al. Weight-adjusted intravenous reslizumab in severe asthma with inadequate response to fixed-dose subcutaneous mepolizumab. *Am J Respir Crit Care Med.* 2018;197(1):38-46.
164. Rup B, Pallardy M, Sikkema D, et al. Standardizing terms, definitions and concepts for describing and interpreting unwanted immunogenicity of biopharmaceuticals: recommendations of the Innovative Medicines Initiative ABIRISK consortium. *Clin Exp Immunol.* 2015;181(3):385-400.
165. Smith SG, Chen R, Kjarsgaard M, et al. Increased numbers of activated group 2 innate lymphoid cells in the airways of patients with severe asthma and persistent airway eosinophilia. *J Allergy Clin Immunol.* 2016;137(1):75-86.e8.
166. Mukherjee M, Bulir DC, Radford K, et al. Sputum autoantibodies in patients with severe eosinophilic asthma. *J Allergy Clin Immunol.* 2018;141(4):1269-1279.
167. Mukherjee M, Lim HF, Thomas S, et al. Airway autoimmune responses in severe eosinophilic asthma following low-dose Mepolizumab therapy. *Allergy Asthma Clin Immunol.* 2017;13:2.
168. Gibson PG, Yang IA, Upham JW, et al. Effect of azithromycin on asthma exacerbations and quality of life in adults with persistent uncontrolled asthma (AMAZES): a randomised, double-blind, placebo-controlled trial. *Lancet.* 2017;390(10095):659-668.
169. Matera MG, Page CP, Calzetta L, Rogliani P, Cazzola M. Pharmacology and therapeutics of bronchodilators revisited. *Pharmacol Rev.* 2020;72(1):218-252.
170. Bonta PI, Chanez P, Annema JT, Shah PL, Niven R. Bronchial thermoplasty in severe asthma: best practice recommendations from an expert panel. *Respiration.* 2018;95(5):289-300.
171. *Mepolizumab for treating severe refractory eosinophilic asthma. Technology appraisal guidance [TA431].* National Institute for Health and Clinical Excellence. Published: 25/01/2017. <https://www.nice.org.uk/guidance/ta431/chapter/1-recommendations>.

172. Wang FP, Liu T, Lan Z, Li SY, Mao H. Efficacy and safety of anti-interleukin-5 therapy in patients with asthma: a systematic review and meta-analysis. *PLoS One*. 2016;11(11):e0166833.
173. Ramirez-Carrozzi V, Sambandam A, Zhou M, et al. Combined blockade of the IL-13 and IL-33 pathways leads to a greater inhibition of type 2 inflammation over inhibition of either pathway alone. *J Allergy Clin Immunol*. 2017;139(2):705-708.
174. Zhang F-Q, Han X-P, Zhang F, et al. Therapeutic efficacy of a co-blockade of IL-13 and IL-25 on airway inflammation and remodeling in a mouse model of asthma. *Int Immunopharmacol*. 2017;46:133-140.
175. Venkataramani S, Low S, Weigle B, et al. Design and characterization of Zweimab and Doppelmab, high affinity dual antagonistic anti-TSLP/IL13 bispecific antibodies. *Biochem Biophys Res Commun*. 2018;504(1):19-24.
176. Roth-Walter F, Adcock IM, Benito-Villalvilla C, et al. Comparing biologicals and small molecule drug therapies for chronic respiratory diseases: An EAACI Taskforce on Immunopharmacology position paper. *Allergy*. 2019;74(3):432-448.
177. Khindri S, Cahn A, Begg M, et al. A multicentre, randomized, double-blind, placebo-controlled, crossover study to investigate the efficacy, safety, tolerability, and pharmacokinetics of repeat doses of inhaled nemiralisib in adults with persistent. *Uncontrolled Asthma. J Pharmacol Exp Ther*. 2018;367(3):405-413.
178. Calbet M, Ramis I, Calama E, et al. Novel inhaled Pan-JAK inhibitor, LAS194046, reduces allergen-induced airway inflammation, late asthmatic response, and pSTAT activation in brown Norway rats. *J Pharmacol Exp Ther*. 2019;370(2):137-147.
179. Zak M, Hanan EJ, Lupardus P, et al. Discovery of a class of highly potent Janus Kinase 1/2 (JAK1/2) inhibitors demonstrating effective cell-based blockade of IL-13 signaling. *Bioorg Med Chem Lett*. 2019;29(12):1522-1531.
180. Sze E, Bhalla A, Nair P. Mechanisms and therapeutic strategies for non-T2 asthma. *Allergy*. 2020;75(2):311-325.
181. Menzies-Gow A, Bafadhel M, Busse WW, et al. An expert consensus framework for asthma remission as a treatment goal. *J Allergy Clin Immunol*. 2020;145(3):757-765.
182. Netea MG, Joosten LAB, Latz E, et al. Trained immunity: A program of innate immune memory in health and disease. *Science*. 2016;352(6284):aaf1098.
183. van de Veen W, Akdis M. The use of biologics for immune modulation in allergic disease. *J Clin Invest*. 2019;130(4):1452-1462.
184. Lunjani N, Satitsuksanoa P, Lukasik Z, Sokolowska M, Eiwegger T, O'Mahony L. Recent developments and highlights in mechanisms of allergic diseases: Microbiome. *Allergy*. 2018;73(12):2314-2327.
185. Chung KF. Potential role of the lung microbiome in shaping asthma phenotypes. *Ann Am Thorac Soc*. 2017;14(Supplement_5):S326-S331.
186. Michalovich D, Rodriguez-Perez N, Smolinska S, et al. Obesity and disease severity magnify disturbed microbiome-immune interactions in asthma patients. *Nat Commun*. 2019;10(1):5711.
187. Venter C, Meyer RW, Nwaru BI, et al. EAACI position paper: Influence of dietary fatty acids on asthma, food allergy, and atopic dermatitis. *Allergy*. 2019;74(8):1429-1444.
188. Pfaar O, Agache I, Blay F, et al. Perspectives in allergen immunotherapy: 2019 and beyond. *Allergy*. 2019;74(Suppl 108):3-25.
189. Agache I, Lau S, Akdis CA, et al. EAACI Guidelines on Allergen Immunotherapy: House dust mite-driven allergic asthma. *Allergy*. 2019;74(5):855-873.
190. Jutel M, Van de Veen W, Agache I, Azkur KA, Akdis M, Akdis CA. Mechanisms of allergen-specific immunotherapy and novel ways for vaccine development. *Allergol Int*. 2013;62(4):425-433.
191. Ravin KA, Loy M. The eosinophil in infection. *Clin Rev Allergy Immunol*. 2016;50(2):214-227.
192. Jacobsen EA, Helters RA, Lee JJ, Lee NA. The expanding role(s) of eosinophils in health and disease. *Blood*. 2012;120(19):3882-3890.
193. Marichal T, Mesnil C, Bureau F. Homeostatic eosinophils: characteristics and functions. *Front Med*. 2017;4:101.
194. Shah K, Ignacio A, McCoy KD, Harris NL. The emerging roles of eosinophils in mucosal homeostasis. *Mucosal Immunol*. 2020. <https://doi.org/10.1038/s41385-020-0281-y>
195. Wen T, Rothenberg ME. The regulatory function of eosinophils. *Microbiol Spectr*. 2016;4(5).
196. Jiménez-Saiz R, Anipindi VC, Galipeau H, et al. Microbial regulation of enteric eosinophils and its impact on tissue remodeling and Th2 immunity. *Front Immunol*. 2020;11:155.
197. Mesnil C, Raulier S, Paulissen G, et al. Lung-resident eosinophils represent a distinct regulatory eosinophil subset. *J Clin Invest*. 2016;126:3279-3295.
198. Pellizzari G, Hoskin C, Crescioli S, et al. IgE re-programs alternatively-activated human macrophages towards pro-inflammatory anti-tumoural states. *EBioMedicine*. 2019;43:67-81.
199. Jensen-Jarolim E, Bax HJ, Bianchini R, et al. AllergoOncology - the impact of allergy in oncology: EAACI position paper. *Allergy*. 2017;72(6):866-887.
200. Crawford G, Hayes MD, Seoane RC, et al. Epithelial damage and tissue $\gamma\delta$ T cells promote a unique tumor-protective IgE response. *Nat Immunol*. 2018;19(8):859-870.
201. Ferastraoar D, Rosenstreich D. IgE deficiency and prior diagnosis of malignancy: Results of the 2005-2006 National Health and Nutrition Examination Survey. *Ann Allergy Asthma Immunol*. 2018;121(5):613-618.
202. Grecis RK. Immunity to helminths: resistance, regulation, and susceptibility to gastrointestinal nematodes. *Annu Rev Immunol*. 2015;33:201-225.
203. Chen F, Liu Z, Wu W, et al. An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat Med*. 2012;18(2):260-266.
204. Minutti CM, Jackson-Jones LH, García-Fojeda B, et al. Local amplifiers of IL-4R α -mediated macrophage activation promote repair in lung and liver. *Science*. 2017;356(6342):1076-1080.
205. DeAngelis RA, Markiewski MM, Kourtzelis I, et al. A complement-IL-4 regulatory circuit controls liver regeneration. *J Immunol*. 2012;188(2):641-648.
206. ISAR Study Group. International severe asthma registry: mission statement. *Chest*. 2019;157(4):805-814.
207. Senna G, Guerriero M, Paggiaro PL, et al. SANI-Severe Asthma Network in Italy: a way forward to monitor severe asthma. *Clin Mol Allergy*. 2017;15:9.
208. Taillé C, Pison C, Nocent C, et al. Patients in the IDEAL cohort: a snapshot of severe asthma in France. *Rev Mal Respir*. 2019;36(2):179-190.
209. Jeffery MM, Shah ND, Karaca-Mandic P, Ross JS, Rank MA. Trends in omalizumab utilization for asthma: evidence of suboptimal patient selection. *J Allergy Clin Immunol Pract*. 2018;6(5):1568-1577.
210. Llanos JP, Bell CF, Packnett E, et al. Real-world characteristics and disease burden of patients with asthma prior to treatment initiation with mepolizumab or omalizumab: a retrospective cohort database study. *J Asthma Allergy*. 2019;12:43-58.
211. Inselman JW, Jeffery MM, Maddux JT, Shah ND, Rank MA. Trends and disparities in asthma biologic use in the United States. *J Allergy Clin Immunol Pract*. 2020;8(2):549-554.e1.
212. Singh H, Peters JI, Kaur Y, Maselli DJ, Diaz JD. Long-term evaluation of response to omalizumab therapy in real life by a novel multimodal approach: the Real-life Effectiveness of Omalizumab Therapy (REALITY) study. *Ann Allergy Asthma Immunol*. 2019;123(5):476-482.

213. Abrams EM, Becker AB, Szefer SJ. Current state and future of biologic therapies in the treatment of asthma in children. *Pediatr Allergy Immunol Pulmonol*. 2018;31(3):119-131.
214. Akdis CA, Arkwright PD, Brügger MC, et al. Type 2 immunity in the skin and lungs. *Allergy*. 2020 Jul;75(7):1582-1605. <https://doi.org/10.1111/all.1431>
215. Cusack RP, Sahadevan A, Lane SJ. Qualitative effects of omalizumab on concomitant IgE-mediated disease in a severe asthmatic population: a real life observational study. *QJM*. 2016;109(9):601-604.
216. Singhanian A, Rupani H, Jayasekera N, et al. Altered epithelial gene expression in peripheral airways of severe asthma. *PLoS One*. 2017;12(1):e0168680.
217. Gafar F, Boudewijn IM, Cox CA, et al. Predictors of clinical response to extrafine and non-extrafine particle inhaled corticosteroids in smokers and ex-smokers with asthma. *Respir Res*. 2018;19(1):256.
218. Castro M, Rabe KF, Corren J, et al. Dupilumab improves lung function in patients with uncontrolled, moderate-to-severe asthma. *ERJ Open Res*. 2020;6(1):00204-2019.
219. Sposato B, Camiciottoli G, Bacci E, et al. Mepolizumab effectiveness on small airway obstruction, corticosteroid sparing and maintenance therapy step-down in real life. *Pulm Pharmacol Ther*. 2020;61:101899.
220. Mathioudakis AG, Custovic A, Deschildre A, et al. Research priorities in pediatric asthma: results of a global survey of multiple stakeholder groups by the Pediatric Asthma in Real Life (PeARL) think tank. *J Allergy Clin Immunol Pract*. 2020;8(6):1953-1960.e9
221. Szefer S. Asthma across the lifespan: time for a paradigm shift. *J Allergy Clin Immunol*. 2018;142:773-780.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Agache I, Akdis C, Akdis M, et al. EAACI Biologicals Guidelines—Recommendations for severe asthma. *Allergy*. 2021;76:14–44. <https://doi.org/10.1111/all.14425>