

<https://helda.helsinki.fi>

Heterogeneous genetic basis of age at maturity in salmonid fishes

Waters, Charles D.

2021-03

Waters , C D , Clemento , A , Aykanat , T , Garza , J C , Naish , K A , Narum , S & Primmer , C R 2021 , ' Heterogeneous genetic basis of age at maturity in salmonid fishes ' , Molecular Ecology , vol. 30 , no. 6 , pp. 1435-1456 . <https://doi.org/10.1111/mec.15822>

<http://hdl.handle.net/10138/339510>

<https://doi.org/10.1111/mec.15822>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

DR. CHARLES D. WATERS (Orcid ID : 0000-0003-4606-3202)

DR. ANTHONY CLEMENTO (Orcid ID : 0000-0003-3598-7151)

DR. JOHN CARLOS GARZA (Orcid ID : 0000-0002-7325-6803)

DR. KERRY-ANN A NAISH (Orcid ID : 0000-0002-3275-8778)

PROF. CRAIG PRIMMER (Orcid ID : 0000-0002-3687-8435)

Article type : Original Article

Heterogeneous genetic basis of age at maturity in salmonid fishes

Charles D. Waters^{1,7,8}, Anthony Clemento^{2,3}, Tutku Aykanat⁴, John Carlos Garza^{2,3*}, Kerry A. Naish^{1*}, Shawn Narum^{5*}, Craig R. Primmer^{4,6*}

¹School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA, USA

²Institute of Marine Sciences, University of California, Santa Cruz, CA, USA

³Santa Cruz Laboratory, Southwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Santa Cruz, CA, USA

⁴Organismal and Evolutionary Biology Research Programme, University of Helsinki, Finland

⁵Hagerman Genetics Laboratory, Columbia River Inter-Tribal Fish Commission, Hagerman, ID, USA

⁶Institute of Biotechnology, University of Helsinki, Finland

⁷Present address: Auke Bay Laboratories, Alaska Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, Juneau, AK, USA

⁸Corresponding author

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/MEC.15822](https://doi.org/10.1111/MEC.15822)

This article is protected by copyright. All rights reserved

* Shared senior authorship

Emails: Charlie.Waters@noaa.gov; Anthony.Clemento@noaa.gov; tutku.aykanat@helsinki.fi;
Carlos.Garza@noaa.gov; knaish@uw.edu; nars@critfc.org; craig.primmer@helsinki.fi

Abstract

Understanding the genetic basis of repeated evolution of the same phenotype across taxa is a fundamental aim in evolutionary biology and has applications in conservation and management. However, the extent to which interspecific life-history trait polymorphisms share evolutionary pathways remains under-explored. We address this gap by studying the genetic basis of a key life-history trait, age at maturity, in four species of Pacific salmonids (genus *Oncorhynchus*) that exhibit intra- and interspecific variation in this trait – Chinook Salmon, Coho Salmon, Sockeye Salmon, and Steelhead Trout. We tested for associations in all four species between age at maturity and two genome regions, *six6* and *vgl3*, that are strongly associated with the same trait in Atlantic Salmon (*Salmo salar*). We also conducted a genome-wide association analysis in Steelhead to assess whether

additional regions were associated with this trait. We found the genetic basis of age at maturity to be heterogeneous across salmonid species. Significant associations between *six6* and age at maturity were observed in two of the four species, Sockeye and Steelhead, with the association in Steelhead being particularly strong in both sexes ($p = 4.46 \times 10^{-9}$ after adjusting for genomic inflation). However, no significant associations were detected between age at maturity and the *vgl3* genome region in any of the species, despite its strong association with the same trait in Atlantic Salmon. We discuss possible explanations for the heterogeneous nature of the genetic architecture of this key life-history trait, as well as the implications of our findings for conservation and management.

Keywords

age at maturity, heterogeneous genetic basis, life-history trait polymorphisms, salmonid, sexual conflict

Introduction

The repeated evolution of the same phenotype across taxa raises fundamental questions on whether evolution has acted on ancestral polymorphisms that are shared between species or on different loci with similar outcomes (Elmer & Meyer, 2011). Well characterized examples of parallel evolution within individual species, such as adaptation to freshwater in three-spined stickleback fish, reveal the role of both mechanisms (Jones et al., 2012; Liu, Ferchaud, Gronkjaer, Nygaard, & Hansen, 2018). In some cases, polymorphic traits that influence fitness are shared across taxa. Three main but non-exclusive pathways for the retention of polymorphic traits during speciation events have been identified (Jamie & Meier, 2020): evolution on standing ancestral variation (Cortez et al., 2014), introgression (Giska et al., 2019), and novel mutations (Mundy, 2005). However, the extent to which interspecific trait polymorphisms share evolutionary pathways remains relatively uncharacterized (Jamie & Meier, 2020). This knowledge can inform how evolution may act to maintain such variation within new (i.e. daughter) species (Ayala & Campbell, 1974; Charlesworth, Nordborg, & Charlesworth, 1997; Guerrero & Hahn, 2017), how genetic architecture may predict the retention of such polymorphisms, and how related species may adapt to shared selection regimes (Jamie & Meier, 2020).

The age at which an individual reaches sexual maturity is a polymorphic life-history trait in many species that can have significant fitness consequences. For example, maturing later provides more opportunities to grow, potentially conferring larger body size and higher energy reserves at reproduction, which in turn can enable higher fecundity and increased offspring survival (Stearns, 1992). However, delayed maturity also increases the risk of mortality before reproduction and translates into longer generation times (Charnov & Gillooly, 2004; Kozlowski, 1992; Roff, Mostow, & Fairbairn, 2002; Stearns, 1992). The large variation in size and age at maturity observed within and among populations of many species indicates that there is often no single, optimal maturation strategy, and considerable research effort has focused on determining the ecological factors and evolutionary mechanisms leading to the maintenance of such variation across different species (Flatt & Heyland, 2011; Stearns, 1989). Yet, the genetic basis of variation underlying different maturation strategies across taxa and species has remained poorly understood until recently (Perry et al., 2014; Roff, 2011).

Fishes display more diversity in reproductive traits than any other vertebrate group (Mank & Avise, 2006). Within fishes, species in the family *Salmonidae* exhibit a wide range of this diversity (Schaffer, 2004), and the potential trade-offs between age at maturity (and therefore size) and survival are well established, particularly in anadromous species and populations (e.g. Fleming, 1996; Hankin, Nicholas, & Downey, 1993; Healey, 1991). As growth rate is high during the marine migratory phase, variation in age may result in dramatic size differences, with the size of individuals potentially doubling with each additional year spent at sea. Larger, older individuals of both sexes have been shown to have higher reproductive success once on the spawning grounds (Fleming, 1996; Janowitz-Koch et al., 2019; Mobley et al., 2019), but they also have a higher probability of mortality during the marine migration phase (Czorlich, Aykanat, Erkinaro, Orell, & Primmer, 2018), thus representing a classic evolutionary trade-off.

In addition to its fundamental importance to ecology and evolution, understanding the drivers of variation in size and age at maturity has applied relevance in fisheries management (Conover & Munch, 2002), aquaculture (Taranger et al., 2010), and conservation (Kindsvater, Mangel, Reynolds, & Dulvy, 2016). For example, shared genetic architectures that are well characterized in one taxon may inform the conservation of evolutionary processes in related species, and selective breeding

Accepted Article

programs (e.g. aquaculture operations) in new species may take advantage of trait-linked markers previously identified in other species. Variation in age at maturity is particularly relevant to the conservation and management of salmonids, one of the most socio-economically important family of fishes. First, variation in breeding age buffers populations from diversity loss following catastrophic environmental events via the ‘portfolio effect’ (Greene, Hall, Guilbault, & Quinn, 2010; Satterthwaite, Carlson, & Criss, 2017; Schindler et al., 2010). Second, such life-history variation reduces inter-annual variability in adult returns and the frequency of fishery closures, which benefits commercial, recreational, and subsistence fisheries and associated communities (Brown & Godduhn, 2015; Copeland, Ackerman, Wright, & Byrne, 2017; Greene et al., 2010; Schindler et al., 2010). Third, as age at maturity is positively correlated with size and larger individuals may be preferentially targeted by fisheries, fisheries induced evolution can result in reductions in population variation and therefore viability. Combined, these factors highlight why knowledge of age structure, as well as the genetic basis of age at maturity, is important for modeling the effects of fisheries harvest and developing appropriate management strategies aimed at maintaining population diversity in fitness-related traits (Bowersox, Corsi, McCormick, Copeland, & Campbell, 2019; Hankin & Healey, 1986; Larsen et al., 2019; Ricker, 1980). Such knowledge can further allow development of more precise monitoring tools for assessing the evolutionary consequences of anthropogenic effects (e.g. Czorlich et al. 2018).

There is a pressing need to understand the mechanisms that drive variation in age at maturity in salmonids due to the trait’s ecological, evolutionary, and economic importance, particularly as the age structures of many populations have been changing in recent decades (Cline, Ohlberger, & Schindler, 2019; DeFilippo et al., 2019; Lewis, Grant, Brenner, & Hamazaki, 2015; Ohlberger, Ward, Schindler, & Lewis, 2018). Age at maturity has been shown to have high heritability in multiple species ($h^2 = 0.21-0.57$; Carlson & Seamons, 2008; Gjerde, 1984; Hankin et al., 1993), suggesting significant genetic variation for this trait that may promote local adaptation (Vähä, Erkinaro, Niemela, & Primmer, 2008). The trait has also been shown to differ considerably between the sexes, with females maturing, on average, later than males (Fleming, 1996; Schaffer, 2004).

Recently, there have been significant advances in understanding the genetic basis of age at maturity in one salmonid species, Atlantic Salmon (*Salmo salar*), with a particular focus on the

number of years spent in the marine migration phase (i.e. sea age). Several studies have identified a large effect locus on chromosome Ssa25, with variation near a strong candidate gene, *vgll3*, explaining almost 40% of the variation in sea age at maturity in wild-caught individuals from over 50 European populations, including several divergent lineages (Ayllon et al., 2015; Barson et al., 2015). Subsequent studies have also identified associations between the same genomic region and age for this species (Ayllon et al., 2019; Christensen, Gutierrez, Lubieniecki, & Davidson, 2017). Further, Barson et al. (2015) identified a second region on chromosome Ssa09 encompassing the candidate gene *six6* that was initially highly significant in a genome-wide association (GWA) analysis, although this signal was no longer statistically significant following correction for population structure. However, a recent study in an aquaculture population of Atlantic Salmon identified strong associations with early maturation and both the chromosome Ssa09 (*six6*) and chromosome Ssa25 (*vgll3*) regions (Sinclair-Waters et al., 2020), suggesting that the population stratification correction in Barson et al. (2015) may have been overly conservative. Interestingly, GWA studies of North American-origin Atlantic Salmon aquaculture stocks have failed to find strong associations between sea age and variants in these regions (Boulding, Ang, Elliott, Powell, & Schaeffer, 2019; Mohamed et al., 2019), possibly due to a lack of variation in these regions (Sinclair-Waters et al., 2020). Given the associations identified in Atlantic Salmon and that both *six6* and *vgll3* have also been linked with pubertal timing in humans (e.g. Cousminer et al., 2013; Cousminer, Widen, & Palmert, 2016; Perry et al., 2014) and other mammals (Cánovas et al., 2014), these same loci may play a role in the maintenance of variation in age at maturity phenotypes in other salmonid species.

Pacific salmonids include numerous species in North America (genus *Oncorhynchus*), most of which exhibit anadromy (Behnke, 2002), and six of which show variation in the number of years spent in the marine environment prior to reproduction. Four species – Chinook Salmon (*O. tshawytscha*), Coho Salmon (*O. kisutch*), Sockeye Salmon (*O. nerka*), and Steelhead Trout (*O. mykiss*, the anadromous form of Rainbow Trout) – have been the focus of intense management and research. As a result, individual-level life-history data and tissue samples for genetic analyses are available for multiple populations in each of these species, making them a promising target for genotype-phenotype association studies. Chinook Salmon and Steelhead Trout exhibit the greatest variation in age at maturity (typically two to six years), Sockeye Salmon commonly mature at three to

five years of age, and Coho Salmon between two and four years (Quinn, 2005), although populations can exhibit different age structures due to genetic (McKinney et al., 2019) and environmental factors such as local rearing and growing conditions (Hankin et al., 1993; Harstad, Larsen, & Beckman, 2014; Harstad et al., 2018; Taylor, 1990). Pacific salmonids therefore provide an excellent opportunity to further assess the influence of *six6* and *vgl3* on age at maturity and improve our understanding of the extent to which genetic architectures are shared across species.

Here, we used a targeted sequencing approach to test for associations between the *six6* and *vgl3* genome regions and age at maturity in four Pacific salmonid species native to North America – Chinook Salmon, Coho Salmon, Sockeye Salmon, and Steelhead Trout - to determine whether these loci are associated with variation in age at maturity across related species. Samples were collected from two to six populations (hatchery and wild) of varying phylogenetic, phenotypic, and geographic backgrounds per species, and SNPs were identified in the *six6* and *vgl3* genes. Associations between these SNPs and age at maturity were then quantified using logistic and cumulative proportional odds models that accounted for potential confounding effects. For Steelhead Trout, we also tested for associations in other genomic regions with whole genome resequencing of groups with differing sea age to determine whether additional regions were associated with this trait in this species. The results improve our understanding of the adaptive genetic variation underlying this key trait across multiple species and provide additional tools to monitor age structure in salmonid populations of commercial, ecological, and evolutionary importance.

Materials and Methods

Sample collections

Populations of Chinook Salmon, Coho Salmon, Sockeye Salmon, and Steelhead Trout (hereafter referred to as Chinook, Coho, Sockeye, and Steelhead) for which individual-based collections of genetic samples and ages at maturity existed were identified for this study. Two to six populations, ranging from California to Alaska, USA, were sampled per species and included those of hatchery and wild origin and variable age structures (Table 1, Tables S1-S4). Genomic DNA from most samples had been previously isolated. When needed, genomic DNA was extracted from tissue samples using DNeasy Blood & Tissue kits (Qiagen, Valencia, CA, USA) following the animal tissue

protocol. Ages had been previously determined from coded wire and passive integrated transponder tags, growth rings on scales, molecular pedigree analyses, or a combination of these methods (Table 1). Ages of Chinook, Coho, and Sockeye correspond to total age at maturity since these species are semelparous. Ages of Steelhead correspond to age at first reproduction since this species is iteroparous. Individual-based data for time spent in freshwater and at sea was not available for all populations, so total age at maturity was examined. There were five Steelhead samples from the Eel River population that matured at age six. Due to the low sample size, these individuals were combined with the age-five fish for association analyses. When possible, existing molecular pedigrees were utilized to ensure the sampling of unrelated individuals, as relatedness may bias results of association analyses (Korte & Farlow, 2013). Detailed population descriptions can be found in the Supplemental Information.

Primer development for six6 and vgl3

Sequences for the *six6* and *vgl3* genes in Atlantic Salmon were obtained from SalmoBase (v. 1, Samy et al., 2017) and aligned to the Coho Salmon (GenBank: GCA_002021735.1) and Rainbow Trout (GCF_002163495.1; Pearse et al., 2019) genomes using *Bowtie2* (v. 2.3.1, Langmead & Salzberg, 2012). The Chinook and Sockeye Salmon genomes were not explicitly considered during primer development, as they had not yet been released at the time of development. Local alignments, rather than end-to-end, were conducted in *Bowtie2* to account for potential differences in gene sequences between species, and up to ten alignments were reported (k=10) so that multiple alignments, including those to homeologous regions, could be investigated. Next, SAM output files from *Bowtie2* were converted to sorted, indexed BAM files using *samtools* (v. 1.3.1, Li et al., 2009). Alignments of the *six6* and *vgl3* genes to the Coho Salmon and Rainbow Trout genomes were then visualized using the Integrated Genome Browser tool (Freese, Norris, & Loraine, 2016). The preferred alignment of each gene was determined by the length and mapping quality (i.e. number of mismatches and indels) of the reported alignments, as well as *a priori* knowledge of chromosomal homologies between species obtained from comparative mapping (Kodama, Briec, Devlin, Hard, & Naish, 2014; Pearse et al., 2019).

Regions where a portion of the best alignments of the *six6* and *vgl3* Atlantic Salmon gene sequences matched, or nearly matched, the Rainbow Trout and Coho Salmon genomes were candidates for primer development. For each gene, six candidate regions spanning approximately 580-780 base pairs (bp) were identified. Next, *Primer3* (Koressaar & Remm, 2007; Untergasser et al., 2012) was used to design PCR primers within 50-100 bp of the sequence tails; minimum, maximum, and optimal primer lengths were set to 21, 25, and 23 bp, respectively. The T7 (5' TAATACGACTCACTATAGGG 3') and T3 (5' ATTAACCCTCACTAAAG 3') universal promoter sequences were then added to all forward and reverse primers, respectively.

PCR amplification

The six candidate regions for each gene were amplified using PCR in a subset of samples ($n=8$ per species) to evaluate amplification success, including possible targeting of gene duplications. PCR reactions were performed in a total volume of 15 μ L, each comprising 1.5 μ L of genomic DNA, 7.5 μ L of 2 \times Qiagen Multiplex PCR Master Mix, 0.3 μ L each of the newly designed forward and reverse primers at 10 μ M concentration, and 5.4 μ L of pure water. The PCR protocol included: initial denaturation at 95°C for 15 minutes, ten cycles of [95°C for 30 seconds (s), 58°C for 90 s, 72°C for 90 s], 25 cycles of [95°C for 30 s, 62°C for 90 s, 72°C for 90 s], and a final extension at 72°C for five minutes. PCR products were cleaned using the ExoSAP-IT PCR Product Cleanup kit (Applied Biosystems) following the manufacturer's protocol and then visualized using agarose gel electrophoresis. Primers that yielded single, high-quality bands of PCR product that matched the expected fragment lengths were deemed suitable for screening in a larger number of samples for single nucleotide polymorphism (SNP) identification.

SNP identification and sample screening

Gene regions were amplified in 16 Chinook, 16 Coho, 16 Sockeye, and 16 Steelhead of varying age classes to identify SNPs. For this screening, 10 μ L PCR reactions were performed, each comprising 1 μ L of genomic DNA, 5 μ L of 2 \times Qiagen Multiplex PCR Master Mix, 0.2 μ L each of 10 μ M forward and reverse primers, and 3.6 μ L of pure water. The PCR protocol and PCR product

cleaning were the same as in the initial amplification. Sanger sequencing (Macrogen, Inc.) was then used to sequence the cleaned PCR products.

Geneious (v. 10.1.3; <https://www.geneious.com>) was used to align sequences for each gene region and species using default parameters. SNPs were identified from the alignments, and we focused on SNPs with intermediate frequencies to maximize variability and the potential to detect genotype-phenotype associations. For Chinook, Coho, and Steelhead, consensus sequences were extracted for regions that contained SNPs; these sequences were then utilized for genotyping in two ways. First, a subset of populations - Cle Elum Chinook, Big Beef Creek Coho, and Forks Creek Steelhead - were genotyped using Sanger sequencing, and alignments were scored in *Geneious*. This initial data was then used to develop primers for high-throughput screening of all study samples using amplicon sequencing methods (Baetscher, Clemento, Ng, Anderson, & Garza, 2018; Campbell, Harmon, & Narum, 2015). All sequences of the *six6* and *vgll3* gene regions obtained from high-throughput screening were aligned and scored as in Baetscher et al. (2018). All Sockeye samples were genotyped directly from Sanger sequences rather than amplicons due to relatively small sample sizes.

SNPs were retained for final analyses if the minor allele frequency was greater than 10%, and individuals were included if they were genotyped successfully at one or more of the SNPs identified within the two genes. When *six6* or *vgll3* contained multiple, highly correlated SNPs (Pearson's $r > 0.95$) that passed filtering, only one "target" SNP was tested. However, when possible, missing genotypes for the target SNP were inferred using genotypes of the linked SNP(s). Steelhead samples from the Eel River, Forks Creek, and Warm Springs populations were the only ones genotyped for the *vgll3* SNPs, as there was not sufficient material remaining to genotype the Dworshak, Pahsimeroi, and Wallowa populations.

We also examined 10 SNPs from the *six6* genome region in Steelhead that had been independently identified using a Pool-seq approach, described in full below (Schlötterer, Tobler, Kofler, & Nolte, 2014). Primers were developed with *Primer3* based on sequences surrounding each target SNP identified from the Pool-seq data. Primers were tested for amplification success in a multiplex reaction using the GT-seq protocol (Campbell et al., 2015). A total of 10 target SNPs were successfully developed (Table S13B) and produced reliable results for further genotyping following standard methods from Campbell et al. (2015). These 10 SNPs included six markers upstream of the

six6 gene on chromosome Omy25 that were presumably in regulatory regions from existing genome annotation, three markers in the *six6* gene sequence, and one SNP downstream of the *six6* gene.

Lastly, all samples were genotyped at suites of putatively neutral SNPs (i.e. presumably not influenced by selection, n=84-242 per species) that had been previously developed for other research and monitoring purposes (Table 1; Tables S14-S17). The putatively neutral loci served as “controls” during association testing, as they were expected to have little or no association with age at maturity, and were also used to account for the possible confounding effects of population structure and relatedness in the association analyses.

Association testing

Associations between age at maturity and SNPs within the *six6* and *vgll3* genome regions were tested using logistic regression and cumulative proportional odds models in R using the *stats* and *ordinal* packages, respectively (Christensen, 2019; R Core Team, 2019). Logistic regression models were employed for Coho and Sockeye since age at maturity was a binary trait for these species (Coho: ages two and three; Sockeye: ages four and five; Table 1). Cumulative proportional odds models were used for Chinook and Steelhead, as age at maturity in these species was an ordinal trait (Chinook and Steelhead: ages two to six; Table 1).

Age at maturity was modeled as a function of SNP, sex, and their interaction. To account for population stratification, a principal component analysis was conducted for each species using putatively neutral loci and the R package *adegenet* (v. 2.1.3, Jombart, 2008; Jombart & Ahmed, 2011). The PCs that reflected population structure (PC1 for Coho and Sockeye; PCs 1-4 for Chinook and Steelhead; Figures S1-S18), as well as their interactions with the SNP, were then included as covariates in the models. Relatedness between samples could not be explicitly included in the models. However, molecular pedigrees were utilized during study design to maximize the sampling of unrelated individuals for most populations (Table 1). Molecular estimates of pairwise relatedness between samples within each population were also generated to determine whether relatedness might have influenced the results (Supplemental Information; Figures S19-S33).

The general form of each full model for Coho and Sockeye was

$$y = \beta_0 + \beta_1 \text{SNP} + \beta_2 \text{Sex} + \beta_3 \text{PC1} + \beta_4 \text{SNP} * \text{Sex} + \beta_5 \text{SNP} * \text{PC1} + \varepsilon ,$$

where y is the binary age at maturity (0 = younger age class, 1 = older age class), SNP is the genotype, Sex is the sex of the individual, and PC1 is the coordinate of the individual on principal component 1. Two full models were constructed for each species, one where the target SNP effect was modelled as additive (i.e. genotype treated as numeric, A/A=1, A/T=2, T/T=3) and a second where the SNP effect was non-additive (i.e. genotype treated as categorical instead of numeric). Models for Coho were evaluated using only males, as females were nearly invariant for age, and thus did not include sex as a covariate. The logistic regression models for Coho and Sockeye estimated the probability of maturing at the younger or older age class for each genotype. Here, the probability of maturing at the older age class was considered to be the probability of delaying maturation.

The general form of each full model for Chinook and Steelhead was

$$y = \beta_0 + \beta_1\text{SNP} + \beta_2\text{Sex} + \beta_3\text{PC1} + \beta_4\text{PC2} + \beta_5\text{PC3} + \beta_6\text{PC4} + \beta_7\text{SNP*Sex} + \beta_8\text{SNP*PC1} + \beta_9\text{SNP*PC2} + \beta_{10}\text{SNP*PC3} + \beta_{11}\text{SNP*PC4} + \varepsilon ,$$

where y is the standardized age at maturity (i.e. youngest age class of each population = 1; Table 1; Table S5), SNP is the genotype, Sex is the sex of the individual, and PCs 1-4 are the coordinates of individuals along the first four principal components. Total age and a binomial age classification were also tested (Tables S5-S6). Four full models were constructed for these species, where the target SNP effect was modelled as additive or non-additive, and, within these models, the threshold structure was either constant or allowed to vary by sex since maturation thresholds may differ between the sexes. The cumulative proportional odds models estimated the probability of maturing at or below a certain age class for each genotype (e.g. probability of maturing at or before age three, probability of maturing at or before age 4, and so forth). For all species, interactions between the SNP and other covariates were included in the full models to determine if SNP effects varied by Sex or location on the PCs (i.e. population). Additional interactions (e.g. Sex*PC1) were not included, as SNP effects were of primary interest.

Backward model selection was performed on all full models within a species (additive versus non-additive for all species; constant versus flexible threshold structure by sex for Chinook and Steelhead) using the *dredge* function of the *MumIn* package (Barton, 2016), and a preferred model was identified using corrected Akaike Information Criterion (AICc) scores. The significance of the SNP effect was then quantified by comparing the likelihood of the preferred model that included

terms with the SNP to the likelihood of a model without the SNP effects. When AICc favored the non-additive SNP model, SNP effects were subsequently partitioned into additive and non-additive components and quantified using the same model comparison approach.

The significance of SNP effects may have been spuriously inflated by cryptic relatedness and residual population structure (Price et al., 2006) despite attempts to sample unrelated individuals and the incorporation of PCs into the models. To monitor and account for such genomic inflation, all polymorphic, putatively neutral loci (n=84 to 242 loci per species, Table 1) were evaluated using the same parsimonious model structures as the *six6* and *vgll3* SNPs. The significance of the *six6* and *vgll3* SNPs were then ranked among those obtained from the putatively neutral SNPs. Genomic inflation factors (i.e. lambda) were also estimated by dividing the median of the observed chi-square values by the median of the expected chi-square distribution of the same degrees of freedom (Devlin & Roeder, 1999; Marchini, Cardon, Phillips, & Donnelly, 2004). Chi-square test statistics of the *six6* and *vgll3* SNPs were then divided by the inflation factors, which were then used to obtain “corrected” *p*-values. It should be noted, however, that lambda values may be overly conservative due to the relatively few number of loci used to estimate them.

Whole genome resequencing collections, library preparation, and sequencing

In order to assess whether other genome regions may harbor loci associated with age at maturity, whole genome resequencing was conducted in Steelhead. Specifically, whole genome resequencing of pooled sample collections was completed with six pools of returning adult male Steelhead that exhibited variable ocean age at maturation phenotypes and originated from three different locations (Table S9A). Individual fish for each pooled sample were collected at a fish trap at Bonneville Dam on the Columbia River (on the border between Oregon and Washington, USA), and the origin of their specific hatchery was identified through parentage analyses (see Hess et al., 2016 for details). Ocean age at maturation was determined by examination of scale annuli collected from each fish at Bonneville Dam, and total age was determined from parentage assignment results. Samples were grouped by ocean age at maturity phenotypes (one or two years in the ocean, or 1- and 2-ocean) within each population, with an average sample size of 48 male fish per pool (Table S9A). DNA was extracted individually with Chelex beads.

Phenotype pools within each population were prepared and sequenced using a standardized Pool-seq protocol described in Micheletti, Hess, Zendt, and Narum (2018). This pooled approach provides estimated allele frequencies for variants across the genome but not individual genotypes. Library preparation included normalizing individual DNA quantity using picogreen fluorescence on a Tecan M200 (Tecan Trading, AG, Switzerland). To ensure a similar contribution of each individual to a pool, individual DNA concentrations were not allowed to deviate more than 20% of the average DNA concentration of all individuals. For a given population, samples were fragmented with NEBNext Ultra dsDNA Fragmentase (New England Biolabs, Ipswich, MA, USA), pooled together, and filtered using Minelute purification (Qiagen, Venlo, Netherlands). Fragment end repair was performed with NEBNext Ultra End Prep (New England Biolabs, Ipswich, MA, USA), and fragments between 400-500 bp were selected for purification using a 25× AMPure beads solution (Beckman Coulter Inc, Indianapolis, IN). Fragments were then amplified with a NEBNext Ultra Q5 PCR protocol, cleaned with AMPure beads, and finally quantified with SYBR quantitative PCR (Thermofisher Scientific, Waltham, MA, USA). Amplified fragments were normalized and then sequenced with high-output runs on an Illumina NextSeq 500 (Illumina, San Diego, CA, USA) with paired-end 150bp reads (2×150bp).

Whole genome resequencing bioinformatics and statistical analyses

For each Pool-seq library, raw 150 bp paired-end reads (2×150 bp) were processed using the PoolParty pipeline (Micheletti & Narum, 2018) that integrates several existing resources into a single pipeline. Briefly, this included multiple steps that started with trimming reads (to a minimum of 50 bp) with a quality score less than 20 using the trim-fastq.pl script part of Popoolation2 (Kofler, Pandey, & Schlotterer, 2011). Trimmed reads were then aligned to the *O. mykiss* genome assembly (GCF_002163495.1) using *bwa-mem* (Li, 2013) with default parameters. PCR duplicates were identified and removed using *SAMblaster* (Faust & Hall, 2014). SAMtools view module (Li, 2011) was used to sort BAM files, which were then combined using the SAMtools mpileup module that extracts SNP and coverage information for each pool. Best practices were implemented following Kofler et al. (2011) to filter out SNPs within 5 bp of indel regions from further analysis and those with minor allele frequency < 0.05. Variant positions were kept with a minimum of 15× depth of coverage

and a maximum of 250× depth of coverage, which was intended to filter out regions that may be paralogs (and thus have high coverage) or regions that are likely represented by a small number of individuals (low coverage).

Filtered allele frequency data were then used to calculate fixation index (F_{ST}) between the collections comprising different ocean ages at maturity (1- or 2-ocean) within each population with a sliding window of 8,000 bases with a step size of 100 bases. Genomic regions with statistically significant differentiation were determined using a local score technique that accounts for linkage disequilibrium of SNPs (Fariello et al., 2017). These local score analyses iteratively determined the tuning parameter (ξ) based on mean $\log_{10} p$ -values for each comparison, and significant regions were displayed (Bonferroni corrected $\alpha = 0.05$) in the form of Manhattan plots using the R package *qqman* (Turner, 2014).

We also implemented a Cochran–Mantel–Haenszel (CMH) test that computes significance between groups of interest (Kofler et al., 2011); in this case, between pairs of 1-ocean and 2-ocean males from each of the three hatchery locations.

Genomic regions associated with differentiation in ocean age were deemed significant if they were shared between both local score analyses (analogous to a Bonferroni corrected $\alpha = 0.05$) and the CMH test (Bonferroni corrected $\alpha = 0.05$). These regions were then investigated for variant annotations using *SnpEff* (Cingolani et al., 2012) with *.gff files available for the genome assembly for Rainbow Trout (GCF_002163495.1).

Results

Primer development for six6 and vgl3 and PCR amplification

The preferred alignments of the Atlantic Salmon *six6* and *vgl3* gene sequences to the Coho Salmon and Rainbow Trout genomes were located on the chromosomes predicted by comparative mapping (Tables S10, S12). Gene sequences also aligned to other chromosomes, including homeologous chromosomes (e.g. Co26 in Coho and Omy22 in Rainbow Trout for *vgl3*), but the alignments were shorter and had lower quality (i.e. more mismatches and indels; Table S10). Six candidate regions for PCR amplification and SNP discovery were identified throughout each gene based on these alignments (Table S11A-B), and most regions amplified successfully.

SNP identification and sample screening

Three regions from *six6* and *vgll3* were then screened in a larger number of samples (n=16 of varying ages per species) to identify SNPs and to generate consensus sequences (Table S11C-D) for further SNP development using amplicon sequencing methods. After genotyping and filtering for minor allele frequency, one to two target SNPs were identified for association analyses within each gene for Chinook, Coho, and Sockeye (Tables S1-S4, S13A). Five SNPs were identified within the *vgll3* gene for Steelhead, and there were 10 SNPs within the *six6* gene identified from the Pool-seq approach (Table S13B). For species-gene combinations with multiple target SNPs, we only present results for the SNP with the strongest association; results for other SNPs are in Table S6. Samples were also genotyped at putatively neutral loci (n=84 to 242 loci per species, Table 1; Tables S1-S4, S14-S17).

Association testing in species with two age classes

Age at maturity in male Coho Salmon was not significantly associated with *six6* or *vgll3* (Table 2; Figure 1; Table S7). In contrast, *six6* was significantly associated with age in Sockeye Salmon, before and after correcting for potential genomic inflation ($p=0.008$, Table 2; Figure 1; Table S7). When partitioned, the non-additive SNP effect was significant while the additive effect was marginally non-significant (Table 2). The probability of delaying maturation (i.e. mature at age five rather than at age four) was 0.34 and 0.53 for the two alternative homozygous genotypes in female Sockeye, while those for male Sockeye were 0.48 and 0.66, respectively (Figure 1). That is, female and male Sockeye Salmon with the C/C genotype were 1.56 and 1.38 times more likely to mature at age five than individuals with the A/A genotype (Figure 1; Table S8A). There was no significant difference in maturation probability between the two sexes. The association of *six6* with age at maturity in Sockeye was greater than any of the 172 putatively neutral SNPs (Table 2).

Vgll3 was also significantly associated with age at maturity in Sockeye Salmon, although the effect became non-significant after accounting for genomic inflation (Table 2; Table S7). Further, the probabilities of delaying maturation were not significantly different between the two homozygous

genotypes (Figure 1; Table S8A). Eleven neutral SNPs exhibited stronger associations with age than *vgll3* in Sockeye Salmon.

Association testing in species with four age classes

Both *six6* and *vgll3* were significantly associated with standardized age at maturity in Chinook Salmon (Table 2; Figure 2; Table S7). However, the SNP effects became non-significant after accounting for potential genomic inflation. The cumulative probabilities of maturation were not significantly different between the two homozygous genotypes in both gene regions, although the maturation probabilities of male Chinook with the T/T genotype appeared to be lower, albeit non-significant, for the younger age classes than the G/G genotype in *vgll3* (Figure 2; Table S8B). In addition, the *p*-values for 10 and 22 putatively neutral loci were smaller (i.e. more significant) than the *six6* and *vgll3* genes, respectively. The specific effects of *vgll3* in Chinook Salmon may vary by population, as suggested by the interactions of the SNP with PCs 2-4. The influence of *vgll3* decreased as the location of individuals along PC2, PC3, and PC4 increased (Table S7). However, there was no apparent geographic gradient in population separation along PCs 2-4 (Figures S2-S5), so extrapolating these effects to other populations may be difficult. The overall effects of *six6* and *vgll3* did not differ significantly between the sexes of Chinook Salmon.

Six6 was highly associated with age at maturity in Steelhead Trout ($p = 4.46 \times 10^{-9}$ after adjusting for genomic inflation), with additive effects being greater than non-additive effects (Table 2). The probabilities of maturing at younger ages were higher for the G/G genotype than the A/A genotype (Figure 3). Specifically, female Steelhead with the G/G genotype were 2.53 and 1.34 times more likely to mature by age one and by age two, respectively, than fish with A/A genotype; male Steelhead with G/G were 2.96 and 1.34 times more likely to mature by age one and by age two than those with A/A (Figure 3; Table S8B). While the maturation thresholds varied by sex, the effect of *six6* on age did not differ significantly between females and males (Table S7). There was a significant interaction between *six6* and PC2, which separated the Forks Creek population from all others (Figure S17); the probability of delayed maturation increased slightly as the position along PC2 increased (Table S7). The other nine SNPs within *six6* were also significantly associated with age at maturity

(Table S6), and all 10 *six6* SNPs exhibited stronger associations with age at maturity than the 242 putatively neutral SNPs tested (Table 2).

Similar to Sockeye Salmon, *vgll3* was significantly associated with age in Steelhead Trout, but the effect became non-significant after correcting for genomic inflation (Table 2). The cumulative maturation probabilities of the three *vgll3* SNP genotypes were very similar (Figure 3; Table S8B), indicating that age was not strongly correlated with variation at *vgll3*. Last, 47 of the 242 neutral SNPs exhibited stronger associations with age than *vgll3* (Table 2).

Whole genome resequencing in Steelhead Trout

Whole genome resequencing provided a means of assessing whether other genome regions may harbor loci associated with age at maturity. After filtering, the six male libraries representing paired collections (one versus two years in the ocean) from three populations had sequence coverage greater than 15× for a range of 48.6-67.0% of the genome (mean 62.2%; Table S9A). The mean depth of coverage for the six libraries was 23.0× with a range of 20.2-25.6×. There were a total of 2,850,964 SNPs identified across all six libraries, and allele frequencies were estimated for each variant in each of the six collections.

Tests for differences in allele frequencies of each paired collection of males (one versus two years in the ocean) within each of the three populations revealed several SNP markers on chromosome Omy25 that were significantly different for each pair based on F_{ST} and Fisher's exact tests. Additionally, CMH tests that compared pools of different ages at maturity for each of the three population pairs identified a region on chromosome Omy25 as consistently significant (Table S9B). Finally, local score results that account for multiple SNPs in physical linkage demonstrated that the signal on chromosome Omy25 remained as the strongest signal ($p < 0.001$ after Bonferroni correction; Figure 4; Table S9B). However, two regions on chromosomes Omy15 and Omy26 were also identified from local score results but with much lower levels of significance (Table S9B). The highly significant region on chromosome Omy25 included markers that were located within and upstream of *six6*, with a total of 216 SNPs between positions 61.27-61.32Mb (Figure 4).

Discussion

We examined whether two genes known to influence age at maturity in Atlantic Salmon (genus *Salmo*), *six6* and *vgll3*, contribute to the phenotypic variation in this trait within four species of Pacific salmonid fishes (genus *Oncorhynchus*). We also examined the broader genetic architecture of this trait in one species, Steelhead Trout, to determine the relative contribution of these genes to phenotypic variation compared to other loci. Two to six populations representing different phylogenetic, phenotypic, and geographic backgrounds were sampled per species. We found significant associations between *six6* and age at maturity in both Sockeye Salmon and Steelhead Trout after correcting for genomic inflation, with the association in Steelhead being particularly strong. Indeed, the odds of maturing by certain ages in this species differed by factors of 1.34 to 2.96 between the two alternative homozygotes, and those in Sockeye differed by factors of 1.38 to 1.56. In contrast, while *vgll3* has been found to be strongly associated with age at maturity in Atlantic Salmon, no significant associations were detected in the four Pacific salmonid species following correction for genomic inflation. Our examination of the relative contribution of these loci in Steelhead using whole genome resequencing also revealed the role of additional loci on separate chromosomes, although the significance of the associations were lower than *six6*. We sought to expand our understanding of the extent to which the genetic architecture of ecologically and evolutionarily important polymorphic traits may be conserved across species radiations. Combined, our results indicate that there is a heterogeneous genetic basis for age at maturity in salmonid fishes, including shared evolutionary pathways for age at maturity involving polymorphisms in the *six6* genomic region in at least two Pacific salmonid species, but no clear evidence for *vgll3* being linked with this trait in the genus *Oncorhynchus*.

The presence of significant associations in both Pacific and Atlantic salmonids suggests that the influence of *six6* on age at maturity has been at least partially conserved over 21 million years of evolution (Lien et al., 2016). Given that significant associations were observed between *six6* and age at maturity in two populations of Sockeye and six populations of Steelhead, this study provides evidence that this genome region harbors broad, ecologically significant variation in salmonid fishes. The *six6* region has frequently been identified as an outlier in numerous evolutionary genetic studies in Atlantic Salmon (summarized in Prichard et al. 2018) and has been associated with landscape factors related to flow rate (Pritchard et al., 2018) as well as the run-timing life-history trait

(Cauwelier, Gilbey, Sampayo, Stradmeyer, & Middlemas, 2018). Further, a RAD locus associated with spawning site selection in Sockeye Salmon (Veale & Russello, 2017) was identified to be located very close to the *six6* genome region (Pritchard et al., 2018). However, associations between *six6* and age at maturity have been inconclusive until recently. Early studies in Atlantic Salmon suggested that strong associations were inflated, or even entirely driven, by population structure (Barson et al., 2015; Johnston et al., 2014). Yet, a recent study detected a strong association between *six6* and early maturation (one year post-smoltification, also known as ‘grilsing’) in >11,000 male Atlantic Salmon from a single aquaculture strain, where population stratification is unlikely to be an issue (Sinclair-Waters et al., 2020). Another recent study identified a strong association between *six6* and age and size at first maturation for inland lineage Steelhead (Willis et al., 2020). The results reported here strengthen the evidence that this locus is relevant to polymorphisms in age at maturity in additional salmonid species.

The molecular functions of *six6* uncovered thus far in vertebrates provide insight into the range of potential functional mechanisms that could link genotype and phenotype in salmonids. The sine oculis homeobox (SIX) protein family consists of evolutionarily conserved transcription factors found in organisms from flies to vertebrates (reviewed by Kumar, 2009). Specific functions of *six6* include playing a role in eye development (Seo, Drivenes, Ellingsen, & Fjose, 1998) and circadian timing in both mammals and fish (Clark et al., 2013; Watanabe et al., 2012). *Six6* has also been identified as a key fertility regulator in mammals (Larder, Clark, Miller, & Mellon, 2011) and has been linked with human pubertal timing (Hou et al., 2017; Perry et al., 2014). Further, *six6* has reported roles in the establishment of the brain-pituitary-gonad (BPG) axis in vertebrates (Jean, Bernier, & Gruss, 1999; Seo et al., 1998). Gene expression studies in Atlantic Salmon support the known roles of *six6* in BPG axis signaling in the hypothalamus and pituitary, as well as eye development, and also suggest links with the Hippo signaling pathway (Kurko et al., 2020). The Hippo pathway is known to regulate cell fate commitment and organ (including gonadal) growth (Kjaerner-Semb et al., 2018; Li et al., 2015). Interestingly, *vgll3* is also known to be involved in the Hippo pathway (Kjaerner-Semb et al., 2018; Kurko et al., 2020), suggesting that genes in this pathway may be relevant to several ecologically important traits in salmonid fishes. However, linking genes of this pathway more directly with the ecological traits of interest awaits further study.

The *vgll3* gene has been linked to age at maturity in Atlantic Salmon in a number of independent studies (Ayllon et al., 2015; Ayllon et al., 2019; Barson et al., 2015; Debes et al., 2019; Verta et al., 2019), as well as with pubertal timing in humans and other mammals (e.g. Cánovas et al., 2014; Cousminer et al., 2013; Cousminer et al., 2016; Perry et al., 2014). The conserved nature of this gene region-phenotype association led us to hypothesize that the same region may be associated with age at maturity in Pacific salmonids. However, our results did not provide conclusive evidence to support this hypothesis. Although *vgll3* genotypes were non-randomly associated with age at maturity in Chinook, Sockeye, and Steelhead, the associations became non-significant after correcting for genomic inflation. Additionally, in each species, a number of randomly chosen SNPs across the genome exhibited even stronger associations with age, suggesting that any effect of *vgll3* is, at best, minor in the Pacific salmonid populations assessed in this study.

There are several possible biological explanations for the heterogeneous results observed between the four Pacific salmonid species analyzed here, as well as between Pacific and Atlantic salmonids. First, the age at maturity trait may have evolved a different genetic architecture in some or all of the species and populations investigated. For example, sample populations of Chinook Salmon and Steelhead Trout spanned large geographic and environmental gradients (Table 1) and represented very diverse lineages (Nielsen, 1999; Waples, Teel, Myers, & Marshall, 2004). Lineage divergence in interior and coastal populations of Chinook and Steelhead reflects past glaciation events that may have resulted in the evolution of different architectures underlying the trait. Further, our sample populations of Chinook Salmon and Steelhead Trout varied in their age at outmigration (e.g. ocean-type and stream-type Chinook) and adult migration timing (e.g. Spring-run and Winter-run Chinook, Summer-run and Winter-run Steelhead; see population descriptions in Supplemental Information). Variability in these other, potentially-correlated life-history traits may have affected the underlying genetic basis of age at maturity, which could be additionally influenced by possible differences in genotype by environment interactions at trait-associated loci. Indeed, population-specific effects of loci associated with age at maturity have been recently documented in Atlantic Salmon (Boulding et al., 2019) and Chinook Salmon (McKinney et al., 2019), as well as for loci associated with adult migration timing in Steelhead Trout (Willis et al., 2020). However, a more complete understanding of differences in genetic architectures between populations, as well as potential genotype by

environment interactions, can only be addressed with surveys of populations across the species' ranges that encompass all life-history strategies.

Different rearing environments may have also affected variation in age at maturity and its underlying genetic architecture, similar to the differences in lineages and other life-history traits described above. Specifically, the populations of Chinook and Steelhead sampled for this study included those of hatchery and wild origin. Four and five hatchery populations were sampled for Chinook and Steelhead, respectively, while one wild population was sampled for each species (Table 1). Hatchery practices may inadvertently lead to younger ages at maturity (Hankin, Fitzgibbons, & Chen, 2009; Thorpe, 2004), particularly for males (Harstad et al., 2014; Larsen et al., 2013). In addition to affecting the phenotype, different selection regimes and environmental conditions experienced in hatcheries might also influence the genetic architecture underlying age at maturity (McKinney, Nichols, & Ford, 2020). To explore to possible effects of origin on the associations of *six6* and *vgll3* with age, we ran the analyses for Chinook and Steelhead with the populations separated by origin (hatchery or wild; results summarized in Table S18). When only hatchery-origin populations were included, the overall findings for Chinook and Steelhead remained unchanged when compared to those when all populations were analyzed together. That is, *six6* was strongly associated with age at maturity in hatchery-origin Steelhead while all other associations were not significant after correcting for possible genomic inflation (Table S18). When the one wild population was analyzed for each species, two of the four results changed. Specifically, *vgll3* was significantly associated with age in the one wild population of Chinook (adjusted $p = 0.009$), while *six6* was not significantly associated with age in the one wild population of Steelhead (adjusted $p = 0.065$). The associations of *six6* in Chinook and *vgll3* in Steelhead remained unchanged. The different results may be a consequence of origin, or they may reflect population-specific effects as described above. The results could also be influenced by the reduced sample sizes for the wild origin models. A future study that samples multiple populations of each origin could better address the question regarding hatchery rearing and its potential effects on age at maturity and its genetic architecture.

Another possible biological explanation for the heterogeneous results is that this study quantified associations between *six6* and *vgll3* and total age at maturity rather than sea age, which was the focal trait in Atlantic Salmon (Ayllon et al., 2015; Barson et al., 2015), because freshwater and

sea age data were not available for every study population. If either gene strictly influences sea age rather than total age, then the strength of associations detected here could have been weakened by variation in freshwater age, particularly since Chinook, Sockeye, and Steelhead are all known to exhibit variation in their freshwater residency time.

Differences in the degree to which age at maturity is under sexual conflict and how it is resolved may have also contributed to the mixed results across species. Age at maturity is under sexual conflict in many salmonid species, whereby the optimal maturation age differs for males and females. One common evolutionary solution to resolve sexual conflict in other taxa is for genes controlling traits subjected to sexual conflict to reside on the sex chromosomes, which more readily allows for sex-specific expression patterns to evolve (Mank, 2017). Although they have a sex-determining locus (*sdY*; Yano et al., 2013), salmon do not have morphologically differentiated sex chromosomes, and therefore other evolutionary solutions are needed. In Atlantic Salmon, the sex-dependent dominance observed at the *vgl3* locus has been suggested to contribute to the partial resolution of sexual conflict (Barson et al., 2015). In Chinook Salmon, it was recently reported that age at maturity in males is associated with male-specific haplotypes from a region on Ots17 (McKinney et al., 2020; McKinney et al., 2019). While this region may include the sex determining locus *sdY*, McKinney et al. (2020) also showed that the sex determination region can be translocated between two chromosomes, Ots17 and Ots18. Age at maturity loci have also been linked to the sex chromosome in Coho Salmon, the sister species to Chinook Salmon, in QTL mapping studies (Kodama, Hard, & Naish, 2018). Linkage of a large effect gene or suite of haplotypes controlling age at maturity with the sex determining region may be an effective strategy for intra-locus sexual conflict resolution, but the specific gene(s) involved are still unknown. Further, it remains to be determined whether similar mechanisms may occur in additional populations and species of salmonids and if other factors, such as selection acting on correlated traits, may also affect the degree of sexual conflict and its resolution.

A final possible biological explanation is that only one of the four species studied here, Steelhead Trout, exhibits similar age at maturity characteristics compared to Atlantic Salmon, as both species are iteroparous and exhibit more than two maturation ages. Interestingly, Steelhead Trout was also the species where the strongest single locus association with age at maturity was observed, albeit

at *six6*. Yet, *six6* was also significant in Sockeye Salmon, a species not known to be iteroparous (Quinn, 2005), and this association was detected in two populations compared to the larger number of populations surveyed here for Steelhead. The age structure in Coho Salmon was the simplest within species studied here – only the males were polymorphic for this trait, and there were only two age classes. However, a broader range of species and populations needs to be studied before it can be concluded whether or not certain life-history characteristics (e.g. iteroparity) may be important in determining the genetic architecture of age at maturity.

There may also be technical explanations for the heterogeneous results we observed that are associated with study design. Specifically, we relied on populations with extensive, individual-based phenotypic and genetic collections. As such, Coho and Sockeye Salmon were represented by only two populations each, while Chinook Salmon and Steelhead Trout were each represented by five to six populations. The power to detect associations may have been reduced in species with few populations, although we note that a significant association with *six6* was detected in Sockeye Salmon. Additionally, environmental and life-history variation across populations (e.g. hatchery or wild origin, inland or coastal lineage, Summer or Winter run) may have inhibited our ability to detect genotype-phenotype associations, and the effects of such “noise” might vary between *six6* and *vgll3* if the genes exhibit different genotype by environment interactions. Efforts to reduce the influence of such population-level differences were implemented here by incorporating multiple principal components – the dominant axes of genetic variation between populations – in the models, a widely employed method in association analyses (Price et al., 2006). Ages were also standardized across populations, where the youngest age class in each population (all of which spent only one year in the ocean) was assigned a value of one (see Table 1 for summary of total and standardized ages), to account for inherent differences in age structures that may be caused by geography, lineage, rearing environment (hatchery or wild), or a combination of these and other factors. Yet, the effects of population-level differences may not have been entirely eliminated. Additional genetic markers to better estimate population differences in the principal components analyses (discussed below), as well as more population replicates for each environmental (e.g. hatchery or wild) and life-history (e.g. Summer or Winter run) variant, could help address this issue in a future study.

Differences in results between species might also be associated with the genotyping. Specifically, surveys were not conducted over the entire genome. Instead, we used a relatively small number of putatively neutral SNPs for estimating genome inflation and correcting for population structure, which may have influenced the statistical power of the analyses. Similarly, relatedness could not be explicitly incorporated into the models. However, we took advantage of known pedigrees where possible to select unrelated individuals, and molecular estimates of pairwise relatedness suggest that fish within populations were predominately unrelated. The fact that a full genome-wide analysis was conducted in only one of the four species means that additional loci that influence age at maturity were not taken into account in three species, although it appears that *six6* has had a significant role in explaining the divergence between phenotypes in Steelhead compared to the other two loci detected on chromosomes Omy15 and Omy26.

Despite these caveats, the findings presented here provide intriguing insight into the evolution of a key trait that is polymorphic across Atlantic and Pacific salmonids. While the genetic basis of age at maturity appears to be heterogeneous both between and within salmonid genera, our results suggest that phenotypic polymorphisms associated with *six6* may represent shared evolutionary processes across the two genera, although at this stage we cannot identify the exact mechanisms involved (Jamie & Meier, 2020). It is interesting to examine the relationships within Pacific salmonids within this context (Crete-Lafreniere, Weir, & Bernatchez, 2012; Macqueen & Johnston, 2014). Steelhead is the most divergent (i.e. outgroup) of the species studied here. This means that the common ancestor for Steelhead Trout and Sockeye Salmon, the two species that share the trait association with *six6*, predates the divergence of all four species, and so the role of the locus has persisted throughout their radiation. If *six6* is indeed not associated with age at maturity in the remaining species examined, then it implies that alternative pathways have evolved. Broadly, the evolution of quantitative traits depends on the relative role of large and small effect loci, their interactions, and the nature of selection acting upon them (Barton & Keightley, 2002; Kardos & Luikart, 2020; Oomen, Kuparinen, & Hutchings, 2020). For example, simulation studies have suggested that the evolution of large effect genes may only be favored under conditions where there is strong selection toward different (local) optima combined with moderate levels of gene flow (Yeaman & Whitlock, 2011). Rates of straying (i.e. gene flow) in salmonids vary between species, populations, and even life-history types, with those of Coho

and Sockeye Salmon being, on average, lower than Atlantic Salmon, Chinook Salmon, and Steelhead Trout (Keefer & Caudill, 2014). Varying levels of gene flow may therefore have played a role in the differential evolution of the trait and the strength of genetic associations observed between species in this study. It has also been shown that the molecular basis of ecologically relevant traits is heterogeneous and does not always involve conserved mechanisms (Kronforst et al., 2012). As a transcription co-factor, *vgl3* expression likely influences a number of other genes (Kurko et al., 2020; Simon, Faucheux, Zider, Theze, & Thiebaud, 2016), and thus it is possible that variation in a different gene or genes in the same pathway could be more directly linked with age at maturity in some of the species investigated here. Indeed, *six6* may also be such a gene.

The discovery of genetic variation underlying ecologically important phenotypic traits across species has significant potential to inform their conservation and management, because architectures that predict evolution in one might be applied to the other. Age at maturity has been declining in many species of Pacific salmonids over recent decades (Bowersox et al., 2019; Cline et al., 2019; Lewis et al., 2015; Ohlberger et al., 2018), leading to lower reproductive potential of females (Ohlberger et al., 2020) and decreased population resilience through reduced phenotypic diversity (Greene et al., 2010; Schindler et al., 2010). Although *six6* is a candidate gene significantly associated with age at maturity in Sockeye and Steelhead, markers developed from this study need to be validated more broadly to determine population and sex-specific effects across the geographic distribution of each species, and the relative contributions of other loci to phenotypic variance need to be identified. In populations where markers from *six6* effectively predict phenotypic variation for age at maturity, it is possible to use this information to better understand mechanisms underlying observed phenotypic changes collected across temporal and spatial replicates, including monitoring effects of climate change and anthropogenic effects such as fisheries-induced selection and supportive breeding. Monitoring of both adaptive and neutral (e.g. Schwartz, Luikart, & Waples, 2007) genetic variation might inform management actions such as harvest regulations aimed at conserving a broad portfolio of life-history variation that can be vital for species to persist in dynamic ecosystems (e.g. Schindler et al., 2010). While acknowledging that gene-targeted conservation can be problematic (Kardos & Luikart, 2020; Shafer et al., 2015; Waples & Lindley, 2018), there are scenarios where large effect loci can be expected to provide direct applications for conservation management.

Acknowledgments

We thank Meri Lindqvist, Ville Aukee, Yann Czorlich, Victoria Pritchard, Jan Laine, Ryan Bare, Jordan Watson, Samuel May, Isadora Jimenez-Hidalgo, Eleni Petrou, Carolyn Tarpey, Cassandra Columbus, Ellen Campbell, Elena Correa, Vanessa Morman, and Steven Micheletti for samples, project development, and laboratory assistance. We also thank Wes Larson, Ingrid Spies, and two anonymous reviewers for providing insightful comments on this manuscript. This work was supported by the Academy of Finland (project numbers 284941, 286334, 314254, and 314255), the European Research Council under the European Articles Union's Horizon 2020 research and innovation program (grant agreement number 742312), the University of Helsinki, Washington Sea Grant (award numbers R/HCE-4 and R/E/I-26), and the School of Aquatic and Fishery Sciences at the University of Washington. The findings and conclusions in this paper are those of the authors and do not necessarily represent the views of the National Marine Fisheries Service. Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

References

- Ayala, F. J., & Campbell, C. A. (1974). Frequency-dependent selection. *Annual Review of Ecology and Systematics*, 5, 115-138.
- Ayllon, F., Kjaerner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M. F., Dahle, G., . . . Wargelius, A. (2015). The vgll3 locus controls age at maturity in wild and domesticated Atlantic salmon (*Salmo salar L.*) males. *PLOS Genetics*, 11(11), 15. doi:10.1371/journal.pgen.1005628
- Ayllon, F., Solberg, M. F., Glover, K. A., Mohammadi, F., Kjaerner-Semb, E., Fjellidal, P. G., . . . Wargelius, A. (2019). The influence of vgll3 genotypes on sea age at maturity is altered in farmed mowi strain Atlantic salmon. *BMC Genetics*, 20, 8. doi:10.1186/s12863-019-0745-9
- Baetscher, D. S., Clemento, A. J., Ng, T. C., Anderson, E. C., & Garza, J. C. (2018). Microhaplotypes provide increased power from short-read DNA sequences for relationship inference. *Molecular Ecology Resources*, 18(2), 296-305. doi:10.1111/1755-0998.12737
- Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., . . . Primmer, C. R. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, 528(7582), 405-408. doi:10.1038/nature16062
- Barton, K. (2016). Package 'MuMIn.' CRAN Repos at <https://cran.r-project.org/web/packages/MuMIn/MuMIn.pdf>.
- Barton, N. H., & Keightley, P. D. (2002). Understanding quantitative genetic variation. *Nature Reviews Genetics*, 3(1), 11-21. doi:10.1038/nrg700
- Behnke, R. J. (2002). *Trout and Salmon of North America*. New York: Free Press.
- Boulding, E. G., Ang, K. P., Elliott, J. A. K., Powell, F., & Schaeffer, L. R. (2019). Differences in genetic architecture between continents at a major locus previously associated with sea age at sexual maturity in European Atlantic salmon. *Aquaculture*, 500, 670-678. doi:10.1016/j.aquaculture.2018.09.025
- Bowersox, B. J., Corsi, M. P., McCormick, J. L., Copeland, T., & Campbell, M. R. (2019). Examining life history shifts and genetic composition in a hatchery steelhead population, with implications for fishery and ocean selection. *Transactions of the American Fisheries Society*, 148(6), 1056-1068. doi:10.1002/tafs.10199

- Brown, C. L., & Godduhn, A. (2015). Socioeconomic effects of declining salmon runs on the Yukon River. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 398, 170 p. Fairbanks, AK.
- Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources*, 15(4), 855-867. doi:10.1111/1755-0998.12357
- Cánovas, A., Reverter, A., DeAtley, K. L., Ashley, R. L., Colgrave, M. L., Fortes, M. R. S., . . . Thomas, M. G. (2014). Multi-tissue omics analyses reveal molecular regulatory networks for puberty in composite beef cattle. *PLOS One*, 9(7), 17. doi:10.1371/journal.pone.0102551
- Carlson, S. M., & Seamons, T. R. (2008). A review of quantitative genetic components of fitness in salmonids: implications for adaptation to future change. *Evolutionary Applications*, 1(2), 222-238. doi:10.1111/j.1752-4571.2008.00025.x
- Cauwelier, E., Gilbey, J., Sampayo, J., Stradmeyer, L., & Middlemas, S. J. (2018). Identification of a single genomic region associated with seasonal river return timing in adult Scottish Atlantic salmon (*Salmo salar*), using a genome-wide association study. *Canadian Journal of Fisheries and Aquatic Sciences*, 75(9), 1427-1435. doi:10.1139/cjfas-2017-0293
- Charlesworth, B., Nordborg, M., & Charlesworth, D. (1997). The effects of local selection, balanced polymorphism and background selection on equilibrium patterns of genetic diversity in subdivided populations. *Genetics Research*, 70(2), 155-174. doi:10.1017/s0016672397002954
- Charnov, E. L., & Gillooly, J. F. (2004). Size and temperature in the evolution of fish life histories. *Integrative and Comparative Biology*, 44(6), 494-497. doi:10.1093/icb/44.6.494
- Christensen, K. A., Gutierrez, A. P., Lubieniecki, K. P., & Davidson, W. S. (2017). TEAD3, implicated by association to grilising in Atlantic salmon. *Aquaculture*, 479, 571-578. doi:10.1016/j.aquaculture.2017.06.026
- Christensen, R. H. B. (2019). ordinal-Regression models for ordinal data. R package version 2019.12-10. <https://CRAN.R-project.org/package=ordinal>.
- Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., . . . Ruden, D. M. (2012). A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:

SNPs in the genome of *Drosophila melanogaster* strain w(1118); iso-2; iso-3. *Fly*, 6(2), 80-92.
doi:10.4161/fly.19695

Clark, D. D., Gorman, M. R., Hatori, M., Meadows, J. D., Panda, S., & Mellon, P. L. (2013). Aberrant development of the suprachiasmatic nucleus and circadian rhythms in mice lacking the homeodomain protein six6. *Journal of Biological Rhythms*, 28(1), 15-25.

doi:10.1177/0748730412468084

Cline, T. J., Ohlberger, J., & Schindler, D. E. (2019). Effects of warming climate and competition in the ocean for life-histories of Pacific salmon. *Nature Ecology & Evolution*, 3(6), 935-942.

doi:10.1038/s41559-019-0901-7

Copeland, T., Ackerman, M. W., Wright, K. K., & Byrne, A. (2017). Life history diversity of Snake River steelhead populations between and within management categories. *North American Journal of Fisheries Management*, 37(2), 395-404. doi:10.1080/02755947.2016.1264506

Cortez, D., Marin, R., Toledo-Flores, D., Froidevaux, L., Liechti, A., Waters, P. D., . . . Kaessmann, H. (2014). Origins and functional evolution of Y chromosomes across mammals. *Nature*, 508(7497), 488-+. doi:10.1038/nature13151

Cousminer, D. L., Berry, D. J., Timpson, N. J., Ang, W., Thiering, E., Byrne, E. M., . . . Early Growth Genetics, E. G. G. (2013). Genome-wide association and longitudinal analyses reveal genetic loci linking pubertal height growth, pubertal timing and childhood adiposity. *Human Molecular Genetics*, 22(13), 2735-2747. doi:10.1093/hmg/ddt104

Cousminer, D. L., Widen, E., & Palmert, M. R. (2016). The genetics of pubertal timing in the general population: recent advances and evidence for sex-specificity. *Current Opinion in Endocrinology Diabetes and Obesity*, 23(1), 57-65. doi:10.1097/med.0000000000000213

Crete-Lafreniere, A., Weir, L. K., & Bernatchez, L. (2012). Framing the *Salmonidae* family phylogenetic portrait: A more complete picture from increased taxon sampling. *PLOS One*, 7(10), 19. doi:10.1371/journal.pone.0046662

Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P., & Primmer, C. R. (2018). Rapid sex-specific evolution of age at maturity is shaped by genetic architecture in Atlantic salmon. *Nature Ecology & Evolution*, 2(11), 1800-1807. doi:10.1038/s41559-018-0681-5

- Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., Parre, N., . . . Primmer, C. R. (2019). Large single-locus effects for maturation timing are mediated via condition variation in Atlantic salmon. *bioRxiv*. doi:https://doi.org/10.1101/780437
- DeFilippo, L. B., Schindler, D. E., Ohlberger, J., Schaberg, K. L., Foster, M. B., Ruhl, D., & Punt, A. E. (2019). Recruitment variation disrupts the stability of alternative life histories in an exploited salmon population. *Evolutionary Applications*, *12*(2), 214-229. doi:10.1111/eva.12709
- Devlin, B., & Roeder, K. (1999). Genomic control for association studies. *Biometrics*, *55*(4), 997-1004. doi:10.1111/j.0006-341X.1999.00997.x
- Elmer, K. R., & Meyer, A. (2011). Adaptation in the age of ecological genomics: insights from parallelism and convergence. *Trends in Ecology & Evolution*, *26*(6), 298-306. doi:10.1016/j.tree.2011.02.008
- Fariello, M. I., Boitard, S., Mercier, S., Robelin, D., Faraut, T., Arnould, C., . . . SanCristobal, M. (2017). Accounting for linkage disequilibrium in genome scans for selection without individual genotypes: The local score approach. *Molecular Ecology*, *26*(14), 3700-3714. doi:10.1111/mec.14141
- Faust, G. G., & Hall, I. M. (2014). SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics*, *30*(17), 2503-2505. doi:10.1093/bioinformatics/btu314
- Flatt, T., & Heyland, A. (2011). *Mechanisms of Life History Evolution: the genetics and physiology of life history traits and trade-offs*: Oxford University Press.
- Fleming, I. A. (1996). Reproductive strategies of Atlantic salmon: Ecology and evolution. *Reviews in Fish Biology and Fisheries*, *6*(4), 379-416. doi:10.1007/bf00164323
- Freese, N. H., Norris, D. C., & Loraine, A. E. (2016). Integrated genome browser: visual analytics platform for genomics. *Bioinformatics*, *32*(14), 2089-2095. doi:10.1093/bioinformatics/btw069
- Giska, I., Farello, L., Pimenta, J., Seixas, F. A., Ferreira, M. S., Marques, J. P., . . . Melo-Ferreira, J. (2019). Introgression drives repeated evolution of winter coat color polymorphism in hares. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(48), 24150-24156. doi:10.1073/pnas.1910471116

- Gjerde, B. (1984). Response to individual selection for age at sexual maturity in Atlantic salmon. *Aquaculture*, 38(3), 229-240. doi:10.1016/0044-8486(84)90147-9
- Greene, C. M., Hall, J. E., Guilbault, K. R., & Quinn, T. P. (2010). Improved viability of populations with diverse life-history portfolios. *Biology Letters*, 6(3), 382-386. doi:10.1098/rsbl.2009.0780
- Guerrero, R. F., & Hahn, M. W. (2017). Speciation as a sieve for ancestral polymorphism. *Molecular Ecology*, 26(20), 5362-5368. doi:10.1111/mec.14290
- Hankin, D. G., Fitzgibbons, J., & Chen, Y. M. (2009). Unnatural random mating policies select for younger age at maturity in hatchery Chinook salmon (*Oncorhynchus tshawytscha*) populations. *Canadian Journal of Fisheries and Aquatic Sciences*, 66(9), 1505-1521. doi:10.1139/f09-085
- Hankin, D. G., & Healey, M. C. (1986). Dependence of exploitation rates for maximum yield and stock collapse on age and sex structure of Chinook salmon (*Oncorhynchus tshawytscha*) stocks. *Canadian Journal of Fisheries and Aquatic Sciences*, 43(9), 1746-1759. doi:10.1139/f86-219
- Hankin, D. G., Nicholas, J. W., & Downey, T. W. (1993). Evidence for inheritance of age at maturity in Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences*, 50(2), 347-358. doi:10.1139/f93-040
- Harstad, D. L., Larsen, D. A., & Beckman, B. R. (2014). Variation in minijack rate among hatchery populations of Columbia river basin Chinook salmon. *Transactions of the American Fisheries Society*, 143(3), 768-778. doi:10.1080/00028487.2014.886621
- Harstad, D. L., Larsen, D. A., Miller, J., Adams, I., Spangenberg, D. K., Nance, S., . . . Beckman, B. R. (2018). Winter-rearing temperature affects growth profiles, age of maturation, and smolt-to-adult returns for yearling summer Chinook salmon in the Upper Columbia River Basin. *North American Journal of Fisheries Management*, 38(4), 867-885. doi:10.1002/nafm.10186
- Healey, M. C. (1991). Life history in Chinook salmon (*Oncorhynchus tshawytscha*). In C. Groot & L. Margolis (Eds.), *Pacific salmon life histories* (pp. 311-393). Vancouver, Canada: University of British Columbia Press.
- Hess, J. E., Ackerman, M. W., Fryer, J. K., Hasselman, D. J., Steele, C. A., Stephenson, J. J., . . . Narum, S. R. (2016). Differential adult migration-timing and stock-specific abundance of

steelhead in mixed stock assemblages. *ICES Journal of Marine Science*, 73(10), 2606-2615.
doi:10.1093/icesjms/fsw138

Hou, H. Y., Uuskula-Reimand, L., Makarem, M., Corre, C., Saleh, S., Metcalf, A., . . . Wilson, M. D. (2017). Gene expression profiling of puberty-associated genes reveals abundant tissue and sex-specific changes across postnatal development. *Human Molecular Genetics*, 26(18), 3585-3599. doi:10.1093/hmg/ddx246

Jamie, G. A., & Meier, J. I. (2020). The persistence of polymorphisms across species radiations. *Trends in Ecology and Evolution*, 35(9), 795-808. doi:10.1016/j.tree.2020.04.007

Janowitz-Koch, I., Rabe, C., Kinzer, R., Nelson, D., Hess, M. A., & Narum, S. R. (2019). Long-term evaluation of fitness and demographic effects of a Chinook Salmon supplementation program. *Evolutionary Applications*, 12(3), 456-469. doi:10.1111/eva.12725

Jean, D., Bernier, G., & Gruss, P. (1999). Six6 (Optx2) is a novel murine Six3-related homeobox gene that demarcates the presumptive pituitary/hypothalamic axis and the ventral optic stalk. *Mechanisms of Development*, 84(1-2), 31-40. doi:10.1016/s0925-4773(99)00068-4

Johnston, S. E., Orell, P., Pritchard, V. L., Kent, M. P., Lien, S., Niemela, E., . . . Primmer, C. R. (2014). Genome-wide SNP analysis reveals a genetic basis for sea-age variation in a wild population of Atlantic salmon (*Salmo salar*). *Molecular Ecology*, 23(14), 3452-3468. doi:10.1111/mec.12832

Jombart, T. (2008). *adeget*: a R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24(11), 1403-1405. doi:10.1093/bioinformatics/btn129

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics*, 27(21), 3070-3071. doi:10.1093/bioinformatics/btr521

Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., . . . Whole Genome Assembly, T. (2012). The genomic basis of adaptive evolution in threespine sticklebacks. *Nature*, 484(7392), 55-61. doi:10.1038/nature10944

Kardos, M., & Luikart, G. (2020). The genetic architecture of fitness drives population viability during rapid environmental change. *BioRxiv*:660803.

Accepted Article
Keefler, M. L., & Caudill, C. C. (2014). Homing and straying by anadromous salmonids: a review of mechanisms and rates. *Reviews in Fish Biology and Fisheries*, 24(1), 333-368.

doi:10.1007/s11160-013-9334-6

Kjaerner-Semb, E., Ayllon, F., Kleppe, L., Sorhus, E., Skaftnesmo, K., Furmanek, T., . . . Edvardsen, R. B. (2018). Vgll3 and the Hippo pathway are regulated in Sertoli cells upon entry and during puberty in Atlantic salmon testis. *Scientific Reports*, 8, 11. doi:10.1038/s41598-018-20308-1

Kodama, M., Briec, M. S. O., Devlin, R. H., Hard, J. J., & Naish, K. A. (2014). Comparative mapping Between coho salmon (*Oncorhynchus kisutch*) and three other salmonids suggests a role for chromosomal rearrangements in the retention of duplicated regions following a whole genome duplication event. *G3-Genes Genomes Genetics*, 4(9), 1717-1730.

doi:10.1534/g3.114.012294

Kodama, M., Hard, J. J., & Naish, K. A. (2018). Mapping of quantitative trait loci for temporal growth and age at maturity in coho salmon: Evidence for genotype-by-sex interactions.

Marine Genomics, 38, 33-44. doi:10.1016/j.margen.2017.07.004

Kofler, R., Pandey, R. V., & Schlotterer, C. (2011). PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics*, 27(24), 3435-3436. doi:10.1093/bioinformatics/btr589

Koressaar, T., & Remm, M. (2007). Enhancements and modifications of primer design program Primer3. *Bioinformatics*, 23(10), 1289-1291. doi:10.1093/bioinformatics/btm091

Korte, A., & Farlow, A. (2013). The advantages and limitations of trait analysis with GWAS: a review. *Plant Methods*, 9, 9. doi:10.1186/1746-4811-9-29

Kozłowski, J. (1992). Optimal allocation of resources to growth and reproduction - implications for age and size at maturity. *Trends in Ecology & Evolution*, 7(1), 15-19. doi:10.1016/0169-5347(92)90192-e

Kronforst, M. R., Barsh, G. S., Kopp, A., Mallet, J., Monteiro, A., Mullen, S. P., . . . Hoekstra, H. E. (2012). Unraveling the thread of nature's tapestry: the genetics of diversity and convergence in animal pigmentation. *Pigment Cell & Melanoma Research*, 25(4), 411-433.

doi:10.1111/j.1755-148X.2012.01014.x

Kumar, J. P. (2009). The sine oculis homeobox (SIX) family of transcription factors as regulators of development and disease. *Cellular and Molecular Life Sciences*, 66(4), 565-583.

doi:10.1007/s00018-008-8335-4

Kurko, J., Debes, P. V., House, A. H., Aykanat, T., Erkinaro, J., & Primmer, C. R. (2020).

Transcription profiles of age-at-maturity-associated genes suggest cell fate commitment regulation as a key factor in the Atlantic salmon maturation process. *G3-Genes Genomes Genetics*, 10(1), 235-246. doi:10.1534/g3.119.400882

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4), 357-U354. doi:10.1038/nmeth.1923

Larder, R., Clark, D. D., Miller, N. L. G., & Mellon, P. L. (2011). Hypothalamic dysregulation and infertility in mice lacking the homeodomain protein six6. *Journal of Neuroscience*, 31(2), 426-438. doi:10.1523/jneurosci.1688-10.2011

Larsen, D. A., Harstad, D. L., Fuhrman, A. E., Knudsen, C. M., Schroder, S. L., Bosch, W. J., . . .

Beckman, B. R. (2019). Maintaining a wild phenotype in a conservation hatchery program for Chinook salmon: The effect of managed breeding on early male maturation. *PLOS One*, 14(5), 32. doi:10.1371/journal.pone.0216168

Larsen, D. A., Harstad, D. L., Strom, C. R., Johnston, M. V., Knudsen, C. M., Fast, D. E., . . .

Beckman, B. R. (2013). Early Life History Variation in Hatchery- and Natural-Origin Spring Chinook Salmon in the Yakima River, Washington. *Transactions of the American Fisheries Society*, 142(2), 540-555. doi:10.1080/00028487.2012.750626

Lewis, B., Grant, W. S., Brenner, R. E., & Hamazaki, T. (2015). Changes in size and age of Chinook salmon *Oncorhynchus tshawytscha* returning to Alaska. *PLOS One*, 10(6), 17.

doi:10.1371/journal.pone.0130184

Li, C. Y., Kan, L. J., Chen, Y., Zheng, X. D., Li, W. N., Zhang, W. X., . . . Chen, D. H. (2015). Ci

antagonizes Hippo signaling in the somatic cells of the ovary to drive germline stem cell differentiation. *Cell Research*, 25(10), 1152-1170. doi:10.1038/cr.2015.114

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and

population genetical parameter estimation from sequencing data. *Bioinformatics*, 27(21), 2987-2993. doi:10.1093/bioinformatics/btr509

- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*. doi:1303.3997v2
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Genome Project Data, P. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, 25(16), 2078-2079. doi:10.1093/bioinformatics/btp352
- Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., . . . Davidson, W. S. (2016). The Atlantic salmon genome provides insights into rediploidization. *Nature*, 533(7602), 200-+. doi:10.1038/nature17164
- Liu, S. L., Ferchaud, A. L., Gronkjaer, P., Nygaard, R., & Hansen, M. M. (2018). Genomic parallelism and lack thereof in contrasting systems of three-spined sticklebacks. *Molecular Ecology*, 27(23), 4725-4743. doi:10.1111/mec.14782
- Macqueen, D. J., & Johnston, I. A. (2014). A well-constrained estimate for the timing of the salmonid whole genome duplication reveals major decoupling from species diversification. *Proceedings of the Royal Society B-Biological Sciences*, 281(1778), 8. doi:10.1098/rspb.2013.2881
- Mank, J. E. (2017). Population genetics of sexual conflict in the genomic era. *Nature Reviews Genetics*, 18(12), 721-730. doi:10.1038/nrg.2017.83
- Mank, J. E., & Avise, J. C. (2006). The evolution of reproductive and genomic diversity in ray-finned fishes: insights from phylogeny and comparative analysis. *Journal of Fish Biology*, 69(1), 1-27. doi:10.1111/j.1095-8649.2006.01132.x
- Marchini, J., Cardon, L. R., Phillips, M. S., & Donnelly, P. (2004). The effects of human population structure on large genetic association studies. *Nature Genetics*, 36(5), 512-517. doi:10.1038/ng1337
- McKinney, G. J., Nichols, K. M., & Ford, M. J. (2020). A mobile sex-determining region, male-specific haplotypes, and rearing environment influence age at maturity in Chinook salmon. *bioRxiv*. doi:https://doi.org/10.1101/2020.04.23.056093
- McKinney, G. J., Seeb, J. E., Pascal, C. E., Schindler, D. E., Gilk-Baumer, S. E., & Seeb, L. W. (2019). Y-chromosome haplotypes drive variation in size and age at maturity in male Chinook salmon. *bioRxiv*. doi:https://doi.org/10.1101/691063

- Micheletti, S. J., Hess, J. E., Zendt, J. S., & Narum, S. R. (2018). Selection at a genomic region of major effect is responsible for evolution of complex life histories in anadromous steelhead. *Bmc Evolutionary Biology*, 18, 11. doi:10.1186/s12862-018-1255-5
- Micheletti, S. J., & Narum, S. R. (2018). Utility of pooled sequencing for association mapping in nonmodel organisms. *Molecular Ecology Resources*, 18(4), 825-837. doi:10.1111/1755-0998.12784
- Mobley, K. B., Granroth-Wilding, H., Ellmen, M., Vaha, J. P., Aykanat, T., Johnston, S. E., . . . Primmer, C. R. (2019). Home ground advantage: Local Atlantic salmon have higher reproductive fitness than dispersers in the wild. *Science Advances*, 5(2), 8. doi:10.1126/sciadv.aav1112
- Mohamed, A. R., Verbyla, K. L., Al-Mamun, H. A., McWilliam, S., Evans, B., King, H., . . . Kijas, J. W. (2019). Polygenic and sex specific architecture for two maturation traits in farmed Atlantic salmon. *Bmc Genomics*, 20, 12. doi:10.1186/s12864-019-5525-4
- Mundy, N. I. (2005). A window on the genetics of evolution: MC1R and plumage colouration in birds. *Proceedings of the Royal Society B-Biological Sciences*, 272(1573), 1633-1640. doi:10.1098/rspb.2005.3107
- Nielsen, J. L. (1999). The evolutionary history of steelhead (*Oncorhynchus mykiss*) along the US Pacific Coast: Developing a conservation strategy using genetic diversity. *ICES Journal of Marine Science*, 56(4), 449-458. doi:10.1006/jmsc.1999.0452
- Ohlberger, J., Schindler, D. E., Brown, R. J., Harding, J. M. S., Adkison, M. D., Munro, A. R., . . . Spaeder, J. (2020). The reproductive value of large females: consequences of shifts in demographic structure for population reproductive potential in Chinook salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, 77(8). Early online <https://doi.org/10.1139/cjfas-2020-0012>
- Ohlberger, J., Ward, E. J., Schindler, D. E., & Lewis, B. (2018). Demographic changes in Chinook salmon across the Northeast Pacific Ocean. *Fish and Fisheries*, 19(3), 533-546. doi:10.1111/faf.12272

- Oomen, R. A., Kuparinen, A., & Hutchings, J. A. (2020). Consequences of single-locus and tightly linked genomic architectures for evolutionary responses to environmental change. *Journal of Heredity*, *111*(4), 319-332.
- Pearse, D. E., Barson, N. J., Nome, T., Gao, G. T., Campbell, M. A., Abadia-Cardoso, A., . . . Lien, S. (2019). Sex-dependent dominance maintains migration supergene in rainbow trout. *Nature Ecology & Evolution*, *3*(12), 1731-+. doi:10.1038/s41559-019-1044-6
- Perry, J. R. B., Day, F., Elks, C. E., Sulem, P., Thompson, D. J., Ferreira, T., . . . Early Growth Genetics, E. G. G. C. (2014). Parent-of-origin-specific allelic associations among 106 genomic loci for age at menarche. *Nature*, *514*(7520), 92-+. doi:10.1038/nature13545
- Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., & Reich, D. (2006). Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics*, *38*(8), 904-909. doi:10.1038/ng1847
- Pritchard, V. L., Makinen, H., Vaha, J. P., Erkinaro, J., Orell, P., & Primmer, C. R. (2018). Genomic signatures of fine-scale local selection in Atlantic salmon suggest involvement of sexual maturation, energy homeostasis and immune defence-related genes. *Molecular Ecology*, *27*(11), 2560-2575. doi:10.1111/mec.14705
- Quinn, T. P. (2005). *The Behavior and Ecology of Pacific Salmon and Trout*. Seattle: University of Washington Press.
- R Core Team. (2019). R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org>. Retrieved from <http://www.R-project.org>
- Ricker, W. E. (1980). *Causes of the decrease in age and size of Chinook salmon (Oncorhynchus tshawytscha)*. Canadian Technical Report of Fisheries and Aquatic Sciences No. 944, 25 p. Nanaimo, British Columbia.
- Roff, D. A. (2011). Alternative strategies: The evolution of switch points. *Current Biology*, *21*(8), R285-R287. doi:10.1016/j.cub.2011.03.016
- Roff, D. A., Mostowy, S., & Fairbairn, D. J. (2002). The evolution of trade-offs: Testing predictions on response to selection and environmental variation. *Evolution*, *56*(1), 84-95.

- Samy, J. K. A., Mulugeta, T. D., Nome, T., Sandve, S. R., Grammes, F., Kent, M. P., . . . Vage, D. I. (2017). SalmoBase: an integrated molecular data resource for Salmonid species. *Bmc Genomics*, *18*, 5. doi:10.1186/s12864-017-3877-1
- Satterthwaite, W. H., Carlson, S. M., & Criss, A. (2017). Ocean size and corresponding life history diversity among the four run timings of California Central Valley Chinook salmon. *Transactions of the American Fisheries Society*, *146*(4), 594-610. doi:10.1080/00028487.2017.1293562
- Schaffer, W. M. (2004). Life histories, evolution, and salmonids. In A. P. Hendry & S. C. Stearns (Eds.), *Evolution Illuminated: Salmon and Their Relatives*: Oxford University Press.
- Schindler, D. E., Hilborn, R., Chasco, B., Boatright, C. P., Quinn, T. P., Rogers, L. A., & Webster, M. S. (2010). Population diversity and the portfolio effect in an exploited species. *Nature*, *465*(7298), 609-612. doi:10.1038/nature09060
- Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals-mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics*, *15*(11), 749-763. doi:10.1038/nrg3803
- Schwartz, M. K., Luikart, G., & Waples, R. S. (2007). Genetic monitoring as a promising tool for conservation and management. *Trends in Ecology & Evolution*, *22*(1), 25-33. doi:10.1016/j.tree.2006.08.009
- Seo, H. C., Drivenes, O., Ellingsen, S., & Fjose, A. (1998). Expression of two zebrafish homologues of the murine Six3 gene demarcates the initial eye primordia. *Mechanisms of Development*, *73*(1), 45-57. doi:10.1016/s0925-4773(98)00028-8
- Shafer, A. B. A., Wolf, J. B. W., Alves, P. C., Bergstrom, L., Bruford, M. W., Brannstrom, I., . . . Zielinski, P. (2015). Genomics and the challenging translation into conservation practice. *Trends in Ecology & Evolution*, *30*(2), 78-87. doi:10.1016/j.tree.2014.11.009
- Simon, E., Fauchaux, C., Zider, A., Theze, N., & Thiebaud, P. (2016). From vestigial to vestigial-like: the *Drosophila* gene that has taken wing. *Development Genes and Evolution*, *226*(4), 297-315. doi:10.1007/s00427-016-0546-3
- Sinclair-Waters, M., Odegard, J., Korsvoll, S. A., Moen, T., Lien, S., Primmer, C. R., & Barson, N. J. (2020). Beyond large-effect loci: large-scale GWAS reveals a mixed large-effect and

- polygenic architecture for age at maturity of Atlantic salmon. *Genetics Selection Evolution*, 52(1), 11. doi:10.1186/s12711-020-0529-8
- Stearns, S. C. (1989). Trade-offs in life-history evolution. *Functional Ecology*, 3(3), 259-268. doi:10.2307/2389364
- Stearns, S. C. (1992). *The Evolution of Life Histories*. London: Oxford University Press.
- Taylor, E. B. (1990). Environmental correlates of life-history variation in juvenile Chinook salmon, *Oncorhynchus tshawytscha* (Walbaum). *Journal of Fish Biology*, 37(1), 1-17. doi:10.1111/j.1095-8649.1990.tb05922.x
- Thorpe, J. E. (2004). Life history responses of fishes to culture. *Journal of Fish Biology*, 65, 263-285. doi:10.1111/j.0022-1112.2004.00556.x
- Turner, S. D. (2014). qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *bioRxiv*. doi:https://doi.org/10.1101/005165
- Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., & Rozen, S. G. (2012). Primer3-new capabilities and interfaces. *Nucleic Acids Research*, 40(15), 12. doi:10.1093/nar/gks596
- Vähä, J. P., Erkinaro, J., Niemela, E., & Primmer, C. R. (2008). Temporally stable genetic structure and low migration in an Atlantic salmon population complex: implications for conservation and management. *Evolutionary Applications*, 1(1), 137-154. doi:10.1111/j.1752-4571.2007.00007.x
- Veale, A. J., & Russello, M. A. (2017). An ancient selective sweep linked to reproductive life history evolution in sockeye salmon. *Scientific Reports*, 7, 10. doi:10.1038/s41598-017-01890-2
- Vertä, J.-P., Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., . . . Primmer, C. R. (2019). Regulatory divergence in *vgl3* underlies variation in age at maturity in male Atlantic salmon. *bioRxiv*. doi:https://doi.org/10.1101/777300
- Waples, R. S., & Lindley, S. T. (2018). Genomics and conservation units: The genetic basis of adult migration timing in Pacific salmonids. *Evolutionary Applications*, 11(9), 1518-1526. doi:10.1111/eva.12687

- Waples, R. S., Teel, D. J., Myers, J. M., & Marshall, A. R. (2004). Life-history divergence in Chinook salmon: Historic contingency and parallel evolution. *Evolution*, 58(2), 386-403.
doi:10.1111/j.0014-3820.2004.tb01654.x
- Watanabe, N., Itoh, K., Mogi, M., Fujinami, Y., Shimizu, D., Hashimoto, H., . . . Suzuki, T. (2012). Circadian pacemaker in the suprachiasmatic nuclei of teleost fish revealed by rhythmic period2 expression. *General and Comparative Endocrinology*, 178(2), 400-407.
doi:10.1016/j.ygcen.2012.06.012
- Willis, S. C., Hess, J. E., Fryer, J. K., Whiteaker, J. M., Brun, C., Gerstenberger, R., & Narum, S. R. (2020). Steelhead (*Oncorhynchus mykiss*) lineages and sexes show variable patterns of association of adult migration timing and age-at-maturity traits with two genomic regions. *Evolutionary Applications*, 21. doi:10.1111/eva.13088
- Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., & Guiguen, Y. (2013). The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids. *Evolutionary Applications*, 6(3), 486-496.
doi:10.1111/eva.12032
- Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-selection balance. *Evolution*, 65(7), 1897-1911. doi:10.1111/j.1558-5646.2011.01269.x

Data Accessibility

All genotypic and phenotypic data are provided in Supplementary Tables S1-S4. The R script and input files for association analyses between age at maturity and *six6* and *vgll3* are in the Dryad Digital Repository at <https://doi.org/10.5061/dryad.x0k6djhhq>. Additional raw sequence data that support the findings of this study will be openly available in the NCBI SRA Database at <https://www.ncbi.nlm.nih.gov/sra>, accession PRJNA650380.

Author Contributions

C.R.P., K.A.N., J.C.G., and S.N. designed the study; C.D.W., A.C., T.A., and S.N. analyzed the data, C.D.W., J.C.G., K.A.N., S.N., and C.R.P. drafted the paper, and all authors contributed to the revision of the manuscript and gave final approval for publication.

Figure captions

Figure 1. Predicted probabilities of delaying maturation with 95% confidence intervals for male Coho Salmon and Sockeye Salmon of both sexes for the genotypes of target SNPs in the *six6* and *vgll3* genome regions, obtained from the logistic regression models. When the most parsimonious model included PCs, the mean PC value was used when calculating probabilities. Only male Coho Salmon were analyzed, as females were nearly invariant in age. Probabilities for Sockeye Salmon at *vgll3* apply to both sexes since sex was not included in the most parsimonious model.

Figure 2. Cumulative probabilities of maturation with 95% confidence intervals for each standardized age class of male and female Chinook Salmon for the genotypes of target SNPs in the *six6* and *vgll3* gene regions, obtained from a cumulative proportional odds model. Mean PC values were used when calculating probabilities. As standardized age four was the oldest age, cumulative probabilities of maturation equal one for this class (i.e. all fish must mature by standardized age four).

Figure 3. Cumulative probabilities of maturation with 95% confidence intervals for each standardized age class of male and female Steelhead Trout for the genotypes of target SNPs in the *six6* and *vgll3* gene regions, obtained from a cumulative proportional odds model. Mean PC values were used when calculating probabilities. As standardized age four was the oldest age, cumulative probabilities of maturation equal one for this class (i.e. all fish must mature by standardized age four).

Figure 4. Genome-wide association test for Steelhead Trout including a) Manhattan plot of genomic regions associated with ocean-age at maturity in paired collections (1-ocean vs. 2-ocean) of male Steelhead from three populations (Local score results from CMH test of each pair of collections per location); b) Manhattan plot (p -values from Fishers exact test) zoomed to the significant region on chromosome Omy25 with green SNPs from the *six6* gene and gray SNPs intergenic, dashed red line is Bonferroni-corrected significance threshold; c) gene diagram for *six6*.

Table 1. Populations of hatchery (H) and wild (W) Chinook Salmon, Coho Salmon, Sockeye Salmon, and Steelhead Trout that were analyzed in this study with sample sizes of females (F) and males (M) for each age class. Ages denote total age at maturity, with standardized ages (i.e. youngest age class of each population = 1) reported in parentheses for Chinook and Steelhead. Standardized ages were analyzed for these two species to account for inherent differences in age structure of the populations. The number of putatively neutral, polymorphic loci genotyped to serve as controls and assess potential confounding effects is also listed for each species. Population locations are given by State; CA - California, OR - Oregon, WA - Washington, AK - Alaska.

Population	Location	Years collected	Age structure and sample sizes	Aging methods	Genotyping methods	Reference
Chinook salmon (neutral loci = 90)						
Cle Elum, WA (H)	47°18'N, 120°96'W	1998-2014	Age 3 (1): F = 3, M = 45 Age 4 (2): F = 49, M = 51 Age 5 (3): F = 22, M = 16	Tags and scales	Primary = GT seq Secondary = Sanger	Waters et al., 2018
Feather River, CA (H)	39°31'N, 121°33'W	2009-2017	Age 2 (1): F = 14, M = 21 Age 3 (2): F = 43, M = 45 Age 4 (3): F = 45, M = 46 Age 5 (4): F = 4, M = 8	Genetic pedigree	GT seq	Clemento, 2013
Chickamin River stock, Little Port Walter, AK (H)	56°38'N, 134°64'W	2013-2016	Age 4 (1): F = 0, M = 47 Age 5 (2): F = 41, M = 43	Tags	GT seq	Templin, 2001
Unuk River stock, Little Port Walter, AK (H)	56°38'N, 134°64'W	2014-2016	Age 3 (1): F = 0, M = 43 Age 4 (2): F = 4, M = 39 Age 5 (3): F = 46, M = 47 Age 6 (4): F = 54, M = 40	Tags	GT seq	Templin, 2001
Sacramento River, CA (W)	40°36'N, 122°26'W	2012-2017	Age 2 (1): F = 48, M = 46 Age 3 (2): F = 48, M = 48 Age 4 (3): F = 2, M = 8	Genetic pedigree	GT seq	Myers et al., 1998
Coho salmon (neutral loci = 84)						
Big Beef Creek, WA (W)	47°39'N, 122°46'W	2008-2009	Age 2: F = 0, M = 96 Age 3: F = 40, M = 54	Genetic pedigree	Primary = GT seq Secondary = Sanger	Kodama et al., 2012
Klamath River, OR (W)	41°55'N, 122°26'W	2009-2012	Age 2: F = 6, M = 44 Age 3: F = 42, M = 41	Genetic pedigree	GT seq	Starks, 2014
Sockeye salmon (neutral loci = 172)						

A Creek, AK (W)	59°34'N, 159°07'W	2008-2010	Age 4: F = 47, M = 25	Genetic pedigree	Sanger	Lin et al., 2008; Peterson et al., 2014; May et al., 2020
C Creek, AK (W)	59°34'N, 159°09'W	2008-2010	Age 5: F = 19, M = 17 Age 4: F = 20, M = 23 Age 5: F = 13, M = 22	Genetic pedigree	Sanger	Lin et al., 2008; Peterson et al., 2014; May et al., 2020
Steelhead trout (neutral loci = 242)						
Dworshak, ID (H)	46°30'N, 116°19'W	2015-2016	Age 3 (1): F = 0, M = 30 Age 4 (2): F = 92, M = 73	Genetic pedigree and scales	GT seq	Hess et al., 2016
Eel River, CA (W)	39°23'N, 123°07'W	2012-2015	Age 2 (1): F = 4, M = 14 Age 3 (2): F = 41, M = 41 Age 4 (3): F = 43, M = 29 Age 5 (4): F = 14, M = 1 Age 6 (4): F = 5, M = 0	Genetic pedigree	GT seq	Unpublished data
Forks Creek, WA (H)	46°33'N, 123°35'W	2000-2009	Age 2 (1): F = 1, M = 30 Age 3 (2): F = 55, M = 60 Age 4 (3): F = 55, M = 45 Age 5 (4): F = 27, M = 14	Genetic pedigree	Primary = GT seq Secondary = Sanger	Naish et al., 2013
Pahsimeroi, ID (H)	44°41'N, 114°02'W	2015-2016	Age 3 (1): F = 42, M = 44 Age 4 (2): F = 74, M = 38 Age 5 (3): F = 29, M = 18	Genetic pedigree and scales	GT seq	Hess et al., 2016
Wallowa, OR (H)	45°25'N, 117°18'W	2015-2016	Age 2 (1): F = 6, M = 3 Age 3 (2): F = 20, M = 19 Age 4 (3): F = 3, M = 7 Age 5 (4): F = 0, M = 30	Genetic pedigree and scales	GT seq	Hess et al., 2016
Warm Springs, CA (H)	38°43'N, 122°59'W	2009-2011	Age 2 (1): F = 52, M = 75 Age 3 (2): F = 47, M = 49	Genetic pedigree	GT seq	Abadía-Cardoso et al., 2013

Table 2. Results of association analyses between age at maturity and the *six6* and *vgll3* genes for each species based on the most parsimonious models. Multiple SNPs within the *six6* and *vgll3* gene regions were tested for Chinook and Steelhead. Results reported here are for the SNPs with the most highly-associated additive components; those for the other SNPs are reported in Table S6. SNP rank indicates the relative significance of the *six6* and *vgll3* SNP associations compared to all putatively neutral SNPs that were tested in the same models. Significant *p*-values are in bold.


Gene	Species	N, {Age classes} ¹ , N _{pop}	Parsimonious model ²	AICc with SNP	Overall (adjusted ³) SNP <i>p</i> -values	Lambda	Additive (non-additive ⁴) component <i>p</i> -values	SNP Rank	Model	Notes
<i>six6</i>	Chinook	1042, {1-4}, 5	<i>six6</i> _{allelic} + PC1 + PC2 + PC3 + PC4 + <i>six6</i> _{allelic} *PC4	2453.2	0.002 (0.075)	2.46	N/A	11 / 91	Ordinal, nominal = sex	one of two SNPs tested
	Coho	233, {2-3}, 2	<i>six6</i> _{allelic} +PC1	313.1	0.104 (0.319)	2.65	N/A	21 / 85	Binomial	Only males ⁵
	Sockeye	176, {4-5}, 2	<i>six6</i> _{geno} + Sex	226.9	0.003 (0.008)	1.22	0.093 (0.003)	1 / 173	Binomial	
	Steelhead	1207,{1-4}, 6	<i>six6</i> _{geno} + PC1 + PC2 + PC3 + PC4 + Sex + <i>six6</i> _{geno} *PC1 + <i>six6</i> _{geno} *PC2 + <i>six6</i> _{geno} *Sex	2633.4	1.835E-19 (4.456E-9)	1.94	2.664E-16 (0.002)	1 / 243	Ordinal, nominal = sex	SNP Omy25_61294400
<i>vgll3</i>	Chinook	1046, {1-4}, 5	<i>vgll3</i> _{geno} + PC1 + PC2 + PC3 + PC4 + Sex + <i>vgll3</i> _{geno} *PC2 + <i>vgll3</i> _{geno} *PC3+ <i>vgll3</i> _{geno} *PC4 + <i>vgll3</i> _{geno} *Sex	2460.7	8.286E-6 (0.147)	2.86	0.028 (3.425E-5)	23 / 91	Ordinal, nominal = sex	one of two SNPs tested
	Coho	203, {2-3}, 2	<i>vgll3</i> _{allelic} + PC1	274.9	0.856 (0.900)	2.11	N/A	83 / 85	Binomial	Only males ⁵
	Sockeye	160, {4-5}, 2	<i>vgll3</i> _{geno}	203.2	0.047 (0.095)	1.30	0.804 (0.071)	12 / 173	Binomial	
	Steelhead	656,{1-4}, 3	<i>vgll3</i> _{allelic} + PC1 + PC2 + PC4 + Sex + <i>vgll3</i> _{allelic} *PC1 + <i>vgll3</i> _{allelic} *PC2	1483.4	0.016 (0.185)	2.14	N/A	48 / 243	Ordinal	SNP <i>vgll3</i> .1_309


¹age classes for Coho and Sockeye Salmon represent total age at maturity, while those for Chinook and Steelhead have been standardized (i.e. youngest age class of each population = 1) to account for inherent differences in age structure between populations.


²geno and allelic refer to non-additive and additive modelling of genotypes, respectively. Null models do not include any terms with the SNP.

³ p -values after correcting for genomic inflation (i.e. lambda).

⁴additive vs. non-additive effects were partitioned when a non-additive model was preferred by model selection; these p -values are unadjusted for possible genomic inflation.







This article is protected by copyright. All rights reserved

—●— A/A —●— A/G —●— G/G

—●— A/A —●— A/T —●— T/T

