





Clinical practice guideline on the management of septic shock and sepsis-associated organ dysfunction in children: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine

Marius Rehn^{1,2,3}  | Michelle S. Chew⁴ | Klaus T. Olkkola⁵ | Martin Ingi Sigurðsson^{6,7}  | Arvi Yli-Hankala^{8,9}  | Morten Hylander Møller^{10,11} 

¹Pre-hospital Division, Air Ambulance Department, Oslo University Hospital, Oslo, Norway

²The Norwegian Air Ambulance Foundation, Drøbak, Norway

³Faculty of Health Sciences, University of Stavanger, Stavanger, Norway

⁴Department of Anaesthesia and Intensive Care, Medicine and Health, Linköping University, Linköping, Sweden

⁵Department of Anaesthesiology, Intensive Care and Pain Medicine, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁶Department of Anaesthesia and Intensive Care Medicine, Landspítali University Hospital, Reykjavík, Iceland

⁷Faculty of Medicine, University of Iceland, Reykjavík, Iceland

⁸Department of Anaesthesia, Tampere University Hospital, Tampere, Finland

⁹Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland

¹⁰Department of Intensive Care, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark

¹¹Collaboration for Research in Intensive Care (CRIC), Copenhagen, Denmark

Correspondence

Marius Rehn, Pre-hospital Division, Air Ambulance Department, Oslo University Hospital, Postbox 4950 Nydalen, 0424 Oslo, Norway.
Email: marius.rehn@norskluftambulanse.no

Funding information

Funding was provided solely from the SSAI and institutional and/or departmental sources.

Abstract

Background: The Clinical Practice Committee of the Scandinavian Society of Anaesthesiology and Intensive Care Medicine endorses the clinical practice guideline *Surviving Sepsis Campaign International Guidelines for the Management of Septic Shock and Sepsis-Associated Organ Dysfunction in Children*. The guideline can serve as a useful decision aid for clinicians managing children with suspected and confirmed septic shock and sepsis-associated organ dysfunction.

KEYWORDS

AGREE II, children, clinical practice guideline, paediatrics, sepsis, septic shock

1 | BACKGROUND

Sepsis remains a leading global cause of mortality and morbidity in children.¹ Early deaths in paediatric sepsis are often caused by refractory shock, whereas later deaths are more attributable to multiple organ dysfunction syndrome.² Anaesthesiologists and intensivists are regularly involved in the identification, resuscitation

and management of children with sepsis. In February 2020, the *Surviving Sepsis Campaign International Guidelines for the Management of Septic Shock and Sepsis-Associated Organ Dysfunction in Children* was published.³

The Scandinavian Society of Anaesthesiology and Intensive Care Medicine (SSAI) Clinical Practice Committee (CPC) decided to appraise this guideline for possible endorsement to guide Scandinavian

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. *Acta Anaesthesiologica Scandinavica* published by John Wiley & Sons Ltd on behalf of Acta Anaesthesiologica Scandinavica Foundation.

anaesthesiologists and intensivists in the identification, resuscitation and management of children with sepsis.

2 | METHODS

The SSAI CPC assessed the guideline using the Appraisal of Guidelines for REsearch and Evaluation (AGREE) II tool,⁴ as per the outlined process for endorsement of non-SSAI guidelines.⁵

3 | RESULTS

3.1 | Quality appraisal (AGREE II)

Five SSAI CPC members completed the appraisal. One member co-authored the guideline and was excluded from the evaluation, as per the SSAI endorsement process.⁵

The individual domain totals were: (1) Scope and purpose 91%; (2) Stakeholder involvement 89%; (3) Rigor of development 85%; (4) Clarity of presentation 82%; (5) Applicability 63%; (6) Editorial independence 95%; and (7) Overall assessment 85%.

The breakdown of the individual appraisers (de-identified) is available in the Supplementary Material.

4 | DISCUSSION

This clinical practice guideline on the management of septic shock and sepsis-associated organ dysfunction in children received overall high ratings on all domains with acceptable agreement between the SSAI CPC appraisers. We believe this guideline can serve as a useful decision aid for clinicians managing children with suspected and confirmed septic shock and sepsis-associated organ dysfunction.

5 | CONCLUSION

The SSAI CPC endorses the clinical practice guideline *Surviving Sepsis Campaign International Guidelines for the Management of Septic Shock and Sepsis-Associated Organ Dysfunction in Children*.³

CONFLICT OF INTEREST

No Clinical Practice Committee member had direct conflicts of interest. MHM was a co-author of the guideline assessed and did not participate in the AGREE II assessment, as per the SSAI endorsement process. No other authors had indirect conflicts of interest.

ORCID

Marius Rehn  <https://orcid.org/0000-0001-9519-241X>

Martin Ingi Sigurðsson  <https://orcid.org/0000-0001-7054-0844>

Arvi Yli-Hankala  <https://orcid.org/0000-0002-5029-9181>

Morten Hylander Møller  <https://orcid.org/0000-0002-6378-9673>

REFERENCES

1. Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kissoon N. The global burden of paediatric and neonatal sepsis: a systematic review. *Lancet Respir Med*. 2018;6:223-230.
2. Weiss SL, Balamuth F, Hensley J, et al. The Epidemiology of hospital death following pediatric severe sepsis: when, why, and how children with sepsis die. *Pediatr Crit Care Med*. 2017;18:823-830.
3. Weiss SL, Peters MJ, Alhazzani W, et al. Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children. *Pediatr Crit Care Med*. 2020;21:e52-e106.
4. Brouwers MC, Kho ME, Browman GP, et al; Consortium ANS. AGREE II: advancing guideline development, reporting and evaluation in health care. *CMAJ*. 2010;182:E839-E842.
5. Rehn M, Chew MS, Olkkola KT, Orn Sverrisson K, Yli-Hankala A, Møller MH. Endorsement of clinical practice guidelines by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine. *Acta Anaesthesiol Scand*. 2019;63:161-163.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Rehn M, Chew MS, Olkkola KT, Sigurðsson MI, Yli-Hankala A, Møller MH. Clinical practice guideline on the management of septic shock and sepsis-associated organ dysfunction in children: Endorsement by the Scandinavian Society of Anaesthesiology and Intensive Care Medicine. *Acta Anaesthesiol Scand*. 2021;65:1365-1366. <https://doi.org/10.1111/aas.13958>