



**Manchester
Metropolitan**
University

Robinson, Orin J, Socolar, Jacob B, Stuber, Erica F, Auer, Tom, Berryman, Alex J, Boersch-Supan, Philipp H, Brightsmith, Donald J, Burbidge, Allan H, Butchart, Stuart HM, Davis, Courtney L, Giacomo, Adrian S Di, Dokter, Adriaan M, Farnsworth, Andrew, Fink, Daniel, Hochachka, Wesley M, Howell, Paige E, La Sorte, Frank A, Lees, Alexander ORCID logo ORCID: <https://orcid.org/0000-0001-7603-9081>, Marsden, Stuart ORCID logo ORCID: <https://orcid.org/0000-0002-0205-960X>, Martin, Robert, Martin, Rowan O, Masello, Juan F, Miller, Eliot T, Moodley, Yoshan, Musgrove, Andy, Noble, David G, Ojeda, Valeria, Quillfeldt, Petra, Royle, J Andrew, Ruiz-Gutierrez, Viviana, Tella, José L, Yorio, Pablo, Youngflesh, Casey and Johnston, Alison (2022) Extreme uncertainty and unquantifiable bias do not inform population sizes. *Proceedings of the National Academy of Sciences of USA*, 119 (10). ISSN 0027-8424

Downloaded from: <https://e-space.mmu.ac.uk/629273/>

Version: Published Version

Publisher: National Academy of Sciences

DOI: <https://doi.org/10.1073/pnas.2113862119>

Please cite the published version

<https://e-space.mmu.ac.uk>



Extreme uncertainty and unquantifiable bias do not inform population sizes

Orin J. Robinson^{a,1}, Jacob B. Socolar^a, Erica F. Stuber^b, Tom Auer^a, Alex J. Berryman^c, Philipp H. Boersch-Supan^{d,e}, Donald J. Brightsmith^f, Allan H. Burbidge^g, Stuart H. M. Butchart^{c,h}, Courtney L. Davis^a, Adriaan M. Dokter^a, Adrian S. Di Giacomoⁱ, Andrew Farnsworth^a, Daniel Fink^a, Wesley M. Hochachka^a, Paige E. Howellⁱ, Frank A. La Sorte^a, Alexander C. Lees^{a,k}, Stuart Marsden^l, Robert Martin^c, Rowan O. Martin^m, Juan F. Maselloⁿ, Eliot T. Miller^a, Yoshan Moodley^o, Andy Musgrove^d, David G. Noble^d, Valeria Ojeda^p, Petra Quillfeldtⁿ, J. Andrew Royle^q, Viviana Ruiz-Gutierrez^a, José L. Tella^r, Pablo Yorio^{s,t}, Casey Youngflesh^u, and Alison Johnston^{a,v}

Species-specific population estimates are fundamental for many aspects of ecology, evolution, and conservation, yet they are lacking for most species. Aiming to fill this gap, Callaghan et al. (1) estimated global bird population sizes by modeling the relationship between eBird reporting rates and independent estimates and extrapolating globally. While we applaud their intention, we caution that their modeling framework is prone to yield extremely uncertain and biased estimates that cannot support robust inferences about species abundance distributions or other applications in ecology, evolution, or conservation (1, 2).

Their methods yield extremely large posterior uncertainties for total global bird abundance (3.9 billion to 2,080 billion; figure 2 of ref. 1), and 96% of individual species had posterior uncertainty spanning three or more orders of magnitude. Glaucous Gull (*Larus hyperboreus*) was listed as the fifth-commonest bird globally; it is difficult to be confident in this conclusion given that the 95% credible interval (CI) for Glaucous Gull overlapped the CIs

for ~67% of all bird species. This uncertainty in species ordering makes it impossible to use these estimates for reliable conservation prioritization as suggested (1).

The tremendous uncertainty associated with the estimates of population size results from the inadequacy of the 10 measures used to account for imperfect detection of birds in eBird data (1), for which there is extreme inter- and intraspecific variation in the observation process across regions, time, and habitat (3). eBird reporting rates also depend heavily on species' overlap with the activity of eBird users, which also varies by region, time, and habitat.

In addition to high uncertainty, the approach also led to biased population estimates for many species. Abundance estimates (1) fell outside minimum–maximum ranges provided by BirdLife International for 81% of the 2,423 species with available estimates (27% below the minimum, 54% above the maximum) (4).^{*} Even the large uncertainty intervals repeatedly failed to cover known true values. For example, Swift Parrot (*Lathamus discolor*), which

^aCornell Laboratory of Ornithology, Cornell University, Ithaca, NY 14850; ^bUS Geological Survey Utah Cooperative Fish and Wildlife Research Unit, Department of Wildland Resources, Utah State University, Logan, UT 84322; ^cBirdLife International, Cambridge CB2 3QZ, United Kingdom; ^dBritish Trust for Ornithology, Thetford IP24 2PU, United Kingdom; ^eDepartment of Geography, University of Florida, Gainesville, FL 32611; ^fSchubot Avian Health Center, Department of Veterinary Pathobiology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, TX 77843; ^gBiodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Bentley, WA 6983, Australia; ^hDepartment of Zoology, University of Cambridge, Cambridge CB2 3EJ, United Kingdom; ⁱCentro de Ecología Aplicada del Litoral, Consejo Nacional de Investigaciones Científicas y Técnicas, 3400 Corrientes, Argentina; ^jDivision of Migratory Bird Management, US Fish and Wildlife Service, Hadley, MA 01035; ^kDepartment of Natural Sciences, Manchester Metropolitan University, Manchester M1 5GD, United Kingdom; ^lDepartment of Natural Sciences, Manchester Metropolitan University, Manchester M1 5GD, United Kingdom; ^mWorld Parrot Trust, Hayle TR27 4HY, United Kingdom; ⁿDepartment of Animal Ecology & Systematics, Justus Liebig University Giessen, D-35392 Giessen, Germany; ^oDepartment of Biological Sciences, University of Venda, Thohoyandou 0950, Republic of South Africa; ^pInstituto de Investigaciones en Biodiversidad y Medio Ambiente, Universidad Nacional del Comahue, Consejo Nacional de Investigaciones Científicas y Técnicas, 8400 Bariloche, Argentina; ^qEastern Ecological Science Center, US Geological Survey, Laurel, MD 20708; ^rDepartment of Conservation Biology, Estación Biológica de Doñana (CSIC), Sevilla, 41092 Spain; ^sCentro para el Estudio de Sistemas Marinos, Consejo Nacional de Investigaciones Científicas y Técnicas, U9120ACD Puerto Madryn, Argentina; ^tWildlife Conservation Society Argentina, Buenos Aires, C1426AKC Argentina; ^uDepartment of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90032; and ^vCentre for Research into Ecological & Environmental Modelling, School of Mathematics and Statistics, University of St Andrews, St Andrews KY16 9AJ, United Kingdom

Author contributions: O.J.R., J.B.S., and E.F.S. designed research; O.J.R., J.B.S., and E.F.S. performed research; and O.J.R., J.B.S., E.F.S., T.A., A.J.B., P.H.B.-S., D.J.B., A.H.B., S.H.M.B., C.L.D., A.M.D., A.S.D.G., A.F., D.F., W.M.H., P.E.H., F.A.L., A.C.L., S.M., R.M., R.O.M., J.F.M., E.T.M., Y.M., A.M., D.G.N., V.O., P.Q., J.A.R., V.R.-G., J.L.T., P.Y., C.Y., and A.J. wrote the paper.

The authors declare no competing interest.

This article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: ojr7@cornell.edu.

This article contains supporting information online at <http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2113862119/-DCSupplemental>.

Published March 1, 2022.

^{*}Population size estimates are given both on the “Summary” tab and the “Data table and detailed info” tab of the factsheet for each species at <http://datazone.birdlife.org/species/search>. They are also available through the IUCN Red List API at <https://api.v3.iucnredlist.org/> and the IUCN Red List Advanced Search at <https://www.iucnredlist.org/search>.

was recently assessed at 280 individuals (5), had a CI from 4,520 to 40 million (1). Spoon-billed Sandpiper (*Eurynorhynchus pygmeus*) has a population size of 490 individuals (95% CI 360 to 620) (6) but had a CI from 6,050 to 47 million (1). San Andres Vireo (*Vireo caribaeus*), used in model training with a population size of 2,500 to 10,000 mature individuals (4), was estimated as extinct (1).

Regional differences in reporting rates create bias, because, as noted by Callaghan et al., the 7% of species used to train the model were heavily biased toward Europe and North America (1). Density imputation based on spatially uneven calibration estimates biases the population estimates to an unknown and

inestimable extent, with downstream influence on the shapes of species abundance distributions and ecological conclusions.

For species with sufficient data quantity and quality, citizen and community science data can produce reliable density estimates (3), and methods for such analyses are constantly improving (7, 8). However, no method currently exists to estimate global population sizes across species while accounting sufficiently for known sources of variation in eBird reporting rates. Meaningful global population estimates would represent a tremendous advance for ecology, evolutionary biology, and conservation but will require considerably more nuanced analysis of globally available data.

-
- 1 C. T. Callaghan, S. Nakagawa, W. K. Cornwell, Global abundance estimates for 9,700 bird species. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2023170118 (2021).
 - 2 A. V. Harry, J. M. Braccini, Caution over the use of ecological big data for conservation. *Nature* **595**, E17–E19 (2021).
 - 3 US Fish and Wildlife Service, "Final report: Bald eagle population size: 2020 update" (Division of Migratory Bird Management, US Fish and Wildlife Service, Washington, DC, 2020).
 - 4 BirdLife International, IUCN red list for birds. <http://datazone.birdlife.org/species/search>. Accessed 29 June 2021.
 - 5 G. Olah, D. Stojanovic, M. H. Webb, R. S. Waples, R. Heinsohn, Comparison of three techniques for genetic estimation of effective population size in a critically endangered parrot. *Anim. Conserv.* **24**, 491–498 (2021).
 - 6 R. E. Green et al., New estimates of the size and trend of the world population of the Spoon-billed Sandpiper using three independent statistical models. *Wader Study* **128**, 22–35 (2021).
 - 7 A. Johnston et al., Analytical guidelines to increase the value of community science data: An example using eBird to estimate species distributions. *Divers. Distrib.* **27**, 1265–1277 (2021).
 - 8 D. Fink et al., eBird status and trends (data version 2019, released 2020, Cornell Lab of Ornithology, Ithaca, NY).