

ARTICLE OPEN



Polygenic risk modeling for prediction of epithelial ovarian cancer risk

© The Author(s) 2021

Polygenic risk scores (PRS) for epithelial ovarian cancer (EOC) have the potential to improve risk stratification. Joint estimation of Single Nucleotide Polymorphism (SNP) effects in models could improve predictive performance over standard approaches of PRS construction. Here, we implemented computationally efficient, penalized, logistic regression models (lasso, elastic net, stepwise) to individual level genotype data and a Bayesian framework with continuous shrinkage, “select and shrink for summary statistics” (S4), to summary level data for epithelial non-mucinous ovarian cancer risk prediction. We developed the models in a dataset consisting of 23,564 non-mucinous EOC cases and 40,138 controls participating in the Ovarian Cancer Association Consortium (OCAC) and validated the best models in three populations of different ancestries: prospective data from 198,101 women of European ancestries; 7,669 women of East Asian ancestries; 1,072 women of African ancestries, and in 18,915 *BRCA1* and 12,337 *BRCA2* pathogenic variant carriers of European ancestries. In the external validation data, the model with the strongest association for non-mucinous EOC risk derived from the OCAC model development data was the S4 model (27,240 SNPs) with odds ratios (OR) of 1.38 (95% CI: 1.28–1.48, AUC: 0.588) per unit standard deviation, in women of European ancestries; 1.14 (95% CI: 1.08–1.19, AUC: 0.538) in women of East Asian ancestries; 1.38 (95% CI: 1.21–1.58, AUC: 0.593) in women of African ancestries; hazard ratios of 1.36 (95% CI: 1.29–1.43, AUC: 0.592) in *BRCA1* pathogenic variant carriers and 1.49 (95% CI: 1.35–1.64, AUC: 0.624) in *BRCA2* pathogenic variant carriers. Incorporation of the S4 PRS in risk prediction models for ovarian cancer may have clinical utility in ovarian cancer prevention programs.

European Journal of Human Genetics (2022) 30:349–362; <https://doi.org/10.1038/s41431-021-00987-7>

INTRODUCTION

Rare variants in known high and moderate penetrance susceptibility genes (*BRCA1*, *BRCA2*, *BRIP1*, *PALB2*, *RAD51C*, *RAD51D* and the mis-match repair genes) account for about 40% of the inherited component of EOC disease risk [1, 2]. Common susceptibility variants, reviewed in Kar et al. and Jones et al., explain about 6% of the heritability of EOC [1, 3]. Polygenic risk scores (PRS) provide an opportunity for refined risk stratification in the general population and in carriers of rare moderate or high risk alleles.

A PRS is calculated as the weighted sum of the number of risk alleles carried for a specified set of variants. The best approach to identify the variant set and their weights to optimize the predictive power of a PRS is unknown. A common approach involves selecting a set of variants that reach a threshold for association based on the *p*-value for each variant with or without pruning to remove highly correlated variants [4, 5]. More complex machine learning approaches that do not assume variant independence have also been used [6, 7], but these methods have produced only modest gains in predictive power for highly polygenic phenotypes [6, 8]. Penalized regression approaches such as the lasso, elastic net and the adaptive lasso have also been used with individual level data [9], but a major drawback is the computational burden required to fit the models [9, 10].

We present novel, computationally efficient PRS models using two approaches: (1) penalized regression models including the

lasso, elastic net and minimax concave penalty (MCP) for use with individual genotype data; and (2) a Bayesian regression model with continuous shrinkage priors for use where only summary statistics are available—referred to as the “select and shrink with summary statistics” (S4) method. We compare these models with two commonly used methods, stepwise regression with *p*-value thresholding and LDPred.

MATERIALS (SUBJECTS) AND METHODS

Model development study population

EOC is a highly heterogeneous phenotype with five major histotypes for invasive disease—high-grade serous, low-grade serous, endometrioid, clear cell, and mucinous histotype. The mucinous histotype is the least common and its origin is the most controversial with up to 60% of diagnosed cases of mucinous ovarian cancer often being misdiagnosed metastasis from non-ovarian sites [11]. Therefore, in this study, we performed PRS modeling and association testing for all cases of invasive, non-mucinous EOC. We used genotype data from 23,564 invasive non-mucinous EOC cases and 40,138 controls with >80% European ancestries from 63 case-control studies included in the Ovarian Cancer Association Consortium (OCAC) for model development. The distribution of cases by histotype was high-grade serous (13,609), low-grade serous (2,749), endometrioid (2,877), clear cell (1,427), and others (2,902). Sample collection, genotyping, and

A full list of authors and their affiliations appears at the end of the paper.

Received: 7 December 2020 Revised: 9 August 2021 Accepted: 27 September 2021

Published online: 14 January 2022

quality control have been previously described [12]. Genotype data were imputed to the Haplotype Reference Consortium reference panel using 470,825 SNPs that passed quality control. Of the 32 million SNPs imputed, 10 million had imputation $r^2 > 0.3$ and were included in this analysis.

Model validation study populations

We validated the best-fitting PRS models developed in the OCAC data in 657 prevalent and incident cases of invasive, non-mucinous EOC and 198,101 female controls of European ancestries from the UK Biobank. Samples were genotyped using either the Affymetrix UK BiLEVE Axiom Array or Affymetrix UK Biobank Axiom Array (which share 95% marker content), and then imputed to a combination of the Haplotype Reference Consortium, the 1000 Genomes phase 3 and the UK10K reference panels [13]. We restricted analysis to genetically confirmed females of European ancestries. We excluded individuals if they were outliers for heterozygosity, had low genotyping call rate <95%, had sex chromosome aneuploidy, or if they were duplicates (cryptic or intended) [12]. All SNPs selected in the model development phase were available in the UK Biobank.

We investigated transferability of the best-fitting PRS models to populations of non-European ancestries using genotype data from females of East Asian and African ancestries genotyped as part of the OCAC OncoArray Project [14, 15]. Women of East Asian ancestries—2,841 non-mucinous invasive EOC and 4,828 controls—were identified using a criterion of >80% Asian ancestries. This included samples collected from studies in China, Japan, Korea,

and Malaysia as well as samples collected from women of Asian ancestry in studies conducted in the US, Europe and Australia [14]. Similarly, women of African ancestries—368 cases of non-mucinous invasive EOC and 704 controls—mainly from studies conducted in the US, were identified using a criterion of >80% African ancestries as described previously [15].

We also assessed the performance of the best-fitting PRS models in women of European ancestries (>80% European ancestries) with the pathogenic *BRCA1* and *BRCA2* variants from the Consortium of Investigators of Modifiers of *BRCA1/2* (CIMBA). We used genotype data from 18,915 *BRCA1* (2,053 invasive EOC cases) and 12,337 *BRCA2* (717 invasive EOC cases) pathogenic variant carriers from 63 studies contributing to CIMBA [16]. Genotyping, data quality control measures, intercontinental ancestries assessment and imputation to the HRC reference panel are as described for the OCAC study population.

STATISTICAL ANALYSIS

Polygenic risk models

For all PRS models, we created scores as linear functions of the allele dosage in the general form $PRS_i = \sum_j^p x_{ij}\beta_j$ where genotypes are denoted as x (taking on the minor allele dosages of 0, 1, and 2), with x_{ij} representing the i th individual for the j th SNP (out of p SNPs) on an additive log scale and β_j represents the weight—the log of the odds ratio—of the j th SNP. We used different approaches to select and derive the optimal weights, β_j , in models as described below.




Fig. 1 PRS model development using penalized regression and LDpred Bayesian approach. Shown in the left panel is the two-stage approach with five-fold cross validation used for individual level genotype data while the right panel shows the LDpred approach used for summary level data.

Penalized logistic regression models

A penalized logistic regression model for a set of SNPs aims to identify a set of regression coefficients that minimize the regularized loss function given by

$$plr(x; \lambda, \kappa) = \begin{cases} x - \lambda \text{sign}(x) / (1 - \kappa) & \text{if } |x| < \lambda / \kappa \text{ and } |(x)| > \lambda \\ x & \text{if } |x| \geq \frac{\lambda}{\kappa} \\ 0 & \text{if } |(x)| < \lambda \end{cases}$$

where x is the effect estimate of a SNP, λ is the tuning parameter and κ is the threshold (penalty) for different regularization paths. λ and κ are parameters that need to be chosen during model development to optimize performance. The lasso, elastic net, MCP, and p -value thresholds are instances of the function with different κ values. We minimized the winner's curse effect on inflated effect estimates for rare SNPs by penalizing rarer SNPs more heavily than common SNPs. Details are provided in the Supplementary Methods.

We used a two-stage approach to reduce computational burden without a corresponding loss in predictive power. The first stage was a SNP selection stage using a sliding windows approach, with 5.5 Mb data blocks and a 500 kb overlap between blocks. SNP selection was performed for each block and selected SNPs were collated. Single SNP association analyses were then run, and all SNPs with a χ^2 test statistic of less than 2.25 were excluded. The 2.25 cutoff was arbitrary and selected to maximize computational efficiency without loss in predictive power. Penalized regression models were applied to the remaining SNPs using λ values of 3.0 and κ values of 0.0, 0.2, 0.4, 0.6, 0.8 and 1.0. SNPs selected in any of these models were included in subsequent analyses. In the second stage, we fit penalized regression models to the training dataset with λ values ranging from 3.0 to 5.5 in increments of 0.1 iterated over κ values from -3.0 to 1 in increments of 0.1. The lasso model ($\kappa = 0$) for each value of λ was fitted first, to obtain a unique maximum. From the fitted maximum the κ value was changed, and the model refitted.

We applied this two-stage approach with five-fold cross-validation (Fig. 1). In each iteration, the data set was split into five, with one part constituting the test data and the other four constituting the training data. The variants and their weights from the two-stage penalized logistic regression modeling in the training data were used to calculate the area under the receiver operating characteristic curve (AUC) in the test data in each iteration. AUC estimates for each combination of λ and κ were obtained. We repeated this process for each cross-validation iteration to obtain a mean AUC for each combination of λ and κ . Finally, we selected the tuning and threshold parameters from the lasso, elastic net and MCP models with the maximum mean cross-validated AUC and fitted penalized logistic regression models with these parameters to the entire OCAC dataset to obtain SNP weights for PRS scores.

Stepwise logistic regression with variable P -value threshold

This model is a general PLR model with $\kappa = 1$. As with the other PLR models, we investigated various values for λ values (corresponding to a variable P -value threshold for including a SNP in the model). However, we observed that the implementation of this model on individual level data was more difficult than for other κ values because the model would sometimes converge to a local optimum rather than the global optimum. Therefore, we applied an approximate conditional and joint association analysis using summary level statistics correcting for estimated LD between SNPs, and utilizing a reference panel of 5,000 individual level genotype OCAC data as described in Yang et al. [17]. Details are provided in the Supplementary Methods.

LDPred

LDPred is a Bayesian approach that shrinks the posterior mean effect size of each marker based on a point-normal prior and LD

information from an external reference panel. We derived seven candidate PRSs assuming the fractions of associated variants were 0.001, 0.003, 0.01, 0.03, 0.1, 0.3, and 1.0 respectively using the default parameters as detailed in Vilhjálmsdóttir et al. [18] and an LD reference panel of 503 samples of European ancestries from the 1000 Genomes phase 3 release with effect estimates from the OCAC model development data.

Select and shrink using summary statistics (S4)

The S4 algorithm is similar to the PRS-CS algorithm [19]—a Bayesian method that uses summary statistics and between-SNP correlation data from a reference panel to generate the PRS scores by placing a continuous shrinkage prior on effect sizes. We adapted this method with penalization of rarer SNPs by correcting for the standard deviation resulting in the selection of fewer SNPs. We varied three parameters, a , b , ϕ , which control the degree of shrinkage of effect estimates. Φ , the overall shrinkage parameter, is influenced by values of a which controls shrinkage of effect estimates around 0 and b which control shrinkage of larger effect estimates. We generated summary statistics for each cross-validation training set and selected the parameters that gave the best results on average from the cross-validation and applied these to the set of summary statistics for the complete OCAC data set to obtain the final set of weights.

PRS based on meta-analysis of OCAC-CIMBA summary statistics

We conducted a meta-analysis of the EOC associations in *BRCA1* variant carriers, *BRCA2* variant carriers and the participants participating in OCAC (see Supplementary Methods) and constructed two PRS models. An S4 PRS was generated by applying the a , b and ϕ parameters from the S4 model described above. A stepwise PRS was generated by selecting all SNPs that were genome-wide significant ($p < 5 \times 10^{-8}$) in the meta-analysis, along with any independent signals in the same region with $p < 10^{-5}$ from the histotype specific analyses for low-grade serous, high-grade serous, endometrioid, clear cell ovarian cancer and non-mucinous invasive EOC.

Polygenic risk score performance

The best lasso, elastic net, stepwise and S4 models from the model development stage were validated using two independent data sources: the UK Biobank data and *BRCA1/BRCA2* pathogenic variant carriers from the CIMBA. In the UK Biobank data, we evaluated discriminatory performance of the models using the AUC and examined the association between standardized PRS and risk of non-mucinous EOC using logistic regression analysis. For the CIMBA data, we assessed associations for each version of the PRS and invasive non-mucinous EOC risk using weighted Cox regression methods [20]. PRSs in the CIMBA data were scaled to the same PRS standard deviations as the OCAC data, meaning that per standard deviation hazard ratios estimated on CIMBA data are comparable to PRS associations in the OCAC and UK Biobank data. The regression models were adjusted for birth cohort (<1920, 1920–1929, 1930–1939, 1940–1949, ≥ 1950) and the first four ancestries informative principal components (calculated separately by iCOGS/OncoArray genotyping array) and stratified by Ashkenazi Jewish ancestries and country. Absolute risks by PRS percentiles adjusting for competing risks of mortality from other causes were calculated as described in the Supplementary Material.

Transferability of PRS scores to non-European ancestries

We implemented two straightforward approaches to disentangle the role of ancestries on polygenic risk scoring. We selected homogenous ancestral samples by using a high cut-off criterion of 80% ancestries and we standardized the PRSs by mean-centering within each population. These approaches led to a more uniform

distribution of PRSs within each ancestral population. Further adjustments using principal components of ancestries did not attenuate risk estimates.

RESULTS

Model development

The results for the models based on individual level genotype data are shown in Table 1. The elastic net model had the best predictive accuracy ($AUC = 0.586$). The optimal value of λ obtained from regularization paths for the MCP model was 3.3 meaning the best MCP model was equivalent to the lasso model. The best-fitting model based on summary statistics was the S4 ($AUC = 0.593$) and the LDPred model had the poorest performance of the methods tested ($AUC = 0.552$). Therefore, the LDPred model was not considered for further validation in other datasets. All SNPs selected and the associated weights for each model are provided in Supplementary Tables 1–6.

Model validation in women of European ancestries

Overall the PLR models performed slightly better in the UK Biobank data than the model development data (Table 2). Of the models developed using the OCAC model development data, the association was strongest with the S4 PRS. In *BRCA1* and *BRCA2* variant carriers, prediction accuracy was generally higher among *BRCA2* carriers than *BRCA1* carriers. Consistent with results from the general population in the UK Biobank, the S4 PRS model also had the strongest association and predictive accuracy for invasive

EOC risk in both *BRCA1* and *BRCA2* carriers. Sensitivity analyses were conducted in which the unadjusted models for *BRCA1* and *BRCA2* carriers were progressively adjusted for birth cohort and 6 principal components. There was little difference in HR estimates and association *P*-values going from the unadjusted model to the model adjusting for six principal components (Supplementary Table 7). The PRS models developed using the OCAC-CIMBA meta-analysis results had better discriminative ability in the UK Biobank than the PRS models developed using only OCAC data. Compared with the S4 PRS using only OCAC data, the S4 PRS model derived from the meta-analysis had fewer SNPs, a stronger association with invasive EOC risk and better predictive accuracy. Similarly, the stepwise model from the OCAC-CIMBA meta-analysis performed better than the stepwise model from only OCAC data, but included more SNPs.

The observed distribution of the OR estimates within centiles of the PRS distribution were consistent with ORs from predicted values under the assumption that all SNPs interact multiplicatively (Fig. 2), with all 95% confidence intervals intersecting with the theoretical estimates for women of European ancestries. Compared with women in the middle quintile, women of European ancestry (UK Biobank) in the top 95th percentile of the lasso derived PRS model had a 2.23-fold increased odds of non-mucinous EOC (95% CI: 1.64–3.02) (Table 3).

Absolute risk of developing ovarian cancer by PRS percentiles

We estimated cumulative risk of EOC within PRS percentiles for women in the general population (Fig. 3), by applying the odds

Table 1. Performance of different PRS models in five-fold cross-validation of OCAC data.

Model	Number of SNPs ^a	Tuning parameter for best performance	AUC	OR per 1 SD of PRS	95% CI
(a) Models based on individual level genotype data					
Lasso	1403	$\lambda = 3.3$	0.583	1.35	1.30–1.39
Elastic net	10,797	$\lambda = 3.3, \kappa = -2.2$	0.586	1.36	1.31–1.40
MCP	1403	$\lambda = 3.3$	0.583	1.35	1.30–1.39
(b) Models based on summary statistics					
LDPred	5,291,719	$\rho = 0.001$	0.552	1.21	1.13–1.29
Stepwise	22	$\lambda = 5.4$	0.572	1.30	1.26–1.34
Select and Shrink (OCAC)	27,240	$a = 2.75, b = 2, \varphi = 3e-6$	0.593	1.39	1.34–1.44

AUC area under the receiver operating characteristic (ROC) curve AUC, OR odds ratio, SD standard deviation, PRS polygenic risk score, CI confidence interval, NA not applicable.

^aNumber of SNPs in PRS model run on full OCAC data set after selection of model parameters.

Table 2. External validation of PRS models in European populations using data from UK Biobank and CIMBA.

Model (data set)	SNPs	UK Biobank			CIMBA <i>BRCA1</i> carriers ^a			CIMBA <i>BRCA2</i> carriers ^a		
		AUC	OR	95% CI	AUC	HR	95% CI	AUC	HR	95% CI
(a) PRS models based on OCAC data										
Lasso (OCAC)	1403	0.587	1.37	1.27–1.48	0.573	1.27	1.21–1.34	0.627	1.48	1.33–1.63
Elastic net (OCAC)	10,797	0.588	1.36	1.26–1.47	0.583	1.32	1.26–1.39	0.617	1.47	1.33–1.63
Stepwise (OCAC)	22	0.588	1.35	1.26–1.46	0.563	1.21	1.16–1.26	0.605	1.39	1.26–1.54
Select and shrink (OCAC)	27,240	0.588	1.38	1.28–1.48	0.592	1.36	1.29–1.43	0.624	1.49	1.35–1.64
(b) PRS models based on meta-analysis of OCAC and CIMBA data										
Stepwise (OCAC-CIMBA) ^b	36	0.595	1.39	1.29–1.50	NA	NA	NA	NA	NA	NA
Select and shrink (OCAC-CIMBA)	18,007	0.596	1.42	1.32–1.54	NA	NA	NA	NA	NA	NA

AUC area under the receiver operating characteristic curve, OR odds ratio, HR hazards ratio.

^aEstimates are from unadjusted models.

^bResults in CIMBA are overfitted as the CIMBA data was used for model development.




Fig. 2 Association between the PLR PRS models and non-mucinous ovarian cancer by PRS percentiles. Shown are estimated odds ratios (OR) and confidence intervals for women of European ancestries by percentiles of polygenic risk scores derived from lasso (A), elastic net (B), stepwise (C) and S4 (D) models relative to the middle quintile.

ratio from the PRS models to age-specific population incidence and mortality data for England in 2016. For *BRCA1* and *BRCA2* pathogenic variant carriers, we applied the estimated hazard ratios from PRS models to age-specific incidence rates obtained from Kuchenbaecker et al. [21]. For women in the general population, the estimated cumulative risks of EOC by age 80 for women at the 99th centile of the PRS distribution were 2.24%, 2.18%, 2.54%, and 2.81% for the lasso, elastic net, stepwise and S4 models, respectively. In comparison, the absolute risks of EOC by age 80 for women at the 1st centile were 0.76%, 0.78%, 0.64%, and 0.56% for the lasso, elastic net, stepwise and S4 models, respectively.

The absolute risks of developing EOC in *BRCA1* and *BRCA2* pathogenic variant carriers were considerably higher than for women in the general population (Figs. S1 and S2). The estimated absolute risk of developing ovarian cancer by age 80 for *BRCA1* carriers at the 99th PRS centiles were 63.2%, 66.3%, 59.0%, and 68.4% for the lasso, elastic net, stepwise and S4 models, respectively. The corresponding absolute risks for women at the 1st PRS centile were 27.7%, 25.6%, 30.8%, and 24.2%. For *BRCA2* carriers the absolute risks for women at the 99th centile were

36.3%, 36.3%, 33.0%, and 36.9%; and 7.10%, 7.12%, 8.24%, and 6.92% at the 1st centile for the lasso, elastic net, stepwise and S4 models, respectively.

PRS distribution and ancestries

To investigate the transferability of the PRS to other populations, we applied the scores to women of African ($N = 1,072$) and Asian ($N = 7,669$) ancestries genotyped as part of the OncoArray project. In general, the distributions of the raw PRS were dependent on both the statistical methods used in SNP selection and ancestral group. PRS models that included more variants had less dispersion, such that the elastic net models had the least between individual variation in all ancestral groups (standard deviation = 0.15, 0.19, and 0.22 for individuals of Asian, African and European ancestries respectively), while the distributions from the stepwise models were the most dispersed (standard deviation = 0.23, 0.27, and 0.30 for individuals of Asian, African and European ancestries respectively). As expected, given the variation in variant frequencies by population, the distribution of polygenic scores was significantly different across the three ancestral groups, with the least dispersion among women of Asian ancestries and the most

Table 3. Association between polygenic risk scores and non-mucinous EOC by PRS percentiles and ancestry.

Percentile	UK Biobank				East Asian				African			
	Controls (n)	Cases (n)	OR (95% CI)	Controls (n)	Cases (n)	OR (95% CI)	Controls (n)	Cases (n)	OR (95% CI)	Controls (n)	Cases (n)	OR (95% CI)
(a) Lasso												
0-5	9880	12	0.42 (0.22-0.72)	278	106	0.65 (0.51-0.83)	35	19	0.89 (0.47-1.65)			
5-10	9870	24	0.83 (0.52-1.27)	271	112	0.71 (0.55-0.90)	41	13	0.52 (0.25-1.01)			
10-20	19,733	53	0.92 (0.66-1.27)	487	280	0.98 (0.82-1.18)	81	26	0.53 (0.31-0.88)			
20-40	39,468	104	0.90 (0.69-1.18)	993	541	0.93 (0.80-1.08)	154	60	0.64 (0.42-0.99)			
40-60	39,457	115	1	967	566	1	133	81	1			
60-80	39,425	147	1.28 (1.00-1.64)	941	593	1.08 (0.93-1.25)	136	78	0.94 (0.64-1.39)			
80-90	19,699	87	1.52 (1.14-2.00)	466	301	1.10 (0.92-1.32)	63	44	1.15 (0.71-1.84)			
90-95	9842	51	1.78 (1.27-2.46)	214	169	1.35 (1.07-1.69)	34	20	0.97 (0.51-1.78)			
95-100	9830	64	2.23 (1.64-3.02)	211	173	1.40 (1.12-1.76)	27	27	1.64 (0.90-3.00)			
(b) Elastic net												
0-5	9876	17	0.67 (0.39-1.09)	277	107	0.72 (0.56-0.92)	35	19	0.90 (0.47-1.64)			
5-10	9876	17	0.67 (0.39-1.09)	271	112	0.78 (0.61-0.99)	41	13	0.52 (0.25-1.01)			
10-20	19,740	45	0.89 (0.62-1.26)	497	270	1.02 (0.85-1.22)	81	26	0.53 (0.31-0.88)			
20-40	39,453	120	1.19 (0.91-1.55)	967	567	1.10 (0.95-1.28)	154	60	0.64 (0.42-0.96)			
40-60	39,471	101	1	1000	533	1	133	81	1			
60-80	39,413	159	1.58 (1.23-2.03)	926	608	1.23 (1.06-1.43)	136	78	0.94 (0.64-1.39)			
80-90	19,695	91	1.80 (1.36-2.40)	457	310	1.27 (1.06-1.52)	63	44	1.15 (0.71-1.84)			
90-95	9841	52	2.07 (1.47-2.87)	226	157	1.30 (1.04-1.64)	34	20	0.97 (0.51-1.78)			
95-100	9839	55	2.18 (1.56-3.02)	207	177	1.60 (1.28-2.01)	27	27	1.64 (0.90-3.00)			
(c) Stepwise												
0-5	9880	13	0.39 (0.21-0.67)	254	130	0.90 (0.71-1.14)	40	14	0.75 (0.37-1.44)			
5-10	9874	19	0.57 (0.34-0.91)	268	115	0.76 (0.59-0.96)	43	11	0.55 (0.26-1.10)			
10-20	19,742	44	0.67 (0.47-0.93)	494	273	0.98 (0.81-1.17)	80	27	0.72 (0.42-1.21)			
20-40	39,470	102	0.77 (0.60-1.00)	970	564	1.03 (0.89-1.19)	142	72	1.09 (0.73-1.63)			
40-60	39,440	132	1	979	564	1	146	68	1			
60-80	39,414	158	1.20 (0.95-1.51)	951	583	1.08 (0.94-1.25)	130	84	1.39 (0.93-2.07)			
80-90	19,697	88	1.33 (1.02-1.75)	456	311	1.21 (1.01-1.44)	61	46	1.62 (1.00-2.61)			
90-95	9853	41	1.24 (0.86-1.75)	236	147	1.10 (0.87-1.38)	35	19	1.17 (0.61-2.17)			
95-100	9834	60	1.82 (1.33-2.46)	220	164	1.32 (1.04-1.65)	27	27	2.15 (1.17-3.95)			
(d) Select and shrink												
0-5	9957	16	0.54 (0.31-0.89)	279	105	0.63 (0.49-0.81)	38	16	0.71 (0.36-1.33)			
5-10	9888	15	0.51 (0.29-0.85)	254	129	0.85 (0.67-1.08)	41	13	0.53 (0.26-1.03)			
10-20	19,812	51	0.87 (0.62-1.20)	489	278	0.96 (0.80-1.14)	81	26	0.54 (0.32-0.90)			
20-40	39,435	113	0.97 (0.75-1.25)	1013	521	0.86 (0.75-1.00)	156	58	0.62 (0.41-0.94)			
40-60	39,512	117	1	961	572	1	134	80	1			
60-80	39,316	158	1.36 (1.07-1.73)	950	584	1.03 (0.89-1.20)	137	77	0.94 (0.63-1.40)			

Table 3 continued

Percentile	UK Biobank			East Asian			African		
	Controls (n)	Cases (n)	OR (95% CI)	Controls (n)	Cases (n)	OR (95% CI)	Controls (n)	Cases (n)	OR (95% CI)
80–90	19,718	77	1.32 (0.98–1.76)	434	333	1.29 (1.08–1.54)	61	46	1.26 (0.79–2.02)
90–95	9791	45	1.55 (1.09–2.17)	233	150	1.08 (0.86–1.36)	30	24	1.34 (0.73–2.45)
95–100	9775	65	2.25 (1.65–3.03)	215	169	1.32 (1.05–1.66)	26	28	1.80 (0.99–3.31)

OR odds ratio, CI confidence interval.

variation in women of European ancestries. The difference in PRS distribution was minimized after correction for ancestry by standardizing the PRS to have unit standard deviation using the control subjects for each ancestral group.

High PRSs were significantly associated with risk of non-mucinous EOC in both Asian and African ancestries (Table 4), although the effects were weaker than in women of European ancestries. For example, with the lasso model, the odds ratio per unit standard deviation increment in polygenic score was 1.16 (95% CI: 1.11–1.22) in women of East Asian ancestries, 1.28 (95% CI: 1.13–1.45) in women of African ancestries and 1.37 (95% CI: 1.27–1.48) in women of European ancestries (p for heterogeneity <0.0001). Variability in effect sizes among ancestral groups was highest for the stepwise model ($I^2 = 92%$) versus 84% and 83% for elastic net and lasso derived polygenic scores respectively. The best discriminative model among women of East Asian and African ancestries were the elastic net PRS (AUC = 0.543) and the S4 PRS derived from OCAC-CIMBA meta-analysis (AUC = 0.596) respectively. Women of African ancestries in the top 5% of the PRS had about two-fold increased risk compared to women in the middle quintile (lasso OR: 1.64, 95% CI: 0.90–3.00; elastic net OR: 1.64, 95% CI: 0.90–3.00; stepwise OR: 2.15, 95% CI: 1.17–3.95; S4 OR: 1.80, 95% CI: 0.99–3.31) (Table 3). Effect estimates were smaller in women of East Asian ancestries with women in the top 5% of the PRS, having about a 1.5 fold increased risk compared to women in the middle quintile (lasso OR: 1.40, 95% CI: 1.12–1.76; elastic net OR: 1.60, 95% CI: 1.28–2.01; stepwise OR: 1.32, 95% CI: 1.04–1.65; S4 OR: 1.32, 95% CI: 1.05–1.66) (Table 3).

DISCUSSION

Genetic risk profiling with PRSs has led to actionable outcomes for cancers such as breast and prostate [22, 23]. Previous PRS scores for invasive EOC risk in the general population and *BRCA1/BRCA2* pathogenic variant carriers have been based on genetic variants for which an association with EOC risk had been established at nominal genome-wide significance [20, 24, 25]. Here, we explored the predictive performance of computationally efficient, penalized, regression methods in modeling joint SNP effects for EOC risk prediction in diverse populations and compared them with common approaches. By leveraging the correlation between SNPs which do not reach nominal genome-wide thresholds and including them in PRS models, the PRSs derived from penalized regression models provide stronger evidence of association with risk of non-mucinous EOC than previously published PRSs in both the general population and in *BRCA1/BRCA2* pathogenic variant carriers.

Recently, Barnes et al. derived a PRS score using 22 SNPs that were significantly associated with high-grade serous EOC risk (PRS_{HGS}) to predict EOC risk in *BRCA1/BRCA2* pathogenic variant carriers [20]. To make effect estimates obtained in this analysis comparable to the effect estimates obtained from the PRS_{HGS}, we standardized all PRSs using the standard deviation from unaffected *BRCA1/BRCA2* carriers and provide estimates which are directly comparable to the PRS_{HGS} in Supplementary Table 9. All PRS models in this analysis except the Stepwise (OCAC only) had higher effect estimates [20]. The AUC estimates from the adjusted PLR methods implemented in this analysis, are higher than the corresponding PRS_{HGS} estimates for *BRCA1* carriers (0.604). In *BRCA2* carriers, the AUC estimates for the lasso and S4 models did slightly better than the PRS_{HGS} AUC estimate (0.667), while the stepwise did slightly worse and the elastic net estimate was comparable. The AUC estimates for women in the general population, as estimated from the UK Biobank, are slightly higher than estimates from previously published PRS models for overall EOC risk by Jia et al. (AUC = 0.57) and Yang et al. (AUC = 0.58) [25, 26].

The level of risk for women above the 95th percentile of the PRS is similar to that conferred by pathogenic variants in moderate penetrance genes such as *FANCM* (RR = 2.1, 95% CI = 1.1–3.9) and




Fig. 3 Cumulative risk of ovarian cancer between birth and age 80 by PRS percentiles and PRS models. Shown are the cumulative risk of ovarian cancer risk in UK women by polygenic risk score percentiles. The lasso (A) and elastic net (B) penalized regression models were applied to individual level genotype data, while the stepwise (C) and S4 (D) models were applied to summary level statistics. Note that the median and the mean risk differ because the distribution of the relative risk in the population is left-skewed (the log relative risk is a Normal distribution).

Table 4. External validation of PRS models in East Asian and African Populations.

Model	East Asian ancestries			African ancestries		
	AUC	OR	95% CI	AUC	OR	95% CI
Lasso	0.541	1.16	(1.11–1.22)	0.576	1.28	(1.13–1.45)
Elastic net	0.543	1.17	(1.12–1.23)	0.574	1.29	(1.14–1.47)
Stepwise (OCAC)	0.528	1.11	(1.06–1.16)	0.581	1.34	(1.18–1.52)
Select and shrink (OCAC)	0.538	1.14	(1.08–1.19)	0.593	1.38	(1.21–1.58)
Stepwise (OCAC-CIMBA)	0.542	1.17	(1.11–1.23)	0.594	1.37	(1.20–1.56)
Select and shrink (OCAC-CIMBA)	0.537	1.14	(1.08–1.19)	0.596	1.41	(1.23–1.61)

PALB2 (RR = 2.91 95% CI = 1.40–6.04) [27, 28]. The inclusion of other risk factors such as family history of ovarian cancer, presence of rare pathogenic variants, age at menarche, oral contraceptive use, hormone replacement therapy, parity, and endometriosis in combination with the PRS could potentially improve risk stratification as implemented in the CanRisk tool (www.canrisk.org), which currently uses a 36-SNP PRS with the potential to use other PRS models [29, 30].

We found that the discrimination of the PRS varied by ancestry with greater discrimination in women of European ancestries than in women of African and East Asian ancestries. The better performance in African than East Asian populations is in contrast to what one would expect given human demographic history, and the performance of PRS for other phenotypes in African populations. This may simply be the play of chance given the small number of samples of African ancestries. Alternatively it reflects the fact that the allele frequencies of the PRS SNPs were more similar between the African and European populations than they were with the East Asian population (Supplementary Tables 10–14).

Further optimization of the models could be achieved by varying the penalization function based on prior knowledge. For

example, varying the penalty function to select more SNPs from genomic regions with known susceptibility variants given that susceptibility variants tend to cluster together. Alternatively, the penalty functions could be modified to incorporate information about functionally active regions of the genome such as promoters, enhancers, and transcription factor binding sites. However, incorporating functional annotation has resulted in limited gains in prediction accuracy for complex traits such as breast cancer, celiac disease, type 2 diabetes, and rheumatoid arthritis [31].

Machine/deep learning approaches are alternative ways to constructing PRS, but methods such as the neural net, support vector machine, and random forest have been shown to be computationally prohibitive or produce inferior results to other approaches [32, 33]. Other machine learning methods, such as those based on gradient boosting do not perform well in genomic regions where strong genetic interactions are present, for which alternative approaches such as the LDpred may perform better [18]. Our approach has several benefits over alternative machine learning methods, including its simplicity, and intrinsic robustness to minor misspecification of LD or association strength.

In conclusion, our results indicate that using the lasso model for individual level genotype data and the S4 model for summary level data in PRS construction provide an improvement in risk prediction for non-mucinous EOC over more common approaches. Our approach overcomes the computational limitations in the use of penalized methods for large-scale genetic data, particularly in the presence of highly correlated SNPs and when the use of cross-validation for parameter estimation is preferred. In practical terms, the PRS provides sufficient discrimination, particularly for women of European ancestries, to be considered for inclusion in risk prediction and prevention approaches for EOC in the future. Further studies are required to optimize these PRSs in ancestrally diverse populations and to validate their performance with the inclusion of other genetic and lifestyle risk factors.

DATA AVAILABILITY

OncoArray germline genotype data for the OCAC studies have been deposited at the European Genome-phenome Archive (EGA; <https://ega-archive.org/>), which is hosted by the EBI and the CRG, under accession EGAS00001002305. Summary statistics for the Ovarian Cancer Association Consortium are available in the NHGRI-EBI GWAS catalogue (<https://www.ebi.ac.uk/gwas/home>) under the accession number GCST90016665. A subset of the OncoArray germline genotype data for the CIMBA studies are publicly available through the database of Genotypes and Phenotypes (dbGaP) under accession phs001321.v1.p1. The complete data set will not be made publicly available because of restraints imposed by the ethics committees of individual studies; requests for further data can be made to the Data Access Coordination Committee (<http://cimba.ccg.medschl.cam.ac.uk/>)

REFERENCES

- Jones MR, Kamara D, Karlan BY, Pharoah PDP, Gayther SA. Genetic epidemiology of ovarian cancer and prospects for polygenic risk prediction. *Gynecol Oncol*. 2017;147:705–13.
- Lyra PCM, Rangel LB, Monteiro ANA. Functional landscape of common variants associated with susceptibility to epithelial ovarian cancer. *Curr Epidemiol Rep*. 2020;7:49–57.
- Kar SP, Berchuck A, Gayther SA, Goode EL, Moysich KB, Pearce CL, et al. Common genetic variation and susceptibility to ovarian cancer: current insights and future directions. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol*. 2018;27:395–404.
- Wray NR, Goddard ME, Visscher PM. Prediction of individual genetic risk to disease from genome-wide association studies. *Genome Res*. 2007;17:1520–8.
- Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, International Schizophrenia Consortium, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460:748–52.
- Abraham G, Kowalczyk A, Zobel J, Inouye M. Performance and robustness of penalized and unpenalized methods for genetic prediction of complex human disease. *Genet Epidemiol*. 2013;37:184–95.
- Habier D, Fernando RL, Kizilkaya K, Garrick DJ. Extension of the bayesian alphabet for genomic selection. *BMC Bioinformatics*. 2011;12:186.
- Szymczak S, Biernacka JM, Cordell HJ, González-Recio O, König IR, Zhang H, et al. Machine learning in genome-wide association studies. *Genet Epidemiol*. 2009;33:551–57.
- Privé F, Aschard H, Blum MGB. Efficient implementation of penalized regression for genetic risk prediction. *Genetics*. 2019;212:65–74.
- Mak TSH, Porsch RM, Choi SW, Zhou X, Sham PC. Polygenic scores via penalized regression on summary statistics. *Genet Epidemiol*. 2017;41:469–80.
- Perren TJ. Mucinous epithelial ovarian carcinoma. *Ann Oncol Off J Eur Soc Med Oncol*. 2016;27:i53–7.
- Phelan CM, Kuchenbaecker KB, Tyrer JP, Kar SP, Lawrenson K, Winham SJ, et al. Identification of 12 new susceptibility loci for different histotypes of epithelial ovarian cancer. *Nat Genet*. 2017;49:680–91.
- Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature*. 2018;562:203–9.
- Lawrenson K, Song F, Hazelett DJ, Kar SP, Tyrer J, Phelan CM, et al. Genome-wide association studies identify susceptibility loci for epithelial ovarian cancer in east Asian women. *Gynecol Oncol*. 2019;153:343–55.
- Manichaikul A, Peres LC, Wang X-Q, Barnard ME, Chyn D, Sheng X, et al. Identification of novel epithelial ovarian cancer loci in women of African ancestry. *Int J Cancer*. 2020;146:2987–98.

- Phelan CM, Kuchenbaecker KB, Tyrer JP, Kar SP, Lawrenson K, Winham SJ, et al. Identification of 12 new susceptibility loci for different histotypes of epithelial ovarian cancer. *Nat Genet*. 2017;49:680–91.
- Yang J, Ferreira T, Morris AP, Medland SE, Genetic Investigation of ANthropometric Traits (GIANT) Consortium, DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium, et al. Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat Genet*. 2012;44:369–75. S1–3
- Vilhjálmsón BJ, Yang J, Finucane HK, Gusev A, Lindström S, Ripke S, et al. Modeling linkage disequilibrium increases accuracy of polygenic risk scores. *Am J Hum Genet*. 2015;97:576–92.
- Ge T, Chen C-Y, Ni Y, Feng Y-CA, Smoller JW. Polygenic prediction via Bayesian regression and continuous shrinkage priors. *Nat Commun*. 2019;10:1776.
- Barnes DR, Rookus MA, McGuffog L, Leslie G, Mooij TM, Dennis J, et al. Polygenic risk scores and breast and epithelial ovarian cancer risks for carriers of BRCA1 and BRCA2 pathogenic variants. *Genet Med Off J Am Coll Med Genet*. 2020;15:576–92.
- Kuchenbaecker KB, Hopper JL, Barnes DR, Phillips K-A, Mooij TM, Roos-Blom M-J, et al. Risks of Breast, Ovarian, and Contralateral Breast Cancer for BRCA1 and BRCA2 Mutation Carriers. *JAMA*. 2017;317:2402–16.
- Mavaddat N, Michailidou K, Dennis J, Lush M, Fachal L, Lee A, et al. Polygenic risk scores for prediction of breast cancer and breast cancer subtypes. *Am J Hum Genet*. 2019;104:21–34.
- Schumacher FR, Al Olama AA, Berndt SI, Benlloch S, Ahmed M, Saunders EJ, et al. Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nat Genet*. 2018;50:928–36.
- Kuchenbaecker KB, McGuffog L, Barrowdale D, Lee A, Soucy P, Dennis J, et al. Evaluation of Polygenic Risk Scores for Breast and Ovarian Cancer Risk Prediction in BRCA1 and BRCA2 Mutation Carriers. *J Natl Cancer Inst*. 2017;109.
- Yang X, Leslie G, Gentry-Maharaj A, Ryan A, Intermaggio M, Lee A, et al. Evaluation of polygenic risk scores for ovarian cancer risk prediction in a prospective cohort study. *J Med Genet*. 2018;55:546–54.
- Jia G, Lu Y, Wen W, Long J, Liu Y, Tao R, et al. Evaluating the utility of polygenic risk scores in identifying high-risk individuals for eight common cancers. *JNCI Cancer Spectr*. 2020;4:pkaa021.
- Song H, Dicks EM, Tyrer J, Intermaggio M, Chenevix-Trench G, Bowtell DD, et al. Population-based targeted sequencing of 54 candidate genes identifies PALB2 as a susceptibility gene for high-grade serous ovarian cancer. *J Med Genet*. 2021;58:305–13.
- Yang X, Leslie G, Doroszuk A, Schneider S, Allen J, Decker B, et al. Cancer risks associated with germline PALB2 pathogenic variants: an international study of 524 families. *J Clin Oncol Off J Am Soc Clin Oncol*. 2020;38:674–85.
- Lee A, Mavaddat N, Wilcox AN, Cunningham AP, Carver T, Hartley S, et al. BOA-DICEA: a comprehensive breast cancer risk prediction model incorporating genetic and nongenetic risk factors. *Genet Med Off J Am Coll Med Genet*. 2019;21:1708–18.
- Welcome to CanRisk. [cited 2021 Aug 15]. <https://canrisk.org/>
- Hu Y, Lu Q, Powles R, Yao X, Yang C, Fang F, et al. Leveraging functional annotations in genetic risk prediction for human complex diseases. *PLoS Comput Biol*. 2017;13:e1005589.
- Gola D, Erdmann J, Müller-Myhsok B, Schunkert H, König IR. Polygenic risk scores outperform machine learning methods in predicting coronary artery disease status. *Genet Epidemiol*. 2020;44:125–38.
- Paré G, Mao S, Deng WQ. A machine-learning heuristic to improve gene score prediction of polygenic traits. *Sci Rep*. 2017;7:12665.

ACKNOWLEDGEMENTS

See Supplementary Material.

AUTHOR CONTRIBUTIONS

EO Dareng, JP Tyrer, DR Barnes, MR Jones, X Yang, KK Aben, MA Adank, S Agata, IL Andrusis, H Anton-Culver, NN Antonenkova, G Aravantis, BK Arun, A Augustinsson, J Balmaña, RB Barkardottir, D Barrowdale, MW Beckmann, A Beeghly-Fadiel, J Benitez, M Bermisheva, MQ Bernardini, L Bjorge, NV Bogdanova, B Bonanni, A Borg, JD Brenton, A Budzylowska, R Butzow, SS Buys, H Cai, MA Caligo, I Campbell, R Cannioto, H Cassingham, J Chang-Claude, SJ Chanock, K Chen, Y Chiew, WK Chung, KB Claes, S Colonna, LS Cook, FJ Couch, MB Daly, F Dao, E Davies, M de la Hoya, R de Putter, J Dennis, A DePersia, P Devilee, O Diez, Y Ding, JA Doherty, SM Domchek, T Dörk, A du Bois, M Dürst, DM Eccles, HA Eliassen, C Engel, D Evans, PA Fasching, JM Flanagan, RT Fortner, E Machackova, E Friedman, PA Ganz, J Garber, F Gensini, GG Giles, G Glendon, AK Godwin, MT Goodman, MH Greene, J Gronwald, E Hahnen, CA Haiman, N Håkansson, U Hamann, TV Hansen, HR Harris, M Hartman, F Heitz, MA Hildebrandt,

E Høgdall, CK Høgdall, JL Hopper, R Huang, C Huff, PJ Hulick, DG Huntsman, EN Imyanitov, C Isaacs, A Jakubowska, PA James, R Janavicius, A Jensen, OT Johannsson, EM John, ME Jones, D Kang, BY Karlan, A Karnezis, LE Kelemen, E Khusnutdinova, LA Kiemeny, B Kim, SK Kjaer, I Komenaka, J Kupryjanczyk, AW Kurian, A Kwong, D Lambrechts, MC Larson, C Lazaro, ND Le, G Leslie, J Lester, F Lesueur, DA Levine, J Li, JT Loud, KH Lu, J Lubi_ski, PL Mai, S Manoukian, JR Marks, R Matsuno, K Matsuo, T May, L McGuffog, JR McLaughlin, IA McNeish, N Mebirouk, A Miller, RL Milne, A Minlikeeva, F Modugno, M Montagna, KB Moysich, E Munro, KL Nathanson, SL Neuhausen, H Nevanlinna, J Ngeow Yuen Yie, H Nielsen, L Nikitina-Zake, K Odunsi, K Offit, E Olah, S Olbrecht, OI Olopade, SH Olson, H Olsson, A Osorio, L Papi, SK Park, MT Parsons, H Pathak, I Pedersen, A Peixoto, T Pejovic, P Perez-Segura, JB Permut, B Peshkin, P Peterlongo, A Piskorz, D Prokofyeva, P Radice, J Rantala, MJ Riggan, HA Risch, C Rodriguez-Antona, E Ross, M Rossing, I Runnebaum, DP Sandler, M Santamariña, P Soucy, RK Schmutzler, V Setiawan, K Shan, W Sieh, J Simard, CF Singer, AP Sokolenko, H Song, MC Southey, H Steed, D Stoppa-Lyonnet, R Sutphen, AJ Swerdlow, Y Tan, MR Teixeira, S Teo, KL Terry, M Terry, M Thomassen, PJ Thompson, L Thomsen, DL Thull, M Tischkowitz, L Titus, AE Toland, D Torres, B Trabert, R Travis, N Tung, SS Tworoger, E Valen, AM van Altena, AH van der Hout, E Van Nieuwenhuysen, EJ van Rosenberg, A Vega, D Velez Edwards, RA Vierkant, F Wang, PM Webb, CR Weinberg, JN Weitzel, N Wentzensen, E White, SJ Winham, A Wolk, Y Woo, AH Wu, L Yan, D Yannoukakos, KM Zavaglia, W Zheng, A Ziogas, KK Zorn, K Lawrenson, TA Sellers, SJ Ramus, AN Monteiro, JM Cunningham, EL Goode, JM Schildkraut, A Berchuck, G Chenevix-Trench, SA Gayther, AC Antoniou, PD Pharoah contributed and/or designed the work that led to this submission, acquired data, played important roles in interpreting results, drafted or revised the manuscript, approved the final version and agreed to be accountable for all aspects of the work.

COMPETING INTERESTS

ADF has received a research grant from AstraZeneca, not directly related to the content of this manuscript. MWB conducts research funded by Amgen, Novartis and Pfizer. PAF conducts research funded by Amgen, Novartis and Pfizer. He received Honoraria from Roche, Novartis and Pfizer. AWK reports research funding to her institution from Myriad Genetics for an unrelated project. UM owns stocks in Abcodia Ltd. Rachel A. Murphy is a consultant for Pharmavite. The other authors declare no conflicts of interest.

ETHICS STATEMENT

All study participants provided written informed consent and participated in research or clinical studies at the host institute under ethically approved protocols. The studies and their approving institutes are listed in the Supplementary Material (Ethics Statement).

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41431-021-00987-7>.

Correspondence and requests for materials should be addressed to Paul D. P. Pharoah.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

Eileen O. Dareng^{1,2,43}, Jonathan P. Tyrer^{2,243}, Daniel R. Barnes¹, Michelle R. Jones³, Xin Yang¹, Katja K. H. Aben^{4,5}, Muriel A. Adank⁶, Simona Agata⁷, Irene L. Andrulis^{8,9}, Hoda Anton-Culver¹⁰, Natalia N. Antonenkova¹¹, Gerasimos Aravantinos¹², Banu K. Arun¹³, Annelie Augustinsson¹⁴, Judith Balmaña^{15,16}, Elisa V. Bandera¹⁷, Rosa B. Barkardottir^{18,19}, Daniel Barrowdale¹, Matthias W. Beckmann²⁰, Alicia Beeghly-Fadiel²¹, Javier Benitez^{22,23}, Marina Bermisheva²⁴, Marcus Q. Bernardini²⁵, Line Bjorge^{26,27}, Amanda Black²⁸, Natalia V. Bogdanova^{11,29,30}, Bernardo Bonanni³¹, Ake Borg³², James D. Brenton³³, Agnieszka Budzylowska³⁴, Ralf Butzow³⁵, Sandra S. Buys³⁶, Hui Cai²¹, Maria A. Caligo³⁷, Ian Campbell^{38,39}, Rikki Cannioto⁴⁰, Hayley Cassingham⁴¹, Jenny Chang-Claude^{42,43}, Stephen J. Chanock⁴⁴, Kexin Chen⁴⁵, Yoke-Eng Chiew^{46,47}, Wendy K. Chung⁴⁸, Kathleen B. M. Claes⁴⁹, Sarah Colonna³⁶, GEMO Study Collaborators^{50,51,52}, GC-HBOC Study Collaborators⁵³, EMBRACE Collaborators¹, Linda S. Cook^{54,55}, Fergus J. Couch⁵⁶, Mary B. Daly⁵⁷, Fanny Dao⁵⁸, Eleanor Davies⁵⁹, Miguel de la Hoya⁶⁰, Robin de Putter⁴⁹, Joe Dennis¹, Allison DePersia^{61,62}, Peter Devilee^{63,64}, Orland Diez^{65,66}, Yuan Chun Ding⁶⁷, Jennifer A. Doherty⁶⁸, Susan M. Domchek⁶⁹, Thilo Dörk³⁰, Andreas du Bois^{70,71}, Matthias Dürst⁷², Diana M. Eccles⁷³, Heather A. Eliassen^{74,75}, Christoph Engel^{76,77}, Gareth D. Evans^{78,79}, Peter A. Fasching^{20,80}, James M. Flanagan⁸¹, Renée T. Fortner⁴², Eva Machackova⁸², Eitan Friedman^{83,84}, Patricia A. Ganz⁸⁵, Judy Garber⁸⁶, Francesca Gensini⁸⁷, Graham G. Giles^{88,89,90}, Gord Glendon⁸, Andrew K. Godwin⁹¹, Marc T. Goodman⁹², Mark H. Greene⁹³, Jacek Gronwald⁹⁴, OPAL Study Group⁹⁵, AOCs Group^{38,46}, Eric Hahnen^{53,96}, Christopher A. Haiman⁹⁷, Niclas Håkansson⁹⁸, Ute Hamann⁹⁹, Thomas V. O. Hansen¹⁰⁰, Holly R. Harris^{101,102}, Mikael Hartman^{103,104}, Florian Heitz^{70,71,105}, Michelle A. T. Hildebrandt¹⁰⁶, Estrid Høgdall^{107,108}, Claus K. Høgdall¹⁰⁹, John L. Hopper⁸⁹, Rhea-Yea Huang¹¹⁰, Chad Huff¹⁰⁶, Peter J. Hulick^{61,62}, David G. Huntsman^{111,112,113,114}, Evgeny N. Imyanitov¹¹⁵, KConFab Investigators³⁸, HEBON Investigators¹¹⁶, Claudine Isaacs¹¹⁷, Anna Jakubowska^{94,118}, Paul A. James^{39,119}, Ramunas Janavicius^{120,121}, Allan Jensen¹⁰⁷, Oskar Th. Johannsson¹²², Esther M. John^{123,124}, Michael E. Jones¹²⁵, Daehee Kang^{126,127,128}, Beth Y. Karlan¹²⁹, Anthony Karnezis¹³⁰, Linda E. Kelemen¹³¹, Elza Khusnutdinova^{24,132}, Lambertus A. Kiemeny⁴, Byoung-Gie Kim¹³³, Susanne K. Kjaer^{107,109}, Ian Komenaka¹³⁴, Jolanta Kupryjanczyk³⁴, Allison W. Kurian^{123,124}, Ava Kwong^{135,136,137}, Diether Lambrechts^{138,139}, Melissa C. Larson¹⁴⁰, Conxi Lazaro¹⁴¹, Nhu D. Le¹⁴², Goska Leslie¹, Jenny Lester¹²⁹, Fabienne Lesueur^{51,52,143}, Douglas A. Levine^{58,144}, Lian Li⁴⁵, Jingmei Li¹⁴⁵, Jennifer T. Loud⁹³, Karen H. Lu¹⁴⁶, Jan Lubiński⁹⁴, Phuong L. Mai¹⁴⁷, Siranoush Manoukian¹⁴⁸, Jeffrey R. Marks¹⁴⁹, Rayna Kim Matsuno¹⁵⁰, Keitaro Matsuo^{151,152}, Taymaa May²⁵, Lesley McGuffog¹, John R. McLaughlin¹⁵³, Iain A. McNeish^{154,155}, Noura Mebirouk^{51,52,143}, Usha Menon¹⁵⁶, Austin Miller¹⁵⁷, Roger L. Milne^{88,89,90}, Albina Minlikeeva¹⁵⁸, Francesmary Modugno^{159,160}, Marco Montagna⁷, Kirsten B. Moysich¹⁵⁸, Elizabeth Munro^{161,162}, Katherine L. Nathanson⁶⁹, Susan L. Neuhausen⁶⁷, Heli Nevanlinna¹⁶³,

Joanne Ngeow Yuen Yie^{164,165}, Henriette Roed Nielsen¹⁶⁶, Finn C. Nielsen¹⁰⁰, Liene Nikitina-Zake¹⁶⁷, Kunle Odunsi¹⁶⁸, Kenneth Offit^{169,170}, Edith Olah¹⁷¹, Siel Olbrecht¹⁷², Olufunmilayo I. Olopade¹⁷³, Sara H. Olson¹⁷⁴, Håkan Olsson¹⁴, Ana Osorio^{23,175}, Laura Papi⁸⁷, Sue K. Park^{126,127,128}, Michael T. Parsons¹⁷⁶, Harsha Pathak⁹¹, Inge Sokilde Pedersen^{177,178,179}, Ana Peixoto¹⁸⁰, Tanja Pejovic^{161,162}, Pedro Perez-Segura⁶⁰, Jennifer B. Permuth¹⁸¹, Beth Peshkin¹⁷, Paolo Peterlongo¹⁸², Anna Piskorz³³, Darya Prokofyeva¹⁸³, Paolo Radice¹⁸⁴, Johanna Rantala¹⁸⁵, Marjorie J. Riggan¹⁸⁶, Harvey A. Risch¹⁸⁷, Cristina Rodriguez-Antona^{22,23}, Eric Ross¹⁸⁸, Mary Anne Rossing^{101,102}, Ingo Runnebaum⁷², Dale P. Sandler¹⁸⁹, Marta Santamariña^{175,190,191}, Penny Soucy¹⁹², Rita K. Schmutzler^{53,96,193}, V. Wendy Setiawan⁹⁷, Kang Shan¹⁹⁴, Weiva Sieh^{195,196}, Jacques Simard¹⁹⁷, Christian F. Singer¹⁹⁸, Anna P. Sokolenko¹¹⁵, Honglin Song¹⁹⁹, Melissa C. Southey^{88,90,200}, Helen Steed²⁰¹, Dominique Stoppa-Lyonnet^{16,50,202,203}, Rebecca Sutphen²⁰⁴, Anthony J. Swerdlow^{125,205}, Yen Yen Tan¹⁹⁸, Manuel R. Teixeira^{180,206}, Soo Hwang Teo^{207,208}, Kathryn L. Terry^{74,209}, Mary Beth Terry²¹⁰, The OCAC Consortium¹⁸⁶, The CIMBA Consortium¹, Mads Thomassen¹⁶⁶, Pamela J. Thompson⁹², Liv Cecilie Vestrheim Thomsen^{26,27}, Darcy L. Thull²¹¹, Marc Tischkowitz^{121,212,213}, Linda Titus²¹⁴, Amanda E. Toland²¹⁵, Diana Torres^{99,216}, Britton Trabert²⁸, Ruth Travis²¹⁷, Nadine Tung²¹⁸, Shelley S. Tworoger^{74,181}, Ellen Valen^{26,27}, Anne M. van Altena⁴, Annemieke H. van der Hout²¹⁹, Els Van Nieuwenhuysen¹⁷², Elizabeth J. van Rensburg¹²⁰, Ana Vega^{175,221,222}, Digna Velez Edwards²²³, Robert A. Vierkant¹⁴⁰, Frances Wang^{224,225}, Barbara Wappenschmidt^{53,96}, Penelope M. Webb⁹⁵, Clarice R. Weinberg²²⁶, Jeffrey N. Weitzel²²⁷, Nicolas Wentzensen²⁸, Emily White^{102,228}, Alice S. Whittemore^{123,229}, Stacey J. Winham¹⁴⁰, Alicja Wolk^{98,230}, Yin-Ling Woo²³¹, Anna H. Wu⁹⁷, Li Yan²³², Drakoulis Yannoukakos²³³, Katia M. Zavaglia³⁷, Wei Zheng²¹, Argyrios Ziogas¹⁰, Kristin K. Zorn¹⁴⁷, Zdenek Kleibl²³⁴, Douglas Easton^{1,2}, Kate Lawrenson^{3,235}, Anna DeFazio^{46,47}, Thomas A. Sellers²³⁶, Susan J. Ramus^{237,238}, Celeste L. Pearce^{239,240}, Alvaro N. Monteiro¹⁸¹, Julie Cunningham²⁴¹, Ellen L. Goode²⁴¹, Joellen M. Schildkraut²⁴², Andrew Berchuck¹⁸⁶, Georgia Chenevix-Trench¹⁷⁶, Simon A. Gayther¹³, Antonis C. Antoniou¹ and Paul D. P. Pharoah^{1,2,23}

¹University of Cambridge, Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, Cambridge, UK. ²University of Cambridge, Centre for Cancer Genetic Epidemiology, Department of Oncology, Cambridge, UK. ³Center for Bioinformatics and Functional Genomics, Cedars-Sinai Medical Center, Los Angeles, CA, USA. ⁴Radboud University Medical Center, Radboud Institute for Health Sciences, Nijmegen, The Netherlands. ⁵Netherlands Comprehensive Cancer Organisation, Utrecht, The Netherlands. ⁶The Netherlands Cancer Institute—Antoni van Leeuwenhoek hospital, Family Cancer Clinic, Amsterdam, The Netherlands. ⁷Veneto Institute of Oncology IOV—IRCCS, Immunology and Molecular Oncology Unit, Padua, Italy. ⁸Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Fred A. Litwin Center for Cancer Genetics, Toronto, ON, Canada. ⁹University of Toronto, Department of Molecular Genetics, Toronto, ON, Canada. ¹⁰University of California Irvine, Department of Epidemiology, Genetic Epidemiology Research Institute, Irvine, CA, USA. ¹¹N.N. Alexandrov Research Institute of Oncology and Medical Radiology, Minsk, Belarus. ¹²Agii Anargiri Cancer Hospital, Athens, Greece. ¹³University of Texas MD Anderson Cancer Center, Department of Breast Medical Oncology, Houston, TX, USA. ¹⁴Lund University, Department of Cancer Epidemiology, Clinical Sciences, Lund, Sweden. ¹⁵Vall d'Hebron Institute of Oncology, Hereditary cancer Genetics Group, Barcelona, Spain. ¹⁶University Hospital of Vall d'Hebron, Department of Medical Oncology, Barcelona, Spain. ¹⁷Rutgers Cancer Institute of New Jersey, Cancer Prevention and Control Program, New Brunswick, NJ, USA. ¹⁸Landspitali University Hospital, Department of Pathology, Reykjavik, Iceland. ¹⁹University of Iceland, BMC (Biomedical Centre), Faculty of Medicine, Reykjavik, Iceland. ²⁰University Hospital Erlangen, Friedrich-Alexander-University Erlangen-Nuremberg, Department of Gynecology and Obstetrics, Comprehensive Cancer Center ER-EMN, Erlangen, Germany. ²¹Vanderbilt University School of Medicine, Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Nashville, TN, USA. ²²Biomedical Network on Rare Diseases (CIBERER), Madrid, Spain. ²³Spanish National Cancer Research Centre (CNIO), Human Cancer Genetics Programme, Madrid, Spain. ²⁴Ufa Federal Research Centre of the Russian Academy of Sciences, Institute of Biochemistry and Genetics, Ufa, Russia. ²⁵Princess Margaret Hospital, Division of Gynecologic Oncology, University Health Network, Toronto, ON, Canada. ²⁶Haukeland University Hospital, Department of Obstetrics and Gynecology, Bergen, Norway. ²⁷University of Bergen, Centre for Cancer Biomarkers CCBIO, Department of Clinical Science, Bergen, Norway. ²⁸National Cancer Institute, Division of Cancer Epidemiology and Genetics, Bethesda, MD, USA. ²⁹Hannover Medical School, Department of Radiation Oncology, Hannover, Germany. ³⁰Hannover Medical School, Gynaecology Research Unit, Hannover, Germany. ³¹IEO, European Institute of Oncology IRCCS, Division of Cancer Prevention and Genetics, Milan, Italy. ³²Lund University and Skåne University Hospital, Department of Oncology, Lund, Sweden. ³³Cancer Research UK Cambridge Institute, University of Cambridge, Cambridge, UK. ³⁴Maria Skłodowska-Curie National Research Institute of Oncology, Department of Pathology and Laboratory Diagnostics, Warsaw, Poland. ³⁵University of Helsinki, Department of Pathology, Helsinki University Hospital, Helsinki, Finland. ³⁶Huntsman Cancer Institute, Department of Medicine, Salt Lake City, UT, USA. ³⁷University Hospital, SOD Genetica Molecolare, Pisa, Italy. ³⁸Peter MacCallum Cancer Center, Melbourne, VIC, Australia. ³⁹The University of Melbourne, Sir Peter MacCallum Department of Oncology, Melbourne, VIC, Australia. ⁴⁰Roswell Park Cancer Institute, Cancer Pathology & Prevention, Division of Cancer Prevention and Population Sciences, Buffalo, NY, USA. ⁴¹Division of Human Genetics, The Ohio State University, Department of Internal Medicine, Columbus, OH, USA. ⁴²German Cancer Research Center (DKFZ), Division of Cancer Epidemiology, Heidelberg, Germany. ⁴³University Medical Center Hamburg-Eppendorf, Cancer Epidemiology Group, University Cancer Center Hamburg (UCCH), Hamburg, Germany. ⁴⁴National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Division of Cancer Epidemiology and Genetics, Bethesda, MD, USA. ⁴⁵Tianjin Medical University Cancer Institute and Hospital, Department of Epidemiology, Tianjin, China. ⁴⁶The University of Sydney, Centre for Cancer Research, The Westmead Institute for Medical Research, Sydney, NSW, Australia. ⁴⁷Westmead Hospital, Department of Gynaecological Oncology, Sydney, NSW, Australia. ⁴⁸Columbia University, Departments of Pediatrics and Medicine, New York, NY, USA. ⁴⁹Ghent University, Centre for Medical Genetics, Gent, Belgium. ⁵⁰INSERM U830, Department of Tumour Biology, Paris, France. ⁵¹Institut Curie, Paris, France. ⁵²Mines ParisTech, Fontainebleau, France. ⁵³Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Familial Breast and Ovarian Cancer, Cologne, Germany. ⁵⁴University of New Mexico, University of New Mexico Health Sciences Center, Albuquerque, NM, USA. ⁵⁵Alberta Health Services, Department of Cancer Epidemiology and Prevention Research, Calgary, AB, Canada. ⁵⁶Mayo Clinic, Department of Laboratory Medicine and Pathology, Rochester, MN, USA. ⁵⁷Fox Chase Cancer Center, Department of Clinical Genetics, Philadelphia, PA, USA. ⁵⁸Memorial Sloan Kettering Cancer Center, Gynecology Service, Department of Surgery, New York, NY, USA. ⁵⁹Cambridge, Cambridge, UK. ⁶⁰CIBERONC, Hospital Clinico San Carlos, IDISCC (Instituto de Investigación Sanitaria del Hospital Clinico San Carlos), Molecular Oncology Laboratory, Madrid, Spain. ⁶¹NorthShore University Health System, Center for Medical Genetics, Evanston, IL, USA. ⁶²The University of Chicago Pritzker School of Medicine, Chicago, IL, USA. ⁶³Leiden University Medical Center, Department of Pathology, Leiden, The Netherlands. ⁶⁴Leiden University Medical Center, Department of Human Genetics, Leiden, The Netherlands. ⁶⁵Vall d'Hebron Institute of Oncology (VHIO), Oncogenetics Group, Barcelona, Spain. ⁶⁶University Hospital Vall d'Hebron, Clinical and Molecular Genetics Area, Barcelona, Spain. ⁶⁷Beckman Research Institute of City of Hope, Department of Population Sciences, Duarte, CA, USA. ⁶⁸University of Utah, Huntsman Cancer Institute, Department of Population Health Sciences, Salt Lake City, UT, USA. ⁶⁹University of Pennsylvania, Basser Center for BRCA, Abramson Cancer Center, Philadelphia, PA, USA. ⁷⁰Ev. Kliniken Essen-Mitte (KEM), Department of Gynecology and Gynecologic Oncology, Essen, Germany. ⁷¹Dr. Horst Schmidt Kliniken Wiesbaden, Department of Gynecology and Gynecologic Oncology, Wiesbaden, Germany. ⁷²Jena University Hospital—Friedrich Schiller University, Department of Gynaecology, Jena, Germany. ⁷³University of Southampton, Faculty of Medicine, Southampton, UK. ⁷⁴Harvard T.H. Chan School of Public Health, Department of Epidemiology, Boston, MA, USA. ⁷⁵Brigham and Women's Hospital and Harvard Medical School, Channing Division of Network Medicine, Boston, MA, USA. ⁷⁶University of Leipzig, Institute for Medical Informatics, Statistics and Epidemiology, Leipzig, Germany. ⁷⁷University of Leipzig, LIFE—Leipzig Research Centre for Civilization Diseases, Leipzig, Germany. ⁷⁸University of Manchester, Manchester Academic Health Science Centre, Division of Evolution and Genomic Sciences, School of Biological Sciences, Faculty of Biology, Medicine and Health, Manchester, UK. ⁷⁹St Mary's Hospital, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, North West Genomics Laboratory Hub, Manchester Centre for Genomic Medicine, Manchester, UK. ⁸⁰University of California at Los Angeles, David Geffen School of Medicine, Department of Medicine Division of Hematology and Oncology, Los Angeles, CA, USA. ⁸¹Imperial College London, Division of Cancer and Ovarian Cancer Action Research Centre, Department of Surgery and Cancer, London, UK. ⁸²Masaryk Memorial Cancer Institute, Department of Cancer Epidemiology and Genetics, Brno, Czech Republic. ⁸³Chaim Sheba Medical Center, The Susanne Levy Gertner Oncogenetics Unit, Ramat Gan, Israel. ⁸⁴Tel Aviv University, Sackler Faculty of Medicine, Ramat Aviv, Israel. ⁸⁵Jonsson Comprehensive Cancer Centre, UCLA, Schools of Medicine and Public Health, Division of Cancer Prevention & Control Research, Los Angeles, CA, USA. ⁸⁶Dana-Farber Cancer Institute, Cancer Risk and Prevention Clinic, Boston,

MA, USA. ⁸⁷University of Florence, Department of Experimental and Clinical Biomedical Sciences 'Mario Serio', Medical Genetics Unit, Florence, Italy. ⁸⁸Cancer Council Victoria, Cancer Epidemiology Division, Melbourne, VIC, Australia. ⁸⁹The University of Melbourne, Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, Melbourne, VIC, Australia. ⁹⁰Monash University, Precision Medicine, School of Clinical Sciences at Monash Health, Clayton, VIC, Australia. ⁹¹University of Kansas Medical Center, Department of Pathology and Laboratory Medicine, Kansas City, KS, USA. ⁹²Cedars-Sinai Medical Center, Samuel Oschin Comprehensive Cancer Institute, Cancer Prevention and Genetics Program, Los Angeles, CA, USA. ⁹³National Cancer Institute, Clinical Genetics Branch, Division of Cancer Epidemiology and Genetics, Bethesda, MD, USA. ⁹⁴Pomeranian Medical University, Department of Genetics and Pathology, Szczecin, Poland. ⁹⁵QIMR Berghofer Medical Research Institute, Population Health Department, Brisbane, QLD, Australia. ⁹⁶Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Integrated Oncology (CIO), Cologne, Germany. ⁹⁷University of Southern California, Department of Preventive Medicine, Keck School of Medicine, Los Angeles, CA, USA. ⁹⁸Karolinska Institutet, Institute of Environmental Medicine, Stockholm, Sweden. ⁹⁹German Cancer Research Center (DKFZ), Molecular Genetics of Breast Cancer, Heidelberg, Germany. ¹⁰⁰Rigshospitalet, Copenhagen University Hospital, Department of Clinical Genetics, Copenhagen, Denmark. ¹⁰¹Fred Hutchinson Cancer Research Center, Program in Epidemiology, Division of Public Health Sciences, Seattle, WA, USA. ¹⁰²University of Washington, Department of Epidemiology, Seattle, WA, USA. ¹⁰³National University of Singapore and National University Health System, Saw Swee Hock School of Public Health, Singapore. ¹⁰⁴National University Health System, Department of Surgery, Singapore. ¹⁰⁵Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department for Gynecology with the Center for Oncologic Surgery Charité Campus Virchow-Klinikum, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Berlin, Germany. ¹⁰⁶University of Texas MD Anderson Cancer Center, Department of Epidemiology, Houston, TX, USA. ¹⁰⁷Danish Cancer Society Research Center, Department of Virus, Lifestyle and Genes, Copenhagen, Denmark. ¹⁰⁸University of Copenhagen, Molecular Unit, Department of Pathology, Herlev Hospital, Copenhagen, Denmark. ¹⁰⁹University of Copenhagen, Department of Gynaecology, Rigshospitalet, Copenhagen, Denmark. ¹¹⁰Roswell Park Cancer Institute, Center For Immunotherapy, Buffalo, NY, USA. ¹¹¹BC Cancer, Vancouver General Hospital, and University of British Columbia, British Columbia's Ovarian Cancer Research (OVCARE) Program, Vancouver, BC, Canada. ¹¹²University of British Columbia, Department of Pathology and Laboratory Medicine, Vancouver, BC, Canada. ¹¹³University of British Columbia, Department of Obstetrics and Gynecology, Vancouver, BC, Canada. ¹¹⁴BC Cancer Research Centre, Department of Molecular Oncology, Vancouver, BC, Canada. ¹¹⁵N.N. Petrov Institute of Oncology, St. Petersburg, Russia. ¹¹⁶Coordinating center: The Netherlands Cancer Institute, The Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON), Amsterdam, The Netherlands. ¹¹⁷Lombardi Comprehensive Cancer Center, Georgetown University, Washington, DC, USA. ¹¹⁸Pomeranian Medical University, Independent Laboratory of Molecular Biology and Genetic Diagnostics, Szczecin, Poland. ¹¹⁹Peter MacCallum Cancer Center, Parkville Familial Cancer Centre, Melbourne, VIC, Australia. ¹²⁰Vilnius University Hospital Santariskiu Clinics, Hematology, oncology and transfusion medicine center, Dept. of Molecular and Regenerative Medicine, Vilnius, Lithuania. ¹²¹State Research Institute Centre for Innovative Medicine, Vilnius, Lithuania. ¹²²Landsþítali University Hospital, Department of Oncology, Reykjavík, Iceland. ¹²³Stanford University School of Medicine, Department of Epidemiology & Population Health, Stanford, CA, USA. ¹²⁴Stanford Cancer Institute, Stanford University School of Medicine, Department of Medicine, Division of Oncology, Stanford, CA, USA. ¹²⁵The Institute of Cancer Research, Division of Genetics and Epidemiology, London, UK. ¹²⁶Seoul National University College of Medicine, Department of Preventive Medicine, Seoul, Korea. ¹²⁷Seoul National University Graduate School, Department of Biomedical Sciences, Seoul, Korea. ¹²⁸Seoul National University, Cancer Research Institute, Seoul, Korea. ¹²⁹University of California at Los Angeles, David Geffen School of Medicine, Department of Obstetrics and Gynecology, Los Angeles, CA, USA. ¹³⁰UC Davis Medical Center, Department of Pathology and Laboratory Medicine, Sacramento, CA, USA. ¹³¹Medical University of South Carolina, Hollings Cancer Center, Charleston, SC, USA. ¹³²Saint Petersburg State University, Saint Petersburg, Russia. ¹³³Sungkyunkwan University School of Medicine, Department of Obstetrics and Gynecology, Samsung Medical Center, Seoul, Korea. ¹³⁴City of Hope Clinical Cancer Genetics Community Research Network, Duarte, CA, USA. ¹³⁵Cancer Genetics Centre, Hong Kong Hereditary Breast Cancer Family Registry, Happy Valley, Hong Kong. ¹³⁶The University of Hong Kong, Department of Surgery, Pok Fu Lam, Hong Kong. ¹³⁷Hong Kong Sanatorium and Hospital, Department of Surgery, Happy Valley, Hong Kong. ¹³⁸VIB Center for Cancer Biology, Leuven, Belgium. ¹³⁹University of Leuven, Laboratory for Translational Genetics, Department of Human Genetics, Leuven, Belgium. ¹⁴⁰Mayo Clinic, Department of Health Sciences Research, Division of Biomedical Statistics and Informatics, Rochester, MN, USA. ¹⁴¹ONCOBELL-IDIBELL-IGTP, Catalan Institute of Oncology, CIBERONC, Hereditary Cancer Program, Barcelona, Spain. ¹⁴²BC Cancer, Cancer Control Research, Vancouver, BC, Canada. ¹⁴³Inserm U900, Genetic Epidemiology of Cancer team, Paris, France. ¹⁴⁴NYU Langone Medical Center, Gynecologic Oncology, Laura and Isaac Pearlmuter Cancer Center, New York, NY, USA. ¹⁴⁵Genome Institute of Singapore, Human Genetics Division, Singapore, Singapore. ¹⁴⁶University of Texas MD Anderson Cancer Center, Department of Gynecologic Oncology and Clinical Cancer Genetics Program, Houston, TX, USA. ¹⁴⁷Magee-Womens Hospital, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA. ¹⁴⁸Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Unit of Medical Genetics, Department of Medical Oncology and Hematology, Milan, Italy. ¹⁴⁹Duke University Hospital, Department of Surgery, Durham, NC, USA. ¹⁵⁰University of Hawaii Cancer Center, Cancer Epidemiology Program, Honolulu, HI, USA. ¹⁵¹Aichi Cancer Center Research Institute, Division of Cancer Epidemiology and Prevention, Nagoya, Japan. ¹⁵²Nagoya University Graduate School of Medicine, Division of Cancer Epidemiology, Nagoya, Japan. ¹⁵³Samuel Lunenfeld Research Institute, Public Health Ontario, Toronto, ON, Canada. ¹⁵⁴Imperial College London, Division of Cancer and Ovarian Cancer Action Research Centre, Department Surgery & Cancer, London, UK. ¹⁵⁵University of Glasgow, Institute of Cancer Sciences, Glasgow, UK. ¹⁵⁶University College London, MRC Clinical Trials Unit at UCL, Institute of Clinical Trials & Methodology, London, UK. ¹⁵⁷Roswell Park Cancer Institute, NRG Oncology, Statistics and Data Management Center, Buffalo, NY, USA. ¹⁵⁸Roswell Park Cancer Institute, Division of Cancer Prevention and Control, Buffalo, NY, USA. ¹⁵⁹Magee-Womens Research Institute and Hillman Cancer Center, Womens Cancer Research Center, Pittsburgh, PA, USA. ¹⁶⁰University of Pittsburgh School of Medicine, Division of Gynecologic Oncology, Department of Obstetrics, Gynecology and Reproductive Sciences, Pittsburgh, PA, USA. ¹⁶¹Oregon Health & Science University, Department of Obstetrics and Gynecology, Portland, OR, USA. ¹⁶²Oregon Health & Science University, Knight Cancer Institute, Portland, OR, USA. ¹⁶³University of Helsinki, Department of Obstetrics and Gynecology, Helsinki University Hospital, Helsinki, Finland. ¹⁶⁴National Cancer Centre, Cancer Genetics Service, Singapore, Singapore. ¹⁶⁵Nanyang Technological University, Lee Kong Chian School of Medicine, Singapore, Singapore. ¹⁶⁶Odense University Hospital, Department of Clinical Genetics, Odense C, Denmark. ¹⁶⁷Latvian Biomedical Research and Study Centre, Riga, Latvia. ¹⁶⁸Roswell Park Cancer Institute, Department of Gynecologic Oncology, Buffalo, NY, USA. ¹⁶⁹Memorial Sloan Kettering Cancer Center, Clinical Genetics Research Lab, Department of Cancer Biology and Genetics, New York, NY, USA. ¹⁷⁰Memorial Sloan Kettering Cancer Center, Clinical Genetics Service, Department of Medicine, New York, NY, USA. ¹⁷¹National Institute of Oncology, Department of Molecular Genetics, Budapest, Hungary. ¹⁷²University Hospitals Leuven, Division of Gynecologic Oncology, Department of Obstetrics and Gynaecology and Leuven Cancer Institute, Leuven, Belgium. ¹⁷³The University of Chicago, Center for Clinical Cancer Genetics, Chicago, IL, USA. ¹⁷⁴Memorial Sloan-Kettering Cancer Center, Department of Epidemiology and Biostatistics, New York, NY, USA. ¹⁷⁵Centro de Investigación en Red de Enfermedades Raras (CIBERER), Madrid, Spain. ¹⁷⁶QIMR Berghofer Medical Research Institute, Department of Genetics and Computational Biology, Brisbane, QLD, Australia. ¹⁷⁷Aalborg University Hospital, Molecular Diagnostics, Aalborg, Denmark. ¹⁷⁸Aalborg University Hospital, Clinical Cancer Research Center, Aalborg, Denmark. ¹⁷⁹Aalborg University, Department of Clinical Medicine, Aalborg, Denmark. ¹⁸⁰Portuguese Oncology Institute, Department of Genetics, Porto, Portugal. ¹⁸¹Moffitt Cancer Center, Department of Cancer Epidemiology, Tampa, FL, USA. ¹⁸²IFOM—the FIRCC Institute of Molecular Oncology, Genome Diagnostics Program, Milan, Italy. ¹⁸³Bashkir State University, Department of Genetics and Fundamental Medicine, Ufa, Russia. ¹⁸⁴Fondazione IRCCS Istituto Nazionale dei Tumori (INT), Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Research, Milan, Italy. ¹⁸⁵Karolinska Institutet, Clinical Genetics, Stockholm, Sweden. ¹⁸⁶Duke University Hospital, Department of Gynecologic Oncology, Durham, NC, USA. ¹⁸⁷Yale School of Public Health, Chronic Disease Epidemiology, New Haven, CT, USA. ¹⁸⁸Fox Chase Cancer Center, Population Studies Facility, Philadelphia, PA, USA. ¹⁸⁹National Institute of Environmental Health Sciences, NIH, Epidemiology Branch, Research Triangle Park, NC, USA. ¹⁹⁰Fundación Pública Galega Medicina Xenómica, Santiago De Compostela, Spain. ¹⁹¹Instituto de Investigación Sanitaria de Santiago de Compostela, Santiago De Compostela, Spain. ¹⁹²Centre Hospitalier Universitaire de Québec – Université Laval Research Center, Genomics Center, Québec City, QC, Canada. ¹⁹³Faculty of Medicine and University Hospital Cologne, University of Cologne, Center for Molecular Medicine Cologne (CMCC), Cologne, Germany. ¹⁹⁴Hebei Medical University, Fourth Hospital, Department of Obstetrics and Gynaecology, Shijiazhuang, China. ¹⁹⁵Icahn School of Medicine at Mount Sinai, Department of Population Health Science and Policy, New York, NY, USA. ¹⁹⁶Icahn School of Medicine at Mount Sinai, Department of Genetics and Genomic Sciences, New York, NY, USA. ¹⁹⁷Centre Hospitalier Universitaire de Québec-Université Laval Research Center, Genomic Center, Québec City, QC, Canada. ¹⁹⁸Medical University of Vienna, Dept of OB/GYN and Comprehensive Cancer Center, Vienna, Austria. ¹⁹⁹University of Cambridge, Department of Public Health and Primary Care, Cambridge, UK. ²⁰⁰The University of Melbourne, Department of Clinical Pathology, Melbourne, VIC, Australia. ²⁰¹Royal Alexandra Hospital, Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, Edmonton, AB, Canada. ²⁰²Institut Curie, Service de Génétique, Paris, France. ²⁰³Université Paris Descartes, Paris, France. ²⁰⁴University of South Florida, Epidemiology Center, College of Medicine, Tampa, FL, USA. ²⁰⁵The Institute of Cancer Research, Division of Breast Cancer Research, London, UK. ²⁰⁶University of Porto, Biomedical Sciences Institute (ICBAS), Porto, Portugal. ²⁰⁷Cancer Research Malaysia, Breast Cancer Research Programme, Subang Jaya, Selangor, Malaysia. ²⁰⁸University of Malaya, Department of Surgery, Faculty of Medicine, Kuala Lumpur, Malaysia. ²⁰⁹Brigham and Women's Hospital and Harvard Medical School, Obstetrics and Gynecology Epidemiology Center, Boston, MA, USA. ²¹⁰Columbia University, Department of Epidemiology, Mailman School of Public Health, New York, NY, USA. ²¹¹Magee-Womens Hospital, University of Pittsburgh School of Medicine, Department of Medicine, Pittsburgh, PA, USA. ²¹²McGill University, Program in Cancer Genetics, Departments of

Human Genetics and Oncology, Montréal, QC, Canada. ²¹³University of Cambridge, Department of Medical Genetics, Cambridge, UK. ²¹⁴Dartmouth College, Geisel School of Medicine, Hanover, NH, USA. ²¹⁵The Ohio State University, Department of Cancer Biology and Genetics, Columbus, OH, USA. ²¹⁶Pontificia Universidad Javeriana, Institute of Human Genetics, Bogotá, Colombia. ²¹⁷University of Oxford, Cancer Epidemiology Unit, Oxford, UK. ²¹⁸Beth Israel Deaconess Medical Center, Department of Medical Oncology, Boston, MA, USA. ²¹⁹University Medical Center Groningen, University Groningen, Department of Genetics, Groningen, The Netherlands. ²²⁰University of Pretoria, Department of Genetics, Arcadia, South Africa. ²²¹Fundación Pública Galega de Medicina Xenómica, Santiago de Compostela, Spain. ²²²Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Complejo Hospitalario Universitario de Santiago, SERGAS, Santiago de Compostela, Spain. ²²³Vanderbilt University Medical Center, Division of Quantitative Sciences, Department of Obstetrics and Gynecology, Department of Biomedical Sciences, Women's Health Research, Nashville, TN, USA. ²²⁴Duke Cancer Institute, Cancer Control and Population Sciences, Durham, NC, USA. ²²⁵Duke University Hospital, Department of Community and Family Medicine, Durham, NC, USA. ²²⁶National Institute of Environmental Health Sciences, NIH, Biostatistics and Computational Biology Branch, Research Triangle Park, NC, USA. ²²⁷City of Hope, Clinical Cancer Genomics, Duarte, CA, USA. ²²⁸Fred Hutchinson Cancer Research Center, Seattle, WA, USA. ²²⁹Stanford University School of Medicine, Department of Biomedical Data Science, Stanford, CA, USA. ²³⁰Uppsala University, Department of Surgical Sciences, Uppsala, Sweden. ²³¹University of Malaya, Department of Obstetrics and Gynaecology, University of Malaya Medical Centre, Kuala Lumpur, Malaysia. ²³²Hebei Medical University, Fourth Hospital, Department of Molecular Biology, Shijiazhuang, China. ²³³National Centre for Scientific Research 'Demokritos', Molecular Diagnostics Laboratory, INRASTES, Athens, Greece. ²³⁴Institute of Biochemistry and Experimental Oncology, First Faculty of Medicine, Charles University, Prague, Czech Republic. ²³⁵Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Centre, Department of Obstetrics and Gynecology, Los Angeles, CA, USA. ²³⁶Royal Pass Road, Tampa, FL, USA. ²³⁷University of NSW Sydney, School of Women's and Children's Health, Faculty of Medicine, Sydney, NSW, Australia. ²³⁸University of NSW Sydney, Adult Cancer Program, Lowy Cancer Research Centre, Sydney, NSW, Australia. ²³⁹University of Michigan School of Public Health, Department of Epidemiology, Ann Arbor, MI, USA. ²⁴⁰University of Southern California Norris Comprehensive Cancer Center, Department of Preventive Medicine, Keck School of Medicine, Los Angeles, CA, USA. ²⁴¹Mayo Clinic, Department of Health Science Research, Division of Epidemiology, Rochester, MN, USA. ²⁴²Emory University, Department of Epidemiology, Rollins School of Public Health, Atlanta, GA, USA. ²⁴³These authors contributed equally: Eileen O. Dareng, Jonathan P. Tyrer. ✉email: pp10001@medschl.cam.ac.uk

GEMO Study Collaborators

Fabienne Lesueur and Noura Mebirouk

GC-HBOC Study Collaborators

Christoph Engel and Rita K. Schmutzler

EMBRACE Collaborators

Daniel Barrowdale, Eleanor Davies, Diana M. Eccles and D. Gareth Evans

KConFab Investigators

Georgia Chenevix-Trench

HEBON Investigators

Muriel A. Adank, Peter Devilee and Annemieke H. van der Hout

The OCAC Consortium

Eileen O. Dareng, Jonathan P. Tyrer, Michelle R. Jones, Katja K. H. Aben, Hoda Anton-Culver, Natalia N. Antonenkova, Gerasimos Aravantinos, Matthias W. Beckmann, Alicia Beeghly-Fadiel, Javier Benitez, Marina Bermisheva, Marcus Q. Bernardini, Line Bjorge, Natalia V. Bogdanova, James D. Brenton, Agnieszka Budzylowska, Ralf Butzow, Hui Cai, Ian Campbell, Rikki Cannioto, Jenny Chang-Claude, Stephen J. Chanock, Kexin Chen, Yoke-Eng Chiew, Linda S. Cook, Fanny Dao, Joe Dennis, Jennifer A. Doherty, Thilo Dörk, Andreas du Bois, Matthias Dürst, Diana M. Eccles, Heather A. Eliassen, Peter A. Fasching, James M. Flanagan, Renée T. Fortner, Graham G. Giles, Marc T. Goodman, Jacek Gronwald, Christopher A. Haiman, Niclas Håkansson, Holly R. Harris, Florian Heitz, Michelle A. T. Hildebrandt, Estrid Høgdall, Claus K. Høgdall, Ruea-Yea Huang, Chad Huff, David G. Huntsman, Anna Jakubowska, Allan Jensen, Michael E. Jones, Daehee Kang, Beth Y. Karlan, Anthony Karnezis, Linda E. Kelemen, Elza Khusnutdinova, Lambertus A. Kiemeneij, Byoung-Gie Kim, Susanne K. Kjaer, Jolanta Kupryjanczyk, Diether Lambrechts, Melissa C. Larson, Nhu D. Le, Jenny Lester, Douglas A. Levine, Karen H. Lu, Jan Lubiński, Jeffrey R. Marks, Rayna Kim Matsuno, Keitaro Matsuo, Taymaa May, John R. McLaughlin, Iain A. McNeish, Roger L. Milne, Albina Minlikeeva, Francesmary Modugno, Kirsten B. Moysich, Elizabeth Munro, Heli Nevanlinna, Kunle Odunsi, Siel Olbrecht, Sara H. Olson, Håkan Olsson, Ana Osorio, Sue K. Park, Tanja Pejovic, Jennifer B. Permuth, Anna Piskorz, Darya Prokofyeva, Marjorie J. Riggan, Harvey A. Risch, Cristina Rodriguez-Antona, Mary Anne Rossing, Ingo Runnebaum, Dale P. Sandler, V. Wendy Setiawan, Kang Shan, Weiva Sieh, Honglin Song, Melissa C. Southey, Helen Steed, Rebecca Sutphen, Anthony J. Swerdlow, Soo Hwang Teo, Kathryn L. Terry, Pamela J. Thompson, Liv Cecilie Vestrheim Thomsen, Linda Titus, Britton Trabert, Ruth Travis, Shelley S. Tworoger, Ellen Valen, Anne M. van Altena, Els Van Nieuwenhuysen, Digna Velez Edwards, Robert A. Vierkant, Frances Wang, Penelope M. Webb, Clarice R. Weinberg, Nicolas Wentzensen, Emily White, Alice S. Whittemore, Stacey J. Winham, Alicja Wolk, Yin-Ling Woo, Anna H. Wu, Li Yan, Drakoulis Yannoukakis, Wei Zheng, Argyrios Ziogas, Kate Lawrenson, Anna deFazio, Susan J. Ramus, Celeste L. Pearce, Alvaro N. Monteiro, Julie M. Cunningham, Ellen L. Goode, Joellen M. Schildkraut, Andrew Berchuck, Simon A. Gayther and Paul D. P. Pharoah

The CIMBA Consortium

Daniel R. Barnes, Xin Yang, Muriel A. Adank, Simona Agata, Irene L. Andrulis, Banu K. Arun, Annelie Augustinsson, Judith Balmaña, Rosa B. Barkardottir, Daniel Barrowdale, Bernardo Bonanni, Ake Borg, Saundra S. Buys, Maria A. Caligo, Hayley Cassingham, Wendy K. Chung, Kathleen B. M. Claes, Sarah Colonna, Fergus J. Couch, Mary B. Daly, Eleanor Davies, Miguel de la Hoya, Robin de Putter, Allison DePersia, Peter Devilee, Orland Diez, Yuan Chun Ding, Susan M. Domchek, Diana M. Eccles, Christoph Engel, D. Gareth Evans, Eva Machackova, Eitan Friedman, Patricia A. Ganz, Judy Garber, Francesca Gensini, Gord Glendon, Andrew K. Godwin, Mark H. Greene, Eric Hahnen, Ute Hamann, Thomas V. O. Hansen, Mikael Hartman, John L. Hopper, Peter J. Hulick, Evgeny N. Imyanitov, Claudine Isaacs, Paul A. James, Ramunas Janavicius, Oskar Th. Johannsson, Esther M. John, Ian Komenaka, Allison W. Kurian, Ava Kwong, Conxi Lazaro, Goska Leslie, Fabienne Lesueur, Jingmei Li, Jennifer T. Loud, Phuong L. Mai, Siranoush Manoukian, Lesley McGuffog, Noura Mebirouk, Austin Miller, Marco Montagna, Katherine L. Nathanson, Susan L. Neuhausen, Joanne Ngeow Yuen Yie, Henriette Roed Nielsen, Liene Nikitina-Zake, Kenneth Offit, Edith Olah, Olufunmilayo I. Olopade, Laura Papi, Michael T. Parsons, Harsha Pathak, Inge Sokilde Pedersen, Ana Peixoto, Pedro Perez-Segura, Beth Peshkin, Paolo Peterlongo, Paolo Radice, Johanna Rantala, Eric Ross, Marta Santamariña, Penny Soucy, Rita K. Schmutzler, Jacques Simard, Christian F. Singer, Anna P. Sokolenko, Dominique Stoppa-Lyonnet, Yen Yen Tan, Manuel R. Teixeira, Mary Beth Terry, Mads Thomassen, Darcy L. Thull, Marc Tischkowitz, Amanda E. Toland, Diana Torres, Nadine Tung, Annemieke H. van der Hout, Elizabeth J. van Rensburg, Ana Vega, Barbara Wappenschmidt, Jeffrey N. Weitzel, Katia M. Zavaglia, Kristin K. Zorn, Thomas A. Sellers, Georgia Chenevix-Trench and Antonis C. Antoniou