

Interatomic Coulombic Decay Processes after Multiple Valence Excitations in Ne Clusters

D. Iablonskyi^{a,1}, K. Nagaya^b, H. Fukuzawa^a, K. Motomura^a, Y. Kumagai^a, S. Mondal^a, T. Tachibana^a, T. Takashi^a, T. Nishiyama^b, K. Matsunami^b, P. Johnsson^c, P. Piseri^d, G. Sansone^e, A. Dubrouil^e, M. Reduzzi^e, P. Carpeggiani^e, C. Vozzi^e, M. Devetta^e, M. Negro^e, D. Faccialà^e, F. Calegari^e, A. Trabattoni^e, M. Castrovilli^e, Y. Ovcharenko^f, T. Möller^f, M. Mudrich^g, F. Stienkemeier^g, M. Coreno^h, M. Alagiaⁱ, B. Schütte^j, N. Berrah^k, C. Callegari^l, O. Plekan^l, P. Finetti^l, C. Spezzani^l, E. Ferrari^l, E. Allaria^l, G. Penco^l, C. Serpico^l, G. De Ninno^l, B. Diviacco^l, S. Di Mitri^l, L. Giannessi^l, K. C. Prince^{i,l}, M. Yao^b, and K. Ueda^a

^a Institute of Multidisciplinary Research for Advanced Materials, Tohoku University, 980-8577 Sendai, Japan

^b Department of Physics, Graduate School of Science, Kyoto University, 606-8502 Kyoto, Japan

^c Department of Physics, Lund University, P. O. Box 118, 22100 Lund, Sweden

^d Dipartimento di Fisica, Università degli Studi di Milano, 20133 Milano, Italy

^e CNR-IFN, Piazza Leonardo da Vinci 32, 20133 Milan, Italy

^f Institut für Optik und Atomare Physik, TU Berlin, 10623 Berlin, Germany

^g Physikalisches Institut, Universität Freiburg, 79104 Freiburg, Germany

^h CNR-ISM, Area della Ricerca di Roma 1, Monterotondo Scalo, Italy

ⁱ CNR-IOM, Area Science Park, 34149 Trieste, Italy

^j Max-Born-Institut, Max-Born-Strasse 2 A, 12489 Berlin, Germany

^k Department of Physics, University of Connecticut, Storrs, CT 06269, USA

^l Elettra-Sincrotrone Trieste, Area Science Park, 34149 Trieste, Italy

Synopsis We present a comprehensive analysis of autoionization processes in Ne clusters (~ 5000 atoms) after multiple valence excitations by free electron laser radiation. The evolution from 2-body interatomic Coulombic decay (ICD) to 3-body ICD is demonstrated when changing from surface to bulk Frenkel exciton excitation. Super Coster-Kronig type 2-body ICD is observed at Wannier exciton which quenches the main ICD channel.

Previously, Yase *et al* [1] investigated multiple excitation of Wannier type excitons (corresponding to the $2p \rightarrow 3d$ atomic resonance) at 20.26 eV in Ne clusters by the intense extreme ultraviolet free electron laser (EUV-FEL) at SCSS (Spring-8 Compact SASE Source, Japan) and found that the electron emission is dominated by low energy electron emission that originates from a nanoplasma.

In the present experiment, we have extended our observations to multiple excitations of surface and bulk Frenkel type excitons (corresponding to the $2p \rightarrow 3s$ atomic resonance) at 17.12 eV and 17.65 eV, respectively, using the new seeded EUV-FEL, FERMI (Trieste, Italy) [2].

At the lowest surface Frenkel exciton we can clearly see the pure 2-body ICD peak at ~ 11.5 eV, predicted by Kuleff *et al* [3], with its multistep ICD tail which is similar to direct multistep ionization in Ar clusters [4]. The situation changes for the bulk Frenkel exciton, where the broad structure around 5 eV is identified as 3-body ICD of knock-off type (also known as collective autoionization [5]) and becomes dominant over 2-body ICD at high FEL intensities. For the Wannier ex-

citon we can see complete quenching of the main 2-body ICD by super Coster-Kronig type ICD in which one 3d electron relaxes to a 3s orbital and another 3d electron is ejected with ~ 1.8 eV kinetic energy (see Fig. 1).




Figure 1. Electron emission spectra for excitation of different excitons and selected FEL intensities.

References

- [1] S. Yase *et al.* 2013 *Phys. Rev. A* **88** 043203
- [2] E. Allaria *et al.* 2012 *Nat. Photonics* **6** 699
- [3] A. I. Kuleff *et al.* 2010 *Phys. Rev. Lett.* **105** 043004
- [4] C. Bostedt *et al.* 2008 *Phys. Rev. Lett.* **100** 133401
- [5] Y. Ovcharenko *et al.* 2014 *Phys. Rev. Lett.* **112** 073401

¹E-mail: denys@tagen.tohoku.ac.jp

