

LETTER • OPEN ACCESS

Oceanic Rossby waves drive inter-annual predictability of net primary production in the central tropical Pacific

To cite this article: Sebastian Brune *et al* 2022 *Environ. Res. Lett.* **17** 014030

View the [article online](#) for updates and enhancements.

You may also like

- [Generalized Darboux transformation and rogue wave solutions for the higher-order dispersive nonlinear Schrödinger equation](#)
Bo Yang, Wei-Guo Zhang, Hai-Qiang Zhang *et al.*
- [Expected maximum of bridge random walks & Lévy flights](#)
Benjamin De Bruyne, Satya N Majumdar and Grégory Schehr
- [Roadmap on optical rogue waves and extreme events](#)
Nail Akhmediev, Bertrand Kibler, Fabio Baronio *et al.*

ENVIRONMENTAL RESEARCH
LETTERS

LETTER


Oceanic Rossby waves drive inter-annual predictability of net primary production in the central tropical Pacific

OPEN ACCESS

RECEIVED
9 September 2021REVISED
15 December 2021ACCEPTED FOR PUBLICATION
16 December 2021PUBLISHED
5 January 2022

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.

Sebastian Brune^{1,*} , Maria Esther Caballero Espejo², David Marcolino Nielsen^{1,4}, Hongmei Li³, Tatiana Ilyina³ and Johanna Baehr¹¹ Institute of Oceanography, CEN, Universität Hamburg, Hamburg, Germany² Ministerio del Ambiente, Dirección de Adaptación al Cambio Climático y Desertificación, Magdalena del Mar, Lima, Peru³ Max Planck Institute for Meteorology, Hamburg, Germany⁴ International Max Planck Research School on Earth System Modelling, Hamburg, Germany

* Author to whom any correspondence should be addressed.

E-mail: sebastian.brune@uni-hamburg.de**Keywords:** decadal climate prediction, net primary productivity, oceanic Rossby waves, ensemble Kalman filter, MPI-ESM, Pacific Ocean, earth system modelling**Abstract**

In the Pacific Ocean, off-equatorial Rossby waves (RWs), initiated by atmosphere–ocean interaction, modulate the inter-annual variability of the thermocline. In this study, we explore the resulting potential gain in predictability of central tropical Pacific primary production, which in this region strongly depends on the supply of macronutrients from below the thermocline. We use a decadal prediction system based on the Max Planck Institute Earth system model to demonstrate that for the time period 1998–2014 properly initialized RWs explain an increase in predictability of net primary productivity (NPP) in the off-equatorial central tropical Pacific. We show that, for up to 5 years in advance, predictability of NPP derived from the decadal prediction system is significantly larger than that derived from persistence alone, or an uninitialized historical simulation. The predicted signal can be explained by the following mechanism: off-equatorial RWs are initiated in the eastern Pacific and travel towards the central tropical Pacific on a time scale of 2–6 years. On their arrival the RWs modify the depths of both thermocline and nutricline, which is fundamental to the availability of nutrients in the euphotic layer. Local upwelling transports nutrients from below the nutricline into the euphotic zone, effectively transferring the RW signal to the near-surface ocean. While we show that skillful prediction of central off-equatorial tropical Pacific NPP is possible, we open the door for establishing predictive systems for food web and ecosystem services in that region.

1. Introduction

In Earth's climate system, dynamic interaction of atmosphere and ocean leads to the initiation of oceanic waves in a wide range of spatio-temporal scales (Gill 1982). Depending on their characteristics, these waves may carry memory within the Earth's climate system, and can therefore be used in skillful climate predictions. Off-equatorial oceanic planetary waves (Rossby 1939, 1945), commonly known as oceanic Rossby waves (RWs) (RWs; Platzman 1968), could carry memory on the interannual to decadal time scales (2–10 years). Oceanic RWs occur with typical phase speeds of 0.03–0.3 m s⁻¹ (approx. 1000–10 000 km/year) and wavelengths larger than

500 km (Chelton and Schlax 1996, Killworth and Blundell 2003). The vertical displacement due to oceanic RWs varies from only a few centimeters at the ocean's surface (Chelton and Schlax 1996) to several tens of meters at thermocline depth (Kessler 1990). Satellite altimetry confirmed the abundance of oceanic RWs in all of Earth's major ocean basins (Chelton and Schlax 1996). However, the longest memory resulting from oceanic RWs can be expected to be found in the Pacific, where they cross the ocean on time scales of 5–30 years (Capotondi and Alexander 2001, Galanti and Tziperman 2003). Here, oceanic RWs are predominantly initiated in the eastern portion of the Pacific Ocean, involving atmospheric and oceanic processes,

e.g. Ekman pumping connected with surface winds (Capotondi and Alexander 2001), or modulations of the pycnocline in connection with El Niño-Southern Oscillation (ENSO, Vega *et al* 2003). Propagating RWs may have an impact not only on the inter-annual variability of the ocean's dynamical state, such as temperature, salinity, sea surface height, but also on nutrient availability, and related to that, biological primary productivity (Siegel 2001, Uz *et al* 2001).

At ocean depths below 100m, where the absence of Sun light prohibits biological production, variability of nutrient concentration is almost negligible on inter-annual time scales (Uz *et al* 2001). At shallower depths, in the euphotic zone, the availability of nutrients varies more than at larger depths, and may determine the strength of biological net primary productivity (NPP) (Barber 1992), in particular in tropical oceans. Vice versa, biological production leads to a depletion of nutrients in the euphotic zone and as a result a steep gradient in nutrient concentration with depth is formed, the nutricline. Oceanic RWs modulate the depth of the nutricline by a few tens of meters, with corresponding impacts on the nutrient availability above the nutricline, and on NPP and the resulting chlorophyll concentration (Uz *et al* 2001, Killworth *et al* 2004). Compared to climatology, a shallower (deeper) than normal nutricline leads to larger (smaller) than normal primary production. Oceanic RWs impact surface chlorophyll concentration also by a vertical displacement of the chlorophyll maximum (Kawamiya and Oschlies 2001). Signatures of oceanic RWs have therefore not only been observed in sea-level anomalies, but also in ocean color attributed to oceanic chlorophyll (Uz *et al* 2001, Gutknecht *et al* 2010). In addition, RWs could impact the variability of surface chlorophyll and organic particles by lateral advection, without a concurrent change in nutrient availability (Dandonneau *et al* 2003, 2004, Killworth 2004). While there is a general consensus that advection due to oceanic RWs directly modulates upper ocean chlorophyll concentrations on the inter-annual time scale, the degree of the impact of oceanic RWs on nutrient availability above the nutricline and a co-varying NPP remains unclear (Killworth *et al* 2004, Charria *et al* 2008). It still remains to be shown how the inter-annual memory potentially carried by oceanic RWs can be used for skillful decadal predictions of nutrient availability and NPP. Here, we set out to precisely fill in this gap, by investigating RWs in a decadal predictions system.

On the basis of comprehensive Earth system models (ESMs), decadal prediction systems have been developed (see, e.g., Boer *et al* 2016) with the aim to predict Earth's climate, in particular the dynamical state of atmosphere and ocean, on inter-annual to decadal time scales. In the slipstream of maturing decadal prediction systems, some studies assessed

the inter-annual prediction of biogeochemical parameters. In a groundbreaking study, Séférian *et al* (2014) found predictability of NPP in the tropical Pacific with an ESM for up to three years in advance, attributed, amongst other drivers, to the poleward advection and persistence characteristics of nutrients, driven directly by ENSO. Ham *et al* (2021) found that in the southern tropical Pacific, ENSO variability has a major impact on the advection of nutrients and thus on surface chlorophyll concentration, limiting skillful predictions to the seasonal time scale. For the North Atlantic sub-polar gyre, Li *et al* (2016) show that the CO₂ uptake by the ocean could be predicted 4–7 years in advance. On the global scale, Park *et al* (2019) found skill in their ESM-based seasonal predictions of chlorophyll concentrations, also mainly connected to ENSO, and Fransner *et al* (2020) showed that oceanic biochemical predictability is closely tied to the representation of the upper ocean dynamical state in their ESM.

Despite the success story of decadal prediction systems, the role of oceanic RWs in the predictability of tropical nutrient availability and NPP has not been comprehensively studied so far. Bridging the gap between dynamical and biogeochemical prediction remains an ongoing challenge. In this study, we use a comprehensive Earth system model to pick up this challenge with the goal to successfully predict NPP in the tropical Pacific several years ahead and to pinpoint the role of off-equatorial oceanic RWs in carrying and distributing inter-annual memory in the tropical Pacific.

2. Model simulations, observational references and prediction skill

We use the Max Planck Institute (MPI) for meteorology ESM at low resolution (MPI-ESM-LR, Giorgetta *et al* 2013) with the atmospheric component ECHAM6 (Stevens *et al* 2013) in T63L47 configuration, equivalent to 2.8° horizontal resolution, and the oceanic component Max Planck Institute Ocean Model (MPIOM) (Jungclaus *et al* 2013) in GR15L40 configuration with 1.5° nominal horizontal resolution. In MPI-ESM, oceanic biogeochemistry is represented by the Hamburg ocean carbon cycle model (HAMOCC, Ilyina *et al* 2013). All simulated values of NPP and nitrate concentration used in this study are calculated by HAMOCC, sea surface temperatures (SST) are taken from MPIOM. We use the vertical mass transport simulated in MPIOM, averaged over 200–450 m depths, to analyze the characteristics of oceanic RWs.

Our prediction system consists of 10-year 8-member ensemble hindcasts with MPI-ESM-LR, initialized from a weakly coupled data assimilation experiment at the 1st of January every year from

1989 to 2014 (Brune *et al* 2018). The 8-member MPI-ESM-LR assimilation covers the time period 1958–2014, where we incorporate ERA-40/ERA-Interim re-analyses of vorticity, divergence, temperature and sea level pressure (Uppala *et al* 2005, Dee *et al* 2011) via atmospheric nudging and observational profiles of oceanic temperature and salinity (EN4, Good *et al* 2013) using a non-localized 8-member oceanic ensemble Kalman filter (EnKF, Brune *et al* 2015). In this weakly coupled assimilation setup, the ocean responds mainly to the atmospheric nudging with weak changes added by the oceanic EnKF. We compare the initialized ensemble hindcasts with an 8-member ensemble of uninitialized MPI-ESM-LR historical simulations. Both the initialized and uninitialized simulations are run under identical prescribed external conditions, e.g. green house gas concentration, solar radiation, and volcanic aerosols, according to CMIP5 (Taylor *et al* 2012). Here, we extend the ‘historical’ time period (until 2005) with the RCP4.5 scenario (after 2005).

NPP observational reference data are estimated from ocean color measurements obtained by the Sea-viewing Wide Field-of-view Sensor (SeaWiFS) instrument of the OrbView-2 satellite for September 1997 to December 2002 and the Moderate Resolution Imaging Spectroradiometer (MODIS) of the Aqua satellite from January 2003 to December 2014 (Behrenfeld and Falkowski 1997, <http://science.oregonstate.edu/ocean.productivity/index.php>). Reference data for SST are taken from the Hadley Centre Global Sea Ice and Sea Surface Temperature data set (HadISST, Rayner *et al* 2003). For supplemental analysis (not shown) we use AVISO as our reference data for sea surface height, this altimeter products were produced by Ssalto/Duacs and distributed by Aviso+, with support from Cnes (www.aviso.altimetry.fr).

Inter-annual prediction skill is measured using the correlation coefficient and the root mean square error between the respective yearly means of simulations and reference data. Prior to skill analysis, linear trends are removed. Significance of correlations is calculated at the 95% level by analyzing both tails of the 1000-member bootstrapped probability distribution (not shown) of the respective simulations with a *p*-value of 0.025 (Efron 1979, Wilks 2011).

In the central tropical Pacific, ENSO has a large direct impact, also on decadal time scales (C-Mode in Takahashi *et al* (2011)), which may mask the oceanic RW signal in oceanic surface quantities. In an effort to carve out the oceanic RW signal, we therefore remove the direct impact of ENSO using linear regressions first on the central Pacific Niño3.4 (5°N–5°S, 170°W–120°W) SST index and then on the eastern Pacific Niño1+2 (0°–10°S, 90°W–80°W) SST index (Trenberth and Stepaniak 2001).

3. Results

3.1. Representation of off-equatorial oceanic Rossby waves

In general, Pacific off-equatorial RWs are abundantly simulated in the hindcasts of our MPI-ESM prediction system (figure 1). In accordance with observations (e.g., Chelton and Schlax 1996), simulated RWs emanate in the tropical Pacific from two distinct regions North and South of the equator: in the North Pacific around the South-Eastern edge of the sub-tropical gyre, in the South Pacific around the Peru-Chile current off-shore South America (figure 1(a)). Two distinct RW trains extend from these source regions to the West: in the North Pacific along 15°N, and in the South Pacific along 10°S. Around these latitudes, the RWs modulate the depth of the nutricline (figures 1(b) and (c)) and the thermocline (not shown). The longitudinal extension of source regions and RW trains along these latitudes does not change over the entire time period 1998–2014 (figures 2(a) and (b)). At 15°N (figure 2(a)), the characteristic period of RWs is about 2 years, the travel time from the source region around 140°W across the basin towards the maritime continent is about 2–4 years. At 10°S (figure 2(b)), the characteristic period is 2–6 years, while the travel time from the source region off-shore South America towards the central tropical Pacific is about 2–6 years. Thus, we determine our target regions in the tropical Pacific as those regions, which show a large impact of oceanic RWs with a characteristic time period of 2–6 years.

Filtering the vertical mass transport with a characteristic RW period of 2–6 years reveals in the Pacific two bands on each side of the equator with large variance due to RWs (figure 2(c)). In each of these bands, the easternmost part represents the source region of RWs. They travel westward and thus defining the target regions for our analysis of prediction skill. In the tropical North Pacific, the source region can be broadly defined as a box at 120°W–140°W, 10°N–20°N, and the target region as a box 150°E–170°W, 10°N–20°N (henceforth referred to as North Pacific box). In the tropical South Pacific the source region is located at around 85°W–100°W, 20°S–10°S, and the target region at 160°W–110°W, 15°S–5°S (henceforth referred to as South Pacific box). In both source regions, RWs in our hindcasts are properly initialized in time and space by the assimilation, indicated by high correlations of the simulated sea surface height anomaly with AVISO observations (not shown, $r = 0.80$ and $r = 0.85$ for the Northern and Southern Pacific source regions, respectively). Hence, in the target regions we expect a significant contribution of RWs not only to the variability of the vertical mass transport around the thermocline, but also to the modulation of the nutricline




Figure 1. (a) Tropical Pacific monthly mean vertical mass transport, averaged over 200–450m depth, with the average seasonal cycle removed, snapshot of December 2000, predicted from January 1998, dashed line boxes indicate source regions of Rossby waves, dotted lines show latitudes 15°N, and 10°S; (b, c) monthly mean nitrate concentration along 15°N, and 10°S, respectively, December 2000 snapshot, predicted from January 1998, the steep gradient depicts the nutricline, dashed lines indicate source regions of Rossby waves.

and the availability of nutrients in the euphotic zone, similar to the findings of Capotondi and Alexander (2001).


3.2. RW impact on prediction of nutrient availability and NPP

The predicted nutrient availability for up to 4 years in advance is modulated by the inter-annual variability of the strength of predicted off-equatorial RWs in the target regions. Variability of nitrate concentration is strongly influenced by RWs not only at depths below 100 m (figures 3(a) and (b)), but the RW signal is also transferred from these depths to the euphotic zone, i.e. the signal can be traced over all depths and often reaches the surface (figures 3(a) and (b)).

In both target regions, the year-to-year and the multi-year variability in nutrient concentration and NPP fit well with the RW signal seen below 100 m depth. In the euphotic zone, there is significant correlation between the predicted nutrient concentration and predicted NPP 4 years in advance (detrended and direct ENSO impact removed) in both the North

Pacific box (figure 3(c), $r = 0.57$) and the South Pacific box (figure 3(d), $r = 0.75$). Simultaneously, the predicted NPP is significantly correlated with the observed NPP (detrended and direct ENSO impact removed) for 4 years in advance in both the North Pacific (figure 3(c), $r = 0.68$) and South Pacific box (figure 3(d), $r = 0.76$).

The correlation between predicted and observed NPP depends on the lead year. For the North Pacific box, we find significantly positive correlation skill for NPP in our prediction system for lead times up to 4 years, and for the South Pacific box for lead times up to 5 years (see filled circles in figure 4). In this time frame, the time-lag auto-correlation of observed NPP alone (NPP persistence) hardly leads to any significantly positive correlation skill, indicating a clear benefit of using MPI-ESM over persistence for NPP prediction. Correlation skill of NPP in a corresponding historical simulation, i.e. without transferring observed states of atmosphere and ocean to MPI-ESM, is negative in both Pacific boxes. The evaluation of the root mean square error (RMSE) against





reference corroborates the findings of our correlation analysis. In the North Pacific box, the RMSE is for all lead years smaller for the predicted NPP than for NPP persistence, and for up to lead year 4 the RMSE for predicted NPP is also smaller than for the historical simulation. In the South Pacific box, RMSE for the predicted NPP is for all lead years smaller than for NPP persistence of NPP from the historical simulation. For predictions up to 5 years in advance, there is a substantial gain in prediction skill when assimilating observed atmospheric and oceanic states into MPI-ESM prior to starting the hindcasts. Thus the proper representation of oceanic RWs, allowed by the assimilation of observed atmospheric and ocean

states at initialization, increases the predictability of NPP at the inter-annual time scales regions with a large impact of RWs.


3.3. Comparison of NPP and SST predictability

In our prediction system, correlation skill of NPP in both the North Pacific and South Pacific boxes is consistently higher than that of SST for lead years 2–5 (figure 4). An important reason for this lies in different susceptibilities of NPP and SST to the local atmospheric surface conditions in the tropical Pacific. Near-surface ocean temperatures are directly coupled to the local atmospheric conditions, which actively



impact SST and add a chaotic component to its prediction, eventually lowering prediction skill. In contrast, NPP in the tropical Pacific is not as strongly affected by local atmospheric conditions as SST, since

temperature is not a dominant factor to NPP in the tropical oceans in general (passive impact). In turn, inter-annual predictions of NPP in the North Pacific and South Pacific boxes benefit more from the



inter-annual variability represented by properly initialized oceanic RWs than SST.

Looking beyond the selected regions, NPP is predictable up to lead year 4 over a substantial portion of the off-equatorial tropical Pacific (figure 5). There is a strong indication that the mechanism proposed in our study is generally complemented by other processes, which eventually lead to predictability of NPP on the inter-annual time scale on a much larger geographic scale. In particular, our prediction system can skillfully predict NPP along the equatorward edges of

the Pacific subtropical gyres. While the gyre boundary may act as a wave guide to oceanic RWs, which eventually modulate NPP variability, the correct prediction of the contrasting dominating effects of lateral and vertical nutrient advection outside the gyres, and strong stratification and nutrient recycling inside the gyres (Séférian *et al* 2014, Karl and Church 2017) could also be the reason for skillful NPP predictions in our decadal prediction system.

Our results also reveal that oceanic RWs do not always imply skillful NPP predictions. Oceanic RWs

traveling within the Pacific subtropical gyres do not seem to, if at all, positively impact inter-annual NPP predictions (e.g. figures 5(e) and (g)), most probably due to the dominance of the nutrient recycling regime within the gyres. Also, prediction skill decreases beyond lead year 5, indicating an increase in the unpredictable components in the climate system, counteracting the memory potentially carried by oceanic RWs.

4. Discussion and conclusion

The potential for inter-annual predictability of biogeochemical parameters in the Pacific has been evaluated previously by S  ferian *et al* (2014). While using an early-state decadal prediction system (e.g. assimilating only SST), they found that prediction skill for NPP could be high for up to 3 years in advance. In our study, we use a more comprehensive decadal prediction system, assimilating 3D atmospheric and ocean states in a model-consistent approach, to specifically examine one potential dynamical source of inter-annual predictability in the tropical Pacific, oceanic RWs, while removing the direct impact of another potential source, ENSO. We find that in regions with a potentially large impact of oceanic RWs, the prediction horizon of NPP could be extended by 1–2 years over the 3 years found in S  ferian *et al* (2014) for the tropical Pacific, with a major contribution by the successfully timed initialization and propagation of oceanic RWs. Although our study highlights the impact of the vertical supply of nutrients in connection to oceanic RWs, other dynamical processes are known to impact nutrient supply and primary production. Killworth *et al* (2004) discuss that horizontal advection of nutrients, also in connection with oceanic waves, may have a strong impact on both nutrient supply and chlorophyll production. Also, nutrients could be transported with oceanic currents, in particular along gyre boundaries, and thus impact NPP on the seasonal to inter-annual time scale. In our study we did not directly account for the effect of nutrient recycling, which has been discussed to impact primary production, in particular inside oceanic gyres (S  ferian *et al* 2014, Karl and Church 2017).

Our initialization approach relies on a rather simple weakly coupled assimilation of both atmospheric and oceanic observational products. It remains to be shown, if other state-of-the-art initialization approaches, such as discussed in Brune and Baehr (2020), Merryfield *et al* (2020) or Fransner *et al* (2020), could lead to further improvements in the prediction of NPP.

In our analysis, we rely on satellite-based ocean color, i.e. chlorophyll, observations as our reference for NPP observations. Earlier studies pointed out that

although oceanic RWs could account for up to 30% of the modulation of regional NPP (Kawamiya and Oschlies 2001), the associated impact on satellite-based chlorophyll measurements could be smaller, Uz *et al* (2001) indicate a range of 5%–20%. In addition to in-situ primary production, visible chlorophyll concentrations are also directly dynamically related to transports within the oceanic mixed layer (Kawamiya and Oschlies 2001). We acknowledge that satellite-based instruments not only changed over the time period 1998–2014, but also may encounter sensor decay over their life-time. This contributes to an increased uncertainty on the reference side in our NPP predictions. To tackle this in future studies it could be advisable to revisit NPP references with Sentinel-3 data (Donlon *et al* 2012).

Although we removed the direct impact of ENSO prior to our analysis of NPP predictability, it should be noted that ENSO is indirectly involved in predicting NPP with MPI-ESM. ENSO determines, among other processes, the atmosphere-ocean interaction in the source regions of oceanic RWs. Therefore, the assimilation of the correct ENSO state at the start of the prediction is a key factor to take advantage of the multi-year memory offered by oceanic off-equatorial RWs in the tropical Pacific, and is the main source of our increased prediction skill in NPP in the initialized predictions when compared to uninitialized historical simulations.

From our analysis of tropical Pacific inter-annual prediction skill of NPP for the time period 1998–2014 in MPI-ESM, we conclude:

- In the tropical Pacific, we find high prediction skill of NPP for up to 5 years in advance, with a large contribution from properly initialized and simulated off-equatorial oceanic RWs.
- The prediction skill of NPP represents a significant increase on prediction skill of SST, which is limited to 1–3 years. We argue that this difference arises from the different impact of near surface atmospheric dynamics on nutrient concentration (passive impact) and near surface ocean temperature (active impact).

The connection of dynamical and biogeochemical predictions is encouraging for further studies on the predictability of the food web in the tropical Pacific ocean.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: https://cera-www.dkrz.de/WDCC/ui/cerasearch/entry?acronym = DKRZ_LTA_1075_ds00006 (Brune *et al* 2021).

Acknowledgments

Model simulations were performed at the German Climate Computing Centre (DKRZ). Reference data for net primary productivity calculated from SeaWiFS and MODIS satellite sensors have been retrieved from <http://science.oregonstate.edu/ocean.productivity/index.php>. Sea surface temperature data from HadISST have been retrieved through www.metoffice.gov.uk/hadobs. This research was funded by the German Ministry of Education and Research (BMBF) under the MiKlip projects AODA-PENG (Grant Nos. 01LP1157C, 01LP1516A), by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy—EXC 2037 'CLICCS—Climate, Climatic Change, and Society'—Project No.: 390683824, contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg, and the Cluster of Excellence CliSAP (EXC177), Universität Hamburg, and by Copernicus Climate Change Service, funded by the EU, under Contract C3S-330. We thank three anonymous reviewers for their constructive and helpful comments.

ORCID iD

Sebastian Brune  <https://orcid.org/0000-0001-7794-5465>

References

- Barber R T 1992 *Geologic and Climatic Time Scales of Nutrient Variability* (Boston, MA: Springer US) pp 89–106
- Behrenfeld M J and Falkowski P G 1997 Photosynthetic rates derived from satellite-based chlorophyll concentration *Limnol. Oceanogr.* **42** 1–20
- Boer G J et al 2016 The decadal climate prediction project (DCPP) contribution to CMIP6 *Geosci. Model Dev.* **9** 3751–77
- Brune S and Baehr J 2020 Preserving the coupled atmosphere-ocean feedback in initializations of decadal climate predictions *WIREs Clim. Change* **11** e637
- Brune S, Baehr J, Pohlmann H and Müller W A 2021 MPI-ESM-LR_1.0.02 decadal predictions globalEnKF (available at: http://cera-www.dkrz.de/WDCC/ui/Compact.jsp?acronym=DKRZ_LTA_1075_ds00006)
- Brune S, Düsterhus A, Pohlmann H, Müller W A and Baehr J 2018 Time dependency of the prediction skill for the North Atlantic subpolar gyre in initialized decadal hindcasts *Clim. Dyn.* **51** 1947–70
- Brune S, Nergler L and Baehr J 2015 Assimilation of oceanic observations in a global coupled Earth system model with the SEIK filter *Ocean Model.* **96** 254–64
- Capotondi A and Alexander M A 2001 Rossby waves in the tropical north pacific and their role in decadal thermocline variability *J. Phys. Oceanogr.* **31** 3496–3515
- Charria G, Dadou I, Cipollini P, Drévilion M and Garçon V 2008 Influence of Rossby waves on primary production from a coupled physical–biogeochemical model in the North Atlantic Ocean *Ocean Sci.* **4** 199–213
- Chelton D B and Schlax M G 1996 Global observations of oceanic Rossby waves *Science* **272** 234–8
- Dandonneau Y, Menkes C, Gorgues T and Madec G 2004 Response to comment on oceanic Rossby waves acting as a Hay Rake for ecosystem floating by-products *Science* **304** 390
- Dandonneau Y, Vega A, Loisel H, du Penhoat Y and Menkes C 2003. Oceanic Rossby waves acting as a “Hay Rake” for ecosystem floating by-products *Science* **302** 1548–51
- Dee D P et al 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q. J. R. Meteorol. Soc.* **137** 553–97
- Donlon C et al 2012 The global monitoring for environment and security (GMES) Sentinel-3 mission *Remote Sens. Environ.* **120** 37–57
- Efron B 1979 Bootstrap methods: another look at the Jackknife *Ann. Stat.* **7** 1–26
- Fransner F, Counillon F, Bethke I, Tjiputra J, Samuelsen A, Nummelin A and Olsen A 2020 Ocean biogeochemical predictions - initialization and limits of predictability *Front. Mar. Sci.* **7** 386
- Galanti E and Tziperman E 2003 A midlatitude-ENSO teleconnection mechanism via baroclinically unstable long Rossby waves *J. Phys. Oceanogr.* **33** 1877–88
- Gill A E 1982 *Atmosphere-Ocean Dynamics (Int. Geophysics Series vol 30)* (New York: Academic)
- Giorgetta M A et al 2013 Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5 *J. Adv. Mod. Earth Sys.* **5** 572–97
- Good S A, Martin M J and Rayner N A 2013 EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates *J. Geophys. Res.* **118** 6704–16
- Gutknecht E, Dadou I, Charria G, Cipollini P and Garçon V 2010 Spatial and temporal variability of the remotely sensed chlorophyll a signal associated with Rossby waves in the South Atlantic Ocean *J. Geophys. Res. Oceans* **115** C05004
- Ham Y-G, Joo Y-S and Park J-Y 2021 Mechanism of skillful seasonal surface chlorophyll prediction over the southern Pacific using a global earth system model *Clim. Dyn.* **56** 45–64
- Ilyina T, Six K D, Segsneider J, Maier-Reimer E, Li H and Nuñez-Riboni I 2013 Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations *J. Adv. Mod. Earth Sys.* **5** 287–315
- Jungclaus J H, Fischer N, Haak H, Lohmann K, Marotzke J, Matei D, Mikolajewicz U, Notz D and von Storch J S 2013 Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model *J. Adv. Mod. Earth Sys.* **5** 422–46
- Karl D M and Church M J 2017 Ecosystem structure and dynamics in the North Pacific Subtropical Gyre: new views of an old ocean *Ecosystems* **20** 433–57
- Kawamiya M and Oschlies A 2001 Formation of a basin-scale surface chlorophyll pattern by Rossby waves *Geophys. Res. Lett.* **28** 4139–42
- Kessler W S 1990 Observations of long Rossby waves in the northern tropical Pacific *J. Geophys. Res. Oceans* **95** 5183–5217
- Killworth P D 2004 Comment on “oceanic Rossby waves acting as a Hay Rake for ecosystem floating by-products *Science* **304** 390
- Killworth P D and Blundell J R 2003 Long extratropical planetary wave propagation in the presence of slowly varying mean flow and bottom topography. Part I: the local problem *J. Phys. Oceanogr.* **33** 784–801
- Killworth P D, Cipollini P, Uz B M and Blundell J R 2004 Physical and biological mechanisms for planetary waves observed in satellite-derived chlorophyll *J. Geophys. Res. Oceans* **109** C07002
- Li H, Ilyina T, Müller W A and Sienz F 2016 Decadal predictions of the North Atlantic CO₂ uptake *Nat. Commun.* **7** 11076

- Merryfield W J *et al* 2020 Current and emerging developments in subseasonal to decadal prediction *Bull. Am. Meteorol. Soc.* **101** E869–96
- Park J-Y, Stock C A, Dunne J P, Yang X and Rosati A 2019 Seasonal to multiannual marine ecosystem prediction with a global Earth system model *Science* **365** 284–8
- Platzman G W 1968 The Rossby wave *Q. J. R. Meteorol. Soc.* **94** 225–48
- Rayner N A, Parker D E, Horton E B, Folland C K, Alexander L V, Rowell D P, Kent E C and Kaplan A 2003 Global analyses of sea surface temperature, sea ice and night marine air temperature since the late nineteenth century *J. Geophys. Res.* **108** 4407
- Rossby C-G 1939 Relation between variations in the intensity of the zonal circulation of the atmosphere and the displacements of the semi-permanent centers of action *J. Mar. Res.* **2** 38–55
- Rossby C-G 1945 On the propagation of frequencies and energy in certain types of oceanic and atmospheric waves *J. Atmos. Sci.* **2** 187–204
- S  ferian R, Bopp L, Gehlen M, Swingedouw D, Mignot J, Guilyardi E and Servonnat J 2014 Multiyear predictability of tropical marine productivity *Proc. Natl Acad. Sci.* **111** 11646–51
- Siegel D A 2001 The Rossby rototiller *Nature* **409** 576–7
- Stevens B *et al* 2013 Atmospheric component of the MPI-M earth system model: ECHAM6 *J. Adv. Mod. Earth Sys.* **5** 146–72
- Takahashi K, Montecinos A, Goubanova K and Dewitte B 2011 ENSO regimes: Reinterpreting the canonical and Modoki El Ni  o *Geophys. Res. Lett.* **38** L10704
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteor. Soc.* **93** 485–98
- Trenberth K E and Stepaniak D P 2001 Indices of El Ni  o Evolution *J. Clim.* **14** 1697–701
- Uppala S M *et al* 2005 The ERA-40 re-analysis *Q. J. R. Meteorol. Soc.* **131** 2961–3012
- Uz B M, Yoder J A and Osychny V 2001 Pumping of nutrients to ocean surface waters by the action of propagating planetary waves *Nature* **409** 597–600
- Vega A, Du-Penhoat Y, Dewitte B and Pizarro O 2003 Equatorial forcing of interannual Rossby waves in the Eastern South Pacific *Geophys. Res. Lett.* **30** 1197
- Wilks D 2011 *Statistical Methods in the Atmospheric Sciences (Int. Geophysics Series vol 100)* (New York: Academic)